

AperTO - Archivio Istituzionale Open Access dell'Università di Torino

Characterization of luteal dynamics in lactating Holstein cows for 32 days after synchronization of ovulation and timed artificial insemination

This is the author's manuscript

Original Citation:

Availability:

This version is available <http://hdl.handle.net/2318/1689812> since 2022-02-17T10:42:45Z

Published version:

DOI:10.3168/jds.2017-13293

Terms of use:

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)

1 Characterization of luteal dynamics in lactating Holstein cows for 32 days after synchronization of 2 ovulation and timed artificial insemination

3 A. Ricci,*† P. D. Carvalho,* M. C. Amundson,* and P. M. Fricke*¹

4 *Department of Dairy Science, University of Wisconsin, Madison 53706

5 †Department of Veterinary Science, Università di Torino, Grugliasco 1009, Italy

6 ABSTRACT

7 Approximately 20 to 30% of cows diagnosed not pregnant 32 d after timed artificial insemination (TAI) lack a
8 corpus luteum (CL), and cows submitted to a resynchronization protocol in the absence of a CL have about 10%
9 fewer pregnancies per AI (P/AI) than cows with a CL. An understanding of luteal dynamics after
10 synchronization of ovulation and TAI may help refine strategies for reinseminating cows failing to conceive.
11 Lactating Holstein cows (n = 141) were synchronized for first TAI using a Double-Ovsynch protocol. Thrice
12 weekly from 4 to 32 d after TAI, blood samples were collected for evaluation of plasma progesterone (P4)
13 concentrations, and CL diameter was measured using transrectal ultrasonography. Pregnancy status was de-
14 termined using transrectal ultrasonography 32 d after TAI. Nonsynchronized cows (n = 4) were removed from
15 the study. For cows diagnosed pregnant 32 d after TAI (n = 57), P4 increased from 4 to 15 d and then remained
16 constant until 32 d after TAI, whereas CL volume increased from 4 to 11 d and then remained constant until 32 d
17 after TAI. For cows diagnosed not pregnant 32 d after TAI (n = 80), P4 profiles were evaluated using statistical
18 cluster analysis based on the day after TAI that P4 decreased to <1 ng/mL, resulting in 5 clusters: (1) CL
19 regression 15 d after TAI (1.3%), (2) CL regression 18 to 22 d after TAI (55.0%), (3) CL regression 25 to 27 d
20 after TAI (17.5%), (4) CL regression 29 to 32 d after TAI (5.0%), and (5) CL maintained until 32 d after
21 TAI (21.3%). Plasma pregnancy-associated glycoprotein (PAG) levels at 25 and 32 d after TAI differed among
22 clusters and were below the cut-off value of the assay for the classifica- tion of cows as not pregnant for cows
23 in clusters 2, 3,

24
25
26 and 4, whereas more than half of the cows in cluster 5 had increased plasma PAG levels. We conclude
27 that at least half of the nonpregnant cows that maintained their CL until 32 d after TAI were initially pregnant
28 but underwent early pregnancy loss based on increased plasma PAG levels at 25 and 32 d after TAI.

29 **Key words:** corpus luteum, progesterone, pregnancy loss, pregnancy-associated glycoproteins

31 INTRODUCTION

32 The use of fertility programs for synchronization of ovulation and timed AI (TAI) has increased service
33 rate and pregnancies per AI (P/AI) in high-producing lactating dairy cows at first service (Fricke et al., 2015;

34 Santos et al., 2016b). Nonetheless, 50 to 70% of cows fail to conceive after first service (Brusveen et al.,
35 2009; Carvalho et al., 2014; Stevenson et al., 2014). Because of poor detection and expression of estrus in
36 high-producing dairy cows (Lopez et al., 2004), coupling a nonpregnancy diagnosis with a management
37 strategy to rapidly resubmit cows to AI increases reproductive efficiency by decreasing the interval between
38 AI services (Fricke, 2002; Fricke et al., 2003, 2016).

39 A common reproductive management strategy to decrease the interval between AI services is weekly sub-
40 mission of nonpregnant cows to an Ovsynch protocol for resynchronization of ovulation (**Resynch**) and
41 TAI (Caraviello et al., 2006; Fricke et al., 2014; Santos et al., 2016a). A key factor limiting fertility during
42 the Ovsynch protocol is the response to each of the sequential hormonal treatments (Giordano et al., 2012b;
43 Carvalho et al., 2015b; Fricke et al., 2015). In this regard, initiation of an Ovsynch protocol during early
44 diestrus (d 5–9) results in more P/AI than initiation of an Ovsynch protocol at other stages of the estrous cycle
45 (Vasconcelos et al., 1999). Lactating dairy cows have an estrous cycle length of approximately 23 d (Sartori
46 et al., 2004); therefore, initiation of a Resynch protocol 32 d after AI (i.e., d 9 of the estrous cycle) should
47 result in more P/AI than initiation of a Resynch protocol

9851

25 d (i.e., d 2 of the estrous cycle) or 39 d (i.e., d 16 of the estrous cycle) after AI. Several studies have
compared initiation of a Resynch protocol at different intervals after AI and reported no differences in P/AI
for cows initiating a Resynch protocol 25 versus 32 d after insemination (Fricke et al., 2003; Silva et al., 2009;
Bruno et al., 2014) or 32 versus 39 d after insemination (Bilby et al., 2013; Lopes et al., 2013). A consistent and
intriguing observation in all of these studies is that the proportion of cows without a corpus luteum (**CL**) or with
low progesterone (**P4**) concentrations at initiation of a Resynch protocol does not differ between cows ini-
tiating the Resynch protocol at 25 versus 32 d (Fricke et al., 2003; Silva et al., 2009; Bruno et al., 2014) or at 32
versus 39 d (Bilby et al., 2013; Lopes et al., 2013) after insemination and ranges from 20 to 30% of the cows
diagnosed not pregnant.

An analysis of 42,252 cows in 159 herds across England and Wales reported that the distribution of
interservice intervals in lactating dairy cows is longer than the expected 18 to 24 d, with a significant pro-
portion of cows having extended interservice intervals (Remnant et al., 2015). Longer interservice intervals,
particularly in high-producing dairy cows, may result from successful conception and maintenance of the CL
followed by early pregnancy loss and luteal regres- sion, thereby delaying return to estrus (Diskin et al., 2011).
Pregnancy-associated glycoproteins (**PAG**) be- long to a large family of aspartic proteinases expressed by the
placenta of domestic ruminants including cows, ewes, and goats (Haugejorden et al., 2006; Wallace et al.,

2015), and assays for detecting PAG levels in blood and milk have been developed and commercialized to determine pregnancy status in cattle (Ricci et al., 2015). Further, circulating PAG levels are dramatically affected by pregnancy loss early during pregnancy and have been used to define pregnancy loss in dairy cows (Giordano et al., 2012; Ricci et al., 2015).

A better understanding of luteal dynamics after synchronization of ovulation and TAI may help refine strategies for reinseminating cows failing to conceive. Thus, the objective of our study was to evaluate P4 concentrations and luteal dynamics from 4 to 32 d after TAI and PAG levels at 25 and 32 d after TAI. Our hypothesis was that luteal regression after TAI among cows diagnosed nonpregnant would not be synchronous and that some cows would have extended luteal phases and increased plasma PAG levels.

MATERIALS AND METHODS

All animal handling procedures were approved by the Animal Care and Use Committee of the College of Agricultural and Life Sciences at the University of Wisconsin–Madison.

Synchronization of Ovulation and Timed AI

Cows were housed at the University of Wisconsin–Madison Dairy Cattle Research Center (Arlington, WI) in loose housing with headlocks. Cows were fed ad libitum a TMR formulated to meet or exceed NRC (2001) requirements for high-producing dairy cows. Primiparous cows were housed separately from multiparous cows. The rolling herd average was 13,880 kg and average daily milk production was 41.6 kg/cow per day with 3.8% fat and 3.2% protein for the herd during the study period. Lactating Holstein cows ($n = 141$; 41 primiparous and 100 multiparous) from 53 ± 3 DIM were synchronized for first TAI using a Double-Ovsynch protocol (Souza et al., 2008). Prostaglandin $F_{2\alpha}$ (25 mg/dose of dinoprost tromethamine; Lutalyse) and GnRH (100 μ g/dose of gonadorelin hydrochloride; Factrel) were from Zoetis (Madison, NJ). Briefly, cows received the first GnRH treatment of the Presynch portion of the Double-Ovsynch protocol at 53 ± 3 DIM, followed by treatment with $PGF_{2\alpha}$ 7 d later and a GnRH treatment 72 h after $PGF_{2\alpha}$. Seven days later, cows received an Ovsynch-56 protocol (GnRH at 70 ± 3 DIM, $PGF_{2\alpha}$ 7 d later, GnRH 56 h after $PGF_{2\alpha}$, and AI 16–20 h later), and all cows received a TAI at 80 ± 3 DIM. Three experienced AI technicians performed all inseminations using sires with high genetic merit and proven fertility.

Blood Sampling, P4, and PAG Assays

Blood samples were collected thrice weekly (Monday, Wednesday, and Friday) from 4 to 32 d after TAI (Figure 1) for analysis of plasma P4 concentrations and at 25 and 32 d after TAI for evaluation of plasma PAG levels. Samples were collected by venipuncture of the median coccygeal artery or vein into 10-mL evacuated plasma collection tubes (Vacutainer; BD, Franklin Lakes, NJ). Blood samples were immediately placed on ice and were centrifuged ($1,600 \times g$; 4°C) for 20 min, and plasma was harvested and stored at -20°C in 2-mL Safe-Lock tubes (Eppendorf AG, Hamburg, Germany). Plasma P4 concentrations were assayed using a solid-phase, no-extraction RIA (Coat-a-Count; Diagnostic Products Corp., Los Angeles, CA). The average sensitivity for the 3 assays was 0.027 ng/mL. The average intra-assay coefficient of variation was 5.61%, and the interassay coefficient of variation was 6.91% based on a quality control sample (2.50 ng/mL of P4) that was replicated within each assay. Plasma PAG levels were assayed for samples collected 25 and 32 d after TAI. After completion of sample collection at the end of the experiment, frozen plasma samples were shipped overnight in a cooled container by courier from the University of Wisconsin to Idexx Laboratories (Westbrook, ME) for analysis of plasma PAG levels using a commercial ELISA assay (Idexx Bovine Pregnancy Test, Idexx Laboratories). Trained technicians, who were blind to the pregnancy outcomes of the individual cows, performed the plasma PAG ELISA assays according to the manufacturer's instructions. Briefly, a microtiter plate format was configured by coating an anti-PAG monoclonal antibody onto the plate. The PAG monoclonal antibody was raised against the PAG-55 protein fraction comprising PAG-4, PAG-6, PAG-9, PAG-16, PAG-18, and PAG-19 (Nagapan et al., 2009). After incubation of the diluted test sample in the coated well, captured PAG was detected with a PAG-specific antibody (detector solution) and horseradish peroxidase conjugate. Unbound conjugate was washed away, and 3,3',5,5'-tetramethylbenzidine substrate was added to the wells. Color development was proportional to the amount of PAG in the sample and was measured using a spectrophotometer. Results were calculated from the optical density (OD) of the sample corrected by subtraction of the reference wavelength OD of the sample (S) minus the OD of the negative control (N) at 450 nm (with both values corrected by subtraction of the reference wavelength OD of the negative control), which resulted in an S - N value. Each microplate included negative and positive controls. Pregnancy outcomes from plasma PAG assays for individual cows were determined based on cut-off values set by the PAG ELISA manufacturer. For the plasma PAG ELISA, when the S - N value was <0.300 , the cow was classified as not pregnant; when the S - N value was >0.300 to <1.000 , the cow was classified as recheck; and when the S - N value was ≥ 1.000 , the cow was classified as pregnant.

Ultrasound Evaluation of CL Volume and Pregnancy Diagnosis

Ovarian structures were evaluated using a portable scanner (Ibex Pro, E. I. Medical Imaging, Loveland, CO)

equipped with a 7.5-MHz linear-array transducer thrice weekly (Monday, Wednesday, and Friday) from 4 to 32 d after TAI to measure CL diameter (Figure 1). The ultrasound image was frozen when the CL was visualized to be at its maximal size, and CL diameter was determined by taking the mean of 2 perpendicular diameter measurements [length (L) and width (W)] obtained using the built-in digital calipers of the ultrasound scanner. When present, CL cavities were measured, mapped, and recorded. Total CL volume was calculated by the formula $V = 4/3 \times \pi \times R^3$ using a radius (R) obtained with the formula $R = (L/2 + W/2)/2$ (Carvalho et al., 2015c). When a CL cavity was present, the volume of the cavity was calculated using the same formula, and the volume of the cavity was subtracted from the total CL volume to estimate total luteal tissue volume. Pregnancy diagnosis was performed 32 d after TAI for all cows using transrectal ultrasonography using the same scanner and transrectal probe used for evaluation of ovarian structures. A positive pregnancy diagnosis was based on visualization of a CL on the ovary ipsilateral to the fluid-filled uterine horn containing an embryo with a heartbeat.

Statistical Analyses

All statistical analyses were performed using SAS computational software (version 9.3 for Microsoft Windows; SAS Institute Inc., Cary, NC). Four cows were removed from all analyses because they failed to ovulate after the final GnRH treatment of the Double-Ovsynch protocol. For individual cows diagnosed nonpregnant 32 d after TAI, the day of CL regression was defined as the day P4 first decreased to <1.0 ng/mL. The Cluster procedure of SAS was used to hierarchically cluster




Figure 1. Schematic representation of the sample collection period. Lactating Holstein cows were submitted to a Double-Ovsynch protocol for synchronization of ovulation for first timed AI (TAI). Blood samples were collected for analysis of plasma progesterone (P4) levels, and ovaries were evaluated using transrectal ultrasonography (US) thrice weekly from 4 to 32 d after TAI. Pregnancy-associated glycoprotein (PAG) levels were assessed at 25 and 32 d after TAI.

the observations. The Cluster procedure segregated nonpregnant cows into 5 clusters (Figure 2); however, because cluster 1 included only 1 cow that underwent CL regression 15 d after TAI, this cow was removed from all subsequent analyses. Cows diagnosed pregnant 32 d after TAI were analyzed as a separate cluster.

Progesterone and CL volume were analyzed using the Mixed procedure of SAS. The model contained the fixed effects of cluster, time, parity (primiparous vs. multiparous), and cluster \times time interaction. Circulating PAG levels were also analyzed using the Mixed procedure of SAS. The model contained the fixed effects of cluster, parity (primiparous vs. multiparous), and cluster \times parity interaction. A contrast statement was used to compare pregnant and nonpregnant cows. Continuous data (P4, CL volume) are presented as least squares means \pm standard error of the mean, whereas binomial data are presented as percentages. A significant difference between levels of a classification variable was considered when $P \leq 0.05$, whereas differences between $P > 0.05$ and $P \leq 0.10$ were considered a statistical tendency.

RESULTS AND DISCUSSION

Luteal Dynamics from 4 to 32 d After TAI

Overall, 41.6% (57/137) of cows were diagnosed pregnant 32 d after TAI, and pregnant cows exhibited plasma




Figure 2. Day of corpus luteum (CL) regression for cows ($n = 80$) diagnosed nonpregnant 32 d after timed AI (TAI). Cows were submitted to a Double-Ovsynch protocol for synchronization of ovulation for first TAI. Day of CL regression was determined based on the day progesterone first decreased to <1.0 ng/mL. Cows were segregated into 5 clusters (C1–C5) based on a statistical cluster analysis of the day of CL regression. Cluster 1, n

= 1; cluster 2, n = 44; cluster 3, n = 14; cluster 4, n = 4; cluster 5, n = 17. NR = cows that did not regress their original CL by 32 d after TAI.




Figure 3. Progesterone (P4) and corpus luteum (CL) volume profiles (LSM \pm SEM) of cows (n = 137) diagnosed nonpregnant 32 d after insemination in clusters 2 to 5 and cows diagnosed pregnant 32 d after insemination from 4 to 32 d after timed AI (TAI). Cows were submitted to a Double-Ovsynch protocol for synchronization of ovulation for first TAI. Cluster 2 = cows undergoing CL regression from 18 to 22 d after TAI, n = 44; cluster 3 = cows undergoing CL regression from 25 to 27 d after TAI, n = 14; cluster 4 = cows undergoing CL regression from 29 to 32 d after TAI, n = 4; cluster 5 = cows that did not undergo CL regression by 32 d after TAI, n = 17; pregnant = cows diagnosed pregnant 32 d after TAI, n = 57. Asterisks (*) above the X-axis in each panel denote the day when CL reached the last statistical increase before CL regression. The arrow with broken line indicates the day when P4 concentrations reached the last statistical increase before CL regression

P4 profiles and CL volume consistent with maintenance of CL function until 32 d after TAI (Figure 3). By contrast, the statistical cluster analysis hierarchically clustered the nonpregnant cows (58.4%; 80/137) into 5 clusters based on the day P4 concentrations decreased to <1.0 ng/mL (Figure 2).

Cluster 1 included only 1 cow (0.73%; 1/137) that underwent CL regression 15 d after TAI (data not shown). Short luteal phases can and do occur in lactating dairy cows. The occurrence of short luteal phases is associated with cyclicity status at the time of submission to an Ovsynch protocol, with 6% of cycling cows and 23% of anovular cows having a luteal phase of <11 d (Gümen et al., 2003). When first postpartum ovulations were excluded from the analysis in another experiment, only 0.5% of cycling cows had a short luteal phase (Opsomer et al., 1998). In the present study, all cows were submitted to a Double-Ovsynch protocol, and cows submitted to a Double-Ovsynch protocol had more CL (94 vs. 68%) at initiation of the Breeding-Ovsynch portion of the protocol than cows submitted to a Presynch-Ovsynch protocol (Ayres et al., 2013). Nonetheless, the cow in cluster 1 ovulated to the last GnRH treatment of the Double-Ovsynch protocol but underwent CL regression earlier than all other nonpregnant cows.

Cows in cluster 2 included nonpregnant cows (32%; 44/137) that underwent CL regression from 18 to 22 d after TAI, and cows in cluster 3 included nonpregnant cows (10%; 14/137) that underwent CL regression from 25 to 27 d after TAI (Figure 3). Differences in timing of luteal regression between cows in clusters 2 and 3 may be attributable to differences in follicular wave patterns among cows. The number of follicular waves per cycle affects estrous cycle length and timing of luteolysis in both lactating dairy cows and dairy heifers (Townson et al., 2002; Ginther and Bashir, 2013; Ginther et al., 2013). Townson et al. (2002) reported that lactating dairy cows with 3 follicular waves had longer estrous cycles (24.5 vs. 21.2 d) and luteal phases (19.3 vs. 17.6 d) than cows with 2 follicular waves. Similarly, Ginther and Bashir (2013) reported that Holstein heifers with 3 follicular waves had longer estrous cycles (23.0 vs. 20.4 d) and luteal phases (20.0 vs. 17.8 d) than heifers with 2 follicular waves, and Sartori et al. (2004) reported longer estrous cycles (23.1 vs. 20.7 d) and luteal phases (19.1 vs. 17.4 d) in Holstein heifers with 3 versus 2 follicular waves.

Interovulatory interval and timing of ovulation relative to CL regression can vary widely among dairy heifers and lactating dairy cows. Sartori et al. (2004) compared ovarian function and circulating steroids during the estrous cycle of Holstein heifers and lactating dairy

cows. Lactating cows with typical cycles had an estrous cycle length of approximately 23 d. Interestingly, some cows were classified as having atypical estrous cycles. Cows with atypical cycles underwent CL regression at the expected time of spontaneous luteolysis; however, ovulation occurred approximately 14

d after luteolysis, whereas cows with typical cycles ovulated approximately 4.9 d after luteolysis (Sartori et al., 2004). In the present study, 74% (59/80; i.e., cows in clusters 1, 2, and 3) of nonpregnant cows would be classified as having typical cycles after TAI according to Sartori et al. (2004).

Cows in cluster 4 included cows (3%; 4/137) that underwent CL regression from 29 to 32 d after TAI, and cows in cluster 5 included cows (12%; 17/137) that maintained the CL formed by the last GnRH treatment of the Double-Ovsynch protocol until 32 d after TAI (Figure 3). These cows (26%; 21/80 of the nonpregnant cows—i.e., cows in clusters 4 and 5), however, did not exhibit typical or atypical cycles as defined by Sartori et al. (2004). Rather, cows in cluster 4 had extended luteal phases and underwent delayed CL regression (Sartori et al., 2004; Ginther and Bashir, 2013), whereas cows in cluster 5 had not undergone CL regression by 32 d after TAI (Figure 3).

Differences might exist between the present study and previous studies (Sartori et al., 2004; Ginther and Bashir, 2013) that evaluated the length of the luteal phase. In our study, all cows were inseminated, whereas the studies by Sartori et al. (2004) and Ginther and Bashir (2013) used lactating cows and heifers that were not inseminated. Nonetheless, the luteal regression distribution of cows in our experiment that included cows with extended luteal phases (Figure 2) is similar to the interservice interval distribution from a large epidemiological data set of dairy cows in the United Kingdom (Remnant et al., 2015).

PAG 25 and 32 d After TAI

Pregnancy-associated glycoproteins belong to a large family of aspartic proteinases expressed by the placenta of domestic ruminants, including cows, ewes, and goats (Haugejorden et al., 2006; Telugu et al., 2010; Wallace et al., 2015), and circulating PAG levels in blood can be used as an indirect method for pregnancy diagnosis (Sasser et al., 1986; Fricke et al., 2016). For pregnant cows, PAG levels begin to increase in circulation by approximately 25 d in gestation, with an early peak occurring 7 d later at 32 d after TAI (Giordano et al., 2012a; Ricci et al., 2015; Carvalho et al., 2017). In addition, between 7 and 14 d is required for circulating PAG and pregnancy-specific protein B (PSPB) levels in cows undergoing induced or spontaneous pregnancy loss to return to baseline levels similar to nonpregnant cows (Giordano et al., 2012a; Ricci et al., 2015).

Plasma PAG levels differed among clusters at both 25 ($P = 0.03$) and 32 ($P < 0.01$) d after TAI (Figure 4). At 25 d after TAI, 95% (54/57) of pregnant cows had detectable plasma PAG levels [37% (21/57) and 58% (33/57) of the pregnant cows were categorized as re-check and pregnant, respectively, by the assay], whereas by 32 d after TAI all pregnant cows had plasma PAG levels above the threshold cut-off set

for a positive pregnancy outcome for this ELISA. Although no cows in clusters 1, 2, 3, or 4 had detectable plasma PAG levels at 25 or 32 d after TAI, 41 and 58% of cows in cluster 5 had detectable plasma PAG levels at 25 and 32 d after TAI, respectively. Based on plasma PAG levels, more than half of the cows in cluster 5 maintained their original CL because they became pregnant and then underwent early pregnancy loss. In addition, it is likely that some or most cows in cluster 5 without detectable




Figure 4. Scatter plot distribution of relative pregnancy-associated glycoprotein (PAG) levels in plasma at 25 and 32 d after insemination for individual cows ($n = 137$). Cows were submitted to a Double- Ovsynch protocol for synchronization of ovulation for first timed AI (TAI). Cows were segregated into clusters based on a statistical clusteranalysis of the day of corpus luteum (CL) regression. Cluster 2 (C2) = cows undergoing CL regression from 18 to 22 d after TAI, $n = 44$; cluster 3 (C3) = cows undergoing CL regression from 25 to 27 d after TAI, $n = 14$; cluster 4 (C4) = cows undergoing CL regression from 29 to 32 d after TAI, $n = 4$; cluster 5 (C5) = cows that did not undergo CL regression by 32 d after TAI, $n = 17$; pregnant = cows diagnosed pregnant 32 d after TAI, $n = 57$. Horizontal lines denote the cut-off values of the assay for the classification of cows as not pregnant when $S - N$ values are <0.300 (dotted line) or pregnant when $S - N$ values are ≥ 1.000 (dashed line). Cows were classified as recheck when the $S - N$ value was >0.300 to <1.000 . Results were calculated from the optical density (OD) of the sample corrected by subtraction of the reference wavelength OD of the sample (S) minus the OD of the negative control (N) at 450 nm (with both values corrected by subtraction of the reference wavelength OD of the negative control), which resulted in an $S - N$ value.

PAG levels 25 or 32 d after TAI could have undergone early pregnancy loss shortly after maternal recognition of pregnancy (Wijma et al., 2016), resulting in enough time to clear residual PAG from circulation before 25 to 32 d after TAI. Nonetheless, not all cows in cluster 5 had detectable PAG levels at 25 or 32 d after TAI in the present experiment.

Although pregnancy loss in lactating dairy cows can occur throughout gestation, the greatest incidence of pregnancy loss occurs during early gestation (Santos et al., 2004; Fricke et al., 2016; Wiltbank et al., 2016). Wiltbank et al. (2016) reported a 20 to 50% pregnancy loss rate during the first week of gestation and a 25 to 41% pregnancy loss rate between 8 and 27 d after AI compared with a 12% pregnancy loss rate between 28 and 60 d and an approximately 2% pregnancy loss rate from 60 d until calving. Similarly, Santos et al. (2004) reported a 34% pregnancy loss rate between 2 and 8 d after AI and a 20% pregnancy loss rate between 6 and 28 d after AI compared with a 12.5% pregnancy loss rate between 28 and 42 d, an 8.5% pregnancy loss rate between 42 and 90 d, and a 6% pregnancy loss rate from 90 d

until calving. Cows undergoing early pregnancy loss have extended luteal phases (>27 d; Wijma et al.,¹¹ 2016). In addition, cows undergoing late embryonic loss have increased interferon- τ stimulated genes 18 to 22 d after AI and increased PSPB concentrations in circulation beginning 24 d after AI, whereas cows undergoing early pregnancy loss have increased interferon- τ stimulated genes 18 to 22 d after AI but may not have increased PSPB concentrations (Wijma et al., 2016). In the present study, nonpregnant cows in cluster 5 that maintained their CL until 32 d after TAI would result in a 23% (17/74) rate of early pregnancy loss that occurs before pregnancy diagnosis is routinely conducted on commercial dairy farms.

Luteal Dynamics for Pregnant and Nonpregnant Cows from 4 to 15 d After TAI

To compare luteal dynamics before spontaneous luteolysis among clusters, plasma P4 concentrations and CL volume were statistically analyzed from 4 to 15 d after TAI (Table 1). Overall, P4 concentrations did not differ ($P = 0.18$) among clusters (Table 1) or parities ($P = 0.22$) from 4 to 15 d after TAI; however, pregnant cows tended ($P = 0.06$) to have greater P4 concentrations from 8 to 13 d after TAI (Table 1). Interestingly, mean plasma P4 concentrations reached maximum concentrations at 15 d after TAI for all clusters (Figure 3). Overall, CL volume did not differ ($P = 0.92$) among clusters ($P = 0.25$) or parities from 4 to 15 d after TAI. In addition, CL volume did not differ ($P = 0.96$) between pregnant and nonpregnant cows (Table 1). The day of maximal CL volume occurred earlier (from 6 to 11 d after TAI; Figure 3) than the day of maximal P4 concentrations (15 d after TAI; Figure 3).

There is an unequivocal requirement for P4 during early pregnancy to support embryo development and for establishment and maintenance of pregnancy (Spencer et al., 2016). Observational studies have consistently reported that pregnant cows have greater P4 concentrations after AI than nonpregnant cows (Sterry et al., 2009; Parr et al., 2012). In this regard, supplementation with exogenous P4 increases circulating P4 concentrations and induces changes in the endometrial transcriptome (Forde et al., 2009; Carter et al., 2010), leading to increased embryo elongation between 13 and 16 d of gestation (Carter et al., 2008; Clemente et al., 2009). By contrast, decreasing P4 concentrations after AI decreases expression of interferon- τ stimulated gene 15 at 20 d after AI, PSPB concentrations from 25 to 67 d after AI, and embryonic crown-rump length 46 d after AI (Carvalho et al., 2017). Results from field trials evaluating the effect of supplementation with exogenous P4, however, are equivocal, with some reporting an increase (Monteiro et al., 2014), no change (Stevenson et al., 2007; Monteiro et al., 2015), or a decrease (Parr et al., 2014) in P/AI. In addition, decreasing P4 concentrations after AI did not affect P/AI or pregnancy loss (Carvalho et al., 2017).

Implications for Reinsemination Strategies

Accurate identification of cows returning to estrus from 18 to 24 d after AI is an early, simple, and inexpensive method for identification and reinsemination of nonpregnant cows (Fricke et al., 2016). Data from the present study highlight several issues that limit this strategy as the only method for reinseminating nonpregnant cows. First, estrous cycle duration varies widely among individual cows, with many cows exhibiting extended interservice intervals (Remnant et al., 2015). Data from the present study support that 26.3% of inseminated cows would be expected to exhibit extended luteal phases and have a delayed return to estrus beyond 24 d. Second, only 51.5% of eligible cows were detected in estrus and inseminated in a study in which detection of estrus was performed through continuous monitoring with an activity monitoring system from AI until a pregnancy diagnosis conducted 32 d after AI (Giordano et al., 2015). A phenomenon referred to as the phantom cow syndrome occurs in seasonally calving herds in Australia and New Zealand in which cows inseminated on the mating start date fail to be detected in estrus within 24 d after AI and are subsequently diagnosed not pregnant 35 d after AI (Cavaliere et al., 2003). Data from the present study support that 21.3% of nonpregnant cows maintained their original CL until 32 d after insemination. Thus, around 1 in 5 inseminated cows would not be expected to return to estrus within 32 d after insemination despite the most accurate and aggressive strategies for detection of estrus. Data on return to estrus after insemination support the need for a resynchronization strategy to reinseminate nonpregnant cows failing to be detected in estrus within 32 d after insemination. A key aspect for fertility to a Resynch protocol is the presence or absence of a CL at the onset of the protocol (Carvalho et al., 2014)

Table 1. Plasma progesterone concentrations and corpus luteum (CL) volume from 4 to 15 d after timed AI (TAI) in lactating Holstein cows¹

Item	CL cluster				Pregnant	P-value	
	Cluster 2	Cluster 3	Cluster 4	Cluster 5		Cluster	C1
n	44	14	4	17	57		
Progesterone (ng/mL)							
d 4	0.9 ± 0.3	0.8 ± 0.5	1.4 ± 0.9	0.8 ± 0.5	1.2 ± 0.3	0.39	0.25
d 6	2.3 ± 0.3	1.9 ± 0.5	2.1 ± 0.9	2.1 ± 0.5	2.3 ± 0.3	0.56	0.14
d 8	3.7 ± 0.3	2.9 ± 0.5	3.8 ± 0.9	3.1 ± 0.5	4.0 ± 0.3	0.11	0.06
d 11	5.0 ± 0.3	4.6 ± 0.5	4.8 ± 0.9	4.3 ± 0.5	5.5 ± 0.3	0.22	0.05
d 13	5.5 ± 0.3	5.5 ± 0.5	5.1 ± 0.9	4.9 ± 0.5	6.0 ± 0.3	0.24	0.05
d 15	6.3 ± 0.3	6.5 ± 0.5	7.0 ± 0.9	5.7 ± 0.5	7.1 ± 0.3	0.42	0.23
CL volume (cm ³)							
d 4	3.3 ± 1.1	3.7 ± 1.9	3.7 ± 2.8	3.4 ± 1.4	3.5 ± 0.8	0.95	0.56
d 6	8.5 ± 0.8	6.9 ± 1.5	8.2 ± 2.5	9.4 ± 1.2	8.4 ± 0.7	0.73	0.84
d 8	10.6 ± 0.8	9.7 ± 1.5	9.7 ± 2.5	11.1 ± 1.3	9.8 ± 0.7	0.71	0.39
d 11	11.1 ± 0.8	12.9 ± 1.3	10.6 ± 2.5	12.6 ± 1.2	11.7 ± 0.7	0.83	0.82
d 13	12.0 ± 0.8	11.1 ± 1.4	11.5 ± 2.5	11.1 ± 1.2	12.0 ± 0.7	0.99	0.84
d 15	11.8 ± 0.8	13.7 ± 1.3	10.6 ± 2.5	11.9 ± 1.2	10.8 ± 0.7	0.27	0.13

¹Cluster 2 = cows undergoing luteal regression 18 and 22 d after TAI; cluster 3 = cows undergoing luteal regression 25 and 27 d after TAI; cluster 4 = cows undergoing luteal regression 29 and 32 d after TAI; cluster 5 = cows that did not undergo luteal regression by 32 d after TAI; C1 = preplanned contrast between pregnant and nonpregnant cows (clusters 2, 3, 4, and 5).

Bisinotto et al., 2015; Carvalho et al., 2015a). Further, fertility of cows lacking a CL at the onset of a Resynch protocol can be increased by administering exogenous P4 during the Resynch protocol (Bilby et al., 2013; Bisinotto et al., 2015). An initial strategy to optimize fertility to Resynch protocols was to empirically determine the optimal interval after TAI to initiate Resynch based on assumptions regarding the physiology of the estrous cycle (Fricke et al., 2003). Assuming an estrous cycle duration of 21 to 23 d, initiation of a Resynch protocol 32 d after TAI should ensure that the first GnRH treatment of the Resynch protocol occurs at d 9 to 11 of the estrous cycle, a stage of the cycle when a CL should be present. Nonetheless, the proportion of nonpregnant cows lacking a CL ranges from 20 to 30% among studies (Fricke et al., 2003; Silva et al., 2009; Bilby et al., 2013; Lopes et al., 2013; Bruno et al., 2014). Data from the present study in which P4 profiles and luteal dynamics were defined from 4 to 32 d after TAI support that implementation of a Resynch protocol based on whether or not a nonpregnant cow has a CL is a better strategy for increasing fertility to TAI than basing initiation of a Resynch protocol on a specific day after AI due to variability in CL regression among groups of cows. Nonpregnant cows without a CL can then be submitted to a protocol with exogenous P4 to increase fertility to TAI. We conclude that at least half of the nonpregnant cows that maintained their CL until 32 d after TAI were initially pregnant but underwent early pregnancy loss based on increased plasma PAG levels at 25 and 32 d after TAI.

ACKNOWLEDGMENTS

We thank the University of Wisconsin–Madison dairy herd personnel at the Emmons Blaine Dairy Cattle Research Center (Arlington, WI) for their assistance with data collection and cow handling during the course of this experiment. Our gratitude is extended to Idexx Laboratories (Westbrook, ME) for running the plasma pregnancy-associated glycoprotein ELISA. This research was supported by Idexx Laboratories and the USDA National Institute of Food and Agriculture (Washington, DC) Hatch Project 231440 to P. M. Fricke.

REFERENCES

- Ayres, H., R. M. Ferreira, A. P. Cunha, R. R. Araujo, and M. C. Wiltbank. 2013. Double-Ovsynch in high-producing dairy cows: Effects on progesterone concentrations and ovulation to GnRH treatments. *Theriogenology* 79:159–164.
- Bilby, T. R., R. G. S. Bruno, K. J. Lager, R. C. Chebel, J. G. N. Moraes, P. M. Fricke, G. Lopes Jr., J. O. Giordano, J. E. P. Santos,

- F. S. Lima, J. S. Stevenson, and S. L. Pulley. 2013. Supplemental progesterone and timing of resynchronization on pregnancy outcomes in lactating dairy cows. *J. Dairy Sci.* 96:7032–7042.
- Bisinotto, R. S., L. O. Castro, M. B. Pansani, C. D. Narciso, N. Martinez, L. D. P. Sinedino, T. L. C. Pinto, N. S. Van de Burgwal, H. M. Bosman, R. S. Surjus, W. W. Thatcher, and J. E. P. Santos. 2015. Progesterone supplementation to lactating dairy cows without a corpus luteum at initiation of the Ovsynch protocol. *J. Dairy Sci.* 98:2515–2528.
- Bruno, R. G. S., J. G. N. Moraes, J. A. H. Hernández-Rivera, K. J. Lager, P. R. B. Silva, A. L. A. Scanavez, L. G. D. Mendonça, R. C. Chebel, and T. R. Bilby. 2014. Effect of an Ovsynch56 protocol initiated at different intervals after insemination with or without a presynchronizing injection of gonadotropin-releasing hormone on fertility in lactating dairy cows. *J. Dairy Sci.* 97:185–194.
- Brusveen, D. J., A. H. Souza, and M. C. Wiltbank. 2009. Effects of additional prostaglandin $F_{2\alpha}$ and estradiol-17 β during Ovsynch in lactating dairy cows. *J. Dairy Sci.* 92:1412–1422.
- Caraviello, D. Z., K. A. Weigel, P. M. Fricke, M. C. Wiltbank, M. J. Florent, N. B. Cook, K. V. Nordlund, N. R. Zwald, and C. L. Rawson. 2006. Survey of management practices on reproductive performance of dairy cattle on large US commercial farms. *J. Dairy Sci.* 89:4723–4735.
- Carter, F., N. Forde, P. Duffy, M. Wade, T. Fair, M. A. Crowe, A. C. O. Evans, D. A. Kenny, J. F. Roche, and P. Lonergan. 2008. Effect of increasing progesterone concentration from Day 3 of pregnancy on subsequent embryo survival and development in beef heifers. *Reprod. Fertil. Dev.* 20:368–375.
- Carter, F., F. Rings, S. Mamo, M. Holker, A. Kuzmany, U. Besenfelder, V. Havlicek, J. P. Mehta, D. Tesfaye, K. Schellander, and P. Lonergan. 2010. Effect of elevated circulating progesterone concentration on bovine blastocyst development and global transcriptome following endoscopic transfer of in vitro produced embryos to the bovine oviduct. *Biol. Reprod.* 83:707–719.
- Carvalho, P. D., C. C. Consentini, S. R. Weaver, R. V. Barleta, L. L. Hernandez, and P. M. Fricke. 2017. Temporarily decreasing progesterone after timed artificial insemination decreased expression of interferon-tau stimulated gene 15 (ISG15) in blood leukocytes, serum pregnancy-specific protein B concentrations, and embryo size in lactating Holstein cows. *J. Dairy Sci.* 100:3233–3242.
- Carvalho, P. D., M. J. Fuenzalida, A. Ricci, A. H. Souza, R. Barletta, M. C. Wiltbank, and

- P. M. Fricke. 2015a. Modifications of Ovsynch improve fertility during resynchronization: Evaluation of presynchronization with gonadotropin-releasing hormone 6 d before initiation of Ovsynch and addition of a second prostaglandin $F_{2\alpha}$ treatment. *J. Dairy Sci.* 98:8741–8752.
- Carvalho, P. D., J. N. Guenther, M. J. Fuenzalida, M. C. Amundson, M. C. Wiltbank, and P. M. Fricke. 2014. Presynchronization using a modified Ovsynch protocol or a single gonadotropin-releasing hormone injection 7 d before an Ovsynch-56 protocol for submission of lactating dairy cows to first timed artificial insemination. *J. Dairy Sci.* 97:6305–6315.
- Carvalho, P. D., A. H. Souza, M. C. Wiltbank, and P. M. Fricke. 2015b. Progesterone concentration at each treatment during an Ovsynch protocol affects fertility to timed AI in lactating Holstein cows. *J. Dairy Sci.* 98(Suppl. 1):92.
- Carvalho, P. D., M. C. Wiltbank, and P. M. Fricke. 2015c. Manipulation of progesterone to increase ovulatory response to the first GnRH treatment of an Ovsynch protocol in lactating dairy cows receiving first timed artificial insemination. *J. Dairy Sci.* 98:8800–8813.
- Cavaliere, J., J. Morton, D. P. Nation, G. Hepworth, S. Pino, A. R. Rabeie, and K. L. Macmillan. 2003. Phantom cows: Predisposing factors, causes and treatment strategies that have been attempted to reduce the prevalence within herds. Pages 365–388 in *Proc. Society of Dairy Cattle Veterinarians of the NZVA Annual Conference*. New Zealand Veterinary Association, Wellington, New Zealand.
- Clemente, M., J. de La Fuente, T. Fair, A. Al Naib, A. Gutierrez-Adan, J. F. Roche, D. Rizos, and P. Lonergan. 2009. Progesterone and conceptus elongation in cattle: A direct effect on the embryo or an indirect effect via the endometrium? *Reproduction* 138:507–517.
- Diskin, M. G., M. H. Parr, and D. G. Morris. 2011. Embryo death in cattle: An update. *Reprod. Fertil. Dev.* 24:244–251.
- Forde, N., F. Carter, T. Fair, M. A. Crowe, A. C. O. Evans, T. E. Spencer, F. W. Bazer, R. McBride, M. P. Boland, P. O’Gaora, P. Lonergan, and J. F. Roche. 2009. Progesterone-regulated changes in endometrial gene expression contribute to advanced conceptus development in cattle. *Biol. Reprod.* 81:784–794.
- Fricke, P. M. 2002. Scanning the future—Ultrasonography as a reproductive management tool for dairy cattle. *J. Dairy Sci.* 85:1918–1926.
- Fricke, P. M., D. Z. Caraviello, K. A. Weigel, and M. L. Welle. 2003. Fertility of dairy cows after

- resynchronization of ovulation at three intervals following first timed insemination. *J. Dairy Sci.* 86:3941–3950.
- Fricke, P. M., J. O. Giordano, A. Valenza, G. Lopes Jr., M. C. Amundson, and P. D. Carvalho. 2014. Reproductive performance of lactating dairy cows managed for first service using timed artificial insemination with or without detection of estrus using an activity-monitoring system. *J. Dairy Sci.* 97:2771–2781.
- Fricke, P. M., A. Ricci, J. O. Giordano, and P. D. Carvalho. 2016. Methods for and implementation of pregnancy diagnosis in dairy cows. *Vet. Clin. North Am. Food Anim. Pract.* 32:165–180.
- Fricke, P. M., M. C. Wiltbank, P. D. Carvalho, and J. O. Giordano. 2015. Fertility programs to achieve high 21-d pregnancy rates in high-producing Holstein dairy herds. Pages 15–27 in *Proc. Dairy Cattle Reprod. Council. Conf. Dairy Cattle Reproduction Council, Buffalo, NY.*
- Ginther, O. J., and S. T. Bashir. 2013. Length of the luteal phase and frequency of the four permutations between two or three follicular waves and ipsilateral or contralateral locations of the corpus luteum and preovulatory follicle in heifers. *Theriogenology* 80:114–119.
- Ginther, O. J., S. T. Bashir, H. B. Rakesh, and M. M. Hoffman. 2013. Hormone concentrations temporally associated with contralateral and ipsilateral relationships between the CL and preovulatory follicle during the third follicular wave in heifers. *Theriogenology* 80:738–747.
- Giordano, J. O., J. N. Guenther, G. Lopes Jr., and P. M. Fricke. 2012a. Changes in serum pregnancy-associated glycoprotein, pregnancy-specific protein B, and progesterone concentrations before and after induction of pregnancy loss in lactating dairy cows. *J. Dairy Sci.* 95:683–697.
- Giordano, J. O., M. L. Stangaferro, R. Wijma, W. C. Chandler, and R. D. Watters. 2015. Reproductive performance of dairy cows managed with a program aimed at increasing insemination of cows in estrus based on increased physical activity and fertility of timed artificial inseminations. *J. Dairy Sci.* 98:2488–2501.
- Giordano, J. O., M. C. Wiltbank, J. N. Guenther, R. Pawlisch, S. Bas, A. P. Cunha, and P. M. Fricke. 2012b. Increased fertility in lactating dairy cows resynchronized with Double-Ovsynch compared with Ovsynch initiated 32 d after timed artificial insemination. *J. Dairy Sci.* 95:639–653.
- Gümen, A., J. N. Guenther, and M. C. Wiltbank. 2003. Follicular size and response to Ovsynch versus detection of estrus in anovular and ovular lactating dairy cows. *J. Dairy Sci.* 86:3184–3194.
- Haugejorden, G., S. Waage, E. Dahl, K. Karlberg, J. F. Beckers, and

- E. Ropstad. 2006. Pregnancy associated glycoproteins (PAG) in postpartum cows, ewes, goats and their offspring. *Theriogenology* 66:1976–1984.
- Lopes, G. Jr., J. O. Giordano, A. Valenza, M. M. Herlihy, J. N. Guenther, M. C. Wiltbank, and P. M. Fricke. 2013. Effect of timing of initiation of resynchronization and presynchronization with gonadotropin-releasing hormone on fertility of resynchronized inseminations in lactating dairy cows. *J. Dairy Sci.* 96:3788–3798.
- Lopez, H., L. D. Satter, and M. C. Wiltbank. 2004. Relationship between level of milk production and estrous behavior of lactating dairy cows. *Anim. Reprod. Sci.* 81:209–223.
- Monteiro, P. L. J. Jr., A. B. Nascimento, G. C. S. Pontes, G. O. Fernandes, L. F. Melo, M. C. Wiltbank, and R. Sartori. 2015. Progesterone supplementation after ovulation: Effects on corpus luteum function and on fertility of dairy cows subjected to AI or ET. *Theriogenology* 84:1215–1224.
- Monteiro, P. L. J. Jr., E. S. Ribeiro, R. P. Maciel, A. L. G. Dias, E. Solé Jr., F. S. Lima, R. S. Bisinotto, W. W. Thatcher, R. Sartori, and J. E. P. Santos. 2014. Effects of supplemental progesterone after artificial insemination on expression of interferon-stimulated genes and fertility in dairy cows. *J. Dairy Sci.* 97:4907–4921.
- Nagappan, M., M. McGrath, and R. Schenkel. 2009. Methods for early detection of pregnancy in cows. US pat. no. 7,604,950 B2. Monsanto Technology LLC, assignee.
- NRC. 2001. *Nutrient Requirements of Dairy Cattle*. 7th rev. ed. Natl. Acad. Press, Washington, DC.
- Opsomer, G., M. Coryn, H. Deluyker, and A. de Kruif. 1998. An analysis of ovarian dysfunction in high yielding dairy cows after calving based on progesterone profiles. *Reprod. Domest. Anim.* 33:193–204.
- Parr, M. H., M. A. Crowe, P. Lonergan, A. C. O. Evans, D. Rizos, and M. G. Diskin. 2014. Effect of exogenous progesterone supplementation in the early luteal phase post-insemination on pregnancy per artificial insemination in Holstein-Friesian cows. *Anim. Reprod. Sci.* 150:7–14.
- Parr, M. H., M. P. Mullen, M. A. Crowe, J. F. Roche, P. Lonergan, A. C. O. Evans, and M. G. Diskin. 2012. Relationship between pregnancy per artificial insemination and early luteal concentrations of progesterone and establishment of repeatability estimates for these traits in Holstein-Friesian heifers. *J. Dairy Sci.* 95:2390–2396.
- Remnant, J. G., M. J. Green, J. N. Huxley, and C. D. Hudson. 2015. Variation in the interservice

- intervals of dairy cows in the United Kingdom. *J. Dairy Sci.* 98:889–897.
- Ricci, A., P. D. Carvalho, M. C. Amundson, R. H. Fourdraine, L. Vincenti, and P. M. Fricke. 2015. Factors associated with pregnancy-associated glycoprotein (PAG) levels in plasma and milk of Holstein cows during early pregnancy and their effect on the accuracy of pregnancy diagnosis. *J. Dairy Sci.* 98:2502–2514.
- Santos, J. E. P., W. W. Thatcher, R. C. Chebel, R. L. A. Cerri, and K. N. Galvão. 2004. The effect of embryonic death rates in cattle on the efficacy of estrus synchronization programs. *Anim. Reprod. Sci.* 82–83:513–535.
- Santos, V. G., P. D. Carvalho, C. Maia, B. Carneiro, A. Valenza, P. M. Crump, and P. M. Fricke. 2016a. Adding a second prostaglandin F2 α treatment to but not reducing the duration of a PRID-Synch protocol increases fertility after resynchronization of ovulation in lactating Holstein cows. *J. Dairy Sci.* 99:3869–3879.
- Santos, V. G., P. D. Carvalho, C. Maia, B. Carneiro, A. Valenza, and P. M. Fricke. 2016b. Fertility of lactating Holstein cows after synchronization of ovulation and timed artificial insemination versus artificial insemination after detection of estrus at a similar DIM range. *J. Dairy Sci.* 99(E-Suppl.):499.
- Sartori, R., J. M. Haughian, R. D. Shaver, G. J. M. Rosa, and M. C. Wiltbank. 2004. Comparison of ovarian function and circulating steroids in estrous cycles of Holstein heifers and lactating cows. *J. Dairy Sci.* 87:905–920.
- Sasser, R. G., C. A. Ruder, K. A. Ivani, J. E. Butler, and W. C. Hamilton. 1986. Detection of pregnancy by radioimmunoassay of a novel pregnancy-specific protein in serum of cows and a profile of serum concentrations during gestation. *Biol. Reprod.* 35:936–942.
- Silva, E., R. A. Sterry, D. Kolb, N. Mathialagan, M. F. McGrath, J. M. Ballam, and P. M. Fricke. 2009. Effect of interval to resynchronization of ovulation on fertility of lactating Holstein cows when using transrectal ultrasonography or a pregnancy-associated glycoprotein enzyme-linked immunosorbent assay to diagnose pregnancy status. *J. Dairy Sci.* 92:3643–3650.
- Souza, A. H., H. Ayres, R. M. Ferreira, and M. C. Wiltbank. 2008. A new presynchronization system (Double-Ovsynch) increases fertility at first postpartum timed AI in lactating dairy cows. *Theriogenology* 70:208–215.
- Spencer, T. E., N. Forde, and P. Lonergan. 2016. The role of progesterone and conceptus-derived factors in uterine biology during early pregnancy in ruminants. *J. Dairy Sci.* 99:5941–5950

- Sterry, R. A., E. Silva, D. Kolb, and P. M. Fricke. 2009. Strategic treatment of anovular dairy cows with GnRH. *Theriogenology* 71:534–542.
- Stevenson, J. S., S. L. Hill, R. L. Nebel, and J. M. DeJarnette. 2014. Ovulation timing and conception risk after automated activity monitoring in lactating dairy cows. *J. Dairy Sci.* 97:4296–4308.
- Stevenson, J. S., M. A. Portaluppi, D. E. Tenhouse, A. Lloyd, D. R. Eborn, S. Kacuba, and J. M. DeJarnette. 2007. Interventions after artificial insemination: Conception rates, pregnancy survival, and ovarian responses to gonadotropin-releasing hormone, human chorionic gonadotropin, and progesterone. *J. Dairy Sci.* 90:331–340.
- Telugu, B. P. V. L., M. O. Palmier, S. R. Van Doren, and J. A. Green. 2010. An examination of the proteolytic activity for bovine pregnancy-associated glycoproteins 2 and 12. *Biol. Chem.* 391:259–270.
- Townson, D. H., P. C. W. Tsang, W. R. Butler, M. Frajblat, L. C. Griel, C. J. Johnson, R. A. Milvae, G. M. Niksic, and J. L. Pate. 2002. Relationship of fertility to ovarian follicular waves before breeding in dairy cows. *J. Anim. Sci.* 80:1053–1058.
- Vasconcelos, J. L. M., R. W. Silcox, G. J. M. Rosa, J. R. Pursley, and M. C. Wiltbank. 1999. Synchronization rate, size of the ovulatory follicle, and pregnancy rate after synchronization of ovulation beginning on different days of the estrous cycle in lactating dairy cows. *Theriogenology* 52:1067–1078.
- Wallace, R. M., K. G. Pohler, M. F. Smith, and J. A. Green. 2015. Placental PAGs: Gene origins, expression patterns, and use as markers of pregnancy. *Reproduction* 149:R115–R126.
- Wijma, R., M. L. Stangaferro, M. M. Kamat, S. Vasudevan, T. L. Ott, and J. O. Giordano. 2016. Embryo mortality around the period of maintenance of the corpus luteum causes alterations to the ovarian function of lactating dairy cows. *Biol. Reprod.* 95:112.
- Wiltbank, M. C., G. M. Baez, A. Garcia-Guerra, M. Z. Toledo, P. L. J. Monteiro, L. F. Melo, J. C. Ochoa, J. E. P. Santos, and R. Sartori. 2016. Pivotal periods for pregnancy loss during the first trimester of gestation in lactating dairy cows. *Theriogenology* 86:239–253.