



AperTO - Archivio Istituzionale Open Access dell'Università di Torino

Habitat-dependent effects of personality on survival and reproduction in red squirrels

This is the author's manuscript
Original Citation:
Availability:
This version is available http://hdl.handle.net/2318/1691151 since 2019-02-08T09:04:39Z
Published version:
DOI:10.1007/s00265-018-2546-y
Terms of use:
Open Access
Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)

1 Title: Habitat-dependent effects of personality on survival and reproduction in red squirrels

Francesca Santicchia^{a*}, Candice Gagnaison ^b, Francesco Bisi^a, Adriano Martinoli^a, Erik Matthysen ^c, Sandro

2

3

4

5

Bertolino^d, Lucas Armand Wauters^{a, c}

- 6 ^a Environment Analysis and Management Unit - Guido Tosi Research Group - Department of Theoretical and 7 Applied Sciences, Università degli Studi dell'Insubria, via J. H. Dunant 3, 21100 Varese, Italy 8 ^b UFR Life, Earth and Environmental Sciences, Université de Bourgogne, 6 Boulevard Gabriel, 21000 Dijon, 9 France 10 ^c Department of Biology, Evolutionary Ecology Group, University of Antwerp, Universiteitsplein 1, 2610 11 Wilrijk, Belgium 12 ^d Department of Life Sciences and Systems Biology, Università degli Studi di Torino, Via Accademia Albertina 13 13, 10123 Torino, Italy 14 15 *Corresponding author (Santicchia F.): Department of Theoretical and Applied Sciences, Università degli Studi 16 dell'Insubria, via J.H. Dunant, 3 – 21100 – Varese. Italy. Telephone: (+39) 0332 421538. email: 17 f.santicchia@uninsubria.it 18 **ORCID IDs:** 19 Francesca Santicchia: 0000-0003-4814-9632 Francesco Bisi: 0000-0002-4908-5012 20 21 Lucas Wauters: 0000-0002-4871-5035 Sandro Bertolino: 0000-0002-1063-8281 22 Adriano Martinoli: 0000-0003-0298-0869 23
- 24

25 Abstract

26 Variation in intrinsic (body mass, reproductive condition) and extrinsic factors (habitat quality, spatio-27 temporal variation in food availability) can affect the costs and benefits of personality traits. Relationships 28 between personality and fitness components can vary with changes in population density and/or habitat 29 quality. Here, using capture-mark-recapture data of individually marked Eurasian red squirrels (Sciurus 30 vulgaris), we explored whether indices of boldness and exploration are correlated with local survival in either 31 of the sexes and with a measure of reproductive success in females, under different conditions of food 32 availability. PCA scores derived from the number of captures (trappability) and number of different traps 33 where an individual was captured (trap diversity), in the first one or two years of presence, were used as 34 indices of personality (boldness and pure exploration). The relationships between boldness and local survival 35 differed between habitats in both sexes. Bold squirrels survived better than shy ones in Norway spruce forest, 36 while in the other two study areas (Scots pine, or mixed spruce-fir forest) survival was lower for bolder 37 squirrels. Pure exploration behavior was negatively correlated with local survival in all habitats and in both 38 sexes. Female reproductive success increased with body mass and decreased with the tendency to explore, 39 a relationship consistent across habitat types. Bolder females did not have a reproductive advantage in years 40 of poor food availability. We suggest that costs and benefits of boldness vary with spatio-temporal 41 differences in availability of high-quality food resources which may help to maintain variation in personality 42 in red squirrels in heterogeneous landscapes.

43

44 Keywords:

45 boldness, capture-mark-recapture, exploration, habitat variability, *Sciurus vulgaris*, trappability

46

47 Significance statement

Differences in personality among individuals can affect their survival or reproductive success. We investigated the relationship between personality traits and fitness components in Eurasian red squirrels under different food availability. We scored boldness and exploration using indices derived from capture51 mark-recapture data. Bold red squirrels survived longer than shy ones in spruce forest where food 52 availability varied strongly between years. Instead, in mixed forests with more stable food supplies shy 53 individuals survived longer. Heavy females produced more litters in their lifetime than those with lower 54 body mass. Explorative squirrels were more likely to die young and to produce no or few litters, 55 independent of the habitat they lived in. Our results suggest that variation in personality traits within 56 populations of Eurasian red squirrels was maintained by changing fitness benefits in relation to habitat type 57 in pulsed resource systems.

58

59 Introduction

60 According to evolutionary theory, the way an individual behaves under local conditions should be 61 adaptive because of natural selection favoring animals whose behavior increases fitness, given that this 62 behavior is heritable. Variation in behaviors is maintained because of spatio-temporal changes in intrinsic 63 and/or extrinsic environmental conditions causing variation in selective pressures (Carter et al. 2013; Weiss 64 and Adams 2013). However, from the perspective of adaptive behavior, the fact that animals display variation 65 in individual behaviors that are consistent over time (within or across generations) and/or under different 66 contexts is more difficult to explain (Dall et al. 2004; Réale et al. 2007; Biro and Stamps 2008). Animal 67 personality refers to between-individual differences in behavior that persist through time (Biro and Stamps 68 2008; Carter et al. 2013). These differences in individual personality, also called temperament or 'coping 69 style' (Bergmüller and Taborsky 2010), are often measured through traits such as reaction to potential risks 70 or handling novelty, aggressiveness and sociability (e.g. Réale et al. 2007). If differences in personality are 71 adaptive, they may affect how populations respond to environmental change and how species are able to 72 persist (Dall et al. 2004; Le Coeur et al. 2015).

An individual's willingness to take risks, referred to as boldness, has been investigated in many wild populations showing that this personality trait is related to several ecological aspects, such as propensity to explore and acquire information about the environment, dispersal distance and/or speed (Fraser et al. 2001; Korsten et al. 2010; Minderman et al. 2010; van Overveld et al. 2015). Although recent studies have shown

77 that personality can affect several components of fitness (Dingemanse and Réale 2005; Patterson and Shulte-78 Hostedde 2011; Mutzel et al. 2013; Zhao et al. 2016), it remains unclear whether bold phenotypes have a 79 fitness advantage over shy ones. Several studies on birds and mammals found that bold and/or fast exploring 80 individuals are often the most dominant ones during intraspecific encounters (Dingemanse and de Goede 81 2004; Rödel et al. 2015). For example, fast-exploring males of the great tit (*Parus major*) occupied better 82 quality territories and fast-fast pairs produced offspring in better condition than slow-exploring pairs (Both 83 et al. 2005). In Norway rats (*Rattus norvegicus*) explorative individuals survived longer than less explorative 84 ones (Cavigelli and McClintock 2003).

85 However, in a taxonomically broad meta-analysis Smith and Blumstein (2008), exploring the effects 86 of exploration and boldness on fitness components (survival and/or reproductive success), reported contrasting findings. Some studies showed that bolder animals encounter more risks due to their exploring 87 88 tendency, which resulted in higher susceptibility to predation and higher mortality risk (e.g. Bremner-89 Harrison et al. 2004; Carter et al. 2010; Hall et al. 2015). In the European rabbit (Oryctolagus cuniculus), explorers had significant survival costs during the juvenile life-stage (Rödel et al. 2015). Still a different 90 91 pattern was revealed in Eastern chipmunks (Tamias striatus) where animals with average exploration levels 92 suffered higher mortality than extreme phenotypes (Bergeron et al. 2013).

Variation in personality may be maintained by fitness trade-offs. For example, bold (active and aggressive) female North American red squirrels (*Tamiasciurus hudsonicus*) were more likely to bequeath their territory, which resulted in higher reproductive success through increased overwinter survival of their offspring. However, bold females had higher winter mortality than shy ones (Boon et al. 2008). In addition, spatio-temporal variation in ecological selection pressures can play a role in maintaining animal personalities in populations in heterogeneous environments, both in short- and long-lived species (Réale et al. 2000; Le Coeur et al. 2015; Vetter et al. 2016).

So far, most studies on personality-fitness relationships have been carried out in a single population and we know only a few studies on mammals that have investigated effects of personality on fitness in different habitats (e.g. Boon et al. 2007; Martin and Réale 2008; Monestier et al. 2015).

103 Here we measured personality traits in the Eurasian red squirrel (Sciurus vulgaris) in three different 104 habitats and investigated to what extent boldness and exploration tendency of adults were correlated with 105 phenotypic traits (body size and body mass) and with fitness components. The red squirrel is a good model 106 species to investigate relationships between personality and fitness components under different habitat 107 conditions for several reasons: 1) previous studies on other tree squirrels have shown that personality traits 108 can affect habitat and/or space use, survival and reproductive success (e.g. Boon et al. 2008; Boyer et al. 109 2010; Le Coeur et al. 2015); 2) reliable methods and indices to study personality have already been developed 110 for other sciurids (Boon et al. 2008; Boyer et al. 2010); 3) long-term capture-mark-recapture data of 111 individually-marked squirrels combined with radio-tracking show high capture probabilities for the majority 112 of resident adults and provide detailed information of phenotype, local survival, and reproductive success 113 (of females) (Wauters et al. 2004, 2007, 2008).

114 Our main hypothesis is that costs and benefits of personality types vary according to environmental 115 conditions. Hence, we predict that in different habitats (with spatio-temporal variation in food availability) 116 different personality types will perform better; in other words, bold squirrels may perform better than shy 117 ones in one habitat type, but not in other habitat(s). To test this hypothesis we first checked for correlations 118 between personality indices and body size and body mass. Since in this species heavier males tend to be 119 more aggressive and have higher mating success (Wauters and Dhondt 1989, 1992; Wauters et al. 1990), 120 while in females body mass positively affects the probability to enter estrus and lifetime reproductive success 121 (Wauters and Dhondt 1989, 1995). Next, we explored potential advantages or disadvantages of being bold 122 by studying whether our indices of boldness and exploration were correlated with local survival in both sexes 123 and with a measure of reproductive success in females, and whether these relationships varied among 124 habitats. Finally, we also tested the hypothesis that being bold might be advantageous for reproduction only 125 in years of poor food availability. We used capture-mark-recapture data of individually marked squirrels to 126 obtain data of sex, body size and body mass, and estimates of local survival and female reproductive success. The total number of captures (trappability) and number of different traps where an individual was captured 127

(trap diversity), over a maximum of two years, were used as indices of the personality traits boldness and
exploration, respectively (Boon et al. 2008; see methods for details).

130

131 Materials and methods

132 Study species

133 Eurasian red squirrels have overlapping home ranges, with intensively used core- areas, whose size 134 differs between the sexes, and in relation to season and habitat quality (e.g. Wauters and Dhondt 1992; Lurz 135 et al. 2000; Wauters et al. 2001, 2005). The mating system is promiscuous, although most females only mate 136 with a dominant male of high body mass (Wauters et al. 1990). Reproduction is seasonal, with 1 to 2 litters per year, and is strongly affected by the female's body condition, food abundance and, in the mountains, 137 138 elevation (Wauters and Dhondt 1995; Wauters et al. 2008; Rodrigues et al. 2010). In our study areas, females 139 can be lactating between April and October and may produce one or two litters/year (Boutin et al. 2006; 140 Wauters et al. 2008). Most animals disperse as juveniles and subadults (from 4 to 10 months old, Wauters 141 and Dhondt 1993; Wauters et al. 1993, 2010). Once settled, adult squirrels tend to have a high site-fidelity. 142 There is no sexual size dimorphism (Wauters et al. 2007).

143

144 Study areas and food availability

145 We studied red squirrels in three study areas in the Italian Alps, two in Lombardy and one in Valle d'Aosta, N. Italy. Cedrasco (CED, 46°07' N, 9°48' E, elevation 1100 – 1600 m a.s.l.) is a mixed montane conifer 146 147 forests on the northern slope of the Orobic Alps, N. Italy. The study area extends over 76 ha of mature 148 secondary forest (mean tree density \pm SD = 449 \pm 163 trees ha⁻¹), with large areas of even-age stands 149 dominated by silver fir (Abies alba, tree cover 56%) and Norway spruce (Picea abies, 25%), with sparse Scots 150 pine (Pinus sylvestris, 5%), larch (Larix decidua, 3%) and dead trees (5%). Spruce and fir were interspersed at 151 a fine-grained level and not clumped. At lower elevations, spruce and fir were mixed with some beech (Fagus 152 sylvatica, 7% tree cover). The second study area, Oga (OGA, 46º28'N, 10º22'E, elevation from 1280 m to 1450 m a.s.l.), is part of a mature mixed montane conifer forest (47 ha, mean tree density \pm SD = 765 \pm 251 153

trees ha⁻¹), dominated by Scots pine (*Pinus sylvestris*, 88.7% tree cover) with some Norway spruce (*Picea abies*, 8.8%) and larch (*Larix decidua*, 2.5%). The third study area, in Val di Rhêmes (RHE, 45°39'N, 7°12'E, elevation 1740 – 1890 m a.s.l.) is a mature Norway spruce forest (85%) with some mixed patches of larch (11%) and spruce (69 ha, mean tree density \pm SD = 773 \pm 333 trees ha⁻¹). Snags (dead trees) are common (4%), occurring throughout the study area. Distances between the areas are CED-OGA 58 km, CED-RHE 217 km, OGA-RHE 271 km (Trizio et al. 2005).

Each year between 26 July – 9 August, the new (current year) maturing cones were counted in the canopy of sample trees of all conifer species to estimate food availability. Methods are described in detail elsewhere (Salmaso et al. 2009; Di Pierro et al. 2011). Based on species-specific energy values per cone (kJ/cone) and data on tree species composition and density (Salmaso et al. 2009), these counts allowed us to estimate the average yearly tree seed-energy production in each study area (Table 1).

The three populations had similar sex ratio's and age structure (Wauters et al. 2007, 2008). There were no differences in allelic richness between areas, but squirrels at RHE had slightly lower observed heterozygosity than at CED and OGA (10 microsatellites, Trizio et al. 2005). All had goshawk, red fox and marten as potential squirrel predators.

169

170 Trapping and handling

Trapping was carried out in three periods per year (April-May, June-July, September-October) between April 2000 and October 2006 in OGA and between April 2000 and April 2009 in CED and RHE (Table 1). A trapping session involved the use of 25 (CED), 23 (OGA) or 30 (RHE) ground-placed Tomahawk "squirrel" traps (models 201 and 202, Tomahawk Live Trap, WI, USA). Steep slopes and mountain orography made it impossible to set traps in a perfect grid, hence traps were more or less homogeneously distributed over the study area, with distances of 100-130 m between traps and average trap density of 0.7-0.8 traps ha⁻¹.

In CED and RHE the forest continued beyond the boundaries of the study area. In contrast, the study
 area of OGA was entirely surrounded by hard edges (road and parking area, meadows, river). Traps were pre-

baited with sunflower seeds and hazelnuts 4 to 6 times over a 30 day period, then baited and set for 6-10
days, until no new, unmarked squirrels were trapped for at least 2 consecutive days (Wauters et al. 2008).
Traps were checked two-three times per day.

182 Each trapped squirrel was flushed into a light cotton handling bag with a zipper or a wire-mesh 183 "handling cone" to minimize stress during handling, and individually marked using numbered metal ear-tags (type 1003 S, National Band and Tag, Newport, KY, USA). It was weighed to the nearest 5 g using a spring-184 185 balance (Pesola AG, Baar, Switzerland) and the length of right hind foot (without nail) was measured (0.5 186 mm) with a thin ruler. Foot length is a measure of structural size and remains fixed once the animal has 187 reached adult size (12-18 month old). In contrast, body mass can also vary with season, age or between years 188 with poor or high food availability (Wauters et al. 2007). Therefore, in our models (see "Effects of personality 189 on fitness components") foot length as adult and average body mass over the three trapping sessions in the 190 first year the squirrel was classified as adult (12-18 months old, see also Wauters and Dhondt 1989) were 191 used as measures of morphological parameters. For foot length, variance from repeated measures was low 192 (13% of total variance) compared to variance caused by individual differences in size (87% of total variance), 193 indicating that measurement error was small (Wauters et al. 2007). Sex and age class were determined on 194 the basis of external genitalia and body mass, with juveniles weighing less than 250 g (Wauters and Dhondt 195 1995). Female reproductive condition was determined based on external genitalia and condition of the 196 nipples: pregnant females have enlarged vulva and small nipples, lactating females have swollen nipples and 197 milk excretion can be stimulated, post-lactation females have still large nipples but no longer produce milk 198 (Wauters and Dhondt 1995). It was not possible to record data blind because our study involved capture-199 mark-recapture in the field.

Capture probabilities of squirrels in the three populations were calculated using loglinear robust design in package Rcapture (Rivest and Baillargeon 2014). Robust design applies a closed population model for each trapping session. The Poisson regression used to fit a robust design model has one entry for each possible capture history, including those that are unobserved.

204

205

Personality: trappability and trap diversity indices

206 For each individual we calculated two indices of personality: a trappability and trap diversity index. 207 The first is a measure of tendency to take risks (boldness), the second measures the animal's propensity to 208 explore novel environments. In sciurid rodents, these two indices have been shown to be repeatable 209 measures of personality, but they also tend to be correlated (Boon et al. 2008; Boyer et al. 2010). The indices 210 were estimated using the standardized values, per study area, of the ratio of number of captures on the 211 length of capture period (i.e. the number of capture days from the first to the last trapping session for a given 212 animal; boldness); or the ratio of the number of different traps in which an animal was captured on the 213 number of available traps (exploration). Since recent studies have shown that personality traits can change 214 over time (flexible component of personality, e.g. Dingemanse et al. 2012), we only used capture data of the 215 first two years an individual was present (using only subadults and adults). Since we only had three 216 sessions/year, using only first year data would strongly reduce the variation among individuals in trappability 217 and trap diversity. Standardization was done for each study area separately because of differences in capture 218 histories and number of available traps.

219 Since the two standardized indices were strongly and positively correlated (r = 0.76; n = 141; P < 220 0.0001), they could not be used together as explanatory variables in multivariate models because of lack of 221 independence when estimating their partial effects. Therefore, we used a principal component analysis (PCA) 222 to derive two new variables, the principal components PC1 and PC2 (orthogonally rotated), maximizing the 223 variance of these among individual squirrels. PC1 (loadings: 0.707 trappability index + 0.707 trap diversity 224 index, eigenvalue 1.33) explained 88% of total variance, PC2 12% (loadings -0.707 trappability index + 0.707 225 trap diversity index, eigenvalue 0.49). PC1 has a high score for those animals that are often captured and in 226 many different traps, so it can be considered as a measure of boldness which includes both risk taking and 227 exploration. PC2 has a high score for those animals that are captured few times but nearly always in a 228 different trap, hence indicating a specific exploration component of the animal's personality. The fact that 229 PC2 has an eigenvalue below 1 is inherent to a PC-analysis with two variables. The commonly used threshold

of PC's with eigenvalues above 1 is thus not meaningful here to decide which components to retain for further analyses. On the contrary, the aim of our PC-analysis was to generate uncorrelated dimensions of personality that can jointly be included in statistical analyses. We thus argue that as long as PC2 is repeatable over time, it contains biologically relevant information to include in our analyses. Recent data on arena tests (open field test and mirror image stimulation test) with red squirrels showed that trappability and trap diversity indices of personality are confirmed by behavior in the arena test, as in other sciurids (Boon et al. 2008; Boyer et al. 2010; Le Coeur et al. 2015; our unpubl. data).

237 Our data could potentially contain a bias in estimates of trappability, trap diversity and survival, due 238 to a lower capture probability for individuals living at the edge of the study areas (e.g. Boyer et al. 2010). 239 However, position and size of home ranges of 91 radio-tracked squirrels (for details see Wauters et al. 2005; 240 Romeo et al. 2010; Di Pierro et al. 2011) showed that their large home range overlapped with several traps 241 (5 to 10) both at study area periphery and more central ones. Hence, to eliminate the risk of a potential bias 242 we removed from our dataset those animals only caught in edge traps. Radio-tracking data also confirmed 243 the reliability of our survival estimates, with only two (2%) radio-tagged squirrels still present (alive) after the 244 last capture event.

245

246 *Repeatability of trapping indices and PCA scores*

247 To analyse the link between trapping characteristics and individual personality we needed to verify 248 that trapping characteristics were consistent across years for the same individual (i.e. trapping characteristics 249 as an expression of personality, e.g. Boyer et al. 2010). Using a restricted sample of 49 individuals that were 250 caught in two successive years during the study period, we estimated the repeatability, also called Intra-class 251 Correlation Coefficient (ICC), in the number of captures per year and in trap diversity per year with a Linear 252 Mixed Model (LMM) (Nakagawa and Schielzeth 2010). Repeatability was estimated using the R package rptR 253 v 0.9.2 (CI= 95%, number of parametric bootstraps for interval estimation= 5000, number of permutations 254 used when calculating asymptotic p-values= 1000; Stoffel et al. 2017). In this case, number of captures was 255 log transformed (using natural logarithm) and trap diversity (number of different traps) square root

transformed to meet assumptions of normality (Shapiro-Wilk's test on transformed data, both W > 0.95) (see also Boyer et al. 2010). We included sex, study area and year and second order interactions as fixed effects and squirrel identity as random factor. Length of capture period was constant between the two years. We also estimated repeatability of the PC1 and PC2 scores derived from these trapping indices (see below) using the same LMM model structure as above.

261

262 Effects of personality on fitness components

263 To explore relationships between personality and fitness parameters we estimated local survival of 264 squirrels of both sexes and reproductive success of females. Length of residency was used as a proxy for local 265 survival and was estimated as the number of months between first and last capture, for those squirrels that 266 remained on the study area for at least 6 months (see also Wauters et al. 2007). To avoid potential bias in 267 survival estimates due to selective disappearance or between population differences in age structure, we 268 only used squirrels from extinct cohorts (n = 132). Reproductive success was measured as the total number 269 of litters produced per female throughout its lifetime, using only females that were trapped in at least two 270 of three sessions/year (for models on number of litters). In April-May, females could be scored as pregnant 271 or lactating if they had a spring litter. In June-July, as pregnant-lactating in case of a single intermediate litter 272 or an autumn litter or as post-lactating in case of spring litter. In September-October, as lactating or post-273 lactating (autumn or intermediate litter, respectively). Hence, females that were trapped in at least two of 274 three sessions allowed to distinguish between production of no, a single or two litters/year. Since both 275 parameters are counts, we used them as dependent variables in GLM models with Poisson error distribution 276 and used sex, study area, body mass, foot length (as co-variate of body mass, see Wauters et al. 2007), and 277 the boldness and exploration scores (PC1 and PC2) from the PCA as fixed effects, and the second order 278 interactions of study area with PC1 and PC2 (both models) and of sex with PC1 and PC2 (survival model). 279 Although the standard deviation of PC2 was below 1 (which is inherently so when only 2 variables are 280 included in the PC-analysis), we did include it in these analyses because of its biological interpretation 281 (specific exploration) and the fact that PC2, as well as PC1, appeared repeatable over time (see below). When

282 none of these interactions were statistically significant (all partial P > 0.05) we presented models only with 283 main effects. In contrast, if one of the interactions was significant, further model selection was carried on a 284 subset of the data (by study area) to allow reliable estimates of fixed effects (Online Resource 1 Tables S1 – 285 S3). Since patterns of longevity were sex-biased (higher local survival in males than in females; significant sex 286 by PC2 interaction, Online Resource 1 Table S1) the models were tested for males and females separately 287 (see also Wauters et al. 2004). In both sexes, significant interactions were found between study area and 288 PC1: they always indicated a different relationship for study area RHE in comparison to study areas CED and 289 OGA (Online Resource 1 Tables S1 – S3, see also results). Therefore, in these cases separate models for (a) 290 CED and OGA, and (b) RHE were further tested using only the main effects.

291 We also tested the hypothesis that being bold might be advantageous for reproduction only in years 292 of poor food availability (e.g. Le Coeur et al. 2015). We classified seed-crops in two categories: poor vs. 293 medium-good seed crop (Table 1). In each reproductive period (spring and summer-autumn) we determined 294 if a female produced a litter or not (binary dependent variable, 190 observations of 57 different females) and investigated relationships with study area, food availability (poor vs medium-good) and PC1 score along with 295 296 the interaction of PC1 with food availability, using a mixed effects logistic regression model which included 297 squirrel identity as a random effect. Spring litters were related to the previous year's seed-crop, summer-298 autumn litters to the same year's seed-crop (Boutin et al. 2006; Wauters et al. 2008).

Distributions of data on longevity and reproductive success were skewed. Therefore, we repeated the above models using a negative binomial distribution of the dependent variable. Since results were similar as models with a Poisson distribution we only present results of the latter.

All tests of significance are two-tailed and the significance level was set at 0.05. Except for repeatability, all other statistical analyses were done using SAS/STAT 9.4 software (Copyright © 2011, SAS Institute Inc., Cary, NC, USA).

306 Results

307 Trappability and trap diversity indices

In the three study areas, the capture probabilities at population level (including only the animals used
 in this study) were high and stable over time (mean capture probability per year ± SE: CED 0.98 ± 0.05; OGA
 0.96 ± 0.02; RHE 0.98 ± 0.01; details in Table S4).

311 Within a period of maximum two years (see methods), the number of captures per individual red 312 squirrel varied between 2 and 41 times (median 7, mean \pm SE = 9.4 \pm 0.6 captures), and number of different 313 traps from 1 to 13 (median 5, mean \pm SE = 5.2 \pm 0.2 traps). On average, males were trapped more often than 314 females (males, n = 78, 11.2 \pm 1.0 captures; females, n = 63, 7.2 \pm 0.7 captures; $F_{1,135}$ = 11.5, P = 0.0009), and 315 in more different traps (males, n = 78, 5.8 ± 0.4 traps; females, n = 63, 4.4 ± 0.3 traps; $F_{1,135} = 9.90$, P = 0.002). 316 The number of times a squirrel was captured did not differ significantly between study areas ($F_{2, 135}$ = 1.40, P 317 = 0.25); however, on average squirrels were trapped in more different traps in OGA (n = 40, 5.3 \pm 0.5 traps) and RHE (n = 56, 5.8 ± 0.4 traps) than in CED (n = 45, 4.2 ± 0.4 traps; area effect $F_{2, 135}$ = 4.59; P = 0.012; Fig. 318 319 1). These differences between areas did not influence our results since trapping indices were standardized 320 within study areas.

321 We observed consistent individual variation across years in number of captures (n = 49; repeatability 322 = 65%, 95% CI = 46 – 81%, Likelihood ratio test LRT = 25.9; df= 1; P < 0.001) and in trap diversity (repeatability 323 = 69%, 95% CI = 52 – 83%, Likelihood ratio test LRT = 29.9; df= 1; P < 0.001). Thus, trappability and trap 324 diversity indices are consistent through time suggesting they are reliable estimates of a red squirrel's 325 personality. Similarly, highly significant repeatabilities were found for the PC1 and PC2 scores derived from 326 these indices (PC1: repeatability = 78%, 95% CI = 66 – 88%, Likelihood ratio test LRT = 43.9; df= 1; P < 0.0001; 327 PC2 repeatability = 37%, 95% CI = 14 - 63%, Likelihood ratio test LRT = 6.58; df= 1; P = 0.005). In spite of the 328 fact that PC2 explained only 12% of the total variation, it does seem to be moderately repeatable through 329 time and this reflect a biologically real phenomenon. This, in combination with the fact that PC2 has a 330 straightforward biological interpretation of an aspect of squirrel personality favors its use in the statistical 331 analyses below.

332

333 Phenotypic variation and personality

334 Neither foot length, nor body mass were correlated with the personality scores (Table 2). These 335 patterns were consistent when analyzed by sex (Table 2).

336

337 Fitness components of males

Local survival of male red squirrels was higher in OGA (n = 21, mean \pm SD = 27 \pm 19 months) than in CED (n = 24, 19 \pm 10 months) and RHE (n = 29, 18 \pm 15 months) and relationships between PC scores and survival differed between study areas (Comparison RHE against OGA and CED, all partial *P* < 0.05). In study areas OGA and CED, local survival increased with body mass but not with body size (Table 3a). Both PC scores for personality were negatively related with longevity (Table 3a).

In study area RHE patterns were different. Larger males had a higher local survival and boldness score (PC1) was positively related to survival. The effect of body size was much stronger than of body mass. The specific exploration score (PC2) was negatively related to local survival as in the other study areas (Table 3b). In the full model, the partial effect of PC1 was not significant, but considered as a single factor there was a positive relationship between boldness score and local survival (estimate 0.13 ± 0.04, χ^2 = 12.9; *P* = 0.0003).

348

349 Fitness components of females

350 Also among female red squirrels, local survival was higher in OGA (n = 19, mean \pm SD = 20 \pm 17 351 months) than in CED (n = 15, 14 ± 8 months) and RHE (n = 24, 15 ± 8 months) and the relationship between 352 survival and the two PC scores differed between study areas (comparison RHE against OGA and CED, all partial P < 0.01). In study areas OGA and CED the GLM model gave the same results as for males: there was 353 354 no significant effect of body size on female survival, while heavier females survived longer than those of 355 lower body mass (Table 4a). Local survival was negatively related to both PC scores, but only PC2 was significant (Table 4a). In contrast, in RHE the coefficient of foot length was negative, while that of body mass 356 357 positive, indicating a strong positive effect of body condition (mass rather than size) on local survival of

female red squirrels (Table 4b). As for males in this study area, the boldness score (PC1) was positively related, and the specific exploration score (PC2) negatively related to female survival (Table 4b).

360 The number of litters that females weaned while present on the study area varied between 0 and 7 (n = 57, mean \pm SD = 1.67 \pm 1.54). Heavier females produced more litters than those of poorer body mass 361 (body mass parameter estimate 0.019 \pm 0.005; χ^2 = 13.7; df = 1; P = 0.0002). Tendency to explore was 362 negatively related to the number of litters produced (partial effect PC2 score -0.86 ± 0.24; χ^2 = 12.3; df = 1; P 363 364 = 0.0005). On average, females tended to produce more litters in OGA (n = 17, mean \pm SE = 2.29 \pm 0.54 litters) 365 than in RHE (n = 23, 1.52 ± 0.23 litters) and CED (n = 17, 1.24 ± 0.31 litters; area effect χ^2 = 5.91; df = 2; P = 366 0.052), but pairwise differences were significant only between OGA and CED (P < 0.05). Partial effects of foot 367 length or boldness score (PC1) were not significant (both P = 0.10).

In periods of poor seed-crops (46 observations), there were 16 cases (35%) of offspring production and 30 cases (65%) of females not reproducing. In contrast, in years of medium-good seed-crops (144 observations) females were more likely to produce offspring (81 cases, 56% against 63 cases of females not reproducing; food availability estimate 0.96 ± 0.38 ; $\chi^2 = 6.42$; df = 1; P = 0.011). Bolder females were not more likely to produce offspring than shy ones (PC1 $\chi^2 = 0.10$; df = 1; P = 0.75), and this did not change between periods of poor against medium-good seed-crops (PC1 by food availability interaction $\chi^2 = 0.69$; df = 1; P =0.40). There was no effect of study area ($\chi^2 = 1.94$; df = 2; P = 0.38).

375

376 Discussion

Our analyses of capture-mark-recapture data of individual Eurasian red squirrels showed that scores derived from the number of times a squirrel is trapped (trappability) and from the number of different traps visited (trap diversity), during the first one or two years of presence, are repeatable across years and therefore can be used as indices of an individual's personality. There was no correlation between boldness or exploration scores and body size or body mass of the squirrels, suggesting that personality was not related to body condition, a good proxy of an individual's quality. We found that the relationships between boldness score and local survival/residency of individual squirrels differed between habitats, in males as well as in 384 females. Survival of bold individuals was higher than that of shy individuals in the habitat dominated by Norway spruce (RHE), while in the other two study areas, a Scots pine dominated forest (OGA) and a mixed 385 386 spruce-fir forest (CED) bolder squirrels survived less well. The second PCA score, related to specific exploration behavior, was negatively correlated with local survival in all habitats and in both sexes. For 387 388 females, the relationships between personality and body mass on one hand, and reproductive output 389 (number of litters in lifetime) on the other hand, were consistent across habitat types. Reproductive success 390 increased with female body mass, but decreased with the tendency to explore. Bolder females did not have 391 a reproductive advantage in years of poor food availability.

392

393 Reliability of capture-mark-recapture data for deriving indices of personality and for estimating local survival 394 In this study, we used two indirect indices of personality: trappability which is generally linked to risk 395 taking behavior, and the tendency to visit more different traps (trap diversity) which is considered as a 396 measure of exploration (Boon et al. 2008; Boyer et al. 2010). Similarly to studies on other Sciuridae we found 397 consistent individual differences across years in both indices suggesting they measure personality traits of 398 Eurasian red squirrels (Boon et al. 2008; Boyer et al. 2010; Montiglio et al. 2012; Le Coeur et al. 2015). 399 Moreover, capture probabilities of red squirrels tend to be high and this was the case also in the study areas 400 used here with high average capture probability per session which varied little over seasons or years (see 401 section "Study species", see also Wauters et al. 2008), allowing us to use and compare data over several 402 years.

Males were trapped on average more often and in more different traps than females, which is consistent with space use differences between the sexes with males using larger home ranges than females (Wauters and Dhondt 1992; Romeo et al. 2010; Di Pierro et al. 2011). Both sexes were captured in more different traps in OGA and RHE than in CED, which may have been due to different numbers of available traps. This potential bias among study areas was eliminated by standardizing trappability and trap diversity within each study area. The first component of the PCA (PC1), measuring both risk taking and exploration tendency and thus considered as a general boldness score, explained 88% of total variance and therefore

410 was the most relevant component. Nevertheless, we feel that PC2 is biologically relevant as it allowed us to 411 measure a pure exploration tendency. However, care is needed in using this component because of the low 412 variance explained, but where models show a significant contribution of PC2, it is not more prone to bias or 413 type I error compared to PC1.

414 We used the same capture-mark-recapture data to derive both personality and survival estimates, 415 which assumes the two estimates to be independent. Length of residency is commonly used in studies on 416 sciurids as an estimate of local survival (Gurnell et al. 2004; Wauters et al. 2004, 2007), with the precaution 417 to use only animals that stayed at least 6 month in the study area (eliminating dispersers) as we have done 418 here. Local survival was estimated from the entire trapping history of the individual and is only based on first 419 and last capture, while personality was estimated using all captures in each session, but only over the first 420 two years the individual was present (see methods). In principle, there could be a potential bias in that shy 421 individuals, which are trapped less often and visit fewer traps, would appear to have a shorter lifespan. 422 However, this was not the case in our study. As stated in the methods, estimated length of residency was 423 confirmed by data on radio-tracked animals. Moreover, in two out of three study areas shy animals (captured 424 fewer times) lived longer than bold ones, showing that the potential bias is weak or non-existent.

425

426 *Personality and phenotypic characteristics*

427 Since personality and body growth can be related, body size/mass should be taken into account when testing for effects of personality on reproductive parameters (Rödel and Meyer 2011; Guenther and Trillmich, 428 429 2015). In some mammalian species, heavier animals were bolder and more explorative than conspecifics of 430 lower body mass, especially among pups (Rödel and Meyer 2011; Guenther and Trillmich, 2015; Vetter et al. 431 2016). Bolder animals tend to be more aggressive (proactive-reactive syndrome) (Huntingford 1976; 432 Koolhaas et al. 1999; Bell and Stamps 2004; Sih et al. 2004; Groothuis and Carere 2005) and, in our study 433 species, heavier squirrels were more aggressive than those in poorer condition (Wauters and Dhondt 1989). 434 However, here we found that personality scores of red squirrels were not correlated with body mass or size 435 (foot length). In high-density populations of red squirrels, where encounters between individuals are

436 frequent, body mass tends to increase with age and heavier, older males are more aggressive than lighter, younger ones and these heavier males are more likely to mate (Wauters and Dhondt 1989; Wauters et al. 437 438 1990). Heavier, older females occupy higher quality (exclusive) core areas and have higher reproductive 439 success than animals of lower body mass (Wauters and Dhondt 1992, 1995). We must underline however 440 that in the alpine population studied here, densities were much lower (0.2 to 0.5 squirrels/ha against 0.8 to 441 1.5/ha in the lowland high-density populations) with less frequent encounters (our unpubl. data), which may 442 lead to animals being less aggressive independent of their body size/mass. This will be tested in future 443 research using Mirror Image Simulation (MIS) in arena tests (e.g. Réale 2007; Boon et al. 2008).

444

445 Body condition, personality and survival

446 In agreement with previous studies we showed that individual variation in body mass is a major factor 447 affecting survival: heavier squirrels of both sexes had a higher local survival than animals of lower body mass 448 (see also Wauters and Dhondt 1989; Wauters et al. 2007). This is a general pattern found in many mammals 449 for which there is a direct relationship between body mass and survival (Festa-Bianchet et al. 1997) and in 450 relation to food availability (Boutin and Larsen 1993). In fact, average survival was higher in the Scots pine 451 dominated forest of OGA where spatio-temporal variation in tree-seed availability was lower than in the 452 other two habitats (Table 1; see also Wauters et al. 2008; Salmaso et al. 2009). In our study, personality was 453 correlated to local survival of both male and female squirrels in different ways according to the study area. In Norway spruce dominated forest (RHE), bolder squirrels of both sexes had a higher survival than shy ones, 454 455 although the pattern was more evident in females. Home range sizes of male and female red squirrels in this 456 habitat were much larger than in the Scots pine wood of OGA and the mixed conifer forest of CED (Wauters 457 et al. 2005; Romeo et al. 2010; Di Pierro et al. 2011). A tentative explanation is that when spatio-temporal 458 variation in food availability is high, having larger home ranges will allow better access to limited food 459 resources, and bolder/exploring animals may use larger ranges than shy ones (Bell 2007; van Overveld and 460 Matthysen 2010; Minderman et al. 2010). Alternatively, bold animals may have better resource acquisition 461 abilities when resources are scarce or more widely distributed (Biro and Stamps 2008), since their high

462 activity or exploration (compared to shy individuals) may bring more opportunities to discover essential 463 resources in a heterogeneous environment (Wolf et al. 2007). In contrast, in our other two study areas (OGA 464 and CED), bold animals had lower local survival than shy ones, which is in accordance with Boon et al. (2008) 465 who found that the more active (higher trappability) female American red squirrels (Tamiasciurus 466 hudsonicus) were less likely to survive until the spring of the following year. These habitat related differences 467 could be explained by a higher cost of risk taking behavior in the latter two study areas, potentially due to 468 higher risk of predation (Sih et al. 2004; Boon et al. 2008) or parasitism (Morand et al. 2006; Boyer et al. 2010; 469 Patterson and Schulte-Hostedde 2011). Although Eurasian red squirrels have a poor macroparasite fauna 470 (only one gastrointestinal helminth and two ectoparasites that are common, with prevalence > 10%), there 471 is marked variation in parasite abundance and/or prevalence between individuals, at least partly related to 472 variation in host body mass and habitat type (Romeo et al. 2013; Santicchia et al. 2015a). It should be further 473 investigated whether bolder red squirrels indeed have higher parasite infections in some habitats, as was the 474 case in Siberian chipmunks (*Eutamias sibiricus*; Boyer et al. 2010). We are aware that our study system lacks 475 replication of the ecological variables mentioned here, at least for the highly-variable Norway spruce forest, 476 but this does not weaken our main finding that the relationships between personality traits and survival 477 differs among populations.

478 Another possible trade-off in costs/benefits of being bold was found in introduced Siberian 479 chipmunks in France: bolder animals had a higher survival and reproductive success than shy ones in low 480 resource years, but this was not the case when resources were abundant (Le Coeur et al. 2015). In great tits 481 (Parus major), faster explorers were favored under lower densities but slower explorers under higher 482 densities and temporal variation in local density represented the primary factor explaining personality-483 related variation in viability selection (Nicolaus et al. 2016). Our multi-year data on survival in different 484 habitats produced rather similar patterns, with an advantage for bolder squirrels only in the habitat with 485 higher variability in food abundance and squirrel density (RHE, Table 1; see also Wauters et al. 2008; Bisi et 486 al. 2016).

487 In this study we separated a general boldness-exploration component of personality from a second pure exploration component (see also Boon et al. 2008). Increased predation risk, encountering more 488 489 parasite infective stages and higher energy demands may all be involved in explaining why we always found 490 a negative effect of our pure exploration score (PC2) on local survival (both sexes) and on reproduction 491 (females). However, two points must be made when considering the effects of our exploration score. First, 492 as stated above, it explains only a limited amount of the variation among individual squirrels (12%); hence its 493 relationships should be interpreted with care. Second, we are measuring local survival of squirrels in the 494 study area: this measure might include animals that disappeared from our study areas due to emigration, 495 especially among those at the lower tail of the survival distribution (those with 6-12 months of local survival) 496 (e.g. Ergon and Gardner 2014). Although most dispersal in red squirrels occurs during the juvenile-subadult 497 phase (Wauters and Dhondt 1993; Wauters et al. 2010, 2011), some adults do undertake dispersal 498 movements (Wauters and Dhondt 1993; Lurz et al. 1997). However, in general site fidelity of adult red 499 squirrels is high and in our study areas 91 out of 107 (85%) radio-tracked adults remained within established 500 home ranges, with most cases of dispersal occurring in RHE in response to a poor seed-crop (e.g. Wauters et 501 al. 2005; Romeo et al. 2010; Di Pierro et al. 2011). We tried to remove dispersers as much as possible from 502 our dataset by using only adults that remained at least for 6 months on the study area and, among radio-503 tracked animals, using only those that settled. Apart from this methodological point, our pure exploration 504 score (PC2) was negatively related to the fitness components studied (survival and reproduction). Only a few 505 studies found a small positive effect of exploration tendency on survival (and no relationship between 506 exploration and reproductive success, e.g. Smith and Blumstein 2008), whereas others reported negative 507 effects as in this study (Carter et al. 2010; Hall et al. 2015), although often related to early life-stages (Rödel 508 et al. 2015). We only used subadults and adults in our dataset, but still wider movements by pure explorers 509 might increase predation risk and/or levels of chronic stress compared to less explorative animals. How stress 510 levels may be linked to personality under differential conditions of habitat or population density are 511 challenges for future research (e.g. Dantzer et al. 2013, 2016).

512

513 Body condition, personality and female reproduction

514 We found that heavier females produced more litters than those of poorer body mass, in agreement 515 with previous studies on red squirrels (Wauters and Dhondt 1989, 1995; Wauters et al. 2007). Because of 516 such strong effects of a female's body condition, variation in personality might have only weak relationships 517 with reproductive success. In this study we found no effect of boldness on reproductive success but a 518 negative effect of exploration. Female red squirrels that explore a lot could be less efficient in obtaining and 519 defending exclusive core-areas (intrasexual territoriality, Wauters and Dhondt 1992), and/or spend less time 520 nursing their offspring. This hypothesis could be tested by radio-tracking squirrels, investigating the 521 relationships of personality with degree of core-area overlap and time spent away from the nest during 522 lactation. Another point to consider is that maybe we need more detailed data of reproductive success: not 523 only number of litters produced, but also data on number of young produced or weaned/litter (Wauters and 524 Dhondt 1995; Santicchia et al. 2015b) to reveal (weak) relationships between personality and reproductive 525 success. Negative effects of exploration on reproductive success were also found in other studies. For 526 example, Vetter et al. (2016) demonstrated that, under high food availability, being less aggressive and less 527 explorative positively influenced the post-weaning litter size in wild boar (Sus scrofa).

528 We believe that our study provides insight in how personality traits can produce fitness trade-offs 529 under different environmental conditions. Unfortunately, our sample size was too small to explore if and 530 how personality might relate to trade-offs among different fitness components, for example differential 531 investment in survival or reproduction. Hence, our results suggest the need for long-term studies measuring 532 individual variation in personality traits, phenotypic factors, space use (home range size, degree of overlap, 533 home range quality) and different fitness components combining data from populations living under different 534 abiotic conditions (habitat types, food supplies) to better assess the impacts of personality on an animal's 535 "performance" both before and during its period of reproductive activity.

536

537 Acknowledgments

We thank the Gran Paradiso National Park, the Orobie Valtellinese Regional Park, the province of Sondrio for allowing fieldwork in the three study areas. We like to thank Ambrogio Molinari, Erica Di Pierro, Marco Adamo and Aleksandar Pizzul for help with the fieldwork and Stefan Van Dongen for statistical advice. Constructive comments by two referees and the Associate Editor greatly helped us to improve the manuscript. Data analysis and writing were carried out in 2015-16 by the first author as part of her PhD project. This study is part of the ASPER (Alpine Squirrel Population Ecology Research) project (paper # 28).

544

545 Compliance with ethical standards

546 Funding

This work was supported by the Committee for Research and Exploration of the National Geographic Society, Washington DC, USA (grant number 6997-01) to LAW and SB; and by the Italian Ministry of Instruction, University and Research (MIUR, Ministero dell'Istruzione, dell'Università e della Ricerca, COFIN project number 2003053710-006) to Università degli Studi dell'Insubria.

551 Ethical approval

All applicable international, national and/or institutional guidelines for the care and use of animals were followed. All procedures performed in studies involving animals were in accordance with the current law on animal research in Italy and were carried out under permission of the authorities for wildlife research and management of Lombardy region (Decree n.18064 of 14/07/2000; Decree n.10816 of 10/06/2002; Decree n.1861 of 16/02/2004; Decree n.2456 of 07/03/2006) and Gran Paradiso National Park (Authorization prot. n. 1716/9.3.7 of 06/07/200 and prot. n 1071//9.3.7 of 07/05/2003). No specific approval from ethics committee was required.

559

560 Conflict of interest

561 The authors declare that they have no conflict of interests

- 562
- 563 Data availability

- All data analysed during this study are included in this published article and its supplementary informationfiles.
- 566

567 References

- 568 Bell AM (2007) Future directions in behavioural syndromes research. Proc R Soc Lond B 274:755-761
- 569 Bell AM, Stamps JA (2004) Development of behavioural differences between individuals and populations of
- 570 sticklebacks, *Gasterosteus aculeatus*. Anim Behav 68:1339-1348
- 571 Bergeron P, Montiglio P-O, Réale D, Humphries MM, Gimenez O, Garant D (2013) Disruptive viability selection
- on adult exploratory behaviour in eastern chipmunks. J Evol Biol 26:766-774
- 573 Bergmüller R, Taborsky M (2010) Animal personality due to social niche specialisation. Trends Ecol Evol 574 25:504-511
- 575 Biro PA, Stamps JA (2008) Are animal personality traits linked to life-history productivity? Trends Ecol Evol 576 23:361-368
- 577 Bisi F, von Hardenberg J, Bertolino S, Wauters LA, Imperio S, Preatoni DG, Mazzamuto MV, Provenzale A,
- 578 Martinoli A (2016) Current and future conifer seed production in the Alps: testing weather factors as cues
- 579 behind masting. Eur J Forest Res 135:743-754
- 580 Boon AK, Réale D, Boutin S (2007) The interaction between personality, offspring fitness and food abundance
- in North American red squirrels. Ecol Lett 10:1094-1104
- 582 Boon AK, Réale D, Boutin S (2008) Personality, habitat use, and their consequences for survival in North
- 583 American red squirrels *Tamiasciurus hudsonicus*. Oikos 117:1321-1328
- 584 Both C, Dingemanse NJ, Drent PJ, Tinbergen JM (2005) Pairs of extreme avian personalities have highest
- 585 reproductive success. J Anim Ecol 74:667-674

- 586 Boutin S, Larsen KW (1993) Does food availability affect growth and survival of males and females differently
- in a promiscuous small mammal, *Tamiasciurus hudsonicus*? J Anim Ecol 62:364-370
- 588 Boutin S, Wauters LA, McAdam AG, Humphries MM, Tosi G, Dhondt AA (2006) Anticipatory reproduction and
- population growth in seed predators. Science 314:1928-1930
- 590 Boyer N, Réale D, Marmet J, Pisanu B, Chapuis JL (2010) Personality, space use and tick load in an introduced
- 591 population of Siberian chipmunks *Tamias sibiricus*. J Anim Ecol 79:538-547
- 592 Bremner-Harrison SB, Prodohl PA, Elwood RW (2004) Behavioural trait assessment as a release criterion:
- 593 boldness predicts early death in a reintroduction programme of captive-bred swift fox (*Vulpes velox*). Anim
- 594 Conserv 7:313-320
- 595 Carter AJ, Feeney WE, Marshall HH, Cowlishaw G, Heinsohn R (2013) Animal personality: what are 596 behavioural ecologists measuring? Biol Rev 88:465-475
- 597 Carter AJ, Goldizen AW, Tromp SA (2010) Agamas exhibit behavioral syndromes: bolder males bask and feed
- 598 more but may suffer higher predation. Behav Ecol 21:655-661
- 599 Cavigelli SA, McClintock MK (2003) Fear of novelty in infant rats predicts adult corticosterone dynamics and
- an early death. P Natl Acad Sci USA 100:16131-16136
- Dall SRX, Houston AI, McNamara JM (2004) The behavioural ecology of personality: consistent individual
 differences from an adaptive perspective. Ecol Lett 7:734-739
- Dantzer B, Newman AEM, Boonstra R, Palme R, Boutin S, Humphries MM, McAdam AJ (2013) Density triggers
- maternal hormones that increase adaptive offspring growth in a wild mammal. Science 340:1215-1217
- 605 Dantzer B, Santicchia F, van Kesteren F, Palme R, Martinoli A, Wauters LA (2016) Measurement of fecal
- 606 glucocorticoid metabolite levels in Eurasian red squirrels (Sciurus vulgaris): effects of captivity, sex,
- 607 reproductive condition, and season. J Mammal 97 :1385-1398

- Di Pierro E, Ghisla A, Wauters LA, Molinari A, Martinoli A, Gurnell J, Tosi G (2011) The effects of seed availability on habitat use by a specialist seed predator. Eur J Wildlife Res 57:585-595
- 610 Dingemanse NJ, Bouwman KM, van de Pol M, van Overveld T, Patrick SC, Matthysen E, Quinn JL (2012)
- 611 Variation in personality and behavioural plasticity across four populations of the great tit *Parus major*. J Anim
- 612 Ecol 81:116-126
- Dingemanse NJ, de Goede P (2004) The relation between dominance and exploratory behavior is contextdependent in wild great tits. Behav Ecol 15:1023-1030
- Dingemanse NJ, Réale D (2005) Natural selection and animal personality. Behaviour 142:1165-1190
- 616 Ergon T, Gardner B (2014) Separating mortality and emigration: modelling space use, dispersal and survival
- 617 with robust-design spatial capture-recapture data. Methods Ecol Evol 5:1327-1336
- Festa-Bianchet M, Jorgenson JT, Bérubé CH, Portier C, Wishart WD (1997) Body mass and survival of bighorn
 sheep. Can J Zool 75:1372-1379
- 620 Fraser DF, Gilliam JF, Daley MJ, Le AN, Skalski GT (2001) Explaining leptokurtic movement distributions:
- 621 intrapopulation variation in boldness and exploration. Am Nat 158:124-135
- Groothuis TGG, Carere C (2005) Avian personalities: characterization and epigenesis. Neurosci Biobehav Rev
 29:137-150
- 624 Guenther A, Trillmich F (2015) Within-litter differences in personality and physiology relate to size differences
- among siblings in cavies. Physiol Behav 145:22-28
- Hall ML, van Asten T, Katsis AC, Dingemanse NJ, Magrath MJL, Mulder RA (2015) Animal personality and pace-
- of-life syndromes: do fast-exploring fairy-wrens die young? Front Ecol Evol 3:28
- 628 Huntingford FA (1976) Relationship between anti-predator behavior and aggression among conspecifics in
- 629 the three-spined stickleback, Gasterosteus aculeatus. Anim Behav 24:245-260

- Koolhaas JM, Korte SM, de Boer SF, van der Vegt BJ, van Reenen CG, Hopster H, de Jong IC, Ruis MAW, Blokuis
 HJ (1999) Coping styles in animals: current status in behavior and stress-physiology. Neurosci Biobehav Rev
 23:925-935
- 633 Korsten P, Mueller JC, Hermannstadter C et al (2010) Association between DRD4 gene polymorphism and
- 634 personality variation in great tits: a test across four wild populations. Mol Ecol 19:832-843
- Le Coeur C, Thibault M, Pisanu B, Thibault S, Chapuis J-L, Baudry E (2015) Temporally fluctuating selection on
 a personality trait in a wild rodent population. Behav Ecol 26:1285-1291
- 637 Lurz PWW, Garson PJ, Wauters LA (1997) Effects of temporal and spatial variation in habitat quality on red
- 638 squirrel dispersal behavior. Anim Behav 54:427-435
- 639 Martin JGA, Réale D (2008) Temperament, risk assessment and habituation to novelty in eastern chipmunks,
- 640 *Tamias striatus*. Anim Behav 75:309-318
- 641 Minderman J, Reid JM, Hughes M, Denny MJH, Hogg S, Evans PGH, Whittinghama MJ (2010) Novel
- 642 environment exploration and home range size in starlings *Sturnus vulgaris*. Behav Ecol 21:1321-1329
- 643 Monestier C, Morellet N, Gaillard J-M, Cargnelutti B, Vanpé C, Hewison AJM (2015) Is a proactive mum a good
- 644 mum? A mother's coping style influences early fawn survival in roe deer. Behav Ecol 26:1395-1403
- 645 Montiglio P-O, Garant D, Pelletier F, Réale D (2012) Personality differences are related to long-term stress
- reactivity in a population of wild eastern chipmunks, *Tamias striatus*. Anim Behav 84:1071-1079
- 647 Morand S, Krasnov B R, Poulin R (2006) Micromammals and macroparasites: from evolutionary ecology to 648 management. Springer, Tokyo
- Mutzel A, Dingemanse NJ, Araya-Ajoy YG, Kempenaers B (2013) Parental provisioning behaviour plays a key
 role in linking personality with reproductive success. Proc R Soc B 280:20131019
- Nakagawa S, Schielzeth H (2010) Repeatability for Gaussian and non-Gaussian data: a practical guide for
 biologists. Biol Rev 85:935-956

- 653 Nicolaus M, Tinbergen JM, Ubels R, Both C, Dingemanse NJ (2016) Density fluctuations represent a key
- 654 process maintaining personality variation in a wild passerine bird. Ecol Lett 19:478-486
- 655 Patterson LD, Schulte-Hostedde AI (2011) Behavioural correlates of parasitism and reproductive success in
- 656 male eastern chipmunks, Tamias striatus. Anim Behav 81:1129-1137
- 657 Réale D, Gallant BY, Leblanc M, Festa-Bianchet M (2000) Consistency of temperament in bighorn ewes and
- 658 correlates with behaviour and life history. Anim Behav 60:589-597
- 659 Réale D, Reader SM, Sol D, McDougall PT, Dingemanse NJ (2007) Integrating animal temperament within
- 660 ecology and evolution. Biol Rev 82:291-318
- 661 Rivest L-P, Baillargeon S (2014) Rcapture: loglinear models for capture-recapture experiments. R package
- 662 version 1.4-2, https://cran.reproject.org/package=Rcapture
- 663 Rödel HG, Meyer S (2011) Early development influences ontogeny of personality types in young laboratory 664 rats. Dev Psychobiol 53:601-613
- 665 Rödel HG, Zapka M, Talke S, Kornatz T, Bruchner B, Hedler C (2015) Survival costs of fast exploration during
- 666 juvenile life in a small mammal. Behav Ecol Sociobiol 69:205-217
- 667 Romeo C, Pisanu B, Ferrari N, Basset F, Tillon L, Wauters LA, Martinoli A, Saino N, Chapuis J-L (2013)
- 668 Macroparasite community of the Eurasian red squirrel (Sciurus vulgaris): poor species richness and diversity. 669 Parasitol Res 112:3527-3536

Romeo C, Wauters L, Preatoni D, Tosi G, Martinoli A (2010) Living on the edge: space use of Eurasian red

- 670
- 671 squirrels in marginal high elevation habitat. Acta Oecol 36:604-610
- 672 Salmaso F, Molinari A, Di Pierro E, Ghisla A, Martinoli A, Preatoni D, Cerabolini B, Tosi G, Bertolino S, Wauters
- 673 LA (2009) Estimating and comparing food availability for tree-seed predators in typical pulsed resource
- 674 systems: alpine conifer forests. Plant Biosyst 143:258-267

- Santicchia F, Romeo C, Martinoli A, Lanfranchi P, Wauters LA, Ferrari N (2015) Effects of habitat quality on
 parasite abundance: do forest fragmentation and food availability affect helminth infection in the Eurasian
 red squirrel? J Zool 296:38-44
- 678 Santicchia F, Romeo C, Grilli G, Vezzoso S, Wauters LA, Mazzamuto MV, Martinoli A, Ferrari N (2015) The use
- of uterine scars to explore fecundity levels in invasive alien tree squirrels. Hystrix 26:95-101
- Sih A, Bell AM, Johnson JC, Ziemba RE (2004) Behavioral syndromes: an integrative overview. Q Rev Biol
 79:241-277
- 682 Smith BR, Blumstein DT (2008) Fitness consequences of personality: a meta-analysis. Behav Ecol 19:448-455
- Trizio I, Crestanello B, Galbusera P, Wauters LA, Tosi G, Matthysen E, Hauffe HC (2005) Geographical distance
- and physical barriers shape the genetic structure of Eurasian red squirrels in the Italian Alps. Mol Ecol 14:469-

685 481

van Overveld T, Adriaensen F, Matthysen E (2015) No evidence for correlational selection on exploratory

behaviour and natal dispersal in the great tit. Evol Ecol 29:137-156

- van Overveld T, Matthysen E (2010) Personality predicts spatial responses to food manipulations in free-
- ranging great tits (*Parus major*). Biol Lett 6:187-190
- 690 Vetter SG, Brandstätter C, Macheiner M, Suchentrunk F, Gerritsmann H, Bieber C (2016) Shy is sometimes

691 better: personality and juvenile body mass affect adult reproductive success in wild boars, *Sus scrofa*. Anim

- 692 Behav 115:193-205
- 693 Wauters LA, Bertolino S, Adamo M, Van Dongen S, Tosi G (2005) Food shortage disrupts social organization:
- the case of red squirrels in conifer forests. Evol Ecol 19:375-404
- 695 Wauters LA, Dhondt AA (1989) Body weight, longevity and reproductive success in red squirrels (*Sciurus*
- 696 *vulgaris*). J Anim Ecol 58:637-651

- 697 Wauters LA, Dhondt AA (1992) Spacing behaviour of red squirrels, *Sciurus vulgaris*: variation between
 698 habitats and the sexes. Anim Behav 43:297-311
- Wauters LA, Dhondt AA (1993) Immigration pattern and success in red squirrels. Behav Ecol Sociobiol 33:159167
- Wauters LA, Dhondt AA (1995) Lifetime reproductive success and its correlates in female Eurasian red
 squirrels. Oikos 72:402-410
- Wauters LA, Dhondt AA, De Vos R (1990) Factors affecting male mating success in red squirrels (*Sciurus vulgaris*). Ethol Ecol Evol 2:195-204
- 705 Wauters LA, Githiru M, Bertolino S, Molinari A, Tosi G, Lens L (2008) Demography of alpine red squirrel
- populations in relation to fluctuations in seed crop size. Ecography 31:104-114
- Wauters LA, Matthysen E, Adriaensen F, Tosi G (2004) Within-sex density dependence and population
 dynamics of red squirrels *Sciurus vulgaris*. J Anim Ecol 73:11-25
- 709 Wauters LA, Preatoni D, Martinoli A, Verbeylen G, Matthysen E (2011) No sex bias in natal dispersal of
- 710 Eurasian red squirrels. Mamm Biol 76:369-372
- 711 Wauters LA, Verbeylen G, Preatoni D, Martinoli A, Matthysen E (2010) Dispersal and habitat cuing of Eurasian
- red squirrels in fragmented habitats. Popul Ecol 52:527-536
- 713 Wauters LA, Vermeulen M, Van Dongen S, Bertolino S, Molinari A, Tosi G, Matthysen E (2007) Effects of
- spatio-temporal variation in food supply on red squirrel *Sciurus vulgaris* body size and body mass and its
- consequences for some fitness components. Ecography 30:51-65
- 716 Weiss A, Adams MJ (2013) Differential behavioral ecology. In: Carere C, Maestripieri D (eds) Animal
- personalities: behavior, physiology and evolution. University of Chicago Press, Chicago, pp 96-123
- 718 Wolf M, van Doorn GS, Leimar O, Weissing FJ (2007) Life-history trade-offs favour the evolution of animal
- 719 personalities. Nature 447:581-584

720	Zhao Q-S, Hu Y-B, Liu P-F, Chen L-J, Sun Y-H (2016) Nest site choice: a potential pathway linking personality
721	and reproductive success. Anim Behav 118:97-103
722	
723	
724	
725	
726	
727	
728	
729	
730	
731	
732	
733	
734	
735	
736	
737	
738	

Table 1 Food abundance estimates (conifer seed-crop in 10³ kJ/ha, mean ± SE) and number of red squirrels
 (males, females) trapped and used to measure personality per year in the three study areas. Total number
 of squirrels refers to sample size of different individuals used in the analyses

Year	Study areas					
	CED		OGA		RHE	
	Food	Squirrels	Food	Squirrels	Food	Squirrels
1999	4357 ± 570		1732 ± 419			
2000	159 ± 63 *	4 (3, 1)	391 ± 82 *	16 (10, 6)	497 ± 252 *	11 (8, 3)
2001	3087 ± 440	10 (7, 3)	588 ± 83	20 (10, 10)	913 ± 109	18 (11, 7)
2002	1867 ± 296	14 (8, 6)	735 ± 153	23 (14,9)	1773 ± 154	21 (12, 9)
2003	846 ± 226 *	20 (11, 9)	552 ± 81	21 (13, 8)	1010 ± 130	20 (11, 9)
2004	6142 ± 728	18 (11, 7)	863 ± 111	21 (12, 9)	2402 ± 233	19 (11, 8)
2005	4022 ± 828	16 (10, 6)	352 ± 73 *	14 (8, 6)	22 ± 5 *	21 (10, 11)
2006	6249 ± 907	14 (9, 5)		5 (3, 2)	1363 ± 179	13 (8, 5)
2007	2547 ± 528	9 (7, 2)			1171 ± 133	8 (4, 4)
2008		1 (1, 0)			96 ± 27 *	4 (2, 2)
Mean/Total	3253 ± 849	45 (27, 18)	745 ± 192	40 (21, 19)	1027 ± 273	56 (30, 26)
% CV	66%		63%		75%	

- 742 *years of poor seed-crop

- 7 4 6

Table 2 Pearson's correlation coefficient with significance level for correlations between foot length, body

- mass and personality scores of red squirrels: a) all animals; b) and c) per sex. Sample size between brackets

	Foot length	Boldness score (PC1)	Exploration score (PC2)
(a) Both sexes (n = 141)		
Body mass	<i>r</i> = 0.49, <i>P</i> < 0.0001	<i>r</i> = 0.001, <i>P</i> = 0.98	<i>r</i> = -0.007, <i>P</i> = 0.93
Foot length		<i>r</i> = 0.06, <i>P</i> = 0.49	<i>r</i> = 0.04, <i>P</i> = 0.60
(b) Males (n = 78)			
Body mass	<i>r</i> = 0.63, <i>P</i> < 0.0001	<i>r</i> = 0.09, <i>P</i> = 0.44	<i>r</i> = -0.04, <i>P</i> = 0.73
Foot length		<i>r</i> = 0.08, <i>P</i> = 0.46	<i>r</i> = 0.01, <i>P</i> = 0.90
(c) Females (n = 63)			
Body mass	<i>r</i> = 0.38, <i>P</i> = 0.002	<i>r</i> = -0.04, <i>P</i> = 0.74	<i>r</i> = -0.03, <i>P</i> = 0.02
Foot length		<i>r</i> = -0.001, <i>P</i> = 0.99	<i>r</i> = 0.11, <i>P</i> = 0.39

Table 3 General Linear Models with Poisson distribution of parameters affecting variation in longevity of male767red squirrels. After significant interactions of PC scores and foot length with study area, models were tested768separately for CED and OGA (interactions, all partial P > 0.05) and for RHE (interactions with both CED and769OGA, all partial P < 0.05)

Fixed effects	Estimates	Chi-square (df = 1)	Р
(a) CED & OGA (n = 45)			
Study area		32.6	< 0.0001
Body mass	0.014 ± 0.002	42.7	< 0.0001
Foot length	-0.058 ± 0.048	1.43	0.23
PC1 score	-0.047 ± 0.022	4.15	0.042
PC2 score	-0.376 ± 0.071	28.4	< 0.0001
(b) RHE (n = 29)			
Body mass	-0.0019 ± 0.0031	0.39	0.53
Foot length	0.221 ± 0.041	29.6	< 0.0001
PC1 score	0.037 ± 0.035	1.06	0.30
PC2 score	-0.478 ± 0.061	60.9	< 0.0001

Table 4 General Linear Models with Poisson distribution of parameters affecting variation in longevity offemale red squirrels. After significant interactions of PC scores with study area, models were testedseparately for CED and OGA (interactions, all partial P > 0.05) and for RHE (interactions with both CED andOGA, all partial P < 0.05)

Fixed effects	Estimates	Chi-square (df = 1)	Р
(a) CED & OGA (n = 34)			
Study area		10.9	0.001
Body mass	0.0092 ± 0.0024	14.8	<0.0001
Foot length	-0.005 ± 0.037	0.02	0.89
PC1 score	-0.099 ± 0.056	3.11	0.078
PC2 score	-1.057 ± 0.120	77.6	< 0.0001
(b) RHE (n = 24)			
Body mass	0.0090 ± 0.0025	13.0	0.0003
Foot length	-0.124 ± 0.035	12.5	0.0004
PC1 score	0.158 ± 0.038	17.4	< 0.0001
PC2 score	-0.494 ± 0.115	18.6	< 0.0001

793	Fig. 1
794	a) Trappability and b) trap diversity (mean ± SE) of male (triangles) and female (circles) red squirrels in
795	three study areas in the Italian Alps. Sample size per study area and sex above error flags
796	
797	
798	
799	
800	
801	
802	
803	
804	
805	
806	
807	
808	
809	
810	This is a post-peer-review, pre-copyedit version of an article published in Behavioral Ecology and
811	Sociobiology. The final authenticated version is available online at: http://dx.doi.org/10.1007/s00265-018-
812	2546-у