

AperTO - Archivio Istituzionale Open Access dell'Università di Torino

Splitting Theorems, Symmetry Results and Overdetermined Problems for Riemannian Manifolds

This is a pre print version of the following article:

Original Citation:

Availability:

This version is available <http://hdl.handle.net/2318/1693256> since 2019-02-18T20:29:14Z

Published version:

DOI:10.1080/03605302.2013.795969

Terms of use:

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)

Splitting theorems, symmetry results and overdetermined problems for Riemannian manifolds

Alberto Farina^(1,2) Luciano Mari⁽³⁾ Enrico Valdinoci^(4,5)

(1) – Laboratoire Amiénois de Mathématique Fondamentale et Appliquée
UMR CNRS 7352, Université Picardie “Jules Verne”
33 Rue St Leu, 80039 Amiens (France).

(2) – Institut “Camille Jordan”
UMR CNRS 5208, Université “Claude Bernard” Lyon I
43 Boulevard du 11 novembre 1918, 69622 Villeurbanne cedex (France).

(3) – Departamento de Matemática
Universidade Federal do Ceará
Campus do Pici, 60455-760 Fortaleza-CE (Brazil).

(4) – Dipartimento di Matematica “Federigo Enriques”
Università degli studi di Milano,
Via Saldini 50, I-20133 Milano (Italy).

(5) – Istituto di Matematica Applicata e Tecnologie Informatiche “Enrico Magenes”
Consiglio Nazionale delle Ricerche
Via Ferrata 1, I-27100 Pavia (Italy).

E-mail addresses: alberto.farina@u-picardie.fr, lucio.mari@libero.it, enrico.valdinoci@unimi.it

Abstract

Our work proposes a unified approach to three different topics in a general Riemannian setting: splitting theorems, symmetry results and overdetermined elliptic problems. By the existence of a stable solution to the semilinear equation $-\Delta u = f(u)$ on a Riemannian manifold with non-negative Ricci curvature, we are able to classify both the solution and the manifold. We also discuss the classification of monotone (with respect to the direction of some Killing vector field) solutions, in the spirit of a conjecture of De Giorgi, and the rigidity features for overdetermined elliptic problems on submanifolds with boundary.

Introduction and main results

In this paper, we will study Riemannian manifolds $(M, \langle \cdot, \cdot \rangle)$ with non-negative Ricci curvature that possess a stable, nontrivial solution of a semilinear equation of the type $-\Delta u = f(u)$. Under reasonable growth assumptions on u , we prove both symmetry results for the solution and the rigidity of the underlying manifold. The case of manifolds with boundary will be considered as well, in the framework of overdetermined problems. The main feature of our work is that we give a unified treatment, thereby providing a bridge between three different

topics in a general Riemannian setting: splitting theorems, symmetry results and overdetermined problems. The key role here is played by a refined geometric Poincaré inequality, improving on those in [41, 40, 18, 19], see Proposition 16 below. In the very particular case of Euclidean space, we recover previously known results in the literature.

Firstly, we deal with complete, non-compact, boundaryless Riemannian manifolds of non-negative Ricci curvature, that admit a non-trivial stable solution. By assuming either a parabolicity condition or a bound on the energy growth, we obtain that the manifold splits off a factor \mathbb{R} that completely determines the solution. More precisely, we will prove

Theorem 1. *Let $(M, \langle \cdot, \cdot \rangle)$ be a complete, non-compact Riemannian manifold without boundary, satisfying $\text{Ric} \geq 0$. Suppose that $u \in C^3(M)$ be a non-constant, stable solution of $-\Delta u = f(u)$, for $f \in C^1(\mathbb{R})$. If either*

- (i) *M is parabolic and $\nabla u \in L^\infty(M)$, or*
- (ii) *The function $|\nabla u|$ satisfies*

$$\int_{B_R} |\nabla u|^2 dx = o(R^2 \log R) \quad \text{as } R \rightarrow +\infty. \quad (1)$$

Then,

- *$M = N \times \mathbb{R}$ with the product metric $\langle \cdot, \cdot \rangle = \langle \cdot, \cdot \rangle_N + dt^2$, for some complete, totally geodesic, parabolic hypersurface N . In particular, $\text{Ric}^N \geq 0$ if $m \geq 3$, and $M = \mathbb{R}^2$ or $\mathbb{S}^1 \times \mathbb{R}$, with their flat metric, if $m = 2$;*
- *u depends only on t , has no critical points, and writing $u = y(t)$ it holds $y'' = -f(y)$.*

Moreover, if (ii) is met,

$$\text{vol}(B_R^N) = o(R^2 \log R) \quad \text{as } R \rightarrow +\infty. \quad (2)$$

$$\int_{-R}^R |y'(t)|^2 dt = o\left(\frac{R^2 \log R}{\text{vol}(B_R^N)}\right) \quad \text{as } R \rightarrow +\infty. \quad (3)$$

Basic facts on parabolicity can be found in [23], Sections 5 and 7. We underline that, under a suitable sign assumption on f , in Theorem 20 below we will obtain that every stable solution is constant.

For our purposes, it is convenient to define \mathcal{F}_2 to be the family of complete manifolds M with non-negative Ricci tensor that, for each fixed $f \in C^1(\mathbb{R})$, do not possess any stable, non-constant solution $u \in C^3(M)$ of $-\Delta u = f(u)$ for which

$$\int_{B_R} |\nabla u|^2 dx = o(R^2 \log R) \quad \text{as } R \rightarrow +\infty.$$

Next Proposition 2 and Theorem 3 give a complete classifications of M using this family:

Proposition 2. *Let $(M^m, \langle \cdot, \cdot \rangle)$ be a complete, non-compact manifold with $\text{Ric} \geq 0$. Then,*

- *if $m = 2$, $M \in \mathcal{F}_2$ if and only if M is neither \mathbb{R}^2 nor $\mathbb{S}^1 \times \mathbb{R}$ with their flat metric;*
- *if $m = 3$, $M \in \mathcal{F}_2$ if and only if M does not split off an Euclidean factor.*

Theorem 3. *Let $(M^m, \langle \cdot, \cdot \rangle)$ be a complete, non-compact manifold with $\text{Ric} \geq 0$ and dimension $m \geq 3$. Suppose that $M \notin \mathcal{F}_2$. Then, one and only one of the following possibilities occur:*

(i) $M = N^{m-1} \times \mathbb{R}$, where $N^{m-1} \in \mathcal{F}_2$ is either compact or it is parabolic, with only one end and with no Euclidean factor. Furthermore,

$$\text{vol}(B_R^N) = o(R^2 \log R) \quad \text{as } R \rightarrow +\infty. \quad (4)$$

(ii) either $m = 3$ and $M = \mathbb{R}^3$ or $\mathbb{S}^1 \times \mathbb{R}^2$ with flat metric, or $m \geq 4$ and $M = \bar{N}^{m-2} \times \mathbb{R}^2$, where $\bar{N}^{m-2} \in \mathcal{F}_2$ is either compact or it is parabolic, with only one end and with no Euclidean factor. Moreover,

$$\text{vol}(B_R^{\bar{N}}) = o(R \log R) \quad \text{as } R \rightarrow +\infty. \quad (5)$$

(iii) either $m = 4$ and $M = \mathbb{S}^1 \times \mathbb{R}^3$ with flat metric, or $m \geq 5$ and $M = \hat{N}^{m-3} \times \mathbb{R}^3$, where \hat{N}^{m-3} is compact with $\text{Ric}^{\hat{N}} \geq 0$.

Remark 4. From the topological point of view, it has been recently proved in [27] that a non-compact, 3-manifold with $\text{Ric} \geq 0$ is either diffeomorphic to \mathbb{R}^3 or its universal cover splits off a line (isometrically). This causes extra-rigidity for the manifolds N, \bar{N} in the previous theorem. On the other hand, compact 3-manifolds with $\text{Ric} \geq 0$ have been classified in [24] (Theorem 1.2) via Ricci flow techniques. Namely, they are diffeomorphic to a quotient of either $\mathbb{S}^3, \mathbb{S}^2 \times \mathbb{R}$ or \mathbb{R}^3 by a group of fixed point free isometries in the standard metrics.

The case of manifolds with boundary will be considered here in the light of overdetermined problems. In this spirit, Killing vector fields play a special role, as underlined by the next

Theorem 5. Let $(M, \langle \cdot, \cdot \rangle)$ be a complete, non-compact Riemannian manifold without boundary, satisfying $\text{Ric} \geq 0$ and let X be a Killing field on M . Let $\Omega \subseteq M$ be an open and connected set with C^3 boundary. Suppose that $u \in C^3(\bar{\Omega})$ is a non-constant solution of the overdetermined problem

$$\begin{cases} -\Delta u = f(u) & \text{on } \Omega \\ u = \text{constant} & \text{on } \partial\Omega \\ \partial_\nu u = \text{constant} \neq 0 & \text{on } \partial\Omega. \end{cases} \quad (6)$$

Such that $\langle \nabla u, X \rangle$ is either positive or negative on Ω . Then, if either

(i) M is parabolic and $\nabla u \in L^\infty(\Omega)$, or

(ii) the function $|\nabla u|$ satisfies

$$\int_{\Omega \cap B_R} |\nabla u|^2 dx = o(R^2 \log R) \quad \text{as } R \rightarrow +\infty,$$

the following properties hold true:

- X is never zero, $\Omega = \partial\Omega \times \mathbb{R}^+$ with the product metric $\langle \cdot, \cdot \rangle = \langle \cdot, \cdot \rangle_{\partial\Omega} + dt^2$, $\partial\Omega$ is totally geodesic in M and satisfies $\text{Ric}_{\partial\Omega} \geq 0$.
- the function u depends only on t , it has no critical points, and writing $u = y(t)$ it holds $y'' = -f(y)$;
- for every $t_0 \in \mathbb{R}$, the projected field $X^\perp = X - \langle X, \partial_t \rangle \partial_t$ at $(\cdot, t_0) \in \partial\Omega \times \{t_0\}$ is still a Killing field tangent to the fiber $\partial\Omega \times \{t_0\}$, possibly with singularities or identically zero;

- if (ii) is met, $\partial\Omega$ satisfies $\text{vol}(B_R^{\partial\Omega}) = o(R^2 \log R)$ as $R \rightarrow +\infty$.

Remark 6. For \mathbb{R}^2 , the above theorem generalizes the one-dimensional symmetry result in Theorem 1.2 of [18]. See also [37] for interesting studies on the geometric and topological properties of overdetermined problems in the Euclidean plane.

By the monotonicity Theorem 1.1 in [4], the relation $\langle \nabla u, X \rangle > 0$ on Ω is automatic for globally Lipschitz epigraphs Ω of Euclidean space and for some large class of nonlinearities f including the prototype Allen-Cahn one $f(u) = u - u^3$ (even without requiring the Neumann condition in (6)). However, it is an open problem to enlarge the class of domains $\Omega \subseteq \mathbb{R}^m$ for which $\langle \nabla u, X \rangle > 0$ is met, or to find nontrivial analogues on Riemannian manifolds. In the last section, we move some steps towards this problem by proving some lemmata that may have independent interest. In particular, we obtain the next result:

Proposition 7. *Let $(M, \langle \cdot, \cdot \rangle)$ be a complete Riemannian manifold satisfying $\text{Ric} \geq -(m-1)H^2 \langle \cdot, \cdot \rangle$, for some $H \geq 0$, and let $f \in C^1(\mathbb{R})$ have the properties*

$$\begin{cases} f > 0 & \text{on } (0, \lambda), & f(\lambda) = 0, & f < 0 & \text{on } (\lambda, \lambda + s_0), \\ f(s) \geq \left(\delta_0 + \frac{(m-1)^2 H^2}{4} \right) s & \text{for } s \in (0, s_0), \end{cases}$$

for some $\lambda > 0$ and some small $\delta_0, s_0 > 0$. Let $\Omega \subseteq M$ be an open, connected subset, and suppose that $u \in C^2(\Omega) \cap C^0(\bar{\Omega})$ is a bounded, non-negative solution of

$$\begin{cases} -\Delta u = f(u) & \text{on } \Omega, \\ u > 0 & \text{on } \Omega, \quad \sup_{\partial\Omega} u < \|u\|_{L^\infty(\Omega)}, \end{cases}$$

Suppose that, for each R , $\Omega_R = \{x \in \Omega : \text{dist}(x, \partial\Omega) > R\}$ is non-empty. Then, the following properties hold:

- (I) $\|u\|_{L^\infty(\Omega)} = \lambda$;
- (II) there exists a $R_0 = R_0(m, H, \delta_0) > 0$ such that, for each connected component V_j of Ω_{R_0} , $u(x) \rightarrow \lambda$ uniformly as $\text{dist}(x, \partial\Omega) \rightarrow +\infty$ along V_j . (7)

Remark 8. We underline that $\partial\Omega$ may even have countably many connected components. Moreover, since Ω is possibly non-compact, Ω_{R_0} may have countably many connected components. In this respect, the uniformity guaranteed at point (II) is referred to each single, fixed connected component.

Although, as said, the general problem of ensuring the monotonicity of each solutions of the Dirichlet problem

$$\begin{cases} -\Delta u = f(u) & \text{on } \Omega \\ u > 0 & \text{on } \Omega, \quad u = 0 & \text{on } \partial\Omega \end{cases} \quad (8)$$

is still open, for some ample class of nonlinearities we will be able to construct non-constant solutions of (8) which are strictly monotone, see the next Proposition 41. To do so, we shall restrict the set of Killing fields to the subclass described in the next

Definition 9. *Let $\Omega \subseteq M$ be an open, connected subset with C^3 boundary. A Killing vector field X on $\bar{\Omega}$ is called good for Ω if its flow Φ_t satisfies*

$$\begin{cases} (i) & \Phi_t(\Omega) \subseteq \Omega, \quad \Phi_t(\partial\Omega) \subseteq \bar{\Omega} & \text{for every } t \in \mathbb{R}^+; \\ (ii) & \text{there exists } o \in \partial\Omega \text{ for which } \text{dist}(\Phi_t(o), \partial\Omega) \rightarrow +\infty & \text{as } t \rightarrow +\infty. \end{cases} \quad (9)$$

Observe that property (i) is the somehow minimal requirement for investigating the monotonicity of u with respect to X . The important assumption (ii) enables us to insert arbitrarily large balls in Ω , an essential requirement for our arguments to work. Under the existence of a good Killing field on Ω , Proposition 7 allow us to produce some energy estimate via the method described in [2], leading to the next particularization of Theorem 5 in the three dimensional case:

Theorem 10. *Let $(M, \langle \cdot, \cdot \rangle)$ be a complete Riemannian 3-manifold, with empty boundary and with $\text{Ric} \geq 0$. Let $\Omega \subseteq M$ be an open, connected set with $\partial\Omega \in C^3$, fix $o \in \partial\Omega$ and assume that*

$$\mathcal{H}^2(\partial\Omega \cap B_R) = o(R^2 \log R) \quad \text{as } R \rightarrow +\infty, \quad (10)$$

where $B_R = B_R(o)$ and \mathcal{H}^2 is the 2-dimensional Hausdorff measure. Suppose that Ω has a good Killing field X . Let $f \in C^1(\mathbb{R})$ be such that

$$\begin{cases} f > 0 & \text{on } (0, \lambda), & f(\lambda) = 0, & f < 0 & \text{on } (\lambda, \lambda + s_0), \\ f(s) \geq \delta_0 s & \text{for } s \in (0, s_0), \end{cases}$$

for some $\lambda > 0$ and some small $\delta_0, s_0 > 0$. If there exists a non-constant, positive, solution $u \in C^3(\overline{\Omega})$ of the overdetermined problem

$$\begin{cases} -\Delta u = f(u) & \text{on } \Omega \\ u = 0 & \text{on } \partial\Omega \\ \partial_\nu u = \text{constant} & \text{on } \partial\Omega, \end{cases} \quad (11)$$

such that

$$\begin{cases} \|u\|_{C^1(\Omega)} < +\infty; \\ \langle X, \nabla u \rangle \geq 0 & \text{on } \Omega, \end{cases} \quad (12)$$

then all the conclusions of Theorem 5 hold.

Remark 11. Particularizing to $M = \mathbb{R}^3$ and for globally Lipschitz epigraphs Ω , we recover Theorem 1.8 in [18], see Corollary 46 below.

Setting and notations

Let $(M, \langle \cdot, \cdot \rangle)$ be a smooth connected Riemannian manifold of dimension $m \geq 2$, without boundary. We briefly fix some notation. We denote with K its sectional curvature and with Ric its Ricci tensor. Having fixed an origin o , we set $r(x) = \text{dist}(x, o)$, and we write B_R for geodesic balls centered at o . If we need to emphasize the manifold under consideration, we will add a superscript M , so that, for instance, we will also write Ric^M and B_R^M . The Riemannian m -dimensional volume will be indicated with vol , and its density with dx , while the will write \mathcal{H}^{m-1} for the induced $(m-1)$ -dimensional Hausdorff measure. Throughout the paper, with the symbol $\{\Omega_j\} \uparrow M$ we mean a family $\{\Omega_j\}$, $j \in \mathbb{N}$, of relatively compact, open sets with smooth boundary and satisfying

$$\Omega_j \Subset \Omega_{j+1} \Subset M, \quad M = \bigcup_{j=0}^{+\infty} \Omega_j,$$

where $A \Subset B$ means $\overline{A} \subseteq B$. Such a family will be called an exhaustion of M . Hereafter, we consider

$$f \in C^1(\mathbb{R}), \quad (13)$$

and a solution u on M of

$$-\Delta u = f(u) \quad \text{on } M. \quad (14)$$

Remark 12. To avoid unessential technicalities, hereafter we assume that $u \in C^3(M)$. By standard elliptic estimates (see [21]), $u \in C^3$ is automatic whenever $f \in C_{\text{loc}}^{1,\alpha}(\mathbb{R})$, for some $\alpha \in (0, 1)$, and u is a locally bounded weak solution of (14). Analogously, for an open set $\Omega \subseteq M$ with boundary, we shall restrict to $u \in C^3(\overline{\Omega})$. This condition is automatically satisfied whenever $\partial\Omega$ is, for instance, of class C^3 .

Remark 13. For the same reason, we shall restrict to $f \in C^1(\mathbb{R})$, although our statements could be rephrased for $f \in \text{Lip}_{\text{loc}}(\mathbb{R})$ with some extra-care in the definition of stability. In this respect, we suggest the reader to consult [15] for a detailed discussion.

We recall that u is characterized, on each open subset $U \Subset M$, as a stationary point of the energy functional $E_U : H^1(U) \rightarrow \mathbb{R}$ given by

$$E_U(w) = \frac{1}{2} \int_U |\nabla w|^2 dx - \int_U F(w) dx, \quad \text{where } F(t) = \int_0^t f(s) ds, \quad (15)$$

with respect to compactly supported variations in U . Let J be the Jacobi operator of E at u , that is,

$$J\phi = -\Delta\phi - f'(u)\phi \quad \forall \phi \in C_c^\infty(M). \quad (16)$$

Definition 14. *The function u solving (14) is said to be a **stable solution** if J is non-negative on $C_c^\infty(M)$, that is, if $(\phi, J\phi)_{L^2} \geq 0$ for each $\phi \in C_c^\infty(M)$. Integrating by parts, this reads as*

$$\int_M f'(u)\phi^2 dx \leq \int_M |\nabla\phi|^2 dx \quad \text{for every } \phi \in C_c^\infty(M). \quad (17)$$

By density, we can replace $C_c^\infty(M)$ in (17) with $\text{Lip}_c(M)$. By a result of [20] and [30] (see also [34], Section 3) the stability of u turns out to be equivalent to the existence of a positive $w \in C^1(M)$ solving $\Delta w + f'(u)w = 0$ weakly on M .

Some preliminary computation

We start with a Picone-type identity.

Lemma 15. *Let $\Omega \subseteq M$ be an open, connected set with C^3 boundary (possibly empty) and let $u \in C^3(\overline{\Omega})$ be a solution of $-\Delta u = f(u)$ on Ω . Let $w \in C^1(\overline{\Omega}) \cap C^2(\Omega)$ be a solution of $\Delta w + f'(u)w \leq 0$ such that $w > 0$ on Ω . Then, the following inequality holds true: for every $\varepsilon > 0$ and for every $\phi \in \text{Lip}_c(M)$,*

$$\begin{aligned} \int_{\partial\Omega} \frac{\phi^2}{w + \varepsilon} (\partial_\nu w) d\mathcal{H}^{m-1} &\leq \int_\Omega |\nabla\phi|^2 dx - \int_\Omega f'(u) \frac{w}{w + \varepsilon} \phi^2 dx \\ &\quad - \int_\Omega (w + \varepsilon)^2 \left| \nabla \left(\frac{\phi}{w + \varepsilon} \right) \right|^2 dx \end{aligned} \quad (18)$$

Furthermore, if either $\Omega = M$ or $w > 0$ on $\overline{\Omega}$, one can also take $\varepsilon = 0$ inside the above inequality. The inequality is indeed an equality if w solves $\Delta w + f'(u)w = 0$ on Ω .

Proof. We integrate $\Delta w + f'(u)w \leq 0$ against the test function $\phi^2/(w + \varepsilon)$ to deduce

$$\begin{aligned} 0 \leq & - \int_{\Omega} (\Delta w + f'(u)w) \frac{\phi^2}{w + \varepsilon} dx = - \int_{\partial\Omega} \frac{\phi^2}{w + \varepsilon} (\partial_{\nu} w) d\mathcal{H}^{m-1} \\ & + \int_{\Omega} \langle \nabla \left(\frac{\phi^2}{w + \varepsilon} \right), \nabla w \rangle dx - \int_{\Omega} f'(u)w \frac{\phi^2}{w + \varepsilon} dx. \end{aligned} \quad (19)$$

Since

$$\langle \nabla \left(\frac{\phi^2}{w + \varepsilon} \right), \nabla w \rangle = 2 \frac{\phi}{w + \varepsilon} \langle \nabla \phi, \nabla w \rangle - \frac{\phi^2}{(w + \varepsilon)^2} |\nabla w|^2, \quad (20)$$

using the identity

$$(w + \varepsilon)^2 \left| \nabla \left(\frac{\phi}{w + \varepsilon} \right) \right|^2 = |\nabla \phi|^2 + \frac{\phi^2}{(w + \varepsilon)^2} |\nabla w|^2 - 2 \frac{\phi}{w + \varepsilon} \langle \nabla w, \nabla \phi \rangle \quad (21)$$

we infer that

$$\langle \nabla \left(\frac{\phi^2}{w + \varepsilon} \right), \nabla w \rangle = |\nabla \phi|^2 - (w + \varepsilon)^2 \left| \nabla \left(\frac{\phi}{w + \varepsilon} \right) \right|^2. \quad (22)$$

Inserting into (19) we get the desired (18). \square

Next step is to obtain an integral equality involving the second derivatives of u . This geometric Poincaré-type formula has its roots in the paper [18], which deals with subsets of Euclidean space, and in the previous works [41, 40].

Proposition 16. *In the above assumptions, for every $\varepsilon > 0$ the following integral inequality holds true:*

$$\begin{aligned} & \int_{\Omega} [|\nabla du|^2 + \text{Ric}(\nabla u, \nabla u)] \frac{\phi^2 w}{w + \varepsilon} dx - \int_{\Omega} |\nabla |\nabla u||^2 \phi^2 dx \\ & \leq \int_{\partial\Omega} \frac{\phi^2}{w + \varepsilon} \left[w \partial_{\nu} \left(\frac{|\nabla u|^2}{2} \right) - |\nabla u|^2 \partial_{\nu} w \right] d\mathcal{H}^{m-1} \\ & + \varepsilon \int_{\Omega} \frac{\phi}{w + \varepsilon} \langle \nabla \phi, \nabla |\nabla u|^2 \rangle dx - \frac{1}{2} \int_{\Omega} \phi^2 \langle \nabla |\nabla u|^2, \nabla \left(\frac{w}{w + \varepsilon} \right) \rangle dx \\ & + \int_{\Omega} |\nabla \phi|^2 |\nabla u|^2 dx - \int_{\Omega} (w + \varepsilon)^2 \left| \nabla \left(\frac{\phi |\nabla u|}{w + \varepsilon} \right) \right|^2 dx. \end{aligned} \quad (23)$$

Furthermore, if either $\Omega = M$ or $w > 0$ on $\bar{\Omega}$, one can also take $\varepsilon = 0$. The inequality is indeed an equality if $\Delta w + f'(u)w = 0$ on Ω .

Proof. We start with the Böchner formula

$$\frac{1}{2} \Delta |\nabla u|^2 = \langle \nabla \Delta u, \nabla u \rangle + \text{Ric}(\nabla u, \nabla u) + |\nabla du|^2, \quad (24)$$

valid for each $u \in C^3(\bar{\Omega})$. The proof of this formula is standard and can be deduced from Ricci commutation laws. Since u solves $-\Delta u = f(u)$, we get

$$\frac{1}{2} \Delta |\nabla u|^2 = -f'(u) |\nabla u|^2 + \text{Ric}(\nabla u, \nabla u) + |\nabla du|^2. \quad (25)$$

Integrating (25) on Ω against the test function $\psi = \phi^2 w / (w + \varepsilon)$ we deduce

$$\begin{aligned}
& \int_{\Omega} [|\nabla du|^2 + \text{Ric}(\nabla u, \nabla u)] \psi dx \\
&= \int_{\Omega} f'(u) |\nabla u|^2 \frac{w}{w + \varepsilon} \phi^2 dx + \frac{1}{2} \int_{\Omega} \frac{w \phi^2}{w + \varepsilon} \Delta |\nabla u|^2 dx = \\
&= \int_{\Omega} f'(u) |\nabla u|^2 \frac{w}{w + \varepsilon} \phi^2 dx + \frac{1}{2} \int_{\partial\Omega} \frac{w \phi^2}{w + \varepsilon} \partial_{\nu} |\nabla u|^2 d\mathcal{H}^{m-1} \\
&\quad - \frac{1}{2} \int_{\Omega} \langle \nabla \left(\frac{w \phi^2}{w + \varepsilon} \right), \nabla |\nabla u|^2 \rangle dx \\
&= \int_{\Omega} f'(u) |\nabla u|^2 \frac{w}{w + \varepsilon} \phi^2 dx + \frac{1}{2} \int_{\partial\Omega} \frac{w \phi^2}{w + \varepsilon} \partial_{\nu} |\nabla u|^2 d\mathcal{H}^{m-1} \\
&\quad - \int_{\Omega} \frac{w \phi}{w + \varepsilon} \langle \nabla \phi, \nabla |\nabla u|^2 \rangle dx - \frac{1}{2} \int_{\Omega} \phi^2 \langle \nabla |\nabla u|^2, \nabla \left(\frac{w}{w + \varepsilon} \right) \rangle dx.
\end{aligned} \tag{26}$$

Next, we consider the spectral inequality (18) with test function $\phi |\nabla u| \in \text{Lip}_c(M)$:

$$\begin{aligned}
& \int_{\partial\Omega} |\nabla u|^2 \frac{\phi^2}{w + \varepsilon} (\partial_{\nu} w) d\mathcal{H}^{m-1} \\
&\leq \int_{\Omega} |\nabla(\phi |\nabla u|)|^2 dx - \int_{\Omega} f'(u) \frac{w}{w + \varepsilon} |\nabla u|^2 \phi^2 dx \\
&\quad - \int_{\Omega} (w + \varepsilon)^2 \left| \nabla \left(\frac{\phi |\nabla u|}{w + \varepsilon} \right) \right|^2 dx \\
&= \int_{\Omega} |\nabla \phi|^2 |\nabla u|^2 dx + \int_{\Omega} \phi^2 |\nabla |\nabla u||^2 dx + 2 \int_{\Omega} \phi |\nabla u| \langle \nabla \phi, \nabla |\nabla u| \rangle dx \\
&\quad - \int_{\Omega} f'(u) \frac{w}{w + \varepsilon} |\nabla u|^2 \phi^2 dx - \int_{\Omega} (w + \varepsilon)^2 \left| \nabla \left(\frac{\phi |\nabla u|}{w + \varepsilon} \right) \right|^2 dx.
\end{aligned} \tag{27}$$

Recalling that $\nabla |\nabla u|^2 = 2 |\nabla u| \nabla |\nabla u|$ weakly on M , summing up (27) and (26), putting together the terms of the same kind and rearranging we deduce (23). \square

Proposition 17. *In the above assumptions, if it holds*

$$\liminf_{\varepsilon \rightarrow 0^+} \int_{\Omega} \phi^2 \langle \nabla |\nabla u|^2, \nabla \left(\frac{w}{w + \varepsilon} \right) \rangle dx \geq 0, \tag{28}$$

Then

$$\begin{aligned}
& \int_{\Omega} [|\nabla du|^2 + \text{Ric}(\nabla u, \nabla u) - |\nabla |\nabla u||^2] \phi^2 dx \\
&+ \liminf_{\varepsilon \rightarrow 0^+} \int_{\Omega} (w + \varepsilon)^2 \left| \nabla \left(\frac{\phi |\nabla u|}{w + \varepsilon} \right) \right|^2 dx \leq \\
&\leq \int_{\Omega} |\nabla \phi|^2 |\nabla u|^2 dx + \liminf_{\varepsilon \rightarrow 0^+} \int_{\partial\Omega} \frac{\phi^2}{w + \varepsilon} \left[w \partial_{\nu} \left(\frac{|\nabla u|^2}{2} \right) - |\nabla u|^2 \partial_{\nu} w \right] d\mathcal{H}^{m-1}.
\end{aligned} \tag{29}$$

Proof. We take limits as $\varepsilon \rightarrow 0^+$ in (23) along appropriate sequences. It is easy to see that

$$\varepsilon \int_{\Omega} \frac{\phi}{w + \varepsilon} \langle \nabla \phi, \nabla |\nabla u|^2 \rangle dx = o(1)$$

as $\varepsilon \rightarrow 0$. Indeed, we can apply Lebesgue convergence theorem, since $|\varepsilon/(w + \varepsilon)| \leq 1$, and we have convergence to the integral of the pointwise limit, which is:

$$\int_{\{w=0\}} \phi \langle \nabla \phi, \nabla |\nabla u|^2 \rangle dx = 0,$$

being $\{w = 0\} \subseteq \partial\Omega$. Lebesgue theorem can also be applied for the other terms in a straightforward way, with the exception of the term that needs (28). \square

Next, we need the following formula, that extends works of P. Sternberg and K. Zumbrun in [41, 40].

Proposition 18. *Let u be a C^2 function on M , and let $p \in M$ be a point such that $\nabla u(p) \neq 0$. Then, denoting with $|II|^2$ the second fundamental form of the level set $\Sigma = \{u = u(p)\}$ in a neighbourhood of p , it holds*

$$|\nabla du|^2 - |\nabla |\nabla u||^2 = |\nabla u|^2 |II|^2 + |\nabla_T |\nabla u||^2,$$

where ∇_T is the tangential gradient on the level set Σ .

Proof. Fix a local orthonormal frame $\{e_i\}$ on Σ , and let $\nu = \nabla u / |\nabla u|$ be the normal vector. For every vector field $X \in \Gamma(TM)$,

$$\nabla du(\nu, X) = \frac{1}{|\nabla u|} \nabla du(\nabla u, X) = \frac{1}{2|\nabla u|} \langle \nabla |\nabla u|^2, X \rangle = \langle \nabla |\nabla u|, X \rangle.$$

Moreover, for a level set

$$II = -\frac{\nabla du|_{T\Sigma \times T\Sigma}}{|\nabla u|}.$$

Therefore:

$$\begin{aligned} |\nabla du|^2 &= \sum_{i,j} [\nabla du(e_i, e_j)]^2 + 2 \sum_j [\nabla du(\nu, e_j)]^2 + [\nabla du(\nu, \nu)]^2 \\ &= |\nabla u|^2 |II|^2 + 2 \sum_j [\langle \nabla |\nabla u|, e_j \rangle]^2 + [\langle \nabla |\nabla u|, \nu \rangle]^2 \\ &= |\nabla u|^2 |II|^2 + |\nabla_T |\nabla u||^2 + |\nabla |\nabla u||^2. \end{aligned}$$

proving the proposition. \square

Splitting and structure theorems: the boundaryless case

Our first result deals with the case when $\Omega = M$ has no boundary. It is inspired by the ones proved in [14, 15] for the Euclidean case, and also extends and strengthens some previous work in [16].

Proof of Theorem 1. In our assumption, we consider the integral formula (23) with $M = \Omega$ and $\varepsilon = 0$. Since $\text{Ric} \geq 0$ we deduce

$$\int_M \left[|\nabla du|^2 - |\nabla|\nabla u||^2 \right] \phi^2 dx \leq \int_M |\nabla\phi|^2 |\nabla u|^2 dx - \int_M w^2 \left| \nabla \left(\frac{\phi|\nabla u|}{w} \right) \right|^2 dx. \quad (30)$$

Next, we rearrange the RHS as follows: using the inequality

$$|X + Y|^2 \geq |X|^2 + |Y|^2 - 2|X||Y| \geq (1 - \delta)|X|^2 + (1 - \delta^{-1})|Y|^2,$$

valid for each $\delta > 0$, we obtain

$$\begin{aligned} w^2 \left| \nabla \left(\frac{\phi|\nabla u|}{w} \right) \right|^2 &= w^2 \left| \frac{|\nabla u| \nabla \phi}{w} + \phi \nabla \left(\frac{|\nabla u|}{w} \right) \right|^2 \\ &\geq (1 - \delta^{-1}) |\nabla u|^2 |\nabla \phi|^2 + (1 - \delta) \phi^2 w^2 \left| \nabla \left(\frac{|\nabla u|}{w} \right) \right|^2. \end{aligned} \quad (31)$$

Substituting in (30) yields

$$\int_M \left[|\nabla du|^2 - |\nabla|\nabla u||^2 \right] \phi^2 dx + (1 - \delta) \int_M \phi^2 w^2 \left| \nabla \left(\frac{|\nabla u|}{w} \right) \right|^2 dx \leq \frac{1}{\delta} \int_M |\nabla\phi|^2 |\nabla u|^2 dx. \quad (32)$$

Choose $\delta < 1$. We claim that, for suitable families $\{\phi_\alpha\}_{\alpha \in I \subseteq \mathbb{R}^+}$, it holds

$$\{\phi_\alpha\} \text{ is monotone increasing to } 1, \quad \lim_{\alpha \rightarrow +\infty} \int_M |\nabla\phi_\alpha|^2 |\nabla u|^2 dx = 0. \quad (33)$$

Choose ϕ as follows, according to the case.

In case (i), fix $\Omega \Subset M$ with smooth boundary and let $\{\Omega_j\} \uparrow M$ be a smooth exhaustion with $\Omega \Subset \Omega_1$. Choose $\phi = \phi_j \in \text{Lip}_c(M)$ to be identically 1 on Ω , 0 on $M \setminus \Omega_j$ and the harmonic capacitor on $\Omega_j \setminus \Omega$, that is, the solution of

$$\begin{cases} \Delta\phi_j = 0 & \text{on } \Omega_j \setminus \Omega, \\ \phi_j = 1 & \text{on } \partial\Omega, \quad \phi_j = 0 & \text{on } \partial\Omega_j. \end{cases}$$

Note that $\phi_j \in \text{Lip}_c(M)$ is ensured by elliptic regularity up to $\partial\Omega$ and $\partial\Omega_j$. By comparison and since M is parabolic, $\{\phi_j\}$ is monotonically increasing and pointwise convergent to 1, and moreover

$$\int_{\Omega_j} |\nabla\phi_j|^2 |\nabla u|^2 dx \leq \|\nabla u\|_{L^\infty}^2 \text{cap}(\Omega, \Omega_j) \rightarrow \|\nabla u\|_{L^\infty}^2 \text{cap}(\Omega) = 0,$$

the last equality following since M is parabolic. This proves (33).

In case (ii), we apply a logarithmic cutoff argument. For fixed $R > 0$, choose the following radial $\phi(x) = \phi_R(r(x))$:

$$\phi_R(r) = \begin{cases} 1 & \text{if } r \leq \sqrt{R}, \\ 2 - 2 \frac{\log r}{\log R} & \text{if } r \in [\sqrt{R}, R], \\ 0 & \text{if } r \geq R. \end{cases} \quad (34)$$

Note that

$$|\nabla\phi(x)|^2 = \frac{4}{r(x)^2 \log^2 R} \chi_{B_R \setminus B_{\sqrt{R}}}(x),$$

where χ_A is the indicatrix function of a subset $A \subseteq M$. Choose R in such a way that $\log R/2$ is an integer. Then,

$$\begin{aligned} \int_M |\nabla\phi|^2 |\nabla u|^2 dx &= \int_{B_R \setminus B_{\sqrt{R}}} |\nabla\phi|^2 |\nabla u|^2 dx = \frac{4}{\log^2 R} \sum_{k=\log R/2}^{\log R-1} \int_{B_{e^{k+1}} \setminus B_{e^k}} \frac{|\nabla u|^2}{r(x)^2} dx \\ &\leq \frac{4}{\log^2 R} \sum_{k=\log R/2}^{\log R} \frac{1}{e^{2k}} \int_{B_{e^{k+1}}} |\nabla u|^2 dx. \end{aligned} \tag{35}$$

By assumption,

$$\int_{B_{e^{k+1}}} |\nabla u|^2 dx \leq (k+1)e^{2(k+1)}\delta(k)$$

for some $\delta(k)$ satisfying $\delta(k) \rightarrow 0$ as $k \rightarrow +\infty$. Without loss of generality, we can assume $\delta(k)$ to be decreasing as a function of k . Whence,

$$\begin{aligned} \frac{4}{\log^2 R} \sum_{k=\log R/2}^{\log R} \frac{1}{e^{2k}} \int_{B_{e^{k+1}}} |\nabla u|^2 dx &\leq \frac{8}{\log^2 R} \sum_{k=\log R/2}^{\log R} \frac{e^{2(k+1)}}{e^{2k}} (k+1)\delta(k) \\ &\leq \frac{8e^2}{\log^2 R} \delta(\log R/2) \sum_{k=0}^{\log R} (k+1) \leq \frac{C}{\log^2 R} \delta(\log R/2) \log^2 R = C\delta(\log R/2), \end{aligned} \tag{36}$$

for some constant $C > 0$. Combining (35) and (36) and letting $R \rightarrow +\infty$ we deduce (33).

In both the cases, we can infer from the integral formula that

$$|\nabla u| = cw, \quad \text{for some } c \geq 0, \quad |\nabla u|^2 = |\nabla|\nabla u||^2, \quad \text{Ric}(\nabla u, \nabla u) = 0. \tag{37}$$

Since u is non-constant by assumption, $c > 0$, thus $|\nabla u| > 0$ on M . From Böchner formula, it holds

$$|\nabla u| \Delta |\nabla u| + |\nabla|\nabla u||^2 = \frac{1}{2} \Delta |\nabla u|^2 = \text{Ric}(\nabla u, \nabla u) + |\nabla du|^2 - f'(u) |\nabla u|^2$$

on M . Using (37), we thus deduce that $\Delta |\nabla u| + f'(u) |\nabla u| = 0$ on M , hence $|\nabla u|$ (and so w) both solve the linearized equation $Jv = 0$.

Now, the flow Φ of $\nu = \nabla u / |\nabla u|$ is well defined on M . Since M is complete and $|\nu| = 1$ is bounded, Φ is defined on $M \times \mathbb{R}$. By (37) and Proposition 18, $|\nabla u|$ is constant on each connected component of a level set N , and N is totally geodesic. Therefore, in a local Darboux frame $\{e_j, \nu\}$ for the level surface N ,

$$\begin{aligned} 0 &= |II|^2 \implies \nabla du(e_i, e_j) = 0 \\ 0 &= \langle \nabla|\nabla u|, e_j \rangle = \nabla du(\nu, e_j), \end{aligned} \tag{38}$$

so the unique nonzero component of ∇du is that corresponding to the pair (ν, ν) . Let γ be any integral curve of ν . Then

$$\frac{d}{dt}(u \circ \gamma) = \langle \nabla u, \nu \rangle = |\nabla u| \circ \gamma > 0$$

and

$$\begin{aligned} -f(u \circ \gamma) &= \Delta u(\gamma) = \nabla du(\nu, \nu)(\gamma) = \langle \nabla |\nabla u|, \nu \rangle(\gamma) \\ &= \frac{d}{dt}(|\nabla u| \circ \gamma) = \frac{d^2}{dt^2}(u \circ \gamma), \end{aligned}$$

thus $y = u \circ \gamma$ solves the ODE $y'' = -f(y)$ and $y' > 0$. Note also that the integral curves γ of ν are geodesics. Indeed,

$$\begin{aligned} \nabla_{\gamma'} \gamma' &= \frac{1}{|\nabla u|} \nabla_{\nabla u} \left(\frac{\nabla u}{|\nabla u|} \right) = \frac{1}{|\nabla u|^2} \nabla_{\nabla u} \nabla u - \frac{1}{|\nabla u|^3} \nabla u (|\nabla u|) \nabla u \\ &= \frac{1}{|\nabla u|^2} \nabla du(\nabla u, \cdot)^\sharp - \frac{1}{|\nabla u|^3} \langle \nabla |\nabla u|, \nabla u \rangle \nabla u \\ &= \frac{1}{|\nabla u|} \nabla du(\nu, \cdot)^\sharp - \frac{1}{|\nabla u|} \langle \nabla |\nabla u|, \nu \rangle \nu = \frac{1}{|\nabla u|} \nabla du(\nu, \cdot)^\sharp - \frac{1}{|\nabla u|} \nabla du(\nu, \nu) \nu = 0, \end{aligned}$$

where the first equality in the last line follows from (38). We now address the topological part of the splitting, following arguments in the proof of [34], Theorem 9.3. Since $|\nabla u|$ is constant on level sets of u , $|\nabla u| = \beta(u)$ for some function β . Evaluating along curves $\Phi_t(x)$, since $u \circ \Phi_t$ is a local bijection we deduce that β is continuous. We claim that Φ_t moves level sets of u to level sets of u . Indeed, integrating $d/ds(u \circ \Phi_s) = |\nabla u| \circ \Phi_s = \alpha(u \circ \Phi_s)$ we get

$$t = \int_{u(x)}^{u(\Phi_t(x))} \frac{d\xi}{\alpha(\xi)},$$

thus $u(\Phi_t(x))$ is independent of x varying in a level set. As $\alpha(\xi) > 0$, this also show that flow lines starting from a level set of u do not touch the same level set. Let N be a connected component of a level set of u . Since the flow of ν is through geodesics, for each $x \in N$ $\Phi_t(x)$ coincides with the normal exponential map $\exp^\perp(t\nu(x))$. Moreover, since N is closed in M and M is complete, the normal exponential map is surjective: indeed, by variational arguments, each geodesic from $x \in M$ to N minimizing $\text{dist}(x, N)$ is perpendicular to N . This shows that $\Phi_{|N \times \mathbb{R}}$ is surjective. We now prove the injectivity of $\Phi_{|N \times \mathbb{R}}$. Suppose that $\Phi(x_1, t_1) = \Phi(x_2, t_2)$. Then, since Φ moves level sets to level sets, necessarily $t_1 = t_2 = t$. If by contradiction $x_1 \neq x_2$, two distinct flow lines of Φ_t would intersect at the point $\Phi_t(x_1) = \Phi_t(x_2)$, contradicting the fact that Φ_t is a diffeomorphism on M for every t . Concluding, $\Phi : N \times \mathbb{R} \rightarrow M$ is a diffeomorphism. In particular, each level set $\Phi_t(N)$ is connected. This proves the topological part of the splitting.

We are left with the Riemannian part. We consider the Lie derivative of the metric in the direction of Φ_t :

$$\begin{aligned} (L_\nu \langle, \rangle)(X, Y) &= \langle \nabla_X \nu, Y \rangle + \langle X, \nabla_Y \nu \rangle \\ &= \frac{2}{|\nabla u|} \nabla du(X, Y) + X \left(\frac{1}{|\nabla u|} \right) \langle \nabla u, Y \rangle + Y \left(\frac{1}{|\nabla u|} \right) \langle \nabla u, X \rangle. \end{aligned}$$

From the expression, using that $|\nabla u|$ is constant on N and the properties of ∇du we deduce that

$$(L_\nu \langle, \rangle)(X, Y) = \frac{2}{|\nabla u|} \nabla du(X, Y) = 0.$$

If at least one between X and Y is in the tangent space of N . If, however, X and Y are

normal, (w.l.o.g. $X = Y = \nabla u$), we have

$$\begin{aligned} (L_\nu \langle \cdot, \cdot \rangle)(\nabla u, \nabla u) &= \frac{2}{|\nabla u|} \nabla du(\nabla u, \nabla u) + 2\nabla u \left(\frac{1}{|\nabla u|} \right) |\nabla u|^2 \\ &= \frac{2}{|\nabla u|} \nabla du(\nabla u, \nabla u) - 2\nabla u(|\nabla u|) = 2\nabla du(\nu, \nabla u) - 2\langle \nabla |\nabla u|, \nabla u \rangle = 0. \end{aligned}$$

Concluding, $L_\nu \langle \cdot, \cdot \rangle = 0$, thus Φ_t is a flow of isometries. Since $\nabla u \perp TN$, M splits as a Riemannian product, as desired. In particular, $\text{Ric}^N \geq 0$ if $m \geq 3$, while, if $m = 2$, $M = \mathbb{R}^2$ or $\mathbb{S}^1 \times \mathbb{R}$ with the flat metric.

We next address the parabolicity. Under assumption (i), M is parabolic and so N is necessarily parabolic too. We are going to deduce the same under assumption (ii). To this end, it is enough to prove the volume estimate (2). Indeed, (2) is a sufficient condition on M to be parabolic. The chain of inequalities

$$\left(\int_{-R}^R |y'(t)|^2 dt \right) \text{vol}(B_R^N) \leq \int_{[-R, R] \times B_R^N} |y'(t)|^2 dt dx^N \leq \int_{B_{R\sqrt{2}}} |\nabla u|^2 dx = o(R^2 \log R)$$

gives immediately (2) and (3), since $|y'| > 0$ everywhere. \square

Remark 19. The proof of Theorem 1 is tightly related to some works in [34] and [35], see in particular Theorems 4.5 and 9.3 in [34], and Theorem 4 in [35]. We note that, however, our technique is different from the one used to prove the vanishing Theorem 4.5 in [34]. Namely, this latter is based on showing that $|\nabla u|/w$ is a weak solution of the inequality

$$\Delta_{w^2} \left(\frac{|\nabla u|}{w} \right) \geq 0 \quad \text{on } M, \quad \text{where} \quad \Delta_{w^2} = w^{-2} \text{div}(w^2 \nabla \cdot),$$

and then concluding via a refined Liouville-type result that improves on works of [3] (Theorem 1.8) and [2] (Proposition 2.1). However, this approach seems to reveal some difficulties when dealing with sets Ω having non-empty boundary, thereby demanding a different method. Our technique, which uses from the very beginning the spectral inequality (18), is closer in spirit to the one in [35].

Under suitable sign assumptions on f , Theorem 1 implies a Liouville type result thanks to a Caccioppoli-type estimate. This is the content of the next

Theorem 20. *Let $(M, \langle \cdot, \cdot \rangle)$ be a complete, non-compact manifold with $\text{Ric} \geq 0$ and let $u \in C^3(M)$ be a bounded stable solution of $-\Delta u = f(u)$ on M , with $f \in C^1(\mathbb{R})$ and*

$$f(r) \geq 0 \quad \text{for any } r \in \mathbb{R}. \quad (39)$$

Suppose that either $m \leq 4$ or

$$\text{vol}(B_R) = o(R^4 \log R) \quad \text{as } R \rightarrow +\infty. \quad (40)$$

Then, u is constant.

Proof. The proof is by contradiction. Suppose that u is not constant and set $u^* = \sup_M u$. Then, multiplying the equation by $\phi^2(u^* - u)$ and integrating by parts we get

$$\begin{aligned} 0 &\geq \int_M |\nabla u|^2 \phi^2 dx - 2 \int_M \phi(u^* - u) \langle \nabla u, \nabla \phi \rangle dx \\ &\geq \int_M |\nabla u|^2 \phi^2 dx - 4\|u\|_{L^\infty} \int_M \phi |\nabla u| |\nabla \phi| dx, \end{aligned}$$

thus, by Young inequality, there exists $C = C(\|u\|_{L^\infty}) > 0$ such that

$$\frac{1}{2} \int_M |\nabla u|^2 \phi^2 dx \leq C \int_M |\nabla \phi|^2 dx.$$

Considering a radial $\phi(x) = \phi_R(r(x))$, where $\phi_R(r)$ satisfies

$$\phi_R(r) = 1 \quad \text{on } [0, R], \quad \phi_R(r) = \frac{2R - r}{R} \quad \text{on } [R, 2R], \quad \phi_R(r) = 0 \quad \text{on } [2R, +\infty)$$

we get

$$\int_{B_R} |\nabla u|^2 dx \leq \int_M |\nabla u|^2 \phi^2 dx \leq C \text{vol}(B_{2R}) R^{-2}$$

where $C > 0$ is a constant independent of R . When $m \leq 4$, we have $\text{vol}(B_{2R}) \leq C'R^4$ (for some constant $C' > 0$ independent of R) by Bishop-Gromov volume comparison theorem, thus condition (1) in Theorem 1 is satisfied. On the other hand, (1) is always satisfied when $m \geq 5$ and (40) are in force. Therefore, by Theorem 1, $u = y(t)$ solves

$$-y'' = f(y) \geq 0$$

hence y , being nonconstant, must necessarily be unbounded, a contradiction that concludes the proof. \square

Theorem 1 can be iterated to deduce the structure Theorem 3. To do so, we define the following families:

$\mathcal{F}_1 = \{\text{complete, parabolic manifolds } (M, \langle \cdot, \cdot \rangle) \text{ with } \text{Ric} \geq 0, \text{ admitting no stable, non-constant solutions } u \in C^3(M) \text{ of } -\Delta u = f(u) \text{ with } |\nabla u| \in L^\infty(M), \text{ for any } f \in C^1(\mathbb{R})\}.$

$\mathcal{F}_2 = \{\text{complete manifolds } (M, \langle \cdot, \cdot \rangle) \text{ with } \text{Ric} \geq 0, \text{ admitting no stable, non-constant solutions } u \in C^3(M) \text{ of } -\Delta u = f(u) \text{ for which}$

$$\int_{B_R} |\nabla u|^2 dx = o(R^2 \log R) \quad \text{as } R \rightarrow +\infty,$$

$\text{for any } f \in C^1(\mathbb{R})\}.$

The next result is an immediate consequence of Theorem 1, and improve upon previous works in [12, 16].

Corollary 21. *If $(M, \langle \cdot, \cdot \rangle)$ be complete, non-compact Riemannian manifold. Suppose that M has quasi-positive Ricci curvature, that is, that $\text{Ric} \geq 0$ and $\text{Ric}_x > 0$ for some point $x \in M$. Then, $M \in \mathcal{F}_2$. If $m = 2$, we also have $M \in \mathcal{F}_1$.*

Proof. Otherwise, by Theorem 1, $M = N \times \mathbb{R}$ with the product metric $\langle \cdot, \cdot \rangle = \langle \cdot, \cdot \rangle_N + dt^2$ (if $m = 2$, by Bishop-Gromov volume comparison $\text{vol}(B_R) \leq \pi R^2$, so M is parabolic). Therefore, $\text{Ric}(\partial_t, \partial_t) = 0$ at every point $x = (\bar{x}, t) \in M$, contradicting the quasi-positivity assumption. \square

Remark 22. The above conclusion is sharp. Indeed, \mathbb{R}^2 equipped with its canonical flat metric is parabolic and supports the function $u(x, y) = x$, which is a non-constant, harmonic function, hence a non-constant stable solution of (14) with $f = 0$.

Remark 23. By results in [25, 15], any compact manifold (M, \langle, \rangle) with $\text{Ric} \geq 0$ belongs to $\mathcal{F}_1 \cap \mathcal{F}_2$.

To proceed with the investigation of $\mathcal{F}_1, \mathcal{F}_2$, we need a preliminary computation.

Proposition 24. *Let X be a vector field on (M^m, \langle, \rangle) , and let $u \in C^3(M)$ be a solution of $-\Delta u = f(u)$, for some $f \in C^1(\mathbb{R})$. Set for convenience $T = \frac{1}{2}L_X \langle, \rangle$. Then, the function $w = \langle \nabla u, X \rangle$ solves*

$$\Delta w + f'(u)w = 2\langle \nabla du, T \rangle + [2\text{div}(T) - \text{dTr}(T)](\nabla u). \quad (41)$$

In particular, if X is conformal, that is, $L_X \langle, \rangle = \eta \langle, \rangle$, for some $\eta \in C^\infty(M)$, then

$$\Delta w + f'(u)w = -\eta f(u) + (2 - m)\langle \nabla \eta, \nabla u \rangle.$$

Proof. Fix a local orthonormal frame $\{e_i\}$, with dual coframe $\{\theta^j\}$. Let R_{ijkl} be the components of the $(4, 0)$ curvature tensor, with the standard sign agreement. We have

$$X = X^k e_k, \quad w = u_k X^k, \quad \nabla X = X_i^k \theta^i \otimes e_k, \quad \nabla du = u_{ki} \theta^i \otimes \theta^k.$$

For notational convenience, we lower all the indices with the aid of the metric $\langle, \rangle = (g_{jk})$. Note that, for X , the lowered index is in the first position, that is, $X_{ki} = g_{kr} X_r^i$. According to the definition of T ,

$$T_{ik} = \frac{1}{2}(L_X \langle, \rangle)_{ik} = \frac{1}{2}(X_{ik} + X_{ki}).$$

Then, since $u \in C^3(M)$,

$$\begin{aligned} \Delta w &= (u_k X_k)_{ii} = u_{ki,i} X_k + 2u_{ki} X_{ki} + u_k X_{ki,i} \\ &= u_{ki,i} X_k + 2u_{ki} T_{ki} + u_k X_{ki,i}, \end{aligned} \quad (42)$$

where the equality in the last row follows since ∇du is symmetric, whence only the symmetric part of ∇X survives. From Ricci commutation laws

$$u_{rk,i} = u_{ri,k} + u_t R_{trki}, \quad X_{rk,i} = X_{ri,k} + X_t R_{trki}, \quad (43)$$

Schwartz symmetry for the second derivatives of u , and the equality $X_{ii} = T_{ii}$ we deduce

$$\begin{aligned} u_{ki,i} &= u_{ik,i} = u_{ii,k} + u_t R_{tik i} = (\Delta u)_k + u_t \text{Ric}_{tk}, \\ X_{ki,i} &= (2T_{ki} - X_{ik})_i = 2T_{ki,i} - X_{ik,i} = 2T_{ki,i} - X_{ii,k} - X_t R_{tik i} \\ &= 2T_{ki,i} - T_{ii,k} - X_t \text{Ric}_{tk} \end{aligned} \quad (44)$$

Using (44) in (42) we infer that

$$\begin{aligned} \Delta w &= (\Delta u)_k X_k + u_t \text{Ric}_{tk} X_k + 2u_{ki} T_{ki} + u_k (2T_{ki,i} - T_{ii,k}) - u_k X_t \text{Ric}_{tk} \\ &= -f'(u)u_k X_k + 2u_{ki} T_{ki} + 2u_k T_{ki,i} - u_k T_{ii,k}, \end{aligned} \quad (45)$$

and (41) follows at once. \square

An immediate application of the strong maximum principle ([21, 36]) yields the following corollary

Corollary 25. *In the assumptions of the above theorem, if X is a Killing vector field, then $w = \langle \nabla u, X \rangle$ is a solution of the Jacobi equation*

$$Jw = -\Delta w - f'(u)w = 0.$$

In particular, if $w \geq 0$ on M , then either $w \equiv 0$ on M or $w > 0$ on M . Therefore, if a solution $u \in C^3(M)$ of $-\Delta u = f(u)$ is weakly monotone in the direction of some Killing vector field, then either u is stable and strictly monotone in the direction of X , or u is constant on the flow lines of X .

With the aid of Corollary 25, we can prove the next results:

Lemma 26. *Let $M = N \times \mathbb{R}$ be a Riemannian product with $\text{Ric} \geq 0$.*

- (I) *If M is parabolic, then $M \notin \mathcal{F}_1$.*
- (II) *If $\text{vol}(B_R^N) = o(R^2 \log R)$, then $M \notin \mathcal{F}_2$.*

Proof. Denote the points of M with (x, t) . Choose $f(t) = t - t^3$, and

$$u(x, t) = \tanh\left(\frac{t}{\sqrt{2}}\right).$$

Then, u is a non-constant, globally Lipschitz solution of $-\Delta u = f(u)$, monotonic in the direction of the Killing field ∂_t . Thus, u is stable by Corollary 25, and (I) immediately follows. Since

$$\int_{B_R} |\nabla u|^2 dx \leq \int_{[-R, R] \times B_R^N} |\partial_t u|^2 dt dx^N \leq \|\partial_t u\|_{L^2(\mathbb{R})}^2 \text{vol}(B_R^N),$$

$M \notin \mathcal{F}_2$ provided that $\text{vol}(B_R^N) = o(R^2 \log R)$, which shows (II). □

Proposition 27. *Denote with $\mathcal{P} = \{\text{parabolic manifolds}\}$. Let m be the dimension of the family of manifolds under consideration. Then*

- (i) $\mathcal{F}_1 \subseteq \mathcal{F}_2$ for every $m \geq 2$;
- (ii) $\mathcal{F}_2 \cap \mathcal{P} \subseteq \mathcal{F}_1$ for $m = 2, 3$;
- (iii) $\mathcal{F}_1 = \mathcal{F}_2$ for $m = 2$, and $\mathcal{F}_2 \cap \mathcal{P} = \mathcal{F}_1$ for $m = 3$;
- (iv) $\mathcal{F}_1 \subsetneq \mathcal{F}_2$ for every $m \geq 3$.

Proof. (i). Suppose that $M \in \mathcal{F}_1 \setminus \mathcal{F}_2$. Then, by Remark 23 M is non-compact and thus, by Theorem 1, $M = N \times \mathbb{R}$. Since M is parabolic, by Lemma 26 we conclude that $M \notin \mathcal{F}_1$, a contradiction.

(ii). Let $M \in \mathcal{F}_2 \cap \mathcal{P}$. If by contradiction $M \notin \mathcal{F}_1$, then M is non-compact, $M = N \times \mathbb{R}$ and $\text{Ric}^N \geq 0$ again by Theorem 1. By Bishop-Gromov theorem, $\text{vol}(B_R^N) \leq CR^{m-1}$. If $m = 2, 3$, N satisfies assumption (II) of Lemma 26, and so $M \notin \mathcal{F}_2$, contradiction.

(iii). By definition, $\mathcal{F}_1 \cap \mathcal{P} = \mathcal{F}_1$. Thus, from (i) and (ii), if $m \leq 3$ it holds $\mathcal{F}_1 = \mathcal{F}_1 \cap \mathcal{P} \subseteq \mathcal{F}_2 \cap \mathcal{P} \subseteq \mathcal{F}_1$, hence $\mathcal{F}_2 \cap \mathcal{P} = \mathcal{F}_1$. On the other hand, if $m = 2$, condition $\text{Ric} \geq 0$ and Bishop-Gromov comparison theorem imply that M is parabolic, thus $\mathcal{F}_2 \cap \mathcal{P} = \mathcal{F}_2$ and so $\mathcal{F}_1 = \mathcal{F}_2$.

(iv). In view of Corollary 21, it is enough to produce a non-parabolic manifold with $\text{Ric} \geq 0$ and $\text{Ric} > 0$ somewhere. For instance, we can take a model manifold M_g , that is, \mathbb{R}^m equipped

with a radially symmetric metric ds^2 whose expression, in polar geodesic coordinates centered at some o , reads $ds^2 = dr^2 + g(r)^2 \langle \cdot, \cdot \rangle_{\mathbb{S}^{m-1}}$, $\langle \cdot, \cdot \rangle_{\mathbb{S}^{m-1}}$ being the standard metric on the unit sphere, with the choice

$$g(r) = \frac{r}{2} + \frac{1}{2} \arctan(r)$$

(see [23] or [32] for basic formulae on radially symmetric manifolds). Standard computations show that $\text{Ric} > 0$ outside o and $\text{vol}(\partial B_r) \geq Cr^{m-1}$, hence $[\text{vol}(\partial B_R)]^{-1} \in L^1(+\infty)$ for each $m \geq 3$, which is a sufficient condition for a model to be non-parabolic (see [23], Corollary 5.6.). Therefore, $M_g \notin \mathcal{F}_1$, as required. \square

Proof of Proposition 2. It follows straightforwardly from Theorem 1, Remark 23 and Lemma 26. \square

We are now ready to prove Theorem 3.

Proof of Theorem 3. Since $M \notin \mathcal{F}_2$, by Theorem 1 we have $M = N \times \mathbb{R}$, for some complete, parabolic N with $\text{Ric}^N \geq 0$ satisfying the growth estimate (4). If $N \in \mathcal{F}_2$ and it is non-compact, then by Lemma 26 it has no Euclidean factor and we are in case (i). In particular, N has only one end, for otherwise it would contain a line and would split off an Euclidean factor according to Cheeger-Gromoll splitting theorem (see [9] or [32], Theorem 68). Suppose that (i) is not satisfied, hence $N \notin \mathcal{F}_2$. Then, if $m = 3$ we have, by Theorem 1, that N is flat and $M = N \times \mathbb{R} = C \times \mathbb{R}^2$ for some curve C , thus $M = \mathbb{R}^3$ or $\mathbb{S}^1 \times \mathbb{R}^2$ with a flat metric. On the other hand, when $m \geq 4$, we have $N = \bar{N} \times \mathbb{R}$ and by (4) we deduce (5). The same analysis performed for N can now be repeated verbatim to \bar{N} in order to obtain the desired conclusion. If (ii) does not hold, then also \bar{N} splits off a line, and $\bar{N} = \hat{N} \times \mathbb{R}$. If $m = 4$, \bar{N} is a flat surface and \hat{N} is a curve, and the sole possibility to satisfy (5) is that $\hat{N} = \mathbb{S}^1$ is closed. If $m \geq 5$, again by (5), we deduce that

$$\text{vol}(B_R^{\hat{N}}) = o(\log R) \quad \text{as } R \rightarrow +\infty.$$

By the Calabi-Yau growth estimate (see [8] and [43]) a non-compact manifold with non-negative Ricci curvature has at least linear volume growth, and this forces \hat{N} to be compact, concluding the proof. \square

An extended version of a conjecture of De Giorgi

We consider an extended version (to Riemannian manifolds with $\text{Ric} \geq 0$) of a celebrated conjecture of E. De Giorgi. Let us recall that in 1978 E. De Giorgi [22] formulated the following question :

Let $u \in C^2(\mathbb{R}^m, [-1, 1])$ satisfy

$$-\Delta u = u - u^3 \quad \text{and} \quad \frac{\partial u}{\partial x_m} > 0 \quad \text{on } \mathbb{R}^m. \quad (46)$$

Is it true that all the level sets of u are hyperplanes, at least if $m \leq 8$?

The original conjecture has been proven in dimensions $m = 2, 3$ and it is still open, in its full generality, for $4 \leq m \leq 8$. We refer the reader to [17] for a recent review on the conjecture of De Giorgi and related topics.

In our setting, we replace the (Euclidean) monotonicity assumption $\partial u / \partial x_m > 0$ on \mathbb{R}^m by the natural one: u is monotone with respect to the flow lines of some Killing vector field, and we investigate the geometry of the level set of u as well as the symmetry properties of

u . This supplies a genuine framework for the study of the above conjecture on Riemannian manifolds. Our conclusion will be that the level sets of u are complete, totally geodesics submanifolds of M , which is clearly the analogous in our context of the classic version of De Giorgi's conjecture. Our results apply to Riemannian manifolds with $\text{Ric} \geq 0$. In particular, they recover and improve the results concerning the Euclidean cases of \mathbb{R}^2 and \mathbb{R}^3 and they also give a description of those manifolds supporting a De Giorgi-type conjecture.

Theorem 28. *Let $(M, \langle \cdot, \cdot \rangle)$ be a complete non-compact Riemannian manifold without boundary with $\text{Ric} \geq 0$ and let X be a Killing field on M . Suppose that $u \in C^3(M)$ is a solution of*

$$\begin{cases} -\Delta u = f(u) & \text{on } M, \\ \langle \nabla u, X \rangle > 0 & \text{on } M, \end{cases}$$

with $f \in C^1(\mathbb{R})$. If either

- (i) M is parabolic and $\nabla u \in L^\infty(M)$ or
- (ii) the function $|\nabla u|$ satisfies

$$\int_{B_R} |\nabla u|^2 dx = o(R^2 \log R) \quad \text{as } R \rightarrow +\infty,$$

then, $M = N \times \mathbb{R}$ with the product metric $\langle \cdot, \cdot \rangle = \langle \cdot, \cdot \rangle_N + dt^2$, for some complete, totally geodesic, parabolic submanifold N . In particular, $\text{Ric}^N \geq 0$ if $m \geq 3$, while, if $m = 2$, $M = \mathbb{R}^2$ or $\mathbb{S}^1 \times \mathbb{R}$ with their flat metric.

Furthermore, u depends only on t and writing $u = y(t)$ it holds

$$-y'' = f(y), \quad y' > 0.$$

Proof. Thanks to Corollary 25, u is a non-constant stable solution of the considered equation. The desired conclusion is then a consequence of Theorem 1. \square

Remark 29. We spend few words to comment on possible topological and geometric restrictions coming from the monotonicity assumption. Condition $\langle \nabla u, X \rangle > 0$ implies that both ∇u and X are nowhere vanishing, hence M is foliated by the smooth level sets of u . However, there is no a-priori Riemannian splitting. Similarly, the presence of the nowhere-vanishing Killing vector field X on M does not force, a-priori, any topological splitting of M along the flow lines of X , as the orthogonal distribution $\mathcal{D}_X : x \mapsto X(x)^\perp$ is not automatically integrable for Killing fields. Therefore, the monotonicity requirement alone does not imply, in general, severe geometric restrictions. However, one should be careful that, when \mathcal{D}_X is integrable and X is Killing, the local geometry of M then turns out to be quite rigid. Indeed, coupling the Frobenius integrability condition for \mathcal{D}_X with the skew-symmetry of ∇X coming from the Killing condition, one checks that each leaf of \mathcal{D}_X is totally geodesic. Since $|X|$ is constant along the integral lines of X , locally in a neighbourhood of a small open subset $U \subseteq N$ the metric splits as the warped product

$$\langle \cdot, \cdot \rangle = \langle \cdot, \cdot \rangle_U + h(x)^{-2} dt^2, \quad \text{where } (x, t) \in U \times \mathbb{R},$$

for some smooth $h(x) = |X|(x)^{-1}$. In particular, the Ricci curvature in the direction of $X = \partial_t$ satisfies

$$\text{Ric}(\partial_t, \partial_t) = -\frac{\Delta h(x)}{h(x)}. \quad (47)$$

Further restrictions then come out when one adds the requirement $\text{Ric} \geq 0$. In this case, by (47) $h(x)$ turns out to be a positive, superharmonic function on U . Consequently, if in a (global) warped product $N \times \mathbb{R}$, with metric $\langle \cdot, \cdot \rangle_N + h^{-2} dt^2$, the factor N is parabolic, then condition $\text{Ric} \geq 0$ forces $N \times \mathbb{R}$ to be a Riemannian product, h being constant by the parabolicity assumption. The dimensional case $m = 2$ is particularly rigid. In fact, if M is a complete surface with non-negative Gaussian curvature and possessing a nowhere vanishing Killing vector field X , then M is flat. Indeed, in this case \mathcal{D}_X is clearly integrable, and the integral curves of the local unit vector field E orthogonal to X are geodesics. For $x \in M$, let $\sigma : \mathbb{R} \rightarrow M$ be a unit speed geodesic with tangent vector everywhere orthogonal to X . The sectional curvature along $\sigma(t)$ is

$$0 \leq K(\sigma' \wedge X) = \frac{R(\sigma', X, \sigma', X)(t)}{|X|^2(\sigma(t))} = -h(t)h''(t),$$

so h is a non-negative, concave function on \mathbb{R} , hence h is constant. Therefore, $K = 0$ along σ , and in particular at x , as claimed. Note that the completeness assumption on M is essential, as the example of the punctured paraboloid $M = \{(x, y, z) \in \mathbb{R}^3 : z = x^2 + y^2, z > 0\}$ shows.

Theorem 28 has some interesting consequences. For instance, in the 2-dimensional case we have the following strengthened version:

Corollary 30. *Let $(M, \langle \cdot, \cdot \rangle)$ be a complete non-compact surface without boundary, with Gaussian curvature $K \geq 0$ and let X be a Killing field on M . Suppose that $u \in C^3(M)$ is a solution of*

$$\begin{cases} -\Delta u = f(u) & \text{on } M \\ \langle \nabla u, X \rangle > 0 & \text{on } M \\ \nabla u \in L^\infty(M) \end{cases}$$

with $f \in C^1(\mathbb{R})$.

Then, M is the Riemannian product \mathbb{R}^2 or $\mathbb{S}^1 \times \mathbb{R}$, with flat metric, u depends only on t and, writing $u = y(t)$, it holds

$$y'' = -f(y), \quad y' > 0.$$

Proof. Since $K \geq 0$ and $\dim(M) = 2$, by Bishop-Gromov comparison theorem $\text{vol}(B_R) \leq \pi R^2$, so M is parabolic by Theorem 7.3 in [23]. Therefore, both (i) and (ii) of Theorem 28 are satisfied. This proves the corollary. \square

Some remarks are in order.

Remark 31. (i) The previous result establishes De Giorgi's conjecture for surfaces with non-negative Gaussian curvature. Actually it yields more, indeed, if $(M, \langle \cdot, \cdot \rangle)$ is a complete non-compact manifold without boundary, with $\text{Ric} \geq 0$ and of dimension $m \geq 2$, it is known that any bounded solution of $-\Delta u = f(u)$ also has bounded gradient (see e.g. Appendix 1). Note also that the converse is not true, since $u(x) = x_1$, is an unbounded monotone harmonic function on $(\mathbb{R}^m, \langle \cdot, \cdot \rangle_{\text{can}})$ whose gradient is bounded (here, and in the sequel, $\langle \cdot, \cdot \rangle_{\text{can}}$ denotes the canonical flat metric on \mathbb{R}^m).

(ii) We recover the case of \mathbb{R}^2 , with its canonical flat metric. Apply Corollary 30 to $(\mathbb{R}^2, \langle \cdot, \cdot \rangle_{\text{can}})$ and $X = \partial/\partial x_2$.

(iii) From Theorem 28 we also recover the case of $(\mathbb{R}^3, \langle \cdot, \cdot \rangle_{\text{can}})$. Indeed, any bounded monotone solution of $-\Delta u = f(u)$ in \mathbb{R}^3 satisfies

$$\int_{B_R} |\nabla u|^2 dx = O(R^2) \quad \text{as } R \rightarrow +\infty$$

(see [15, 1]). Hence, the conclusion follows by applying Theorem 28 with $X = \partial/\partial x_3$.

(iv) By Remark 29, the flatness of M is automatic in Corollary 30 from the sole assumptions $K \geq 0$ and X Killing and nowhere vanishing.

(v) If $m \geq 2$ and $M^m = N \times \mathbb{R}$ with the product metric $\langle \cdot, \cdot \rangle = \langle \cdot, \cdot \rangle_N + dt^2$, then it is always possible to construct a solution of (14) which is monotone in the direction of the Killing vector field ∂_t (proceed as in the proof of Lemma 26). Our main Theorem 28 states that the converse holds true if the manifold M has non-negative Ricci curvature and it supports a De Giorgi-type conjecture.

Overdetermined boundary value problems

In this section we study the case of overdetermined elliptic problems on open and connected sets with C^3 boundary. In the situation considered here, the boundary term in (29) may cause extra difficulties. Surprisingly, for solutions monotone in the direction of some Killing vector field, the boundary term indeed can be ruled out, as the next lemma reveals:

Lemma 32. *Let u be such that u and $\partial_\nu u$ are constant on $\partial\Omega$ and $\partial_\nu u \neq 0$ on $\partial\Omega$. Suppose that w is of the form $w = \langle \nabla u, X \rangle$ in a neighbourhood of $\partial\Omega$, for some vector field X . Then*

$$w \partial_\nu \left(\frac{|\nabla u|^2}{2} \right) - |\nabla u|^2 \partial_\nu w = -|\nabla u|^3 \langle \nu, \nabla X(\nu) \rangle \quad \text{on } \partial\Omega. \quad (48)$$

In particular, if X satisfies $\langle \nabla X(\nu), \nu \rangle \geq 0$ on $\partial\Omega$, the boundary terms in (23) and (29) are non-positive.

Proof. Let us define the constant $c = \partial_\nu u$ on $\partial\Omega$. Since u is constant, $\nabla u = (\partial_\nu u)\nu = c\nu$, so $|\nabla u|^2 = c^2$ is constant on $\partial\Omega$. Therefore, its gradient has only normal component:

$$\partial_\nu (|\nabla u|^2)\nu = \nabla (|\nabla u|^2) = 2\nabla du(\nabla u, \cdot)^\sharp.$$

It follows that, in our assumptions,

$$\begin{aligned} |\nabla u|^2 \partial_\nu w &= c^2 \partial_{\nabla u/c} w = c \nabla u \langle \nabla u, X \rangle \\ &= c [\langle \nabla_{\nabla u} \nabla u, X \rangle + \langle \nabla u, \nabla_{\nabla u} X \rangle] = c [\nabla du(\nabla u, X) + c^2 \langle \nu, \nabla X(\nu) \rangle] \\ &= c \nabla du(\nabla u, X) + c^3 \langle \nu, \nabla X(\nu) \rangle = \frac{c}{2} \langle \nabla |\nabla u|^2, X \rangle + |\nabla u|^3 \langle \nu, \nabla X(\nu) \rangle \\ &= \frac{c}{2} \partial_\nu (|\nabla u|^2) \langle \nu, X \rangle + |\nabla u|^3 \langle \nu, \nabla X(\nu) \rangle = \frac{\partial_\nu (|\nabla u|^2)}{2} \langle \nabla u, X \rangle + |\nabla u|^3 \langle \nu, \nabla X(\nu) \rangle \\ &= \frac{\partial_\nu (|\nabla u|^2)}{2} w + |\nabla u|^3 \langle \nu, \nabla X(\nu) \rangle, \end{aligned}$$

as claimed. □

Remark 33. Clearly, any Killing vector field fulfills the requirement $\langle \nabla X(\nu), \nu \rangle \geq 0$, but the class is much more general. For instance, $\langle \nabla X(\nu), \nu \rangle \geq 0$ is met whenever X solves

$$L_X \langle \cdot, \cdot \rangle \geq 0 \quad \text{as a quadratic form.}$$

Examples of such X also include positively conformal vector fields, that is, fields satisfying $L_X \langle \cdot, \cdot \rangle = \eta \langle \cdot, \cdot \rangle$ for a non-negative $\eta \in C^\infty(M)$, and gradients of convex functions $X = \nabla \psi$, being $L_{\nabla \psi} \langle \cdot, \cdot \rangle = 2\nabla d\psi$.

The above Lemma is the key to prove Theorem 5.

Proof of Theorem 5. In our assumptions, by Lemma 32 either $w = \langle \nabla u, X \rangle$ or $w = -\langle \nabla u, X \rangle$ is a positive solution on $\Delta w + f'(u)w = 0$ on Ω . Up to changing the sign of X , we can suppose that $w = \langle \nabla u, X \rangle > 0$ on Ω . In particular, X is nowhere vanishing on Ω . We are going to show that condition (28) is satisfied, namely, that

$$\liminf_{\varepsilon \rightarrow 0^+} \int_{\Omega} \phi^2 \langle \nabla |\nabla u|^2, \nabla \left(\frac{w}{w + \varepsilon} \right) \rangle dx \geq 0. \quad (49)$$

By a density argument, this will be accomplished once we prove that

$$\liminf_{\varepsilon \rightarrow 0^+} \int_K \langle \nabla |\nabla u|^2, \nabla \left(\frac{w}{w + \varepsilon} \right) \rangle dx \geq 0 \quad \forall K \Subset \bar{\Omega}. \quad (50)$$

We first claim that there exists a constant $C = C(K, m, \|u\|_{C^3(K)}) > 0$ such that

$$|\langle \nabla |\nabla u|^2, \nabla w \rangle| \leq C|w| \quad \text{on } K \cap \partial\Omega. \quad (51)$$

First we observe that, since u is constant on $\partial\Omega$, ∇u has only normal component, thus

$$|w| = |\langle \nabla u, X \rangle| = |\nabla u| |\langle \nu, X \rangle|. \quad (52)$$

From the further property that $|\nabla u|^2 = c^2$ is constant along $\partial\Omega$, we deduce that $\nabla |\nabla u|^2$ is parallel to ν and therefore, by Kato inequality,

$$|\langle \nabla |\nabla u|^2, \nabla w \rangle| = |\nabla |\nabla u|^2| |\partial_\nu w| \leq 2|\nabla u| |\nabla du| |\partial_\nu w| \quad (53)$$

on $\partial\Omega$. Using the fact that X is a Killing vector field and $\nabla u = c\nu$ on $\partial\Omega$, the following chain of equalities is true:

$$\begin{aligned} \partial_\nu w &= \nu(\langle \nabla u, X \rangle) = \nabla du(\nu, X) + \langle \nabla u, \nabla_\nu X \rangle \\ &= \nabla du(\nu, X) + c\langle \nu, \nabla_\nu X \rangle = \nabla du(\nu, X). \end{aligned} \quad (54)$$

Now, we use that $\partial_\nu u$ is constant on $\partial\Omega$, whence $\nabla(\partial_\nu u)$ is also parallel to ν :

$$\pm |\nabla(\partial_\nu u)|\nu = \nabla(\partial_\nu u) = \nabla(\langle \nabla u, \nu \rangle) = \nabla du(\nu, \cdot)^\sharp + \langle \nabla u, \nabla \nu \rangle. \quad (55)$$

on $\partial\Omega$, where $\nabla \nu$ is the $(1, 1)$ version of the second fundamental form of $\partial\Omega$, that is, the (opposite of the) Weingarten transformation. Taking the inner product with X we deduce that, on $\partial\Omega$,

$$\begin{aligned} \pm |\nabla(\partial_\nu u)|\langle \nu, X \rangle &= \nabla du(\nu, X) + \langle \nabla u, \nabla_X \nu \rangle = \nabla du(\nu, X) + c\langle \nu, \nabla_X \nu \rangle \\ &= \nabla du(\nu, X) + \frac{c}{2} X(|\nu|^2) = \nabla du(\nu, X), \end{aligned} \quad (56)$$

whence combining with (54) we conclude

$$|\partial_\nu w| = |\nabla du(\nu, X)| = |\nabla(\partial_\nu u)| |\langle \nu, X \rangle|. \quad (57)$$

Inserting the equalities (52) and (57) into (53) we deduce

$$\begin{aligned} |\langle \nabla |\nabla u|^2, \nabla w \rangle| &\leq 2|\nabla u| |\nabla du| |\partial_\nu w| \leq 2|\nabla u| |\nabla du| |\nabla(\partial_\nu u)| |\langle \nu, X \rangle| \\ &= 2|\nabla du| |\nabla(\partial_\nu u)| |w|. \end{aligned}$$

Since $u \in C^3(\overline{\Omega})$, the terms ∇du and $\nabla(\partial_\nu u)$ are bounded on K , and the claimed inequality (51) is proved.

Our next task is to extend the bound in (51) to a whole neighbourhood of $\partial\Omega$. More precisely, we claim that there exist $C > 0$, possibly depending on K , f , u and $\partial\Omega$, such that

$$|\langle \nabla |\nabla u|^2, \nabla w \rangle| \leq C|w| \quad \forall x \in \Omega \cap K. \quad (58)$$

To prove this, we notice that it is enough to prove the bound in a neighbourhood of $K \cap \partial\Omega$. By the compactness of $K \cap \partial\Omega$, it is enough to work locally around any $x_0 \in \partial\Omega$. Towards this aim we note that, since $\partial\Omega$ is C^3 , for any $x_0 \in \partial\Omega$ Fermi coordinates (T, Ψ) can be defined in a collar $T \Subset \Omega$ of x_0 :

$$\Psi : T \longrightarrow [0, \delta) \times U \subseteq \mathbb{R}_0^+ \times \partial\Omega, \quad \Psi(x) = (t, \pi(x)),$$

where U is open in $\partial\Omega$ and contains x_0 . In particular, $\pi(x) \in \partial\Omega$ is the unique point of $\partial\Omega$ realizing $\text{dist}(x, \partial\Omega)$, and the smooth coordinate $t \in [0, \delta)$ satisfies

$$t(x) = \text{dist}(x, \partial\Omega) = \text{dist}(x, \pi(x)).$$

Again since $\partial\Omega$ is smooth enough, up to shrinking further T there exists a bounded domain $D_0 \Subset \Omega$, of class C^3 and containing T , that satisfies

$$t(x) = \text{dist}(x, \partial\Omega) = \text{dist}(x, \partial D_0) \quad \forall x \in T.$$

In the chart Ψ , the function $w \in C^2(\overline{T})$ satisfies a linear elliptic equation, the expression in chart of $\Delta w + f'(u)w = 0$, to which the Hopf-type Lemma 1 of [42] can be applied to deduce

$$w(x) \geq C \text{dist}(x, \partial D_0) = Ct(x) \quad \forall x \in T, \quad (59)$$

for some $C > 0$. Next, since $u \in C^3(\overline{T})$, the function

$$g(x) = |\langle \nabla |\nabla u|^2, \nabla w \rangle| \in \text{Lip}(\overline{T}),$$

whence

$$|g(\pi(x)) - g(x)| \leq C \text{dist}(\pi(x), x) = Ct(x) \quad (60)$$

All in all, combining (59) and (60), and using also (51) we obtain:

$$\frac{g(x)}{|w(x)|} \leq \frac{|g(x) - g(\pi(x))| + |g(\pi(x))|}{|w(x)|} \leq \frac{|g(x) - g(\pi(x))| + C|w(x)|}{|w(x)|} \leq C, \quad (61)$$

for a suitable $C > 0$. This completes the proof of (58). Now we observe that the integrand in (50) may be written as

$$\frac{\varepsilon}{(\varepsilon + w)^2} \langle \nabla |\nabla u|^2, \nabla w \rangle =: \psi_\varepsilon.$$

Notice that ψ_ε is well-defined in Ω since $w > 0$, and

$$\lim_{\varepsilon \rightarrow 0^+} \psi_\varepsilon(x) = 0,$$

for each $x \in \Omega$. Moreover, by (58), on \bar{T}

$$|\psi_\varepsilon(x)| \leq \frac{(\varepsilon + w)}{(\varepsilon + w)w} |\langle \nabla |\nabla u|^2, \nabla w \rangle| \leq C.$$

Then, (50) with $K = T$ follows from Lebesgue convergence theorem.

Applying Proposition 17 with the aid of Lemma 32, the boundary term in (29) vanishes since X is Killing, and we get

$$\begin{aligned} & \int_{\Omega} \left[|\nabla du|^2 + \text{Ric}(\nabla u, \nabla u) - |\nabla |\nabla u|^2| \right] \phi^2 dx + \liminf_{\varepsilon \rightarrow 0^+} \int_{\Omega} (w + \varepsilon)^2 \left| \nabla \left(\frac{\phi |\nabla u|}{w + \varepsilon} \right) \right|^2 dx \\ & \leq \int_{\Omega} |\nabla \phi|^2 |\nabla u|^2 dx. \end{aligned} \quad (62)$$

Hereafter, we can proceed in a way analogous to that in Theorem 1. In particular, the use of appropriate cutoff functions $\{\phi_\alpha\}$ satisfying (33), and the assumption $\text{Ric} \geq 0$, imply

$$|\nabla du|^2 = |\nabla |\nabla u|^2|, \quad \text{Ric}(\nabla u, \nabla u) = 0 \quad \text{on } \Omega, \quad (63)$$

thus inserting into (62) we obtain

$$\liminf_{\varepsilon \rightarrow 0^+} \int_{\Omega} (w + \varepsilon)^2 \left| \nabla \left(\frac{\phi |\nabla u|}{w + \varepsilon} \right) \right|^2 dx \leq \int_{\Omega} |\nabla \phi|^2 |\nabla u|^2 dx. \quad (64)$$

For every small $\delta > 0$, we define $\Omega_\delta = \{x \in \Omega : \text{dist}(x, \partial\Omega) > \delta\}$. By the positivity of the integrand, and since away from $\partial\Omega$ the function w is locally uniformly bounded away from zero,

$$\begin{aligned} \liminf_{\varepsilon \rightarrow 0^+} \int_{\Omega} (w + \varepsilon)^2 \left| \nabla \left(\frac{\phi |\nabla u|}{w + \varepsilon} \right) \right|^2 dx & \geq \lim_{\varepsilon \rightarrow 0^+} \int_{\Omega_\delta} (w + \varepsilon)^2 \left| \nabla \left(\frac{\phi |\nabla u|}{w + \varepsilon} \right) \right|^2 dx \\ & = \int_{\Omega_\delta} w^2 \left| \nabla \left(\frac{\phi |\nabla u|}{w} \right) \right|^2 dx. \end{aligned}$$

Letting $\delta \rightarrow 0$ we thus get

$$\liminf_{\varepsilon \rightarrow 0^+} \int_{\Omega} (w + \varepsilon)^2 \left| \nabla \left(\frac{\phi |\nabla u|}{w + \varepsilon} \right) \right|^2 dx \geq \int_{\Omega} w^2 \left| \nabla \left(\frac{\phi |\nabla u|}{w} \right) \right|^2 dx.$$

In particular, by (64) the RHS of the above inequality is finite and

$$\int_{\Omega} w^2 \left| \nabla \left(\frac{\phi |\nabla u|}{w} \right) \right|^2 dx \leq \int_{\Omega} |\nabla \phi|^2 |\nabla u|^2 dx.$$

An application of Young type inequality (31) transforms the above inequality into

$$(1 - \delta) \int_{\Omega} w^2 \left| \nabla \left(\frac{|\nabla u|}{w} \right) \right|^2 dx \leq \frac{1}{\delta} \int_{\Omega} |\nabla \phi|^2 |\nabla u|^2 dx,$$

for each $\delta \in (0, 1)$. Consequently, choosing again the appropriate cut-offs $\{\phi_\alpha\}$ satisfying (33) as in Theorem 1, we also get $|\nabla u| = cw$ for some constant $c \geq 0$. Since u is non-constant, $c > 0$. The topological part of the splitting needs some extra care. We shall divide into two cases, according to the sign of the constant $\partial_\nu u$ on $\partial\Omega$. Since the discussions are specular, we just consider the case when $\partial_\nu u$ is positive on $\partial\Omega$. Denote with $N \subseteq \Omega$ any level set of u , and with Φ_t the flow of $\nu = \nabla u / |\nabla u|$ on Ω . Observe that, for $x \in \partial\Omega$, the fact that $u \circ \Phi_t$ is strictly increasing implies that $\Phi_t(x) \in \Omega$ for each $t \in \mathbb{R}^+$. From the Sternberg-Zumbrun identity in Proposition 18, N is totally geodesic, $|\nabla u|$ is constant (and non-zero) on N and the only non-vanishing component of ∇du is that corresponding to the pair (ν, ν) . Therefore, integral curves of ν are geodesics. Write $|\nabla u| = \beta(u)$, for some continuous β . We claim that, for each $x \in \Omega$, $\Phi_t(x)$ touches $\partial\Omega$ at a finite, negative time $t_0(x)$. Indeed, consider the rescaled flow Ψ_s of the vector field $Y = \nabla u / |\nabla u|^2$. Clearly, $\Phi(t, x) = \Psi(s(t), x)$, where

$$s(t) = s(0) + \int_0^t |\nabla u|(\Phi_\tau(x)) d\tau = s(0) + \int_0^t \beta(u \circ \Phi_\tau(x)) d\tau$$

is a locally Lipschitz bijection with inverse $t(s)$. From $u(\Psi_s(x)) = u(x) + s$ and from $\partial_\nu u > 0$, we deduce that $\Psi_s(x)$ touches $\partial\Omega$ at a finite, negative value $s_0(x)$. Now, since $\Phi_t(x)$ is a geodesic, and geodesics are divergent as $t \rightarrow -\infty$, then necessarily the correspondent value $t_0(x) = t(s_0(x))$ is finite. Consequently, the flow of ν starting from $\partial\Omega$ covers the whole Ω . Having fixed a connected component Σ of $\partial\Omega$, proceeding as in the proof of Theorem 1 it can be shown that $\Phi : \Sigma \times \mathbb{R}^+ \rightarrow \Omega$ is a C^3 diffeomorphism. Thus, $\Sigma \equiv \partial\Omega$ and we have the desired topological splitting. The proof that each Φ_t is an isometry is identical to the boundaryless case. It thus follows, via a simple approximation, that $\partial\Omega$ is totally geodesic and isometric to any other level set of Ω , and thus Ω splits as a Riemannian product $\partial\Omega \times \mathbb{R}^+$. Setting $u(x, t) = y(t)$, y solves

$$y'(t) = |\nabla u|(x, t) > 0, \quad y''(t) = -f(y(t)).$$

As regards the volume estimate for $\partial\Omega$, it follows exactly along the same lines as those yielding (2):

$$\left(\int_0^R |y'(t)|^2 dt \right) \text{vol}(B_R^{\partial\Omega}) \leq \int_{(0,R) \times B_R^{\partial\Omega}} |y'(t)|^2 dt dx^{\partial\Omega} \leq \int_{B_{R\sqrt{2}} \cap \Omega} |\nabla u|^2 dx = o(R^2 \log R)$$

as $R \rightarrow +\infty$, according to (ii). Lastly, we address the mutual position of X and $\partial_t = \nu$. From the identity $|\nabla u| = cw = c\langle \nabla u, X \rangle$ we deduce that

$$\langle \partial_t, X \rangle = \frac{1}{|\nabla u|} \langle \nabla u, X \rangle = \frac{1}{c}$$

is constant on M . Consequently, the projected vector field

$$X^\perp = X - \langle X, \partial_t \rangle \partial_t$$

is still a Killing field, since so are X and ∂_t . This concludes the proof. The case $\partial_\nu u < 0$ on $\partial\Omega$ can be dealt with analogously, by considering the flow of $\nu = -\nabla u / |\nabla u|$. \square

Clearly, in the above theorem a key role is played by the monotonicity condition $\langle \nabla u, X \rangle > 0$, for some Killing vector field X . As remarked in the Introduction, this condition is automatically satisfied for globally Lipschitz epigraphs $\Omega \subseteq \mathbb{R}^m$, and for $f \in \text{Lip}(\mathbb{R})$ satisfying some mild assumptions, thanks to the following remarkable result by H. Berestycki, L. Caffarelli and L. Nirenberg in [4]:

Theorem 34 ([4], Theorem 1.1). *Let $\Omega \subseteq \mathbb{R}^m$ be an open subset that can be written as the epigraph of a globally Lipschitz function φ on \mathbb{R}^{m-1} , that is,*

$$\Omega = \{(x', x_m) \in \mathbb{R}^m = \mathbb{R}^{m-1} \times \mathbb{R} : x_m > \varphi(x')\}.$$

Let $f \in \text{Lip}(\mathbb{R})$ satisfy the requirements

$$\begin{cases} f > 0 & \text{on } (0, \lambda), & f \leq 0 & \text{on } (\lambda, +\infty), \\ f(s) \geq \delta_0 s & \text{for } s \in (0, s_0), \\ f & \text{is non-increasing on } [\lambda - s_0, \lambda], \end{cases}$$

for some positive λ, δ_0, s_0 . Let $u \in C^2(\Omega) \cap C^0(\overline{\Omega})$ be a bounded, positive solution of

$$\begin{cases} -\Delta u = f(u) & \text{on } \Omega, \\ u > 0 & \text{on } \Omega, \quad u = 0 & \text{on } \partial\Omega. \end{cases} \quad (65)$$

Then, u is monotone in the x_m -direction, that is, $\partial u / \partial x_m > 0$ on Ω .

The proof of this result relies on some techniques which are tightly related to the peculiarities of Euclidean space as a Riemannian manifold. It would be therefore very interesting to investigate the following

problem: determine reasonable assumptions on the manifold $(M, \langle \cdot, \cdot \rangle)$ and on Ω, f which ensure that every bounded, sufficiently smooth solution u of (65), or at least of (6), is monotone in the direction of a Killing vector field X .

In the next section, we prove some preliminary results addressed to the above problem. In doing so, we obtain an improvement of Theorem 5 in the dimensional case $m = 3$.

Further qualitative properties of solutions, and the monotonicity condition

This last section is devoted to move some first steps towards a proof of the monotonicity condition in a manifold setting. In doing so, we extend results in [4], [5] to Riemannian manifolds satisfying $\text{Ric} \geq -(m-1)H^2 \langle \cdot, \cdot \rangle$, for some $H \geq 0$. Although the proofs below are in the same spirit as those in [4] and [5], in order to deal with the lack of symmetry of a general M we shall introduce some different arguments that may have independent interest. In particular, we mention Proposition 7 for its generality. Combining the results of this section will lead us to a proof of Theorem 10. Hereafter, we shall restrict ourselves to a class of nonlinearities f satisfying the following general assumptions:

$$\begin{cases} f > 0 & \text{on } (0, \lambda), & f(\lambda) = 0, & f < 0 & \text{on } (\lambda, \lambda + s_0), \\ f(s) \geq \delta_0 s & \text{for } s \in (0, s_0), \end{cases}$$

for some $\lambda > 0$ and some small $\delta_0, s_0 > 0$. Let $\Omega \subseteq M$ be an open, connected subset with possibly noncompact closure, and let $u \in C^2(\Omega) \cap C^0(\overline{\Omega})$, $u > 0$ on Ω solve

$$-\Delta u = f(u) \quad \text{on } \Omega.$$

For $R_0 > 0$, set

$$\Omega_{R_0} = \{x \in \Omega : \text{dist}(x, \partial\Omega) > R_0\}, \quad \Omega^{R_0} = \{x \in \Omega : \text{dist}(x, \partial\Omega) < R_0\}.$$

Moreover, for notational convenience, for $y \in M$ define $r_y(x) = \text{dist}(y, x)$.

Remark 35. We observe that, even when Ω is connected, Ω_{R_0} may have infinitely many connected components. By a compactness argument, however, such a number is always finite if Ω is relatively compact.

The first lemma ensures that, for suitable f , u is bounded from below by some positive constant on each connected component of Ω_{R_0} . The strategy of the proof is somehow close to the spirit of the sliding method, although this latter cannot be applied due to the lack of a group of isometries acting transitively on M .

Lemma 36. *Let $(M^m, \langle \cdot, \cdot \rangle)$ be a complete Riemannian manifold such that $\text{Ric} \geq -(m-1)H^2 \langle \cdot, \cdot \rangle$, for some $H \geq 0$. Suppose that $f \in C^1(\mathbb{R})$ satisfies*

$$f(s) \geq \left(\delta_0 + \frac{(m-1)^2 H^2}{4} \right) s \quad \text{for } s \in (0, s_0), \quad (66)$$

for some positive, small δ_0, s_0 . Let $u \in C^2(\Omega)$ be a positive solution of $-\Delta u = f(u)$ on Ω . Then, there exists $R_0 = R_0(m, H, \delta_0) > 0$ such that the following holds: for each connected component V_j of Ω_{R_0} , there exists $\varepsilon_j = \varepsilon_j(\delta_0, H, m, V_j) > 0$ such that

$$u(x) \geq \varepsilon_j \quad \text{if } x \in V_j. \quad (67)$$

In particular, if Ω_{R_0} has only finitely many connected components,

$$\inf_{\Omega_{R_0}} u > 0$$

Proof. Let M_H be a space form of constant sectional curvature $-H^2 \leq 0$ and dimension m . In other words, $M_H = \mathbb{R}^m$ for $H = 0$, and M_H is the hyperbolic space of curvature $-H^2$ if $H > 0$. Let $o \in M_H$. For $R > 0$, denote with $\lambda_1(\mathbb{B}_R)$ the first Dirichlet eigenvalue of $-\Delta$ on the geodesic ball $\mathbb{B}_R = B_R(o) \subseteq M_H$. By a standard result (combine for instance [29] and [7]), the bottom of the spectrum of $-\Delta$ on M_H , $\lambda_1(M_H)$, is given by

$$\lambda_1(M_H) = \lim_{R \rightarrow +\infty} \lambda_1(\mathbb{B}_R) = \frac{(m-1)^2 H^2}{4}.$$

Therefore, by (66) we can choose $R_0 = R_0(\delta_0, H, m)$ such that, for $R \geq R_0/2$,

$$\lambda_1(\mathbb{B}_R) s < f(s) \quad \text{for } s \in (0, s_0]. \quad (68)$$

Let $\text{sn}_H(r)$ be a solution of

$$\begin{cases} \text{sn}_H''(r) - H^2 \text{sn}_H(r) = 0 & \text{on } \mathbb{R}^+ \\ \text{sn}_H(0) = 0, \quad \text{sn}_H'(0) = 1 \end{cases}$$

and set $\text{cn}_H(r) = \text{sn}_H'(r)$. Moreover, let z be a first eigenfunction of \mathbb{B}_R . Then, via a symmetrization argument and since the space of first eigenfunctions has dimension 1, z is radial and (up to normalization) solves

$$\begin{cases} z''(r) + (m-1) \frac{\text{cn}_H(r)}{\text{sn}_H(r)} z'(r) + \lambda_1(\mathbb{B}_R) z(r) = 0 & \text{on } (0, R), \\ z(0) = 1, \quad z'(0) = 0, \quad z(R) = 0, \quad z(r) > 0 & \text{on } [0, R). \end{cases}$$

A first integration shows that $z' < 0$ on $(0, R)$, so $z \leq 1$. From assumption $\text{Ric} \geq -(m-1)H^2 \langle \cdot, \cdot \rangle$ and the Laplacian comparison theorem (see for instance [32], Ch. 9 or [34], Section 2), we deduce that

$$\Delta r_y(x) \leq \frac{\text{cn}_H}{\text{sn}_H}(r_y(x))$$

pointwise outside the cut-locus of y and weakly on the whole M . Therefore, for every $y \in \Omega_R$, the function $\varphi_y : M \rightarrow \mathbb{R}$ defined as

$$\varphi_y(x) = \begin{cases} z(r_y(x)) & \text{if } x \in B_R(y) \\ 0 & \text{otherwise} \end{cases}$$

is a Lipschitz, weak solution of

$$\begin{cases} \Delta\varphi_y + \lambda_1(\mathbb{B}_R)\varphi_y \geq 0 & \text{on } B_R(y) \\ 0 < \varphi_y \leq 1 \text{ on } B_R(y), \quad \varphi_y = 0 \text{ on } \partial B_R(y), \quad \varphi_y(y) = 1. \end{cases} \quad (69)$$

Fix any $R \in (R_0/2, R_0)$. Note that, in this way, for each $y \in \Omega_{R_0}$ it holds $B_R(y) \Subset \Omega$, and (68) is met. Let $\{V_j\}$ be the connected components of Ω_{R_0} (possibly, countably many). For each j , choose $y_j \in V_j$ and $\varepsilon_j \in (0, s_0)$ sufficiently small that

$$\varepsilon_j \varphi_{y_j}(x) < u(x) \quad \text{for every } x \in B_R(y_j) \Subset \Omega.$$

This is possible since $u > 0$ on $\overline{B_R(y_j)}$. Let $y \in V_j$. From $\varphi_y \leq 1$, $\varepsilon_j \varphi_y \leq s_0$, thus

$$f(\varepsilon_j \varphi_y) > \lambda_1(\mathbb{B}_R) \varepsilon_j \varphi_y \quad \text{on } B_R(y). \quad (70)$$

We are going to show that, for each $y \in V_j$, $u(y) \geq \varepsilon_j$. Towards this aim, let $\gamma : [0, l] \rightarrow V_j$ be a unit speed curve joining y_j and y , in such a way that $\gamma(0) = y_j$. Define

$$w_t(x) = u(x) - \varepsilon_j \varphi_{y(t)}(x) = u(x) - \varepsilon_j z(\text{dist}(y(t), x)).$$

Then, the curve $w : t \in [0, l] \rightarrow w_t \in C^0(\overline{\Omega})$, where $C^0(\overline{\Omega})$ is endowed with the topology of uniform convergence on compacta, is continuous. Indeed, by the triangle inequality and since γ has speed 1 we have

$$\begin{aligned} \|w(t) - w(s)\|_{L^\infty(\Omega)} &= \varepsilon_j \|z(\text{dist}(\cdot, y(t))) - z(\text{dist}(\cdot, y(s)))\|_{L^\infty(\Omega)} \\ &\leq s_0 \text{Lip}(z) \|\text{dist}(\cdot, y(t)) - \text{dist}(\cdot, y(s))\|_{L^\infty(\Omega)} \\ &\leq s_0 \text{Lip}(z) \text{dist}(y(t), y(s)) \leq \varepsilon_0 \text{Lip}(z) |t - s|. \end{aligned}$$

It follows that the set

$$T = \left\{ t \in [0, l] : i(t) = \inf_{B_R(y(t))} w_t > 0 \right\}$$

is open on $[0, l]$ and contains $t = 0$. We stress that, for each $t \in [0, l]$, by construction $B_R(y(t)) \Subset \Omega$. We claim that $T = [0, l]$. If not, there is a first point $a \leq l$ such that $i(t) > 0$ for $t \in [0, a)$ and $i(a) = 0$. Since $w_a > 0$ on $\partial B_R(y(a))$ by construction, the minimum of w_a is attained on some $x_0 \in B_R(y(a))$. Now, by (69) and (70), w_a solves weakly

$$\Delta w_a = -f(u) + \varepsilon_j \lambda_1(\mathbb{B}_R) \varphi_{y(a)} < -f(u) + f(\varepsilon_j \varphi_{y(a)}) = c(x) w_a,$$

where as usual

$$c(x) = \frac{f(\varepsilon_j \varphi_{y(a)}(x)) - f(u(x))}{u(x) - \varepsilon_j \varphi_{y(a)}(x)}.$$

Now, from $w_a \geq 0$ and $w_a(x_0) = 0$, by the local Harnack inequality for Lipschitz weak solutions (see [21]), $w_a \equiv 0$, contradicting the fact that $w_a > 0$ on $\partial B_R(y(a))$. This proves the claim.

Now, from $w_l > 0$ on $\overline{B_R(y)}$, in particular

$$0 < w_l(y) = u(y) - \varepsilon_j \varphi_y(y) = u(y) - \varepsilon_j,$$

proving the desired (67). \square

In the second Lemma, we specify the asymptotic profile of the solution as $\text{dist}(x, \partial\Omega) \rightarrow +\infty$. First, we shall need some notation. Let $R > 0$, and consider the (radial) solution v_R of

$$\begin{cases} \Delta v_R = -1 & \text{on } \mathbb{B}_R \subseteq M_H, \\ v_R = 0 & \text{on } \partial\mathbb{B}_R, \end{cases}$$

where $\mathbb{B}_R = B_R(o) \subseteq M_H$ as in the previous lemma. Since v_R is radial, integrating the correspondent ODE we see that

$$v_R(r) = \int_r^R \frac{1}{\text{sn}_H(t)^{m-1}} \left[\int_0^t \text{sn}_H(s)^{m-1} ds \right] dt.$$

Denote with $C_H(R) = \|v_R\|_{L^\infty([0,R])} = v_R(0)$, and observe that

$$C_H(R) \downarrow 0^+ \quad \text{as } R \rightarrow 0^+, \quad C_H(R) \uparrow +\infty \quad \text{as } R \uparrow +\infty. \quad (71)$$

Let $R_0, \{V_j\}$ and ε_j be as in the previous lemma, and for $y \in V_j$ set

$$\delta_j(y) = \min \{f(s) : s \in [\varepsilon_j, u(y)]\}. \quad (72)$$

Lemma 37. *In the assumptions of the previous lemma, suppose further that u is bounded above, and that*

$$f > 0 \quad \text{on } (0, \|u\|_{L^\infty}). \quad (73)$$

Then, for every $y \in V_j \subseteq \Omega_{R_0}$,

$$\delta_j(y)C_H([\text{dist}(y, \partial\Omega) - R_0]) \leq \|u\|_{L^\infty}. \quad (74)$$

Proof. Under assumption (73), $\delta_j(y) \geq 0$ for each $y \in V_j$ and each j . Suppose by contradiction that there exists $y \in V_j$ such that

$$\delta_j(y)C_H([\text{dist}(y, \partial\Omega) - R_0]) > \|u\|_{L^\infty},$$

and let $R < \text{dist}(y, \partial\Omega) - R_0$ be such that

$$\delta_j(y)C_H(R) > \|u\|_{L^\infty}. \quad (75)$$

Note that $\delta_j(y) > 0$ and that, with such a choice of R , $B_R(y) \Subset V_j$. By the positivity of $\delta_j(y)$ and since $u(y) > 0$, $\Delta u(y) = -f(u(y)) < 0$. Thus, arbitrarily close to y we can find a point $\bar{y} \in V_j$ such that $u(\bar{y}) < u(y)$, and we can choose \bar{y} in order to satisfy the further relation $y \in B_R(\bar{y}) \Subset V_j$. Define

$$h(x) = \delta_j(y)v_R(r_{\bar{y}}(x)).$$

Since $v'_R < 0$, by the Laplacian comparison theorem it holds

$$\begin{cases} \Delta h \geq -\delta_j(y) & \text{weakly on } B_R(\bar{y}), \\ h = 0 & \text{on } \partial B_R(\bar{y}). \end{cases} \quad (76)$$

Note that $\|h\|_{L^\infty(B_R(\bar{y}))} = \delta_j(y)\|v_R\|_{L^\infty([0,R])} = \delta_j(y)C_H(R)$, and that the norm of h is attained at \bar{y} . For $\tau > 0$, define on $B_R(\bar{y})$

$$w(x) = \tau h(x) - u(x).$$

If τ is small enough, then $w < 0$. Choose τ to be the first value for which τh touches u from below. Hence, $w \leq 0$ and there exists x_0 such that $w(x_0) = 0$. From $h = 0$ on $\partial B_R(\bar{y})$, we deduce that $x_0 \in B_R(\bar{y})$ is an interior point. From our choice of h and \bar{y} ,

$$\tau \delta_j(y) C_H(R) = \tau h(\bar{y}) = w(\bar{y}) + u(\bar{y}) \leq u(\bar{y}) < u(y) \leq \|u\|_{L^\infty(\Omega)}.$$

By assumption (75), we deduce that necessarily $\tau < 1$. Now, from

$$u(x_0) = \tau h(x_0) \leq \tau h(\bar{y}) = w(\bar{y}) + u(\bar{y}) \leq u(\bar{y}) < u(y),$$

there exists a small neighbourhood $U \subset B_R(\bar{y})$ of x_0 such that $u|_U < u(y)$. But then, on U ,

$$\Delta u = -f(u) \leq -\min_U(f \circ u) \leq -\min_{t \in [\varepsilon_j, u(y)]} f(t) = -\delta_j(y). \quad (77)$$

Finally, combining (76) and (77), from $\tau < 1$ w satisfies

$$\begin{cases} \Delta w = \tau \Delta h - \Delta u \geq -\tau \delta_j(y) + \delta_j(y) > 0 & \text{weakly on } V, \\ w \leq 0 & \text{on } V, \quad w(x_0) = 0, \end{cases}$$

which contradicts the maximum principle and proved the desired (74). \square

Putting together the two theorems leads to the proof of Proposition 7.

Proof of Proposition 7. In our assumptions, we can modify the function u in a tiny neighbourhood $T \subseteq \Omega$ of $\partial\Omega$ to produce a function $\bar{u} \in C^2(M)$ such that $\bar{u} = u$ on $\Omega \setminus T$, $\sup_{T \cup (M \setminus \Omega)} \bar{u} < \|u\|_{L^\infty}$. For instance, choose $\varepsilon > 0$ be such that $\sup_{\partial\Omega} u + \varepsilon < u^*$, and let $\psi \in C^\infty(\mathbb{R})$ be such that

$$0 \leq \psi \leq 1, \quad \psi = 0 \text{ on } \left(-\infty, \sup_{\partial\Omega} u + \frac{\varepsilon}{2}\right), \quad \psi = 1 \text{ on } \left(\sup_{\partial\Omega} u + \varepsilon, +\infty\right).$$

Then, $\bar{u}(x) = \psi(u(x))$ (extended with zero on $M \setminus \Omega$) meets our requirements. Denote with $u^* = \sup_M \bar{u} = \sup_\Omega u > 0$. In our assumptions on the Ricci tensor, the strong maximum principle at infinity holds on M (see [33] and Appendix 1 below), thus we can find a sequence $\{x_k\}$ such that

$$\bar{u}(x_k) > u^* - \frac{1}{k}, \quad \frac{1}{k} \geq \Delta \bar{u}(x_k) = -f(\bar{u}(x_k)).$$

For k large enough, by the first condition $x_k \in \Omega \setminus \bar{T}$, thus $u = \bar{u}$ around x_k . Letting $k \rightarrow +\infty$ we deduce that $f(u^*) \geq 0$. Since $f(u(\Omega))$ is connected in \mathbb{R}_0^+ and contains 0, in our assumptions $u^* \leq \lambda$. Applying Lemmata 36 and 37 we infer the existence of a large R_0 such that, for each connected component V_j of Ω_{R_0} and for each $y \in V_j$,

$$\delta_j(y) C_H(\text{dist}(y, \partial\Omega) - R_0) \leq \lambda,$$

where $\delta_j(y)$ is defined in (72). Letting $\text{dist}(y, \partial\Omega) \rightarrow +\infty$ along V_j and using (71), we deduce that $\delta(y) \rightarrow 0$ uniformly as y diverges in V_j . By the very definition of $\delta_j(y)$ and our assumption on f , this implies $u(y) \rightarrow \lambda$ uniformly as $\text{dist}(y, \partial\Omega) \rightarrow +\infty$ along V_j . This also implies that $u^* = \lambda$, and concludes the proof. \square

Remark 38. If u solves $-\Delta u = f(u)$, in the sole assumptions

$$f > 0 \text{ on } (0, \lambda), \quad f(\lambda) = 0, \quad 0 \leq u \leq \lambda \text{ on } \Omega,$$

and $u \not\equiv 0$, $u \not\equiv \lambda$, then $0 < u < \lambda$ on Ω , by the strong maximum principle.

To deal with monotonicity properties of solutions, we shall investigate good Killing fields more closely. We begin with the next simple observation:

Lemma 39. *Let $(M, \langle \cdot, \cdot \rangle)$ be a complete Riemannian manifold, and let $\Omega \subseteq M$ be an open subset with non-empty boundary. Suppose that X is a Killing vector field on M , with associated flow Φ_t . Then, the next two conditions are equivalent:*

(i) *there exists $x \in \partial\Omega$ such that $\text{dist}(\Phi_t(x), \partial\Omega) \rightarrow +\infty$ as $t \rightarrow +\infty$;*

(ii) *$\text{dist}(\Phi_t(y), \partial\Omega) \rightarrow +\infty$ locally uniformly for $y \in \partial\Omega$.*

Proof. Indeed, suppose that (i) holds and let $y \in \partial\Omega$. Since Φ_t is a flow of isometries, $\text{dist}(x, y) = \text{dist}(\Phi_t(x), \Phi_t(y))$. For each $p_t \in \partial\Omega$ realizing $\text{dist}(\Phi_t(y), \partial\Omega)$, by the triangle inequality we thus get

$$\begin{aligned} \text{dist}(\Phi_t(x), \partial\Omega) - \text{dist}(x, y) &\leq \text{dist}(\Phi_t(x), p_t) - \text{dist}(x, y) \\ &\leq \text{dist}(\Phi_t(y), p_t) = \text{dist}(\Phi_t(y), \partial\Omega), \end{aligned}$$

from which (ii) immediately follows. \square

Remark 40. Note that condition (ii) in Definition 9, together with Lemma 39, implies that a good Killing vector field X is nowhere vanishing on $\bar{\Omega}$.

As anticipated, in the presence of a good Killing field on Ω , and for suitable nonlinearities f , we can construct a strictly monotone, non-constant solution of

$$\begin{cases} -\Delta u = f(u) & \text{on } \Omega, \\ u > 0 & \text{on } \Omega, \quad u = 0 & \text{on } \partial\Omega. \end{cases} \quad (78)$$

Proposition 41. *Let $(M, \langle \cdot, \cdot \rangle)$ with Ricci tensor satisfying $\text{Ric} \geq -(m-1)H^2 \langle \cdot, \cdot \rangle$, let $\Omega \subseteq M$ be an open, connected set with C^3 -boundary, and let $f \in C^1(\mathbb{R})$ with the properties*

$$\begin{cases} \text{(I)} & f > 0 \text{ on } (0, \lambda), \quad f(0) = f(\lambda) = 0, \\ \text{(II)} & f(s) \geq \left(\delta_0 + \frac{(m-1)^2 H^2}{4} \right) s \text{ for } s \in (0, s_0), \end{cases} \quad (79)$$

for some $\lambda > 0$ and some small $\delta_0, s_0 > 0$. Suppose that Ω admits a good Killing field X transverse to $\partial\Omega$, with flow $\Phi : \mathbb{R}_0^+ \times \bar{\Omega} \rightarrow \bar{\Omega}$, and suppose further that

$$\Phi(\mathbb{R}_0^+ \times \partial\Omega) \equiv \bar{\Omega}. \quad (80)$$

Then, there exists a non-constant solution $u \in C^2(\Omega) \cap C^0(\bar{\Omega})$ of (78) satisfying $0 < u < \lambda$ and the monotonicity $\langle \nabla u, X \rangle > 0$ on Ω .

Remark 42. Condition (II) in (79) is only required in order to show that the constructed solution is not identically zero.

Remark 43. The validity of (80) and the connectedness of Ω imply that also $\partial\Omega$ is connected.

Remark 44. Property (80) is not automatic for good Killing fields. As a counterexample, consider $M = \mathbb{R}^m$ with coordinates $(x', x_m) \in \mathbb{R}^{m-1} \times \mathbb{R}$, and set

$$\Omega = \mathbb{R}^m \setminus \{x = (x', x_m) \in \mathbb{R}^m : |x'| < 1, x_m \leq -(1 - |x'|^2)^{-1}\}.$$

Clearly, $X = \partial/\partial x_m$ is a good vector field on Ω , transverse to $\partial\Omega$, but $\Phi(\mathbb{R}_0^+ \times \partial\Omega)$ only covers the portion of Ω inside the cylinder $\{(x', x_m) : |x'| < 1\}$.

The proof of the above proposition relies on the classical sliding method in [5], [4]. For the convenience of the reader, we postpone it to Appendix 2.

The control on the asymptotic behaviour of u as $\text{dist}(x, \partial\Omega) \rightarrow +\infty$ ensured by Proposition 7, coupled with the existence of a good Killing field enables us to proceed along the lines in [2, 18] to obtain a sharp energy estimate.

Theorem 45. *Let $(M^m, \langle \cdot, \cdot \rangle)$ be a complete Riemannian manifold with $\text{Ric} \geq -(m-1)H^2 \langle \cdot, \cdot \rangle$, for some $H > 0$, and let $f \in C^1(\mathbb{R})$ with the properties*

$$\begin{cases} f > 0 & \text{on } (0, \lambda), & f(\lambda) = 0, & f < 0 & \text{on } (\lambda, \lambda + s_0), \\ f(s) \geq \left(\delta_0 + \frac{(m-1)^2 H^2}{4} \right) s & \text{for } s \in (0, s_0), \end{cases}$$

for some $\lambda > 0$ and some small $\delta_0, s_0 > 0$. Let $\Omega \subseteq M$ be a connected open set with smooth boundary, and suppose that Ω supports a good Killing field X . Let u be a bounded solution of

$$\begin{cases} -\Delta u = f(u) & \text{on } \Omega, \\ u > 0 & \text{on } \Omega, & u = 0 & \text{on } \partial\Omega \end{cases} \quad (81)$$

with the properties that

$$\begin{cases} \|u\|_{C^1(\Omega)} < +\infty; \\ \langle X, \nabla u \rangle \geq 0 & \text{on } \Omega. \end{cases} \quad (82)$$

Then, there exists a positive $C = C(\|u\|_{C^1(\Omega)})$ such that

$$\int_{\Omega \cap B_R} |\nabla u|^2 dx \leq C \left[\mathcal{H}^{m-1}(\partial B_R) + \mathcal{H}^{m-1}(\partial\Omega \cap B_R) \right] \quad (83)$$

Proof. Set $B_R = B_R(o)$. By Corollary 25, $\langle \nabla u, X \rangle > 0$ on Ω . Indeed, the possibility $\langle \nabla u, X \rangle = 0$ is ruled out by (9) and since $u > 0$ on Ω , $u = 0$ on $\partial\Omega$. Define $u_t(x) = u(\Phi_t(x))$, and note that, by the first assumption in (9), u_t is defined on Ω . Since Φ_t is an isometry and X is Killing,

$$\begin{cases} \partial_t u_t = \langle \nabla u, X \rangle \circ \Phi_t > 0, & -\Delta u_t = f(u_t), \\ |\nabla u_t|^2 = |(\text{d}\Phi_t \circ \text{d}u)^\#|^2 = |\nabla u|^2 \circ \Phi_t, \\ \text{d}(\partial_t u_t) = \text{d}(\langle \nabla u, X \rangle) \circ \text{d}\Phi_t = \nabla \text{d}u(X, \text{d}\Phi_t) + \langle \nabla u, \nabla_{\text{d}\Phi_t} X \rangle. \end{cases} \quad (84)$$

We claim that $\nabla u_t = \text{d}\Phi_{-t}(\nabla u)$. Indeed, for every vector field W and using that Φ_t is an isometry,

$$\langle \nabla u_t, W \rangle = \text{d}u_t(W) = \text{d}u(\text{d}\Phi_t(W)) = \langle \nabla u, \text{d}\Phi_t(W) \rangle = \langle \text{d}\Phi_{-t}(\nabla u), W \rangle.$$

We thus deduce that, from (84) and again the Killing property,

$$\begin{aligned} \langle \nabla u_t, \nabla(\partial_t u_t) \rangle &= \text{d}(\partial_t u_t)(\nabla u_t) = \nabla \text{d}u(X, \text{d}\Phi_t(\nabla u_t)) + \langle \nabla u, \nabla_{\text{d}\Phi_t(\nabla u_t)} X \rangle \\ &= [\nabla \text{d}u(X, \nabla u)] \circ \Phi_t + \langle \nabla u, \nabla_{\nabla u} X \rangle = [\nabla \text{d}u(X, \nabla u)] \circ \Phi_t; \\ \partial_t |\nabla u_t|^2 &= \text{d}(|\nabla u|^2 \circ \Phi_t)(\partial_t) = \langle \nabla |\nabla u|^2, X \rangle \circ \Phi_t = [2\nabla \text{d}u(\nabla u, X)] \circ \Phi_t, \end{aligned}$$

whence

$$\frac{1}{2} \partial_t |\nabla u_t|^2 = \langle \nabla u_t, \nabla(\partial_t u_t) \rangle. \quad (85)$$

Set for convenience

$$E_R(t) = E_{\Omega \cap B_R}(u_t) = \frac{1}{2} \int_{\Omega \cap B_R} |\nabla u_t|^2 dx - \int_{\Omega \cap B_R} F(u_t),$$

where $E_{\Omega \cap B_R}$ and F are as in (15). In our assumptions, since u is bounded, by Proposition 7 we have that $\|u\|_{L^\infty} = \lambda$ and $u(x) \rightarrow \lambda$ uniformly as $\text{dist}(x, \partial\Omega) \rightarrow +\infty$ along each fixed connected component of Ω_{R_0} . Using (ii) of Definition 9, and Lemma 39, we deduce that

$$\|u_t\|_{L^\infty} \leq \lambda, \quad u_t(x) \rightarrow \lambda \quad \text{as } t \rightarrow +\infty, \text{ pointwise on } \Omega \cap B_R.$$

By the first assumption in (82), there exists a uniform constant C such that

$$\|\nabla u_t\|_{L^\infty(\Omega \cap B_R)} \leq C \quad \text{for every } t \in \mathbb{R}^+, \quad (86)$$

whence, by elliptic estimates, up to a subsequence

$$u_t \rightharpoonup \lambda \quad \text{in } C^{2,\alpha}(\Omega \cap B_R). \quad (87)$$

Differentiating under the integral sign with the aid of (85), integrating by parts and using (84), (86) we get

$$\begin{aligned} \frac{dE_R(t)}{dt} &= \int_{\Omega \cap B_R} \langle \nabla u_t, \nabla(\partial_t u_t) \rangle dx - \int_{\Omega \cap B_R} f(u_t)(\partial_t u_t) \\ &= \int_{\partial(\Omega \cap B_R)} (\partial_\nu u_t) \partial_t u_t d\sigma - \int_{\Omega \cap B_R} (\partial_t u_t) [\Delta u_t + f(u_t)] dx \\ &= \int_{\partial(\Omega \cap B_R)} (\partial_\nu u_t) \partial_t u_t d\sigma \geq -C \int_{\partial(\Omega \cap B_R)} \partial_t u_t d\sigma. \end{aligned}$$

Now, integrating on $(0, T)$ and using Tonelli's theorem we obtain

$$\begin{aligned} E_R(T) - E_R(0) &\geq -C \int_0^T \int_{\partial(\Omega \cap B_R)} \partial_t u_t d\sigma dt = C \int_{\partial(\Omega \cap B_R)} [u_T - u_0] d\sigma \\ &\geq -2C\lambda \mathcal{H}^{m-1}(\partial(\Omega \cap B_R)) \\ &\geq -2C\lambda [\mathcal{H}^{m-1}(\partial B_R) + \mathcal{H}^{m-1}(\partial\Omega \cap B_R)]. \end{aligned}$$

Since $F(u_t) \geq 0$, we deduce

$$\int_{\Omega \cap B_R} |\nabla u|^2 dx \leq E_R(0) \leq E_R(T) + 2C\lambda [\mathcal{H}^{m-1}(\partial B_R) + \mathcal{H}^{m-1}(\partial\Omega \cap B_R)].$$

By (87), $E_R(T) \rightarrow 0$ as $T \rightarrow +\infty$, from which the desired estimate (83) follows. \square

Putting together with Theorem 5, we easily prove Theorem 10.

Proof of Theorem 10. By Corollary 25, either $\langle \nabla u, X \rangle > 0$ or $\langle \nabla u, X \rangle = 0$ on Ω . However, from the existence of $o \in \partial\Omega$ with the property (ii) of Definition 9, and since $u > 0$ on Ω and $u = 0$ on $\partial\Omega$, we infer that the second possibility cannot occur. Moreover, by Hopf Lemma $\partial_\nu u > 0$ on $\partial\Omega$. Via Bishop-Gromov volume estimate, assumption $\text{Ric} \geq 0$ implies $\mathcal{H}^2(\partial B_R) \leq 4\pi R^2$ for $R > 0$, thus applying Theorem 45 with $H = 0$ we deduce that, by (10),

$$\int_{\Omega \cap B_R} |\nabla u|^2 \leq C [4\pi R^2 + \mathcal{H}^2(\partial\Omega \cap B_R)] = o(R^2 \log R)$$

as $R \rightarrow +\infty$. Now, the conclusion follows by applying Theorem 5. \square

To conclude, particularizing to the flat case we recover Theorem 1.8 in [18].

Corollary 46. *Let $\Omega \subseteq \mathbb{R}^3$ be an open set with a C^3 boundary. Suppose that Ω can be described as the epigraph of a function $\varphi \in C^3(\mathbb{R}^2)$ over some plane \mathbb{R}^2 , and that φ is globally Lipschitz on \mathbb{R}^2 . Let $f \in C^1(\mathbb{R})$ satisfy*

$$\begin{cases} f > 0 & \text{on } (0, \lambda), & f(\lambda) = 0, & f < 0 & \text{on } (\lambda, \lambda + s_0), \\ f(s) \geq \delta_0 s & \text{for } s \in (0, s_0), \\ f \text{ is non-increasing} & \text{on } [\lambda - s_0, \lambda], \end{cases} \quad (88)$$

for some $\lambda > 0$ and some small $\delta_0, s_0 > 0$. Then, if there exists a non-constant, positive, bounded solution $u \in C^3(\bar{\Omega})$ of the overdetermined problem

$$\begin{cases} -\Delta u = f(u) & \text{on } \Omega \\ u = 0 & \text{on } \partial\Omega \\ \partial_\nu u = \text{constant} & \text{on } \partial\Omega, \end{cases} \quad (89)$$

Ω is an half-space and, up to translations, $\partial\Omega = v^\perp$ for some $v \in \mathbb{S}^2$, $u(x) = y(\langle x, v \rangle)$ and $y'' = -f(y)$ on \mathbb{R}^+ .

Proof. Up to an isometry, we can assume that $\mathbb{R}^3 = \mathbb{R}^2 \times \mathbb{R}$ with coordinates (x', x_3) , and that

$$\Omega = \{(x', x_3) : x' \in \mathbb{R}^2, x_3 > \varphi(x')\}.$$

Let $X = \partial/\partial x_3$ be the translational vector field along the third coordinate direction, and let Φ_t be the associated flow. Clearly, $\text{dist}(\Phi_t(x), \partial\Omega) \rightarrow +\infty$ as $t \rightarrow +\infty$, thus X is a good Killing field for Ω . By Theorem 1.1 in [4], the monotonicity $\langle \nabla u, X \rangle > 0$ is satisfied on Ω . Having fixed an origin $o \in \partial\Omega$, since φ is globally Lipschitz we deduce that

$$\mathcal{H}^2(\Omega \cap B_R) \leq \int_{B_R \subseteq \mathbb{R}^2} \sqrt{1 + |\nabla \varphi(x')|^2} dx' \leq CR^2 \quad \text{as } R \rightarrow +\infty.$$

Therefore, the desired conclusion follows from Theorem 10. \square

Appendix 1: some remarks on L^∞ bounds for u and ∇u .

Under some mild conditions on the nonlinearity $f(t)$, it can be proved that both u and $|\nabla u|$ are globally bounded on M . In this appendix, we collect and comment on two general estimates. We first examine L^∞ bounds for u . Suppose that Δ satisfies the strong maximum principle at infinity (also called the Omori-Yau maximum principle), briefly (SMP). We recall that, by definition, Δ satisfies (SMP) if, for every $w \in C^2(M)$ with $w^* = \sup w < +\infty$, there always exist a sequence $\{x_k\} \subseteq M$ such that

$$w(x_k) > w^* - \frac{1}{k}, \quad |\nabla w|(x_k) < \frac{1}{k}, \quad \Delta w(x_k) < \frac{1}{k}.$$

As it is shown in [33], (SMP) turns out to be an extremely powerful tool in modern Geometric Analysis, and its validity is granted via mild function-theoretic properties of M . In particular, if $r(x)$ denotes the distance from a fixed point, the conditions

$$\begin{aligned} \text{Ric}(\nabla r, \nabla r)(x) &\geq -(m-1)G(r(x)), \\ G(t) &= Ct^2 \log^2 t \quad \text{for } t \gg 1, \end{aligned} \quad (90)$$

where $C > 0$ and G is a smooth, positive and non-decreasing function defined on $[0, +\infty)$, ensure that (SMP) holds for Δ . A proof of this fact can be found, for instance, in [33], Example 1.13. Observe that (90) includes the cases

$$(i) \quad \text{Ric} \geq -(m-1)H^2 \langle \cdot, \cdot \rangle \quad \text{and} \quad (ii) \quad K \geq -H^2, \quad \text{for some constant } H \geq 0,$$

which have originally been investigated by S.T. Yau (case (i), in [10, 11]) and H. Omori (case (ii), in [31]). Under the validity of (SMP), the next general result enables us to obtain L^∞ bounds for wide classes of differential inequalities.

Theorem 47 ([33], Theorem 1.31). *Suppose that Δ satisfies (SMP), and let $u \in C^2(M)$ be a solution of $\Delta u \geq -f(u)$, for some $f \in C^0(\mathbb{R})$. Then,*

$$u^* = \sup_M u < +\infty \quad \text{and} \quad f(u^*) \geq 0$$

provided that there exists a function F , positive on $[a, +\infty)$ for some $a \in \mathbb{R}$, with the following properties:

$$\left\{ \int_a^t F(s) ds \right\}^{-1/2} \in L^1(+\infty), \quad \limsup_{t \rightarrow +\infty} \frac{\int_a^t F(s) ds}{tF(t)} < +\infty, \quad \liminf_{t \rightarrow +\infty} \frac{-f(t)}{F(t)} > 0. \quad (91)$$

Next, we consider L^∞ bounds for ∇u , where $u \in C^3(M)$ is a bounded solution of $-\Delta u = f(u)$ and $f \in C^1(\mathbb{R})$.

Remark 48. Since $\text{Ric} \geq 0$ and M is complete, the property $: u \in L^\infty(M) \Rightarrow \nabla u \in L^\infty(M)$ holds true as a (quite standard) consequence of the Bochner identity and the De Giorgi-Nash-Moser's regularity theory for PDE's (cfr. for instance [26, 38]). Indeed, if u solves $-\Delta u = f(u)$ then, by Bochner formula,

$$\Delta |\nabla u|^2 \geq -2f'(u)|\nabla u|^2 + 2\text{Ric}(\nabla u, \nabla u) \geq -C|\nabla u|^2 \quad \text{on } M.$$

Now, we can run the Moser iteration to get the desired bound (since $f(u)$ is bounded on M). In fact, we have used the well-known facts that any Riemannian manifold with $\text{Ric} \geq 0$ has a scale invariant L^2 Neumann Poincaré inequality and a relative volume comparison property. We conclude this remark by pointing out that the property: $u \in L^\infty(M) \Rightarrow \nabla u \in L^\infty(M)$ holds true also under the less restrictive assumption $\text{Ric} \geq -(m-1)H^2 \langle \cdot, \cdot \rangle$, for some $H \geq 0$ (cfr. for instance [28], Theorem 2.6 and Corollary 2.7). We stress that the techniques to prove this generalized result are different from the ones outlined above, and rely on Ahlfors-Yau type gradient estimates.

As a prototype case, we now prove uniform L^∞ bounds for u and ∇u for the Allen-Cahn equation appearing in De Giorgi's conjecture.

Corollary 49. *Let M be a complete manifold satisfying $\text{Ric} \geq -(m-1)H^2 \langle \cdot, \cdot \rangle$, for some $H \geq 0$, and let $u \in C^2(M)$ be a solution of the Allen-Cahn equation*

$$-\Delta u = u - u^3 \quad \text{on } M.$$

Then, u is smooth, $-1 \leq u(x) \leq 1$ for every $x \in M$ and $|\nabla u| \in L^\infty(M)$.

Proof. By standard elliptic estimates u is smooth on M . In our assumptions on the Ricci curvature, by the remarks above M satisfies (SMP). Set $f(t) = t - t^3$. It is easy to check that $F(t) = t^3$ satisfies the assumptions in (91). Then, by Theorem 47, u is bounded above

and $(u^*)^3 - u^* \leq 0$, which gives $u^* \in [0, 1]$ or $u^* \leq -1$. Analogously, the function $w = -u$ satisfies

$$\Delta w = -\Delta u = u - u^3 = w^3 - w,$$

and applying the same result we deduce that either $w^* \in [0, 1]$ or $w^* \leq -1$. Since $w^* = -u_* := -\inf_M u$ we deduce that either $u_* \in [-1, 0]$ or $u_* \geq 1$. Combining with the above estimates for u^* the L^∞ bound for u follows immediately. The L^∞ bound for ∇u is a direct consequence of Remark 48. \square

Remark 50. In the Euclidean case, all the distributional solutions $u \in L^1_{\text{loc}}(\mathbb{R}^m)$ of the Allen-Cahn equation $-\Delta u = u - u^3$ (and more generally of the vector valued Ginzburg-Landau equation $-\Delta u = u(1 - |u|^2)$) always satisfy the bound $|u| \leq 1$, see Proposition 1.9 in [13]. Hence, by standard elliptic estimates, they are smooth and all their derivatives are bounded too.

Appendix 2: construction of a monotone solution

In this appendix, under the presence of a good Killing field on Ω , we construct a non-constant solution of

$$\begin{cases} -\Delta u = f(u) & \text{on } \Omega, \\ u > 0 & \text{on } \Omega, \quad u = 0 \quad \text{on } \partial\Omega. \end{cases} \quad (92)$$

We recall the geometric assumptions: let $(M, \langle \cdot, \cdot \rangle)$ with Ricci tensor satisfying $\text{Ric} \geq -(m-1)H^2 \langle \cdot, \cdot \rangle$, let $\Omega \subseteq M$ be an open set with C^3 -boundary, and let $f \in C^1(\mathbb{R})$ with the properties

$$\begin{cases} \text{(I)} & f > 0 \quad \text{on } (0, \lambda), \quad f(0) = f(\lambda) = 0, \\ \text{(II)} & f(s) \geq \left(\delta_0 + \frac{(m-1)^2 H^2}{4} \right) s \quad \text{for } s \in (0, s_0), \end{cases} \quad (93)$$

for some $\lambda > 0$ and some small $\delta_0, s_0 > 0$. Suppose that X is a good Killing field on Ω , with flow $\Phi : \mathbb{R}_0^+ \times \bar{\Omega} \rightarrow \bar{\Omega}$.

Proposition 51. *In the above assumptions, suppose further that X is transverse to $\partial\Omega$ and that*

$$\Phi(\mathbb{R}_0^+ \times \partial\Omega) \equiv \bar{\Omega}. \quad (94)$$

Then, there exists a non-constant solution $u \in C^2(\Omega) \cap C^0(\bar{\Omega})$ of (92) such that $0 < u < \lambda$ and $\langle \nabla u, X \rangle > 0$ on Ω .

Proof. Let $\{U_j\} \uparrow \partial\Omega$ be a smooth exhaustion of $\partial\Omega$. By the properties of flows and the transversality of X and $\partial\Omega$, the map Φ restricted to $\mathbb{R}^+ \times U_j$ realizes a diffeomorphism onto its image. We briefly prove it. To show that Φ is injective, suppose that there exist $(t_1, x_1) \neq (t_2, x_2)$ for which $\Phi(t_1, x_1) = \Phi(t_2, x_2)$. Then, by the properties of the flow, necessarily $t_1 < t_2$ (up to renaming). Since $\Phi(t_2, x_2) = \Phi(t_1, \Phi_{t_2-t_1}(x_2))$, the equality and the fact that Φ_{t_1} is a diffeomorphism imply that $x_1 = \Phi_{t_2-t_1}(x_2)$. Hence, the flow line $\Phi_t(x_1)$ intersects twice the boundary $\partial\Omega$, and by property (i) of good Killing fields it holds $\Phi_{[0, t_2-t_1]}(x_1) \subseteq \partial\Omega$, which is impossible since X is transverse to $\partial\Omega$. Next, we show that $d\Phi$ is nonsingular. Indeed, if at a point $\Phi(t, x)$ we have $X_{\Phi(t, x)} = d\Phi(\partial_t) = d\Phi(Z_x)$ for some nonzero $Z_x \in T_x \partial\Omega$, then applying $d\Phi_{-t}$ we would have $X_x = Z_x$, which is impossible again by the transversality of X and $\partial\Omega$. Next, choose a sequence $\{T_k\} \uparrow +\infty$ and define the cylinders $C_{jk} = \Phi([0, T_k] \times U_j)$. By (94), $\Omega = \bigcup_{j,k} C_{jk}$ (this is the only point where

(94) is used). Denote with $\pi_1 : \mathbb{R}_0^+ \times \bar{U}_j \rightarrow \mathbb{R}_0^+$ the projection onto the first factor, and with $\pi = \pi_1 \circ \Phi^{-1} : \bar{C}_{jk} \rightarrow \mathbb{R}_0^+$ its image through Φ . Take a sequence $\{\psi_k\} \subseteq C^\infty(\mathbb{R}_0^+)$ with the following properties:

$$\begin{aligned} 0 \leq \psi_k \leq \lambda \text{ on } \mathbb{R}^+, \quad \psi_k \equiv \lambda \text{ on } [T_k, +\infty), \quad \psi_k(0) = 0 \text{ for each } k, \\ \psi_k \text{ is strictly monotone on } [0, T_k], \quad \psi_k \geq \psi_{k+1} \text{ on } \mathbb{R}_0^+. \end{aligned}$$

For every pair (j, k) , let $u_{jk} \in C^2(C_{jk}) \cap C^0(\bar{C}_{jk})$ be a solution of

$$\begin{cases} -\Delta u_{jk} = f(u_{jk}) & \text{on } C_{jk} \\ u_{jk} = \psi_k \circ \pi & \text{on } \Phi([0, T_k] \times \partial U_j) \\ u_{jk} = 0 & \text{on } \Phi(\{0\} \times U_j) \quad u_{jk} = \lambda \text{ on } \Phi(\{T_k\} \times U_j), \end{cases} \quad (95)$$

constructed via the monotone iteration scheme (see [39]) by using $u \equiv 0$ as a subsolution and $u \equiv \lambda$ as a supersolution. Then, $0 \leq u_{jk} \leq \lambda$ on \bar{C}_{jk} , and the inequality is strict on C_{jk} by the strong maximum principle (see Remark 38).

Step 1: u_{jk} is monotone in t on C_{jk} .

To prove this claim, for $t \in \mathbb{R}^+$ set

$$w_t = u_{jk} \circ \Phi_{-t} - u_{jk} \quad \text{on } V_t = \Phi_t(C_{jk}) \cap C_{jk} = \begin{cases} \emptyset & \text{if } t \geq T_k \\ \Phi(U_j \times (t, T_k)) & \text{if } t \in [0, T_k). \end{cases}$$

Hereafter we omit writing the pair (j, k) . In our assumptions, for every $t > 0$

$$\begin{cases} -\Delta w_t = c_t(x)w_t & \text{on } V_t \\ w_t < 0 & \text{on } \partial V_t, \end{cases} \quad \text{where } c_t(x) = \frac{f(u_{jk} \circ \Phi_{-t}) - f(u_{jk})}{w_t} \quad (96)$$

We now claim that, if t is sufficiently close to T_k , then the operator $L_t = \Delta + c_t(x)$ is non-negative on V_t (as observed by S.R.S. Varadhan and A. Bakelman, see Proposition 1.1 of [5]). Indeed let $S > 0$ be the L^2 -Sobolev constant of $W = \Phi([0, 2T_k] \times U_j)$:

$$S\|\phi\|_{L^{2^*}(W)} \leq \|\nabla\phi\|_{L^2(W)} \quad \text{for every } \phi \in C_c^\infty(W).$$

Then, for every $V \subseteq W$ and every $\phi \in C_c^\infty(V)$, by Cauchy-Schwarz inequality

$$\int c_t \phi^2 \leq \text{Lip}_{[0, \lambda]}(f) \int \phi^2 \leq \text{Lip}_{[0, \lambda]}(f) |V|^{\frac{2}{m}} \left(\int \phi^{\frac{2m}{m-2}} \right)^{\frac{m-2}{m}} \leq \frac{\text{Lip}_{[0, \lambda]}(f) |V|^{\frac{2}{m}}}{S} \int |\nabla\phi|^2,$$

where $\text{Lip}_{[0, \lambda]}(f)$ is the Lipschitz constant of f on $[0, \lambda]$. If $|V|$ is sufficiently small (and the bound does not depend on $t \in (0, T_k]$), it thus follows that

$$\int |\nabla\phi|^2 - \int c_t \phi^2 \geq 0,$$

which means that L_t has non-negative spectrum. Particularizing to $V = V_t$ proves the claim. By a classical result, [6], the non-negativity of L_t on V_t is equivalent to the validity of the maximum principle for L_t on V_t , hence, by (96), $w_t \leq 0$ on V_t . The strong maximum principle then implies the strict inequality $w_t < 0$ on V_t . Now, consider

$$\mathcal{T} = \{t \in [0, T_k] : w_s < 0 \text{ on } V_s, \text{ for each } s \in [t, T_k]\},$$

which by the previous claim is non-empty and contains a left neighbourhood of T_k . We are going to prove that $\bar{t} = \inf \mathcal{T} = 0$. If, by contradiction, $\bar{t} > 0$, then by continuity $w_{\bar{t}} \leq 0$ on $V_{\bar{t}}$. Since, by (96), $(w_{\bar{t}})|_{\partial V_{\bar{t}}} < 0$, the strong maximum principle implies that $w_{\bar{t}} < 0$ on $\bar{V}_{\bar{t}}$. By compactness, let $\varepsilon > 0$ be such that $w_{\bar{t}} < -\varepsilon$ on $\bar{V}_{\bar{t}}$, and by continuity choose $\eta > 0$ sufficiently small in order to satisfy the next requirements:

- the operator $L_{\bar{t}-\eta}$ is non-negative on $V_{\bar{t}-\eta} \setminus V_{\bar{t}} = \Phi((\bar{t} - \eta, \bar{t}] \times U_j)$;
- $w_{\bar{t}-\eta} \leq -\frac{\varepsilon}{2}$ on $V_{\bar{t}}$.

By our construction, $w_{\bar{t}-\eta} < 0$ on $\partial(V_{\bar{t}-\eta} \setminus V_{\bar{t}})$, thus by the maximum principle $w_{\bar{t}-\eta} < 0$ on $V_{\bar{t}-\eta} \setminus V_{\bar{t}}$ and so on $V_{\bar{t}-\eta} = V_{\bar{t}} \cup (V_{\bar{t}-\eta} \setminus V_{\bar{t}})$, contradicting the minimality of \bar{t} . Concluding, $\bar{t} = 0$, hence $w_t > 0$ on \mathbb{R}^+ for every $t \in (0, T_k]$, which proves the monotonicity of u in the t -direction.

Step 2: the limiting procedure.

First, by requirement (ii) in Definition 9 of a good Killing field we argue that $\Omega_{R_0} = \{x \in \Omega : \text{dist}(x, \partial\Omega) \geq R_0\}$ is non-empty for each R_0 . In our assumptions on Ric and on f , by a comparison procedure identical to that performed in Lemma 36 we can find:

- $R > 0$ such that $\lambda_1(\mathbb{B}_R)s < f(s)$ for every $s \in [0, s_0]$, where $\lambda_1(\mathbb{B}_R)$ is the first eigenvalue of a geodesic ball \mathbb{B}_R in a space form M_H ;
- $y \in \Omega_{2R_0}$ and a Lipschitz, weak solution $w \in \text{Lip}(B_R(y))$ of

$$\begin{cases} -\Delta w \leq \lambda_1(\mathbb{B}_R)w < f(w) & \text{on } B_R(y), \\ w|_{\partial B_R(y)} = 0, \quad w > 0 & \text{on } B_R(y), \quad \|w\|_{L^\infty(B_R(y))} < s_0. \end{cases}$$

We arrange the exhaustion $\{U_j\}$ in such a way that $B_R(y) \Subset \Phi(\mathbb{R}^+ \times U_0)$, and for each fixed j we let $k = k_j$ be such that $B_R(y) \Subset C_{jk}$ for every $k \geq k_j$. This latter property is possible by (ii) of Definition 9 of a good Killing field, together with Lemma 39. Since $\partial\Omega \in C^3$, we can smooth the corners of C_{jk} in such a way that $\partial C_{jk} \in C^3$. By uniform elliptic estimates, up to passing to a subsequence $\{u_{jk}\}_k$ converges in $C_{\text{loc}}^{2,\alpha}$ to a solution u_j of

$$-\Delta u_j = f(u_j) \quad \text{on } C_j = \Phi(\mathbb{R}_0^+ \times U_j), \quad u_j = 0 \quad \text{on } \Phi(\{0\} \times U_j), \quad 0 \leq u_j \leq \lambda.$$

Moreover, by comparison $u_{jk} \geq w$ on $B_R(y)$, hence $u_j \geq w$. Letting now $j \rightarrow +\infty$ and using again elliptic estimates we get the existence of the desired u with $0 \leq u \leq \lambda$. From $u_j \geq w$ we deduce that $u \geq w$, thus u is non-zero. By Remark 38 and since $u = 0$ on $\partial\Omega$, $0 < u < \lambda$ on Ω . The monotonicity relation $\langle \nabla u, X \rangle \geq 0$ follows from that of u_{jk} via pointwise convergence. To prove the stronger $\langle \nabla u, X \rangle > 0$, we apply Corollary 25 to get that either $\langle \nabla u, X \rangle \equiv 0$ or $\langle \nabla u, X \rangle > 0$. The first case is ruled out, because it would mean that u is constant on the flow lines of X : starting from a point $x \in \partial\Omega$, this and the positivity of u on Ω would imply that $\Phi_t(x) \in \partial\Omega$ for every $t \in \mathbb{R}^+$, contradicting property (ii) of Definition 9 (or, even, contradicting the transversality of X and $\partial\Omega$). \square

Acknowledgements: The second author is indebted to Jorge Herbert de Lira for a stimulating discussion about Killing fields on Riemannian manifolds. The first and the third authors were supported by the ERC grant EPSILON (*Elliptic Pde's and Symmetry of Interfaces and Layers for Odd Nonlinearities*).

References

- [1] G. Alberti, L. Ambrosio, and X. Cabré. On a long-standing conjecture of E. De Giorgi: symmetry in 3D for general nonlinearities and a local minimality property. *Acta Appl. Math.*, 65(1-3):9–33, 2001. Special issue dedicated to Antonio Avantaggiati on the occasion of his 70th birthday.
- [2] L. Ambrosio and X. Cabré. Entire solutions of semilinear elliptic equations in \mathbb{R}^3 and a conjecture of De Giorgi. *J. Amer. Math. Soc.*, 13(4):725–739 (electronic), 2000.
- [3] H. Berestycki, L. Caffarelli, and L. Nirenberg. Further qualitative properties for elliptic equations in unbounded domains. *Ann. Scuola Norm. Sup. Pisa Cl. Sci. (4)*, 25(1-2):69–94 (1998), 1997. Dedicated to Ennio De Giorgi.
- [4] H. Berestycki, L.A. Caffarelli, and L. Nirenberg. Monotonicity for elliptic equations in unbounded Lipschitz domains. *Comm. Pure Appl. Math.*, 50(11):1089–1111, 1997.
- [5] H. Berestycki and L. Nirenberg. On the method of moving planes and the sliding method. *Bol. Soc. Brasil. Mat. (N.S.)*, 22(1):1–37, 1991.
- [6] H. Berestycki, L. Nirenberg, and S.R.S. Varadhan. The principal eigenvalue and maximum principle for second-order elliptic operators in general domains. *Comm. Pure Appl. Math.*, 47(1):47–92, 1994.
- [7] R. Brooks. A relation between growth and the spectrum of the Laplacian. *Math. Z.*, 178(4):501–508, 1981.
- [8] E. Calabi. On manifolds with non-negative Ricci curvature II. *Notices Amer. Math. Soc.*, 22:A205, 1975.
- [9] J. Cheeger and D. Gromoll. The splitting theorem for manifolds of nonnegative Ricci curvature. *J. Diff. Geom.*, 6:119–128, 1971/72.
- [10] S.Y. Cheng and S.T. Yau. Differential equations on Riemannian manifolds and their geometric applications. *Comm. Pure Appl. Math.*, 28(3):333–354, 1975.
- [11] S.Y. Cheng and S.T. Yau. Hypersurfaces with constant scalar curvature. *Math. Ann.*, 225:195–204, 1977.
- [12] L. Dupaigne and A. Farina. Stable solutions of $-\Delta u = f(u)$ in \mathbb{R}^N . *J. Eur. Math. Soc.*, 12(4):855–882, 2010.
- [13] A. Farina. Finite-energy solutions, quantization effects and Liouville-type results for a variant of the Ginzburg-Landau systems in \mathbb{R}^K . *Differential Integral Equations*, 11(6):875–893, 1998.
- [14] A. Farina. Propriétés qualitatives de solutions d’équations et systèmes d’équations non-linéaires. 2002. Habilitation à diriger des recherches, Paris VI.
- [15] A. Farina, B. Sciunzi, and E. Valdinoci. Bernstein and De Giorgi type problems: new results via a geometric approach. *Ann. Sc. Norm. Super. Pisa Cl. Sci. (5)*, 7(4):741–791, 2008.
- [16] A. Farina, Y. Sire, and E. Valdinoci. Stable solutions of elliptic equations on Riemannian manifolds. *J. Geom. Anal.*, 2012.

- [17] A. Farina and E. Valdinoci. The state of the art for a conjecture of De Giorgi and related problems. In *Recent progress on reaction-diffusion systems and viscosity solutions*, pages 74–96. World Sci. Publ., Hackensack, NJ, 2009.
- [18] A. Farina and E. Valdinoci. Flattening results for elliptic PDEs in unbounded domains with applications to overdetermined problems. *Arch. Ration. Mech. Anal.*, 195(3):1025–1058, 2010.
- [19] A. Farina and E. Valdinoci. Correction of a technical point and some remarks about the paper appeared in arch. ration. mech. anal. 195(2010), no. 3, 1025-1058. *Arch. Ration. Mech. Anal.*, DOI: 10.1007/s00205-012-0568-6, <http://www.springerlink.com/content/p1m68k11856667w7/>
- [20] D. Fischer-Colbrie and R. Schoen. The structure of complete stable minimal surfaces in 3-manifolds of non negative scalar curvature. *Comm. Pure Appl. Math.*, XXXIII:199–211, 1980.
- [21] D. Gilbarg and N. Trudinger. *Elliptic Partial Differential Equations of Second Order*. third ed., Springer-Verlag, 1998.
- [22] E. De Giorgi. Convergence problems for functionals and operators. In *Proceedings of the International Meeting on Recent Methods in Nonlinear Analysis (Rome, 1978)*, pages 131–188, Bologna, 1979. Pitagora.
- [23] A. Grigor’yan. Analytic and geometric background of recurrence and non-explosion of the Brownian motion on Riemannian manifolds. *Bull. Amer. Math. Soc.*, 36:135–249, 1999.
- [24] R.S. Hamilton. Four-manifolds with positive curvature operator. *J. Differential Geom.*, 24(2):153–179, 1986.
- [25] S. Jimbo. On a semilinear diffusion equation on a Riemannian manifold and its stable equilibrium solutions. *Proc. Japan Acad. Ser. A Math. Sci.*, 60(10):349–352, 1984.
- [26] P. Li. Lecture notes on geometric analysis. *Lecture Notes Series No. 6, Research Institute of Mathematics, Global Analysis Research Center, Seoul National University, Korea (1993)*.
- [27] G. Liu. 3-manifolds with nonnegative Ricci curvature. *available at arXiv:1108.1888*.
- [28] P. Mastrolia and M. Rigoli. Diffusion-type operators, Liouville theorems and gradient estimates on complete manifolds. *Nonlinear Anal.*, 72(9-10):3767–3785, 2010.
- [29] H.P. McKean. An upper bound to the spectrum of Δ on a manifold of negative curvature. *J. Diff. Geom.*, 4:359–366, 1970.
- [30] W.F. Moss and J. Piepenbrink. Positive solutions of elliptic equations. *Pac. J. Math.*, 75:219–226, 1978.
- [31] H. Omori. Isometric immersions of Riemannian manifolds. *J. Math. Soc. Japan*, 19:205–214, 1967.
- [32] P. Petersen. *Riemannian Geometry*. Springer-Verlag, 1997.
- [33] S. Pigola, M. Rigoli, and A.G. Setti. Maximum principles on Riemannian manifolds and applications. *Mem. Amer. Math. Soc.*, 174:no.822, 2005.

- [34] S. Pigola, M. Rigoli, and A.G. Setti. *Vanishing and finiteness results in Geometric Analysis. A generalization of the Böchner technique*, volume 266 of *Progress in Math.* Birkäuser, 2008.
- [35] S. Pigola and G. Veronelli. Remarks on L^p -vanishing results in Geometric Analysis. *Internat. J. Math.*, 23(1):18pp., 2012.
- [36] M.H. Protter and H.F. Weinberger. *Maximum principles in differential equations*. (Corrected reprint of the 1967 original), Springer-Verlag, 1984.
- [37] A. Ros and P. Sicbaldi. Geometry and topology of some overdetermined elliptic problems. *Preprint*, 2012.
- [38] L. Saloff-Coste. *Aspects of Sobolev-Type Inequalities*. London Mathematical Society Lecture Notes Series, 2001.
- [39] D.H. Sattinger. Monotone methods in nonlinear elliptic and parabolic boundary value problems. *Indiana Univ. Math. J.*, 21:979–1000, 1971/72.
- [40] P. Sternberg and K. Zumbrun. Connectivity of phase boundaries in strictly convex domains. *Arch. Rational Mech. Anal.*, 141(4):375–400, 1998.
- [41] P. Sternberg and K. Zumbrun. A Poincaré inequality with applications to volume-constrained area-minimizing surfaces. *J. Reine Angew. Math.*, 503:63–85, 1998.
- [42] W. Walter. A theorem on elliptic differential inequalities with an application to gradient bounds. *Math. Z.*, 200(2):293–299, 1989.
- [43] S.T. Yau. Some function-theoretic properties of complete Riemannian manifold and their applications to geometry. *Indiana Univ. Math. J.*, 25(7):659–670, 1976.