

AperTO - Archivio Istituzionale Open Access dell'Università di Torino

Long-term efficacy of docosahexaenoic acid (DHA) for Spinocerebellar Ataxia 38 (SCA38) treatment: An open label extension study

This is a pre print version of the following article:

Original Citation:

Availability:

This version is available <http://hdl.handle.net/2318/1694891> since 2019-08-23T17:03:16Z

Published version:

DOI:10.1016/j.parkreldis.2019.02.040

Terms of use:

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)

1 **Long-term efficacy of Docosahexaenoic acid (DHA) for Spinocerebellar Ataxia 38**
2 **(SCA38) treatment: an open label extension study**

3 Manes M¹, Alberici A¹, Di Gregorio E^{2,3}, Boccone L⁴, Premi E¹, Mitro N⁵, Pasolini MP⁶, Pani C⁴, Paghera
4 B⁷, Orsi L⁸, Costanzi C⁹, Ferrero M³, Tempia F¹⁰, Caruso D⁵, Padovani A¹, Brusco A^{2,3}, Borroni B¹

5
6 ¹Neurology Unit, Department of Clinical and Experimental Sciences, University of Brescia, Brescia,
7 Italy

8 ²Medical Genetics Unit, Città della Salute e della Scienza University Hospital, Turin, Italy

9 ³Department of Medical Sciences University of Turin, Turin, Italy

10 ⁴Ospedale Regionale Microcitemie, AOBrotzu, Cagliari, Italy

11 ⁵Department of Pharmacological and Biomolecular Sciences, University of Milan, Milan, Italy,

12 ⁶Neurophysiology Unit, "Spedali Civili", Brescia, Italy

13 ⁷Department of Nuclear Medicine, University of Brescia, Brescia, Italy

14 ⁸Neurologic Division 1 Department of Neuroscience and Mental Health AOU Città della Salute e della
15 Scienza di Torino, Turin, Italy

16 ⁹Neurology Unit, Cremona Hospital, Cremona, Italy

17 ¹⁰Neuroscience Institute Cavalieri Ottolenghi (NICO) and Department of Neuroscience, University of
18 Turin, Turin, Italy

19

20 Corresponding author:
21 Barbara Borroni, MD
22 Centre for Ageing Brain and Neurodegenerative Disorders,
23 Neurology Unit, Department of Clinical and Experimental Sciences
24 University of Brescia
25 Piazzale Spedali Civili 1, Brescia, Italy
26 Email: bborroni@inwind.it

27
28 Number of characters in the title: xx
29 Number of words in the Abstract: xx
30 Number of words in the body of the manuscript: xxx
31 Number of figures: xx

32

33

Field Code Changed

Formatted: English (United States)

34 **Abstract**

35 **Background:** Spinocerebellar Ataxia 38 (SCA38) is caused by *ELOVL5* gene mutation, with significant
36 reduction of serum docosahexaenoic acid (DHA) levels. DHA supplementation has been proven
37 effective at short-term follow-up.

38 **Objective:** To evaluate long-term safety and efficacy of 600 mg/day oral DHA in SCA38 by a 2-year
39 open label extension study.

40 **Methods:** Nine SCA38 patients underwent standardised clinical assessment at 62 (T1), 82 (T2) and
41 104 (T3) weeks, and compared to pre-treatment scores (T0). Brain 18-Fluorodeoxyglucose Positron
42 Emission Tomography and electroneurography were performed at T0 and T3.

43 **Results:** We found a significant maintenance of clinical symptom improvement at each follow-up time-
44 point ($p<0.001$) as compared to T0, a sustained increase of cerebellar metabolism at T3 as compared
45 to T0 ($p=0.013$), and no worsening of neurophysiological parameters. No side effect was recorded.

46 **Conclusions:** Long-term DHA supplementation is an eligible treatment for SCA38.

47

48 **Introduction**

49 We have recently suggested supplementation with docosahexaenoic acid (DHA) as an effective
50 treatment in Spinocerebellar ataxias 38 (SCA38), an autosomal dominant disorders phenotypically
51 characterized by gait and limb ataxia, followed by dysarthria, dysphagia, and opthalmoparesis, and in
52 which *pes cavus* and hyposmia may be considered distinctive associated features (Borroni et al.; Manes
53 Ann Neurol).

54 SCA38 is caused by mutations in the *ELOVL5* gene (Di Gregorio), which encodes an elongase enzyme
55 involved in the synthesis of very long-chain fatty acids with a high and specific expression in Purkinje
56 cells (Tempia). The mutations lead to a reduction of serum DHA of the omega-3 polyunsaturated fatty
57 acid class (Di Gregorio).

58 Given the reduction of DHA levels in SCA38 patients, we have previously evaluated the safety and
59 efficacy of DHA supplementation, and we carried out a double-blind randomised placebo-controlled
60 study for 16 weeks, followed by an open-label study with overall 40-week DHA treatment. We
61 demonstrated that oral DHA supplementation is a safe and effective treatment, exerting clinical
62 efficacy and ameliorating cerebellar metabolism (Manes et al).

63 Here, we report the results of the subsequent long-term open-label extension study, aimed at
64 evaluating the long-term safety and efficacy of oral DHA supplementation (600 mg/day) on clinical
65 disease course, cerebellar metabolism and neurophysiological parameters up to two years follow-up.

66 **Methods**

67 This is a 104-week open-label extension study of a previous 16-week double blind, randomised,
68 placebo-controlled trial followed by a 40-week open label phase. The patients included in the previous
69 trial have been periodically evaluated at the Centre for Ageing Brain and Neurodegenerative
70 Disorders, Department of Clinical and Experimental Sciences, University of Brescia, Italy up to two-
71 years. We considered 9 patients out of 10 with the *ELOVL5* c.689G>T (p.Gly230Val) variant, as one
72 died during the study (death not related to the disease). Mean age was 48.7±10.8, and the mean age at
73 onset was 38.4±6.8; six patients were females.

74 The study was approved by the ethics committee of the Brescia Hospital, Italy (NP1821) and was
75 conformed to the Declaration of Helsinki principles. Written informed consent was obtained from all
76 patients. Trial was registered at ClinicalTrials.gov (NCT03109626).

77 During this 104-week extension, each patient received 600 mg/day b.i.d. of a fish oil derived DHA
78 (Sofedus srl, Milan, Italy). Inclusion and exclusion criteria were previously published (Manes).
79 Each patient underwent standardised clinical assessment at 62 week (T1), 82 week (T2) and 104 week
80 (T3) DHA supplementation follow-up, as compared to baseline (T0); moreover, we assessed cerebellar
81 metabolism and neurophysiological parameters at T3, as compared to T0 (see **Figure 1**, study design).
82 Standardised clinical assessment included Scale for the Assessment and Rating of Ataxia (SARA, range
83 0-40)⁸ and the International Cooperative Ataxia Rating Scale, (ICARS, range 0-100)⁹. As this was an
84 open label study, to ensure blindness in the clinical assessment scoring, at each time-point (T0, T1, T2
85 and T3) neurological examination was video-recorded and analysed in blindness by AA, who was
86 unaware of time-point, as the videos were presented randomly.

87 Cerebellar metabolism was assessed by brain 18-fluorodeoxyglucose (FDG) Positron Emission
88 Tomography (PET) scan, as previously reported (Manes). Briefly, processing and statistical analyses
89 were carried out running on MATLAB (<http://it.mathworks.com/products/matlab/>) (Mathworks Inc.,
90 Sherborn, Mass., USA) and Statistical Parametric Mapping (SPM,
91 <http://www.fil.ion.ucl.ac.uk/spm/software/SPM12/>), a fully automated, unbiased and operator
92 independent software. Cerebellar metabolism changes were evaluated by non-parametric permutation

Field Code Changed

Field Code Changed

93 test (10,000 permutation, Statistical nonParametric Mapping 13, <http://warwick.ac.uk/snpm>; T0 vs.
94 T3), and the threshold set at $p < 0.05$, Family-Wise Error (FWE) cluster-level corrected¹¹.

95 Electromyography/electroneurography (EMG/ENG) examination was performed according to
96 standard procedures.

97 To assess the effect of DHA treatment over time, we applied one-way mixed ANOVA with TIME as
98 within-subjects factors. Mauchly's test was used to test for assumption of sphericity, while
99 Greenhouse–Geisser epsilon determination was used to correct in case of sphericity violation. Results
100 were expressed as mean values \pm standard error (SE), unless otherwise specified. Statistical analyses
101 were performed using SPSS version 21 (SPSS, Inc., Chicago, IL, USA).

102 Evaluation of *ELOVL5* expression was assessed by quantitative real-time PCR on retrotranscribed
103 mRNA extracted from nine patients' PAX-gene blood at different time points of DHA supplementation
104 (see REFERENCE): 62 week (T1), 82 week (T2) and 104-week (T3). The mean and S.E. was calculated
105 and normalised to baseline (T0). Data analysis was performed by GraphPad using a second degree
106 polynomial regression curve.

Field Code Changed

107 **Results**

108 Nine patients completed the 104-week follow-up. No discontinuations occurred, and no side effects
109 were reported during DHA supplementation at established dosage.

110 We found a significant maintenance of improvement of clinical symptoms over time. Repeated measures
111 ANOVA performed on total SARA score revealed a main effect of time ($F(3,24)=9.82, p<0.001$, partial η^2
112 $=0.55$). Post-hoc analysis showed a significant reduction of SARA score values at T1 ($6.0\pm 1.4, p=0.001$),
113 T2 ($6.4\pm 1.8, p=0.01$) and T3 ($6.4\pm 1.8, p=0.013$) compared to baseline (8.9 ± 1.4) (**Figure 2, panel A**). No
114 significant differences among T1, T2 and T3 time-points were reported.

115 The same main effect of time was shown for ICARS score ($F(1.35,10.86)=8.07, p=0.012$, partial $\eta^2 =0.50$).
116 Post-hoc analysis showed a significant reduction of ICARS score values at T1 ($13.1\pm 3.9, p=0.008$), T2
117 ($13.0\pm 4.0, p=0.012$) and T3 ($13.8\pm 4.3, p<0.001$) compared to baseline (18.7 ± 4.3) (**Figure 2, panel A**).
118 No significant differences among T1, T2 and T3 time-points were reported.

119 A significant difference in cerebellar metabolism between T0 and T3 was observed, with a sustained
120 significant increase of cerebellar metabolism at T3 as compared to T0 in the left exterior cerebellar lobe
121 ($x,y,z= -44, -60, -26; T=10,26; p=0.013$, cluster size=227) (see **Figure 2, panel B**). Importantly, no
122 significant differences in the opposite contrast (T0>T3) were found at the pre-established threshold,
123 ruling out worsening of regional cerebellar metabolism after 104-week DHA treatment.

124 At EMG/ENG, motor and sensory conduction velocities did not significantly worsen during 104-week
125 DHA treatment.

126 We showed a slight but significant reduction of *ELOVL5* expression in blood comparing each time point
127 with baseline, although a high degree of variability was present. An increased expression was also

128 present in all cases at T3 (r^2 : 0.8989)(Figure 3).**Discussion**

129 In a double-blind, randomised, placebo-controlled study followed by a short-term open label phase, we
130 have previously demonstrated that oral DHA is a safe and effective treatment for SCA38 patients
131 (Manes). However, we did not know whether the beneficial effect of this treatment might be extended
132 over time, with no long-term side effects and with no tolerance mechanisms occurring.

133 In the present study, by a long-term follow-up, evaluating patients up to 2 years, we have
134 demonstrated that oral DHA at a dosage of 600 mg/day is the eligible treatment for SCA38. By a
135 comprehensive assessment, we reported that DHA was able to stabilize clinical symptoms, as
136 measured by SARA and ICARS scales, to maintain the significant amelioration of cerebellar
137 metabolism, and to avoid the worsening of neurophysiological measures. Moreover, we have further
138 proven that DHA supplementation was well tolerated, affordable and easy to administer dietary
139 intervention, with no side effect and with a relatively low cost. Clinical findings were in line with the
140 slightly decreased expression of *ELOVL5* in patients' blood. We also noted a final increment of *ELOVL5*
141 expression, which might suggest a compensatory mechanism after long-term administration and
142 possibly an adjustment of the therapy if prolonged for many years..

143 The use of DHA supplementation in SCA38 stemmed from the initial observation that *ELOVL5*
144 mutations lead to an increased amount of the encoded protein with a mislocalization of the aberrant
145 form in the perinuclear area instead of endoplasmic reticulum and by a decrease of its final products,
146 in particular DHA, in patients' serum (Di Gregorio et al., 2014). Thus, DHA supplementation can act by
147 compensating the decrease of very long chain fatty acids and by lowering *ELOVL5* aberrant protein via
148 a transcriptional feedback loop.

149 Given the open-label study design, we tried to minimize possible biases in the clinical evaluation
150 scoring, by video-recording and randomising the clinical assessments and making them evaluating by
151 a neurologist who was unaware of the follow-up time-points. Furthermore, we used an unbiased,

Commented [s1]: Questa può anche essere una figura
supplementare

152 automated, and operator independent software for analysing imaging data, which allowed us to
153 exclude regional worsening of cerebellar metabolism at 2 year follow-up.
154 Even though carried out in a small sample of patients, this study suggests that DHA supplementation is
155 an effective treatment approach in SCA38. Its intake for the whole lifewould be beneficial but now we
156 have no sufficient data to exclude the development of tolerance associated to a prolonged
157 administration. Based on our observation, we might speculate that DHA treatment is even beneficial in
158 the presymptomatic stage of the disease to delay disease onset and slow the progression of symptoms.

159

160 **Acknowledgement**

161 The authors are indebted to the patients and their families for taking part into the study, to Dr. Stefano
162 Gazzina for imaging analyses, and to Dr. Maura Cosseddu for technical support. This work was
163 supported by the Fondazione Telethon grant number GGP14225.

164

165

166 **Author Contributions**

167 B.A., and B.B. contributed to the concept and study design. M.M., A.A., D.G.E., B.L., P.E., M.N., P.M.P., P.C.,
168 P.B., O.L., C.C., F.M., T.F., C.D., P.A., B.A., B.B. contributed to the data acquisition and analysis. M.M., A.A.,
169 B.A., and B.B. drafted the manuscript and figures, and all authors approved the final version.

170

171 **Potential Conflicts of Interest**

172 The authors do not have any potential conflict of interest.

173

174

175

176

177

178

179

180 **References**

- 181 1. Ruano L, Melo C, Silva MC, Coutinho P. The global epidemiology of hereditary ataxia and spastic
182 paraplegia: a systematic review of prevalence studies. *Neuroepidemiology*. 2014;42(3):174-83.
- 183 2. Durr A. Autosomal dominant cerebellar ataxias: polyglutamine expansions and beyond. *Lancet*
184 *Neurol*. 2010 Sep;9(9):885-94.
- 185 3. van Gaalen J, Giunti P, van de Warrenburg BP. Movement disorders in spinocerebellar ataxias.
186 *Mov Disord*. 2011 Apr;26(5):792-800.
- 187 4. Sun YM, Lu C, Wu ZY. Spinocerebellar ataxia: relationship between phenotype and genotype - a
188 review. *Clin Genet*. 2016 Oct;90(4):305-14.
- 189 5. Di Gregorio E, Borroni B, Giorgio E, et al. ELOVL5 mutations cause spinocerebellar ataxia 38.
190 *Am J Hum Genet*. 2014 Aug 07;95(2):209-17.
- 191 6. Borroni B, Di Gregorio E, Orsi L, et al. Clinical and neuroradiological features of spinocerebellar
192 ataxia 38 (SCA38). *Parkinsonism Relat Disord*. 2016 Jul;28:80-6.
- 193 7. Lien EL. Toxicology and safety of DHA. *Prostaglandins Leukot Essent Fatty Acids*. 2009 Aug-
194 Sep;81(2-3):125-32.
- 195 8. Yabe I, Matsushima M, Soma H, Basri R, Sasaki H. Usefulness of the Scale for Assessment and
196 Rating of Ataxia (SARA). *J Neurol Sci*. 2008 Mar 15;266(1-2):164-6.
- 197 9. Trouillas P, Takayanagi T, Hallett M, et al. International Cooperative Ataxia Rating Scale for
198 pharmacological assessment of the cerebellar syndrome. The Ataxia Neuropharmacology Committee
199 of the World Federation of Neurology. *J Neurol Sci*. 1997 Feb 12;145(2):205-11.
- 200 10. Della Rosa PA, Cerami C, Gallivanone F, et al. A standardized [18F]-FDG-PET template for
201 spatial normalization in statistical parametric mapping of dementia. *Neuroinformatics*. 2014
202 Oct;12(4):575-93.
- 203 11. Nichols TE, Holmes AP. Nonparametric permutation tests for functional neuroimaging: a
204 primer with examples. *Hum Brain Mapp*. 2002 Jan;15(1):1-25.
- 205 12. Cooper JM, Korlipara LV, Hart PE, Bradley JL, Schapira AH. Coenzyme Q10 and vitamin E
206 deficiency in Friedreich's ataxia: predictor of efficacy of vitamin E and coenzyme Q10 therapy. *Eur J*
207 *Neurol*. 2008 Dec;15(12):1371-9.

- 208 13. Cooper JM, Schapira AH. Friedreich's ataxia: coenzyme Q10 and vitamin E therapy.
209 Mitochondrion. 2007 Jun;7 Suppl:S127-35.
- 210 14. Moon YA, Hammer RE, Horton JD. Deletion of ELOVL5 leads to fatty liver through activation of
211 SREBP-1c in mice. J Lipid Res. 2009 Mar;50(3):412-23.
- 212 15. Demar JC, Jr., Ma K, Chang L, Bell JM, Rapoport SI. alpha-Linolenic acid does not contribute
213 appreciably to docosahexaenoic acid within brain phospholipids of adult rats fed a diet enriched in
214 docosahexaenoic acid. J Neurochem. 2005 Aug;94(4):1063-76.
- 215 16. Igarashi M, Ma K, Chang L, Bell JM, Rapoport SI, DeMar JC, Jr. Low liver conversion rate of
216 alpha-linolenic to docosahexaenoic acid in awake rats on a high-docosahexaenoate-containing diet. J
217 Lipid Res. 2006 Aug;47(8):1812-22.
- 218 17. Rapoport SI, Chang MC, Spector AA. Delivery and turnover of plasma-derived essential PUFAs
219 in mammalian brain. J Lipid Res. 2001 May;42(5):678-85.
- 220 18. Rapoport SI, Rao JS, Igarashi M. Brain metabolism of nutritionally essential polyunsaturated
221 fatty acids depends on both the diet and the liver. Prostaglandins Leukot Essent Fatty Acids. 2007 Nov-
222 Dec;77(5-6):251-61.
- 223 19. Nguyen LN, Ma D, Shui G, et al. Mfsd2a is a transporter for the essential omega-3 fatty acid
224 docosahexaenoic acid. Nature. 2014 May 22;509(7501):503-6.
- 225 20. Hacioglu G, Seval-Celik Y, Tanriover G, et al. Docosahexaenoic acid provides protective
226 mechanism in bilaterally MPTP-lesioned rat model of Parkinson's disease. Folia Histochem Cytobiol.
227 2012 Jul 04;50(2):228-38.
- 228 21. Yassine HN, Braskie MN, Mack WJ, et al. Association of Docosahexaenoic Acid Supplementation
229 With Alzheimer Disease Stage in Apolipoprotein E epsilon4 Carriers: A Review. JAMA Neurol. 2017
230 Mar 01;74(3):339-47.
- 231 22. Bourre JM. Effects of nutrients (in food) on the structure and function of the nervous system:
232 update on dietary requirements for brain. Part 1: micronutrients. J Nutr Health Aging. 2006 Sep-
233 Oct;10(5):377-85.

- 234 23. Bazan NG. Neuroprotectin D1-mediated anti-inflammatory and survival signaling in stroke,
235 retinal degenerations, and Alzheimer's disease. *J Lipid Res.* 2009 Apr;50 Suppl:S400-5.
- 236 24. Senior K. Lorenzo's oil may help to prevent ALD symptoms. *Lancet Neurol.* 2002 Dec;1(8):468.
- 237 25. Eder K, Kish SJ, Kirchgessner M, Ross BM. Brain phospholipids and fatty acids in Friedreich's
238 ataxia and spinocerebellar atrophy type-1. *Mov Disord.* 1998 Sep;13(5):813-9.
- 239

240 **Legend to figures.**

241

242 **Figure 1. Study design of the open-label extension study with DHA supplementation in SCA38.**

243 Black blocks indicate the time points of clinical assessment; black arrows indicated the time points of
244 brain 18-fluorodeoxyglucose Positron Emission Tomography (FDG-PET) and electroneurography
245 (ENG). T0: baseline; T1: 62 weeks follow-up; T2: 82 weeks follow-up; T3: 104 weeks follow-up. T*:
246 previously published open label follow-up study (Manes et al., 2017) at 40 weeks follow-up.
247 wk: weeks.

248

249 **Figure 2. Clinical assessment and cerebellar metabolism in the open-label extension study.**

250 **Panel A.** SARA and ICARS scores at baseline (T0), after 62 weeks (T1), 82 weeks (T2) and 104 weeks
251 (T3) DHA treatment in SCA38 patients.

252 *p-value<0.05.

253 **Panel B.** Pattern of cerebellar metabolism in SCA38 patients at T0 vs. T3 (T0<T3) (p<0.05, FWE corrected).

254 See results for details. The results are superimposed on a 3D standardized template template.

255 P: posterior, L: left; R: right.

256

257 **Figure 3.** *ELOVL5* expression analysis in DHA-treated patients' blood. Expression was evaluated by

258 Quantitative real-time PCR on retrotranscribed mRNA extracted from PAX-gene blood at different time points

259 (62 week (T1), 82 week (T2) and 104 week (T3)). *ELOVL5* was normalized against *TBP* gene. The average

260 value and S.E. is shown. Statistical analysis showed the data had a best fit with a second degree polynomial

261 regression curve ($r^2= 0.8989$), with an initial reduction of expression and a final increase to levels close to

262 baseline.

263