AperTO - Archivio Istituzionale Open Access dell'Università di Torino

Abstract 5620: Novel phage display-derived neuroblastoma-targeting peptides potentiate the effect of drug nanocarriers in preclinical settings

This is the author's manuscript

Original Citation:

Availability:

This version is available <http://hdl.handle.net/2318/1695386> since 2019-03-25T14:26:50Z

Published version:

DOI:10.1158/1538-7445.AM2013-5620

Terms of use:

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)

Cancer Research

Search...

[Advanced Search](#)[Home](#)[About](#)[Articles](#)[For Authors](#)[Alerts](#)[News](#)

Experimental and Molecular Therapeutics

Abstract 5620: Novel phage display-derived neuroblastoma-targeting peptides potentiate the effect of drug nanocarriers in preclinical settings.

Monica Loi, Daniela Di Paolo, Marco Soster, Chiara Brignole, Alice Bartolini, Laura Emionite, Jessica Sun, Pamela Becherini, Flavio Curnis, Andrea Petretto, Monica Sani, Alessandro Gori, Claudio Gambini, Renato Longhi, Michele Cilli, Theresa M. Allen, Federico Bussolino, Wadih Arap, Renata Pasqualini, Angelo Corti, Mirco Ponzoni, Serena Marchiò, and Fabio Pastorino

DOI: 10.1158/1538-7445.AM2013-5620 Published April 2013

[Article](#)[Info & Metrics](#)

Proceedings: AACR 104th Annual Meeting 2013; Apr 6-10, 2013; Washington, DC

Abstract

Purpose: Molecular targeting of drug delivery nanocarriers is expected to improve their therapeutic index while decreasing their toxicity. The identification of novel peptide ligands specific for cells present in high-risk neuroblastoma, a childhood tumor mostly refractory to current therapies, is needed.

Experimental design: We performed combined in vitro/ex-vivo phage display screenings on human neuroblastoma cell lines and on tumors derived from orthotopic mouse models of human neuroblastoma. Binding validation and homing in vivo of selected phage clones were tested by immunohistochemistry/immunofluorescent analyses. Cell association experiments in vitro with the corresponding synthetic biotin-labeled peptides were performed. In vitro cytotoxicity and in vivo tumor

April 2013
Volume 73,
Issue 8
Supplement
[Table of Contents](#)
[Index by Author](#)

Search this issue

[Sign up for alerts](#)[© Permissions](#)[! Article Alerts](#)[✉ Email Article](#)[🌐 Citation Tools](#)[↪ Share](#)[🐦 Tweet](#)

Advertisement

[▼ Related Articles](#)

accumulation and therapeutic experiments were

Jump to section

peptide-targeted, doxorubicin-loaded,

Article

Info & Metrics

ining proper subtractive protocols, we

identified phage clones specific either for the primary tumor, its metastases, or for the stromal components. Globally, we isolated 121 phage-displayed neuroblastoma-binding peptides; of these, 26 bound the primary tumor, 15 the metastatic mass, 57 and 23 their respective microenvironments. Of these, five phage clones were further validated for their specific binding ex-vivo to biopsies from stage IV neuroblastoma patients and to neuroblastoma tumors derived from mice. All five clones also targeted tumor cells and vasculature in vivo when injected into neuroblastoma-bearing mice. Coupling of the corresponding targeting peptides with doxorubicin-loaded nanocarriers led to a significant inhibition in tumor volume and enhanced survival in preclinical neuroblastoma models.

Conclusions: Our findings demonstrate that novel ligands of neuroblastoma-associated markers are functional in the design of nanocarriers with therapeutic efficacy paving the way to their clinical development.

Citation Format: Monica Loi, Daniela Di Paolo, Marco Soster, Chiara Brignole, Alice Bartolini, Laura Emionite, Jessica Sun, Pamela Becherini, Flavio Curnis, Andrea Petretto, Monica Sani, Alessandro Gori, Claudio Gambini, Renato Longhi, Michele Cilli, Theresa M. Allen, Federico Bussolino, Wadih Arap, Renata Pasqualini, Angelo Corti, Mirco Ponzoni, Serena Marchiò, Fabio Pastorino. Novel phage display-derived neuroblastoma-targeting peptides potentiate the effect of drug nanocarriers in preclinical settings. [abstract]. In: Proceedings of the 104th Annual Meeting of the American Association for Cancer Research; 2013 Apr 6-10; Washington, DC. Philadelphia (PA): AACR; Cancer Res 2013;73(8 Suppl):Abstract nr 5620. doi:10.1158/1538-7445.AM2013-5620

©2013 American Association for Cancer Research

No related articles found.

Google Scholar

► Cited By...

► More in this TOC Section

^ Back to top