

AperTO - Archivio Istituzionale Open Access dell'Università di Torino

In vitro Digestibility Measurement of Feedstuffs in Donkeys Using the DaisyII Incubator

This is the author's manuscript

Original Citation:

Availability:

This version is available <http://hdl.handle.net/2318/1702983> since 2019-05-28T10:47:22Z

Published version:

DOI:<https://doi.org/10.1016/j.jevs.2019.02.002>

Terms of use:

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)

Manuscript Details

Manuscript number	JEVS_2018_283_R2
Title	In vitro Digestibility Measurement of Feedstuffs in Donkeys Using the DaisyII Incubator
Article type	Research paper

Abstract

Successful studies on in vitro digestibility measurement of feedstuffs with fecal inoculum have been reported for horses, while data on donkeys are currently lacking. In this study we evaluated the use of the DaisyII Incubator (Ankom Technology, Fairport, NY, USA) for in vitro digestibility measurement of feedstuffs using donkey feces as source of microbial inoculum. The method was tested using 7 feedstuffs commonly used in donkey diets (alfalfa, bromegrass, ryegrass and timothy hays; wheat bran and wheat straw; barley grains). Feces were obtained from 4 female donkeys and incubations were carried out at one-week intervals for 4 consecutive weeks. Two bags of each feedstuff were incubated in digestion vessels containing a buffer/feces solution (90:10). In vitro apparent dry matter digestibility (DMD), true dry matter digestibility (IVTD), and neutral detergent fiber digestibility (NDFD) were evaluated at 4 incubation times: 30, 48, 60, and 72 h. All digestibility parameters significantly increased from 30 to 72 h of incubation. At 72 h of incubation, the within-laboratory repeatability and reproducibility of the method were 2.7% and 5.0% for DMD and 1.6% and 3.9% for IVTD, respectively. The method was less repeatable and reproducible for NDFD (4.5% and 10.4%, respectively).

Keywords	Equus asinus L.; fecal inoculum; DaisyII Incubator; measurement error; incubation time.
Corresponding Author	Manuela Renna
Order of Authors	Sonia Tassone, Manuela Renna, Salvatore Barbera, EMANUELA VALLE, Riccardo Fortina
Suggested reviewers	Jennifer E. Earing, Mauro Spanghero, Barbara Padalino, Valentina Ragno, Eric Davis

Submission Files Included in this PDF

File Name [File Type]

Cover letter.docx [Cover Letter]
Response to reviewers.docx [Response to Reviewers]
Highlights.doc [Highlights]
Manuscript_R2.docx [Manuscript File]
Figure 1.docx [Figure]
Figure 2.docx [Figure]
Figure captions.docx [Figure]
Table 1.docx [Table]
Table 2.docx [Table]
Conflict of interest statement.doc [Conflict of Interest]

To view all the submission files, including those not included in the PDF, click on the manuscript title on your EVISE Homepage, then click 'Download zip file'.



UNIVERSITÀ DEGLI STUDI DI TORINO

DIPARTIMENTO DI SCIENZE VETERINARIE



Dr. Manuela Renna

Department of Veterinary Sciences

Largo Paolo Braccini, 2 – 10095 Grugliasco, Turin

University of Turin

Italy

Telephone: +39-011.6709053

Email: manuela.renna@unito.it

To Dr. Edward L. Squires
Editor-in-Chief
Journal of Equine Veterinary Science

Re: Revision Requested - JEVS_2018_283_R1 for Journal of Equine Veterinary Science

Dear Dr. Edward L. Squires,

I have read carefully your letter dated January 28th, 2019 and the comments from reviewer 1 regarding our manuscript. I prepared a revised version of the paper according to these comments. All changes made to the R1 version of the manuscript have been highlighted in red font. I hope this modified version will fully satisfy the reviewers and the Editor, and match the standards of the *Journal of Equine Veterinary Science*.

The comments of the reviewer have been addressed one by one, and are listed in the “Response to reviewers” file.

Yours sincerely, Dr. Manuela Renna.

Largo Braccini 2 – 10095 Grugliasco To

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59

RESPONSE TO REVIEWERS

Reviewer 1

This reviewer congratulates the authors on a well written and detailed manuscript. Thank you for thoroughly addressing all comments and for your effort in adding new sections and references. In this reviewer's opinion, the authors' suggestion to change the manuscript from short communication to original research article should be accommodated.

Please see below my minor comments to this version of the manuscript.

AU Response: Thank you for the positive evaluation of the revised version of our manuscript.

Line 109: doubled weighed

AU Response: Changed to: "weighed in duplicate".

Line 227: improve "when" extending

AU Response: Corrected.

Line 237: provides

AU Response: The sentence has been modified.

Lines 238-239: "the absolute difference... will differ" I suggest trying to rephrase and shorten this sentence for clarity

AU Response: We rewrote the sentence as follows: "The estimated repeatability and reproducibility coefficients (Table 2b) are precision measures providing additional information, as they represent the maximum absolute difference that can be expected between repeated measurements on 95% of occasions [17]."

Lines 244-248: since conclusions (i) and (iii) appear to overlap, could they be condensed into one?

AU Response: We rewrote the sentence as follows: "From the obtained results, it can be concluded that an incubation period of 72 h is required to obtain accurate DMD, IVTD, and NDFD estimates in donkeys and to improve the repeatability and reproducibility of the method. For NDFD, the repeatability and reproducibility of the method were comparable to those obtained with the Daisy^{II} Incubator in ruminants."

Reviewer 2

The revision of the manuscript is satisfactory.

AU Response: Thank you for the positive evaluation of the revised version of our manuscript.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59

Highlights

- Daisy^{II} Incubator can be successfully used for digestibility studies in donkeys
- NDFD had lower repeatability and reliability than DMD and IVTD
- Digestibility increased significantly from 30 to 72 h of incubation

1
2
3 1 ***In vitro* Digestibility Measurement of Feedstuffs in Donkeys Using the Daisy^{II} Incubator**
4
5
6 2

7 3 Sonia Tassone ^a, Manuela Renna ^{b, *}, Salvatore Barbera ^a, Emanuela Valle ^b, Riccardo Fortina ^a
8
9
10 4

11
12 5 ^a Department of Agricultural, Forest and Food Sciences, University of Torino, Largo P. Braccini 2,
13
14 6 10095 Grugliasco (TO), Italy

15
16 7 ^b Department of Veterinary Science, University of Torino, Largo P. Braccini 2, 10095 Grugliasco
17
18 8 (TO), Italy
19
20 9

21
22 10 Corresponding author: Dr. Manuela Renna, Department of Veterinary Science, University of
23
24 11 Torino, Largo P. Braccini 2, 10095 Grugliasco (TO), Italy – Tel. +39.011.6709053 – Fax:
25
26 12 +39.011.6708563 - Email: manuela.renna@unito.it
27
28
29 13

30
31 14 E-mail addresses:

32
33 15 ST: sonia.tassone@unito.it
34

35 16 MR: manuela.renna@unito.it
36

37 17 SB: salvatore.barbera@unito.it
38

39 18 EV: emanuela.valle@unito.it
40

41 19 RF: riccardo.fortina@unito.it
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

61
62
63 **21 Abstract**
64

65 22 Successful studies on *in vitro* digestibility measurement of feedstuffs with fecal *inoculum* have been
66
67 23 reported for horses, while data on donkeys are currently lacking. In this study we evaluated the use
68
69 24 of the Daisy^{II} Incubator (Ankom Technology, Fairport, NY, USA) for *in vitro* digestibility
70
71 25 measurement of feedstuffs using donkey feces as source of microbial *inoculum*. The method was
72
73 26 tested using 7 feedstuffs commonly used in donkey diets (alfalfa, bromegrass, ryegrass and timothy
74
75 27 hays; wheat bran and wheat straw; barley grains). Feces were obtained from 4 female donkeys and
76
77 28 incubations were carried out at one-week intervals for 4 consecutive weeks. Two bags of each
78
79 29 feedstuff were incubated in digestion vessels containing a buffer/feces solution (90:10). *In vitro*
80
81 30 apparent dry matter digestibility (DMD), true dry matter digestibility (IVTD), and neutral detergent
82
83 31 fiber digestibility (NDFD) were evaluated at 4 incubation times: 30, 48, 60, and 72 h. All
84
85 32 digestibility parameters significantly increased from 30 to 72 h of incubation. At 72 h of incubation,
86
87 33 the within-laboratory repeatability and reproducibility of the method were 2.7% and 5.0% for DMD
88
89 34 and 1.6% and 3.9% for IVTD, respectively. The method was less repeatable and reproducible for
90
91 35 NDFD (4.5% and 10.4%, respectively).
92
93
94
95
96

97 37 **Key words:** *Equus asinus* L.; fecal *inoculum*; Daisy^{II} Incubator; measurement error; incubation
98
99 38 time.
100

101 39
102
103

104 40
105

106 41
107

108 42
109

110 43
111

112 44
113

114 45
115

116 46
117
118
119
120

121
122
123 **47 1. Introduction**
124

125 48 The use of *in vitro* techniques to study diet digestion and fermentative end products has been
126
127
128 49 successfully applied in horse nutrition. Lowman et al. [1] was the first to demonstrate that equine
129
130 50 feces can be used as source of microbial *inoculum* and that fecal microflora can remain viable for
131
132 51 several hours after excretion. According to Abdouli et al. [2], a two-stage *in vitro* technique
133
134 52 (combining enzymatic pre-digestion and fermentation) can estimate the organic matter digestibility
135
136 53 of both forages and concentrates using horse feces as source of microbial *inoculum*. Ringler et al.
137
138 54 [3] demonstrated that the use of equine fecal *inoculum* in a closed-system fermentation apparatus
139
140
141 55 can yield valid *in vitro* estimates of dry matter (DM) and fiber digestibility. Lattimer et al. [4]
142
143 56 reported that the Daisy^{II} Incubator could be used to predict valid estimates of DM digestibility of
144
145 57 diets using equine feces as *in vitro inoculum* source. Earing et al. [5] confirmed that feces are a
146
147 58 suitable source of microbial *inoculum* for *in vitro* digestibility studies in horses. More recently,
148
149 59 Murray et al. [6] compared samples from different sites of the equine intestine, showing that feces
150
151
152 60 are a suitable alternative to cecal fluid, thus highlighting the potential of this kind of *inoculum* for *in*
153
154 61 *vitro* digestibility studies.
155

156 62 Very few studies are currently available on digestibility measurement of feedstuffs in donkeys,
157
158 63 most of the data resulting from *in vivo* feeding trials [7-9]. *In vitro* digestibility in donkeys was
159
160 64 studied using the neutral cellulase plus gamanase technique developed by Ankom Technology
161
162 65 (Fairport, NY, USA) and via fecal-Near Infrared Spectroscopy [9-11], but with poor success.
163
164
165 66 Since the breeding of lactating donkeys in Western Europe is increasing [12], it becomes needful to
166
167 67 evaluate tools to study the *in vitro* digestibility of feedstuffs used in donkey nutrition. Therefore, the
168
169 68 objectives of this study were (i) to analyse, at different incubation times, the *in vitro* digestibility
170
171 69 measurement of feedstuffs in donkeys using the Daisy^{II} Incubator and donkey feces as source of
172
173 70 microbial *inoculum*, and (ii) to test the within-laboratory repeatability and reproducibility of the
174
175 71 method.
176
177
178
179
180

181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240

72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97

2. Materials and Methods

The protocol was designed according to the guidelines of the current European Directive (2010/63/EU) on the care and protection of animals used for scientific purposes.

Ethical approval for this study was obtained from the Animal Welfare Committee of the Department of Veterinary Science of the University of Torino (Italy).

2.1. Feedstuffs

Seven feedstuffs (alfalfa hay, bromegrass hay, ryegrass hay, timothy hay, wheat bran, wheat straw, and barley grains) were chosen for the trial. Hays and straws are commonly used as forages in donkey nutrition; wheat bran and flaked barley are fed to working and lactating donkeys to increase the palatability and/or energy density of diet [13,14].

Feedstuffs were analysed for their proximate constituents and fiber fractions following the procedures reported by Fortina et al. [15].

2.2. Animals and diet

The feces were collected from 4 healthy (Body Condition Score = 5) female Ragusana donkeys (live weight: 193 ± 24 kg) fed with a constant ration during the whole trial. The ration was based on first cut meadow hay given *ad libitum* [DM 909 g/kg; ash 79 g/kg DM; crude protein (CP) 68 g/kg DM; ether extract 9 g/kg DM; neutral detergent fiber (NDF) 550 g/kg DM; acid detergent fiber (ADF) 313 g/kg DM; lignin 38 g/kg DM]. The donkeys were individually housed in 3×4 m covered pens at the experimental facility of the Department of Veterinary Science, University of Torino. Wood chips were used for bedding and fresh water was available at all times. On a daily basis, the donkeys were turned out in pairs into a sandy paddock, allowing free movement and socialization.

2.3. *In vitro* digestibility

For feces collection and *in vitro* analyses, the procedures described by Earing et al. [5] were followed. Approximately 250 g of feces were collected directly from the rectum of each donkey on a weekly basis, for 4 consecutive weeks (i.e. 4 runs), always at the same hour in the morning. Immediately after collection, the feces were sealed in individual airtight bags expelling air as much as possible to maintain anaerobic conditions [4,5] and transported to the laboratory (approximately 3 min travel from the barn to the laboratory). During transport, the temperature of the feces was maintained at about 39°C using a warm water-containing cooler. At the laboratory, the feces of the 4 donkeys were pooled and a mixture of them was used for the analysis.

The *in vitro* digestibility measures were performed using the Ankom Daisy^{II} Incubator (Ankom Technology Corporation, Fairport, USA). Feed samples (approximately 0.50 g) were weighed in duplicate into Ankom F57 filter bags, which were then heat sealed and put into each of the four vessels with 1800 mL of a mixture (5:1 ratio) of two buffer solutions (A and B), prepared as described by Ankom Technology [16]. About 200 g of feces per vessel were weighed and transferred to a blender jar with 400 mL of the warmed buffer solution. The content was purged with CO₂ for 15 s and blended for 30 s using a high-speed blender (Osterizer cyclo trol eight, Oster, Moncalieri (TO), Italy). The blended content was transferred to the digestion jar and gently mixed. The pH of the mixture was adjusted to 7.0 using additional amounts of the two (unmixed) buffers (A or B) as suggested by Earing et al. [5].

Considering previous experiments on horses [3-5], 4 incubation times were assigned to the vessels: 30, 48, 60, and 72 h. The shortest incubation time (30 h) was included considering that the mean retention time of the gastrointestinal tract in horses ranges from 18-20 h [8] to 45-60 h [7]. As donkeys retain feeds for longer time than other equids [7], the 72 h incubation time was chosen trying to ensure complete nutrient digestion [5].

301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360

123 At the end of the assigned time, the bags were removed from each vessel, rinsed thoroughly with
124 cold tap water and placed in a 50 °C forced-air oven to dry for 24 h. The bags were weighed and
125 then analysed for their NDF content with the Ankom²⁰⁰ Fiber Analyzer (Ankom Technology).

126 The obtained data were used to calculate:

127 - *in vitro* apparent DM digestibility (DMD, g/kg DM) = $1000 \times (DM_{0h} - DM_{residue}) / DM_{0h}$

128 - *in vitro* true digestibility (IVTD, g/kg DM) = $1000 \times (DM_{0h} - NDF_{residue}) / DM_{0h}$

129 - *in vitro* NDF digestibility (NDFD, g/kg NDF) = $1000 \times (NDF_{0h} - NDF_{residue}) / NDF_{0h}$.

130 To test the within-laboratory repeatability of the method, two replicates were carried out for each
131 sample. The within-laboratory reproducibility of the method tested instead the variation in the
132 measurements under changing conditions (i.e. the 4 runs) [17].

134 *2.4. Statistical analysis*

135 Data were analysed using SAS [18].

136 A mixed model repeated measures analysis [18] was carried out, considering incubation time as
137 repeated measure, the effects of run, incubation time and their interaction, and the random effect of
138 feedstuff [19]. Different covariance structures were fitted and, considering the Schwarz Bayesian
139 Information Criterion, the autoregressive structure was chosen [19] to account for intra-feedstuff
140 correlation.

141 For each incubation time, the repeatability and reproducibility of the method were calculated as
142 reported by Spanghero et al. [20]. For this purpose, a generalized linear model (GLM) was applied
143 according to the following equation:

144 $Y_{ijk} = \mu + \alpha_i + \beta_j + (\alpha\beta)_{ij} + \varepsilon_{ijk}$,

145 where: μ = overall mean; α = effect of run (i = 1,4); β = effect of feedstuff (j = 1,7); ε = residual
146 error.

147 The standard deviation (SD) of repeatability and reproducibility were calculated as:

361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420

148 $SD_repeatability (S_r) = \sqrt{\sigma_e^2}$

149 $SD_reproducibility (S_R) = \sqrt{\left\{ \left(\frac{\sigma_\alpha^2 - \sigma_{\alpha\beta}^2}{7 \times 2} \right) + \left(\frac{\sigma_{\alpha\beta}^2 - \sigma_e^2}{2} \right) + \sigma_e^2 \right\}}$,

150 where σ_α^2 , $\sigma_{\alpha\beta}^2$, and σ_e^2 are the variance components of run effect, interaction between run and
151 feedstuff effects, and error, respectively. Repeatability and reproducibility were then expressed as
152 coefficients of variation: $SD / \text{mean} \times 100$ [20].

153 In addition, the repeatability and reproducibility coefficients were also estimated, as reported by
154 Bartlett and Frost [17]:

- 155 • repeatability coefficient = $1.96 \times \sqrt{2} \times S_r$
- 156 • reproducibility coefficient = $1.96 \times \sqrt{2} \times S_R$.

157 A linear regression analysis for incubation time was used to obtain, for each feedstuff, the
158 regression parameters (intercept and slope) of each *in vitro* digestion (4 runs \times 7 feedstuffs).

159 Regression parameters were analysed by Canonical Discriminant Analysis (CDA), which finds
160 linear combinations (canonical functions) of the quantitative variables that provide maximal
161 separation between classes or groups [17].

162 Significance was set at $P < 0.05$.

163 3. Results

164 3.1. Effect of incubation time on estimates of DMD, IVTD, and NDFD

165 The chosen feedstuffs were characterized by a wide range of CP (66-166 g/kg DM), NDF (213-797
166 g/kg DM), ADF (69-550 g/kg DM), and lignin (12-100 g/kg DM) contents (Table 1).

167 The mixed model repeated measure analysis showed that DMD, IVTD, and NDFD were not
168 influenced by run ($P > 0.05$). Incubation time significantly affected DMD, IVTD and NDFD (Table
169 2a). All considered digestibility parameters ranked according to the following order: 72 h $>$ 60 h $>$
170 48 h $>$ 30 h. For each feedstuff, DMD and NDFD trends over incubation time are shown in Fig. 1.

421
422
423 173 DMD (Fig. 1a) and NDFD (Fig. 1b) of hays and straw were lower than that of barley grains. Straw
424
425 174 showed the absolute lowest DMD values (Fig. 1a). Wheat bran and ryegrass hay showed the highest
426
427 NDFD values at 72 h, while alfalfa hay and wheat straw had the lowest; bromegrass hay, timothy
428 175
429 hay, and barley grains showed intermediate values at this incubation time (Fig. 1b).
430 176

431 432 177 433 434 178 *3.2. Repeatability and reproducibility of the method*

435
436 179 For each incubation time, the repeatability and reproducibility of the method are reported in Table
437
438 180 2b. Repeatability and reproducibility, expressed as coefficients of variation, ranged 2.7 - 3.8% and
439
440 181 5.0 - 10.3% for DMD, 1.6 - 1.8% and 2.8 - 5.4% for IVTD, and 4.5 - 11.6% and 10.4 - 18.5% for
441
442 182 NDFD, respectively. Particularly for NDFD, both repeatability and reproducibility improved from
443
444 183 30 to 72 h of incubation.

445
446
447 184 The results of the multivariate statistical analysis are shown in Fig. 2. The regression parameters
448
449 185 (intercept and slope) describe each *in vitro* digestion (4 runs \times 7 feedstuffs) and Fig. 2, for IVTD as
450
451 186 an example, shows the separation among the different feedstuffs by CDA.

452 453 187 454 455 188 **4. Discussion**

456 457 189 458 459 190 *4.1. Effect of incubation time on estimates of DMD, IVTD, and NDFD*

460
461 191 As expected, estimates of DMD, IVTD and NDFD increased while increasing the incubation time
462
463 192 (Table 2a). The obtained results indicate that an incubation period of 72 h or higher may be required
464
465 193 for accurate DMD, IVTD, and NDFD estimates in donkeys. Similar findings were recently obtained
466
467 194 by Franzan et al. [21], studying DM, organic matter, and NDF degradation using horse feces as
468
469 *inoculum* in an *in vitro* fermentation assay. Moreover, comparing *in vivo* digestibility estimates in
470 195
471 horses with those obtained *in vitro* using the Daisy^{II} Incubator at 30, 48, and 72 h of incubation,
472 196
473 Earing et al. [5] showed that, for high-fiber diets (44 - 51% NDF), a 72 h incubation time provided
474 197
475 estimates closer to *in vivo* data. However, the use of a set incubation time for all kind of feedstuffs
476 198
477

481
482
483 199 may be questionable, because more fibrous feedstuffs may need longer incubation times to estimate
484
485 200 accurately DMD and NDFD [4,5]. In the current trial, the effect of incubation time was not
486
487 201 evaluated for each considered feedstuff, because only one sample (in duplicate) per feedstuff was
488
489
490 202 incubated. Further targeted trials are needed to assess the optimum incubation time for different
491
492 203 feedstuffs using the Daisy^{II} Incubator for the digestibility evaluation of feedstuffs in donkeys.
493
494 204 As we did not use multiple samples per each feedstuff, at each incubation time differences among
495
496 205 feedstuffs were not statistically evaluated. Despite lack of targeted statistical evaluation, the single
497
498 206 feedstuffs showed expected trends according to their chemical composition (Fig. 1). High-fiber
499
500 207 feedstuffs showed in fact lower DMD (Fig. 1a) and NDFD (Fig. 1b) values when compared to high-
501
502 208 starch samples. The lowest NDFD values obtained for alfalfa hay and wheat straw at 72 h of
503
504 209 incubation can be the consequence of their high lignin content (Fig. 1b). It is known that the type of
505
506 210 feed, stage of maturity and processing can affect the digestive process and the gastrointestinal
507
508 211 retention time in equids [22-23]. In forages, the cell wall components are the main factors
509
510 212 influencing digestibility. Cell wall components increase with plant maturity [24] and lignification in
511
512 213 particular imposes a barrier to complete cell wall polysaccharide digestion [25-26].
513
514 214 The CDA applied to our data (Fig. 2) allowed a good separation of the feedstuffs; this is also a way
515
516 215 to synthesize the method's action according to the chemical composition of the feedstuffs.
517
518 216

521 217 *4.2. Repeatability and reproducibility of the method*

523 218 Concerning the repeatability and reproducibility of the method, it has to be pointed out that values
524
525 219 can vary among laboratories [27]. Despite the large use of the Daisy^{II} Incubator to measure
526
527 220 digestibility (especially in ruminants), very few studies aimed at verifying the associated
528
529 221 measurement errors [20]. To the best of our knowledge, no published literature is available on the
530
531 222 repeatability and reproducibility of DMD and IVTD obtained with the Daisy^{II} Incubator in
532
533 223 ruminants or equids. The results obtained in our study are in accordance with those commonly
534
535 224 found by Ankom Technology using the Daisy^{II} Incubator [Brian Layton, Ankom Technology;
536
537
538
539
540

541
542
543 225 personal communication]. Regarding fiber digestibility, in ruminants Spanghero et al. [20] showed
544
545 226 that the repeatability and reproducibility of NDFD and digestible NDF using the Daisy^{II} Incubator
546
547
548 227 improve **when** extending the fermentation time from 30 to 48 h. Our data also show a general
549
550 228 improvement of the repeatability and reproducibility of the method, particularly for NDFD, as a
551
552 229 consequence of longer incubations (Table 2b). At the shorter incubation times (30, 48, and 60 h),
553
554 230 low repeatability and reproducibility for NDFD may be the consequence of insufficient degradation
555
556 231 of the fiber fraction of the diet, which instead improved at 72 h of incubation, also allowing a better
557
558 232 precision degree of the method. At 72 h of incubation, the method tested in our study was less
559
560 233 repeatable and reproducible for NDFD than for DMD and IVTD (Table 2b). The repeatability and
561
562 234 reproducibility found by Spanghero et al. [20] for NDFD at 48 h of incubation were equal to 6.83%
563
564 235 and 10.46%; such values are quite similar to those found for NDFD at 72 h of incubation in the
565
566 236 current study (4.5% and 10.4%, respectively). **The estimated repeatability and reproducibility**
567
568 237 **coefficients (Table 2b) are precision measures providing additional information, as they represent**
569
570 238 **the maximum absolute difference that can be expected between repeated measurements on 95% of**
571
572 239 **occasions [17].**

573 240 574 575 241 **5. Conclusion**

576
577 242 We evaluated the use of the Daisy^{II} Incubator (Ankom Technology) for determining DMD, IVTD,
578
579 243 and NDFD of feedstuffs using donkey feces as source of microbial *inoculum*. **From the obtained**
580
581 244 **results, it can be concluded that an incubation period of 72 h is required to obtain accurate DMD,**
582
583 245 **IVTD, and NDFD estimates in donkeys and to improve the repeatability and reproducibility of the**
584
585 246 **method. For NDFD, the repeatability and reproducibility of the method were comparable to those**
586
587 247 **obtained with the Daisy^{II} Incubator in ruminants.** Overall, the trial shows that the method described
588
589 248 by different authors [3-5] for *in vitro* digestibility studies in horses using the Daisy^{II} Incubator could
590
591 249 be successfully extended to donkeys.
592
593
594
595
596
597
598
599
600

601
602
603 250 Further studies should evaluate the among-laboratory repeatability and reproducibility of the
604
605 251 method. In addition, trials aimed at comparing results from *in vitro* and *in vivo* digestibility are
606
607 252 needed to assess the most suitable incubation time of different feedstuffs for donkeys, and to
608
609
610 253 propose equations that fit the shortest *in vitro* incubation time with *in vivo* data.
611

612 254

613 614 255 **Acknowledgements**

615
616 256 Financial support for this work was provided by the University of Torino (ex 60% grant 2016). The
617
618 257 authors gratefully acknowledge Mr. Andrea Gamarra (“Asi Lait”, San Benigno Canavese, Torino,
619
620 258 Italy) for providing the donkeys, and Dr. Giovanni Perona, Dr. Stefano Nurisso, Dr. Francesca
621
622 259 Spada, Dr. Federica Raspa, Dr. Silvia Boggero, Dr. Barbara Negri, and Mrs. Vanda Maria Malfatto
623
624 260 for animal care and analytical help. Dr. Brian Layton (Technical Sales and Service, Ankom
625
626 261 Technology, Macedon NY, USA) is also gratefully acknowledged for his technical support.
627
628

629 262

630 631 263 **References**

- 632
633 264 [1] Lowman RS, Theodorou MK, Hyslop JJ, Dhanoa MS, Cuddeford D. Evaluation of an *in vitro*
634
635 265 batch culture technique for estimating the *in vivo* digestibility and digestible energy content of
636
637 266 equine feeds using equine feces as the source of microbial inoculum. *Anim Feed Sci Technol*
638
639 267 1999; 80:11–27.
640
641 268 [2] Abdouli H, Ben Attia S. Evaluation of a two-stage *in vitro* technique for estimating digestibility
642
643 269 of equine feeds using horse feces as the source of microbial inoculum. *Anim Feed Sci Technol*
644
645 270 2007; 132:155–62.
646
647
648 271 [3] Ringler JE, Cassill BD, Hayes S, Lawrence LM. Effect of incubation time on *in vitro* estimates
649
650 272 of DM, NDF and ADF digestibility obtained using equine feces inoculum. In: *Proceedings of the*
651
652 273 *19th Equine Science Society*; 2005, p. 307–8.
653
654
655
656
657
658
659
660

661
662
663
664 274 [4] Lattimer JM, Cooper SR, Freeman DW, Lalman DL. Effect of yeast culture on in vitro
665 fermentation of a high concentrate or high fiber diet using equine fecal inoculum in a Daisy^{II}
666 275 incubator. J Anim Sci 2007; 85:2484–91.
667
668 276
669
670 277 [5] Earing JE, Cassill BD, Hayes SH, Vanzant ES, Lawrence LM. Comparison of in vitro
671 digestibility estimates using the Daisy^{II} incubator with in vivo digestibility estimates in horses. J
672 278 Anim Sci 2010; 88:3954–63.
673
674 279
675
676 280 [6] Murray JMD, McMullin P, Handel I, Hastie PM. Comparison of intestinal contents from
677 different regions of the equine gastrointestinal tract as inocula for use in an *in vitro* gas production
678 281 technique. Anim Feed Sci Technol 2014; 187:98-103.
680
681 282
682
683 283 [7] Cuddeford D, Pearson RA, Archibald RF, Muirhead RH. Digestibility and gastro-intestinal
684 transit time of diets containing different proportions of alfalfa and oat straw given to
685 284 Thoroughbreds, Shetland ponies, Highland ponies and donkeys. Anim Sci 1995; 61:407–17.
686
687 285
688
689 286 [8] Pearson RA, Archibald RF, Muirhead RH. The effect of forage quality and level of feeding on
690 digestibility and gastrointestinal transit time of oat straw and alfalfa given to ponies and donkeys.
691 Br J Nutr 2001; 85:599–606.
692 287
693
694 288
695
696 289 [9] Kidane NF, Stuth JW, Tolleson DR. Predicting diet quality of donkeys via fecal-NIRS
697 calibrations. Rangeland Ecol Manage 2008; 61:232–9.
698
699 290
700
701 291 [10] Wood SJ. Some factors affecting the digestible energy, requirements and dry matter intake of
702 mature donkeys and comparison with normal husbandry practises. PhD Thesis. University of
703 292 Edinburgh; 2010.
704
705 293
706
707 294 [11] Ankom Technology. Method for Determining Neutral Cellulase plus Gamanase (NCGD).
708 2006. Retrieved from
709
710 295 <https://www.ankom.com/sites/default/files/document-files/NCGD%20Procedure.pdf> (last
711 296 accessed 16.07.2018).
713
714 297
715
716
717
718
719
720

- 721
722
723
298 [12] Valle E, Raspa F, Giribaldi M, Barbero R, Bergagna S, Antoniazzi S, Mc Lean AK, Minero M,
724
725
299 Cavallarini L. A functional approach to the body condition assessment of lactating donkeys as a
726
727
300 tool for welfare evaluation. PeerJ 2017; 5:e3001.
728
729
301 [13] Burden F, Thiemann A. Donkeys are different. J Equine Vet Sci 2015; 35:376–82.
730
731
302 [14] Valle E, Pozzo L, Giribaldi M, Bergero D, Gennero MS, Dezzutto D, McLean A, Borreani G,
732
733
303 Coppa M, Cavallarini L. Effect of farming system on donkey milk composition. J Sci Food Agric
734
735
304 2018; 98:2801-8.
736
737
305 [15] Fortina R, Gasmi-Boubaker A, Lussiana C, Malfatto V, Tassone S, Renna M. Nutritive value
738
739
306 and energy content of the straw of selected *Vicia L. taxa* from Tunisia. Ital J Anim Sci 2015;
740
741
307 14:280–4.
742
743
308 [16] Ankom Technology. In vitro true digestibility using the Daisy^{II} Incubator. 2005. Retrieved
744
745
309 from: [https://www.ankom.com/sites/default/files/document-](https://www.ankom.com/sites/default/files/document-files/Method_3_Invitro_D200_D200I.pdf)
746
747
310 [files/Method_3_Invitro_D200_D200I.pdf](https://www.ankom.com/sites/default/files/document-files/Method_3_Invitro_D200_D200I.pdf) (last accessed 21.12.2018).
748
749
750
751
311 [17] Bartlett JW, Frost C. Reliability, repeatability and reproducibility: analysis of measurement
752
753
312 errors in continuous variables. Ultrasound Obstet Gynecol 2008; 31:466-75.
754
755
313 [18] SAS. The SAS System for Windows, Release 9.4. SAS Institute Inc., Cary, NC, USA. 2018
756
757
314 <http://support.sas.com/documentation> (last accessed 07.12.2018).
758
759
315 [19] Littell RC, Henry PR, Ammerman CB. Statistical analysis of repeated measures data using
760
761
316 SAS procedures. J Anim Sci 1998; 76:1216–31.
762
763
317 [20] Spanghero M, Berzaghi P, Fortina R, Masoero F, Rapetti L, Zanfi C, Tassone S, Gallo A,
764
765
318 Colombini S, Ferlito JC. Technical note: Precision and accuracy of in vitro digestion of neutral
766
767
319 detergent fiber and predicted net energy of lactation content of fibrous feeds. J Dairy Sci 2010;
768
769
320 93(10):4855-9.
770
771
772
773
774
775
776
777
778
779
780

781
782
783
784
785
786
787
788
789
790
791
792
793
794
795
796
797
798
799
800
801
802
803
804
805
806
807
808
809
810
811
812
813
814
815
816
817
818
819
820
821
822
823
824
825
826
827
828
829
830
831
832
833
834
835
836
837
838
839
840

[21] Franzan BC, Franco TW, Stefani G, Pereira MM, de Almeida FQ, Silva VP. Equine fecal inoculum optimization in *in vitro* fermentation assays of dehydrated roughage. R Bras Zootec 2018; 47:e20180006.

[22] Rosenfeld I, Austbø D. Effect of type of grain and feed processing on gastrointestinal retention times in horses. J Anim Sci 2009; 87:3991-6.

[23] Van Weyenberg S, Sales J, Janssens GPJ. Passage rate of digesta through the equine gastrointestinal tract: a review. Livest Sci 2006; 99:3-12.

[24] Peiretti PG, Gai F, Alonzi S, Battelli G, Tassone S. Characterisation of Alpine highland pastures located at different altitudes: forage evaluation, chemical composition, *in vitro* digestibility, fatty acid and terpene contents. Plant Biosyst 2017; 151(1):50-62.


[25] Gomes DI, Detmann E, Valadares Filho SC, Fukushima RS, de Souza MA, Valente TNP, Paulino MF, de Queiroz AC. Evaluation of lignin contents in tropical forages using different analytical methods and their correlations with degradation of insoluble fiber. Anim Feed Sci Tech 2011; 168:206-22.

[26] Krizsan SJ, Huhtanen P. Effect of diet composition and incubation time on feed indigestible neutral detergent fiber concentration in dairy cows. J Dairy Sci 2013; 96:1715-26.

[27] Van Laar H, Van Straalen WM, Van Gelder AH, De Boever JL, D'heer B, Vedder H, Kroes R, de Bot P, Van Hees J, Cone JW. Repeatability and reproducibility of an automated gas production technique. Anim Feed Sci Technol 2006; 127:133-50.

1 **Figure 1.**

2 **(a)**



Substrate

- Alfalfa hay
- Bromegrass hay
- Timothy hay
- Wheat straw
- Barley grain
- Ryegrass hay
- Wheat bran

3

4

5

6


7

8

9

10

11 (b)




12

13

1 **Figure 2.**

2



3

Figure captions

Figure 1. Trends for (a) dry matter digestibility (DMD, g/kg DM) and (b) neutral detergent fiber digestibility (NDFD, g/kg NDF) over incubation time for different feedstuffs. Error bars represent standard deviation.

Figure 2. Canonical Discriminant Analysis of regression parameters of *in vitro* true dry matter digestibility (IVTD) for different feedstuffs.

[Color should be used for all figures in print.]

1 **Table 1. Chemical composition of feedstuffs (g/kg DM, unless otherwise stated).**

	DM, g/kg	Ash	CP	EE	NDF	ADF	Lignin	NFC*
Alfalfa hay	911	91	156	10	616	480	100	127
Bromegrass hay	918	112	89	13	662	406	45	124
Ryegrass hay	870	90	156	32	588	332	25	135
Timothy hay	859	68	82	14	598	361	47	238
Wheat bran	899	59	166	48	675	145	39	51
Wheat straw	925	93	66	15	797	550	77	30
Barley grains	883	28	117	19	213	69	12	623

2 Abbreviations: DM, dry matter; CP, crude protein; EE, ether extract; NDF, neutral detergent fiber;

3 ADF, acid detergent fiber; NFC, non-fiber carbohydrates.

4 * Calculated as: $NFC = 1000 - (\text{ash} + CP + EE + NDF)$.

1 **Table 2. *In vitro* apparent dry matter digestibility (DMD), true dry matter digestibility (IVTD) and neutral detergent fiber digestibility**
2 **(NDFD) obtained using the Daisy^{II} Incubator and donkey feces as source of microbial *inoculum* (n=208). (a) Effect of incubation time on**
3 **estimates of DMD, IVTD and NDFD. (b) Repeatability and reproducibility of the method.**

Incubation time (h)	DMD				IVTD				NDFD			
	30	48	60	72	30	48	60	72	30	48	60	72
(a)												
Mean*	330.9 ^d	432.6 ^c	497.1 ^b	525.2 ^a	510.7 ^d	587.4 ^c	639.6 ^b	658.3 ^a	193.2 ^d	338.4 ^c	412.5 ^b	443.1 ^a
(b)												
Variances												
Residual error	160.7	262.8	261.7	199.6	85.6	97.0	123.3	108.8	505.9	278.1	499.9	389.0
Runs	2185.5	6607.4	18963.6	3417.0	409.5	8495.9	7903.5	5562.9	968.8	22086.0	21907.9	14642.5
Run × Feedstuffs	529.4	1001.8	2642.9	776.0	292.6	859.0	1170.1	475.1	2220.3	1759.3	4389.6	2095.5
Precision parameters												
Repeatability (CV, %)	3.8	3.7	3.3	2.7	1.8	1.7	1.7	1.6	11.6	4.9	5.4	4.5
Reproducibility (CV, %)	6.5	7.4	10.3	5.0	2.8	5.4	5.3	3.9	18.5	14.7	14.7	10.4
Repeatability coefficient*	35.1	44.9	44.8	39.2	25.8	27.3	30.8	28.9	62.4	46.2	62.0	54.7
Reproducibility coefficient*	59.7	89.1	141.8	72.1	39.0	88.7	93.1	71.0	98.9	137.8	168.5	128.2

4 Abbreviations: CV, coefficient of variation.

5 For the effect of incubation time, different superscripts within row and digestibility parameter indicate significant differences (a, b, c, d: $P < 0.05$).

6 * Expressed as g/kg DM for DMD and IVTD; expressed as g/kg NDF for NDFD.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59

Conflict of interest statement

The Authors declare no conflict of interest.