AperTO - Archivio Istituzionale Open Access dell'Università di Torino

Gut microbiota and chronic exercise in diabetic patients: not only bacteria

This is the author's manuscript

Original Citation:

Availability:

This version is available <http://hdl.handle.net/2318/1706982> since 2020-11-22T10:02:09Z

Published version:

DOI:10.23736/S0026-4806.19.06090-7

Terms of use:

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)

Gut microbiota and chronic exercise in diabetic patients: not only bacteria

Journal: Minerva Medica

Paper code: Minerva Med-6090

Submission date: April 6, 2019

Article type: Letter to the Editor

Files:

1. Manuscript

Version: 1

Description: manoscritto

File format: application/msword

Gut microbiota and chronic exercise in diabetic patients: not only bacteria

Davide Giuseppe Ribaldonea,^{1*} Rinaldo Pellicano²

¹Department of Surgical Sciences, University of Turin, Turin, Italy;

²Unit of Gastroenterology, Molinette Hospital, Turin, Italy

Conflicts of interest: none to declare.

Corresponding author: Davide Giuseppe Ribaldone - Department of Surgical Sciences, University of Turin, Corso Bramante, n° 88, 10126 Turin, Italy; tel (0039)0116333918, fax (0039)0116333623, davrib_1998@yahoo.com

Key words: Fungal microbiota - Mycobiome - Microbiome – Metabolic syndrome – Physical exercise

TO THE EDITOR: In a recent interesting paper Pasini *et al.* have shown that exercise controls diabetes also by modifying intestinal mycobiota composition and gut barrier function. In particular, while diabetes was associated with significant gut mycetes overgrowth, exercise improved glycemia and reduced gut mycetes overgrowth. The analysis of the gut microbiota included bacterial species, *Candida Albicans* and *Mycetes spp.* Only *Candida albicans* and *Mycetes spp.* were significantly reduced after exercise ($P = 0.043$ and $P < 0.001$, respectively).¹

We would like to highlight three crucial points regarding the results of this study.

First, each strategy aiming to obtain the homeostasis of the microbiota should consider the mycobiota. This is not always considered in a time of prevalent interest toward the bacteria.²

Second, the study of the gut permeability, as indicator of gut barrier function, by the search for zonulin³ should be mandatory in this type of investigations.

Third, more and more data confirmed the potential involvement of gut microbiota in several extra-intestinal diseases.^{4,5}

References

1. Pasini E, Corsetti G, Assanelli D, Testa C, Romano C, Dioguardi FS, et al. Effects of chronic exercise on gut microbiota and intestinal barrier in human with type 2 diabetes. *Minerva Med.* 2019 Jan;110(1):3–11.
2. Pellicano R, Ribaldone DG, Astegiano M, Dughera L, Battaglia E, Morgando A, et al. Gastroenterology today: between certainties and news. *Minerva Gastroenterol Dietol.* 2018 Nov;64(4):323–32.
3. Caviglia GP, Dughera F, Ribaldone DG, Rosso C, Abate ML, Pellicano R, et al. Serum zonulin in patients with inflammatory bowel disease: a pilot study. *Minerva Med.* 2019 Feb;110(2):95–100.

4. Champion D, Ponzo P, Alessandria C, Saracco GM, Balzola F. The role of microbiota in autism spectrum disorders. *Minerva Gastroenterol Dietol*. 2018 Nov;64(4):333–50.
5. Ribaldone DG, Venero M, Pellicano R. Is microbiome a target for the management of allergy associated diseases in children? *Eur Rev Med Pharmacol Sci*. 2018 Dec;22(23):8061–2.