

AperTO - Archivio Istituzionale Open Access dell'Università di Torino

**PLASTICIZERS AS POSSIBLE METABOLIC DISRUPTORS: MIXTURE EFFECTS ON ADIPOCYTE DIFFERENTIATION AND LIPID ACCUMULATION IN 3T3-L1 CELLS**

**This is the author's manuscript**

*Original Citation:*

*Availability:*

This version is available <http://hdl.handle.net/2318/1708671> since 2019-08-12T10:48:27Z

*Terms of use:*

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)

## PLASTICIZERS AS POSSIBLE METABOLIC DISRUPTORS: MIXTURE EFFECTS ON ADIPOCYTE DIFFERENTIATION AND LIPID ACCUMULATION IN 3T3-L1 CELLS

Erika Cottone, Valentina Pomatto, Paolo Cocci, Erik Russel Nelson, Francesco Alessandro Palermo, Patrizia Bovolin

A subset of Endocrine Disrupting Chemicals (EDCs), named Metabolic Disruptors, have been demonstrated to be able of promoting adiposity and alteration of energy homeostasis. We investigated the adipogenic activity and nuclear receptor interactions of four plasticizers approved for the manufacturing of food-contact materials (FCMs). Differentiating 3T3-L1 mouse preadipocytes were exposed to scalar concentrations of DiNP (Di-iso-nonyl-phthalate), DiDP (Di-iso-decyl-phthalate), DEGDB (Diethylene glycol dibenzoate), or TMCP (Tri-m-cresyl phosphate). Rosiglitazone, a well-known pro-adipogenic peroxisome proliferator activated receptor gamma (PPAR $\gamma$ ) agonist, and the plasticizer BPA were included as reference compounds. All concentrations of plasticizers were able to enhance lipid accumulation. Accordingly, *in silico* and gene-reporter experiments showed binding affinities and receptor activities to the nuclear receptor PPAR $\gamma$  comparable to those of Rosiglitazone. Differently from BPA, the four plasticizers were most effective in enhancing lipid accumulation when added in the mid-late phase of differentiation. qRT-PCR studies showed that all plasticizers were able to increase the expression of CCAAT/enhancer binding protein  $\beta$  (*Cebp $\beta$* ), in the early steps of adipogenesis, and *Ppar $\gamma$ 2*, the adipogenesis master gene, in the middle phase. In addition, TMCP and DEGDB were both able to modulate the expression of Fatty Acid Binding Protein 4/Adipocyte Protein 2 (*Fabp4/Ap2*) transcripts in the late phase of adipogenesis. It is well known that exposure in daily life is not restricted to individual compounds and humans are exposed to complex mixtures of EDCs. We therefore began to investigate the effects of different combinations of these plasticizers in 3T3-L1 cells. Our preliminary results suggest an enhancement of lipid accumulation, without an apparent additive effect. Taken together, our results suggest that exposure to single plasticizers or mixtures at low, environmentally relevant doses can affect adipogenesis and lipid accumulation in fat cells *in vitro*. Future experiments will better characterize the combined molecular and cellular effects of these chemicals in differentiating adipocytes.