

AperTO - Archivio Istituzionale Open Access dell'Università di Torino

Characterization and temporal evolution of the elemental composition of PM₁₀ collected at Ny-Ålesund (The Arctic)

This is the author's manuscript

Original Citation:

Availability:

This version is available <http://hdl.handle.net/2318/1712224> since 2019-09-23T12:13:36Z

Publisher:

Società chimica italiana

Terms of use:

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)

CHARACTERIZATION AND TEMPORAL EVOLUTION OF THE ELEMENTAL COMPOSITION OF PM₁₀ COLLECTED AT NY-ÅLESUND (THE ARCTIC)

E. Conca¹, O. Abollino¹, A. Giacomino², S. Buoso¹, A. Ruo Redda², R. Traversi³, S. Becagli³, M. Grotti⁴, M. Malandrino¹

¹ *Department of Chemistry, University of Turin, Via P. Giuria 5 - 10125 Turin, Italy*

² *Department of Drug Science and Technology, University of Turin, Via P. Giuria 5 - 10125 Turin, Italy*

³ *Department of Chemistry "Ugo Schiff", University of Florence, Via della Lastruccia, 3-13 - 50019 Sesto Fiorentino, Italy*

⁴ *Department of Chemistry and Industrial Chemistry, University of Genoa, Via Dodecaneso 31 - 16146 Genoa, Italy*

In this study, the elemental composition of PM₁₀ samples collected in Ny-Ålesund (Norwegian Arctic) in the sampling campaigns 2010-2013 was investigated, with the purpose to identify the PM sources and to understand short- and long-range transport processes [1]. Enrichment factors were calculated for all elements with respect to the mean values for the Earth's crust reported by Wedepohl [2] and to the mean abundances in sea water reported by Goldberg [3], in order to distinguish elements having natural, anthropic or mixed origin. The results obtained so far evidence a remarkable seasonal trend for most of the investigated elements. For both geogenic and anthropogenic elements, concentrations are generally higher in March and April, when the ground is almost entirely covered by snow and ice, suggesting that long-range transport processes might be taking place. On the other hand, the elements typically deriving from marine aerosol (i.e. Mg and Na) present higher concentrations in late spring and summer, together with Co, Ni and V, typical anthropogenic metals related to ship emissions. From the Kruskal-Wallis and Conover-Iman tests it emerged that, for most of the analytes, the four campaigns are not significantly different; therefore, in the studied period, the composition of PM₁₀ in Ny-Ålesund did not vary remarkably. Principal Component Analysis allowed us to better understand the sources of different elements and to lay the basis for an interpretation of the chemical and physical processes concerning the Arctic atmosphere. Airborne pollution deriving from ship fuels, local vehicle and continental emissions resulted to be the main sources of anthropogenic elements.

[1] A. Bazzano, F. Ardini, M. Grotti, M. Malandrino, A. Giacomino, O. Abollino, D. Cappelletti, S. Becagli, R. Traversi, R. Udisti, Elemental and lead isotopic composition of atmospheric particulate measured in the Arctic region (Ny-Ålesund, Svalbard Islands). *Rend. Fis. Acc. Lincei* 27 (2016) S73-S84.

[2] K.H. Wedepohl, The composition of the continental crust. *Geochim. Cosmochim. Ac.* 59 (1995) 1217-1232.

[3] E.D. Goldberg, Chapter 5. In: J.P. Riley, G. Skirrow, *Chemical Oceanography*. Academic Press, 1965.