

P-01-19

Organoleptic properties of celta pig meat fed with traditional or commercial feed (#346)Sol Zamuz¹, Laura Purriños¹, Alberto Brugiapaglia², Daniel Franco¹, Paulo MuneKata¹, Mirian Pateiro¹, Laura Cutillas-Barreiro¹, Jose M. Lorenzo¹¹ Centro Tecnológico de la Carne, Ourense, Spain; ² University of Turin, Department of Agricultural, Forest and Food Sciences, Grugliasco, Torino, Italy**Introduction**

The organoleptic attributes of meat are usually considered key factors to assess meat quality, among which the sensory acceptability is a major factor to address the consumers preferences, which can ultimately affect their purchasing intention. Celta pig is a native breed of Galicia (NW Spain) highly appreciated by its succulent meat. Moreover, previous studies have shown the influence of diet and slaughter weight in the nutritional quality of meat (Lorenzo et al., 2014). Changes in the physicochemical properties of meat could modify the sensorial characteristic and influence the acceptance of consumers. The aim of this study was to evaluate the effect of feeding (traditional feed (C) and commercial feed (F)) and slaughter weight (low weight (Lw) and high weight (Hw)) on the sensory attributes of Celta pig meat.

Methods

The organoleptic attributes of meat are usually considered key factors to assess meat quality, among which the sensory acceptability is a major factor to address the consumers preferences, which can ultimately affect their purchasing intention. Celta pig is a native breed of Galicia (NW Spain) highly appreciated by its succulent meat. Moreover, previous studies have shown the influence of diet and slaughter weight in the nutritional quality of meat (Lorenzo et al., 2014). Changes in the physicochemical properties of meat could modify the sensorial characteristic and influence the acceptance of consumers. The aim of this study was to evaluate the effect of feeding (traditional feed (C) and commercial feed (F)) and slaughter weight (low weight (Lw) and high weight (Hw)) on the sensory attributes of Celta pig meat.

Results

The results of the QDA of Celta pig loins are shown in Figure 1. Significant differences ($P < 0.05$) in color intensity, marbling, tenderness and juiciness were observed. The C-Hw samples received the highest scores in these attributes. Regarding PCA, the first two components (Figure 2) accumulated 94.55% of total variance (F1 = 83.95% and F2 = 10.60%, first and second principal components, respectively). This outcome is in accordance with Reis et al. (2010), who argued that to select principal components, the accumulated percentage of variance has to be equal or greater than 70%. The PCA indicated that color, tenderness, and flavor were located in the same quadrant with C-Hw. This outcome agrees with the data reported by Cannata et al. (2010) who observed that highly marbled pork loins also received significantly high scores on juiciness and tenderness attributes. In addition, the HCA was applied to data and three groups with different preference

profiles were generated. The PREFMAP indicated that the applied model was significant for Group 1 and 2 ($P > F = 0.004$ and 0.45 , respectively). External preference mapping and contour plot (Figure 3) showed areas that 80 and 100% of consumers were satisfied with fresh loin obtained from C-Hw and C-Lw treatments. The high preference values for these treatments could be attributed to the high scores obtained for tenderness and juiciness attributes. This hypothesis is supported by the data obtained by Aaslyng et al. (2007), who reported that consumer preference was positively associated with tenderness and juiciness in pork loins.

Conclusion

Feeding and slaughter weight influenced the sensorial perception of color, marbling, tenderness and juiciness of Celta pig loins. The high percentage of consumers satisfied with the loins obtained from C-Hw treatment support the use of traditional feed and high weight slaughter to produce Celta pig loin in the conditions used in the present study.

Acknowledgements

Authors are grateful to Xunta de Galicia for the financial support (Project FEADER 2017/038B). Financial support from the Agencia Estatal de Investigación for supporting of Sol Zamuz (PTA2017-14156-I). Paulo E. S. MuneKata acknowledges postdoctoral fellowship support from Ministry of Economy and Competitiveness (MINECO, Spain) "Juan de la Cierva" program (FICI-2016-29486).

References

Aaslyng et al. (2007). The impact of sensory quality of pork on consumer preference. *Meat Science*, 76(1), 61-73.
 Cannata et al. (2010). Effect of visual marbling on sensory properties and quality traits of pork loin. *Meat Science*, 85(3), 428-434.
 ISO 8586 (2012). Sensory analysis - general guidelines for the selection, training and monitoring of selected assessors and expert sensory assessors. International Standards Organisation publications.
 Lorenzo et al. (2014). Effect of the inclusion of chestnut in the finishing diet on volatile compounds during the manufacture of dry-cured "Lacón" from Celta pig breed. *Meat Science*, 96(1), 211-223.
 Reis et al. (2010). Mapa de preferéncia. In V. P. R. Minim (Ed.), *Análise Sensorial: Estudos com consumidores* (pp. 214-257). Viçosa: Ed. UFV.