

AperTO - Archivio Istituzionale Open Access dell'Università di Torino

Investigation of permafrost sensor dynamics

This is the author's manuscript

Original Citation:

Availability:

This version is available <http://hdl.handle.net/2318/1712357> since 2019-09-25T15:35:35Z

Terms of use:

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)

<i>Topic</i>	T9.1 Meteorological traceability and uncertainty
<i>Oral or Poster Presentation</i>	Poster Presentation

INVESTIGATION OF PERMAFROST SENSOR DYNAMICS

P. Pavlasek ^{1,2}, A. Merlone ³, F. Sanna ³, G. Coppa ³, S. Ďuriš ², Z. Ďurišová ², J. Palenčar ²

¹ *Slovak Institute of Metrology, Bratislava, Slovakia*

² *Slovak University of Technology, Bratislava, Slovakia*

³ *Istituto Nazionale di Ricerca Metrologica, Torino, Italy*

E-mail (corresponding author): Pavlasek@smu.gov.sk

Permafrost soils are an important part of the whole climate and environmental system as they play an important role in storing carbon and water. As permafrost makes around 15 % of the global land cover the amount of stored carbon and water is not negligible. These types of soils provide a direct source of water for plants in severe drought areas, during summer and keep surplus water in the soil until the next season. Furthermore the trapped carbon is of high concern as the increasing global temperature could cause the release of this undesired substance.

To be able to determine the current state of the permafrost thickness and its evolution in time it is important to have a precise measurement of permafrost temperature. Commonly the measurements are performed by temperature sensors that are placed in a narrow drill hole several meters deep. The temperature is recorded continuously during the whole year. The problematic point of this measurement is the day/night cycle, during which the temperature elevates due to increasing solar radiation. This creates an oscillating behaviour of measured temperature that is affected by the sensor dynamics that is different from sensor to sensor. In order to be able to have reliable and traceable permafrost temperature data the different sensor dynamics need to be determined and included into the measurement uncertainty.

This paper presents an experimental measurement method by which means it is possible to determine the different permafrost sensor dynamic in the range from -30 °C to +30 °C. The paper furthermore discusses the uncertainty contribution from this sensor property and furthermore compares the dynamics of different tested sensors.

Acknowledgements

This work is being developed within the frame of the EMRP (European Metrology Research Programme) joint research project ENV58 “METEOMET”. Furthermore the work was supported by the Slovak R&D Agency, projects APVV-15- 0295, APVV-15-0164, the Scientific Grant Agency of the Ministry of Education of the Slovak Republic projects VEGA 2/0610/17, VEGA 1/0556/18 VEGA - 1/0098/18 and KEGA039STU-4/2017.