AperTO - Archivio Istituzionale Open Access dell'Università di Torino

Stay with the flow: How macroinvertebrate communities recover during the rewetting phase in Alpine streams affected by an exceptional drought

This is a pre print version of the following article:

Original Citation:

Availability:

This version is available <http://hdl.handle.net/2318/1718418> since 2021-12-23T12:36:55Z

Published version:

DOI:10.1002/rra.3563

Terms of use:

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)

**Stay with the flow: how macroinvertebrate communities
recover during the rewetting phase in Alpine streams
affected by an exceptional drought**

Journal:	<i>River Research and Applications</i>
Manuscript ID	Draft
Wiley - Manuscript type:	Research Article
Date Submitted by the Author:	n/a
Complete List of Authors:	Doretto, Alberto; University of Piemonte Orientale; University of Turin; ALPSTREAM - Alpine Stream Research Center Bona, Francesca; University of Turin; ALPSTREAM - Alpine Stream Research Center Falasco, Elisa; University of Piemonte Orientale; ALPSTREAM - Alpine Stream Research Center Morandini, Daniele; University of Turin Piano, Elena; University of Piemonte Orientale; ALPSTREAM - Alpine Stream Research Center Fenoglio, Stefano; Università del Piemonte Orientale; ALPSTREAM - Alpine Stream Research Center
Keywords:	benthic invertebrates, Alpine streams, water scarcity, recolonization, biodiversity, resilience

SCHOLARONE™
Manuscripts

Stay with the flow: how macroinvertebrate communities recover during the rewetting phase in Alpine streams affected by an exceptional drought

Short running title: Resilience of macroinvertebrates to droughts in Alpine streams

Alberto Doretto^{1,2,3,*}, Francesca Bona^{2,3}, Elisa Falasco^{1,3}, Daniele Morandini², Elena Piano^{1,3}, Stefano Fenoglio^{1,3}

¹DISIT, University of Piemonte Orientale, Viale Teresa Michel 25, I-15121, Alessandria, Italy

²DBIOS, University of Torino, Via Accademia Albertina 13, I-10123, Torino, Italy

³ALPSTREAM – Alpine Stream Research Center, I-12030 Ostan, Italy

*Corresponding author: alberto.doretto@unito.it; alberto.doretto@uniupo.it

Orcid ID

Alberto Doretto: <https://orcid.org/0000-0002-4105-473X>

Acknowledgements

Authors are very grateful to Maria Cristina Bruno, Laura Gruppuso, Marco Baltieri (ATAAI – Associazione per la Tutela degli Ambienti Acquatici e dell’Ittiofauna) and the Monviso Natural Park for their assistance. This work was supported by the project PRIN NOACQUA “Risposte di comunità e processi ecOsistemici in corsi d’ACQUA soggetti a intermittenza idrologica” - code 201572HW8F, funded by the Italian Ministry of Education, University and Research.

For Peer Review

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Abstract

Drought occurrence is affecting an increasing number of lotic ecosystems worldwide due to the combined effects of climatic and anthropogenic pressures. Unlike naturally intermittent rivers, where the drying phase is a part of the annual flow regime, water scarcity in Alpine rivers represent a relatively recent phenomenon and, therefore, a major threat for the biodiversity of these lotic ecosystems, but the response of aquatic communities to this disturbance is still poorly investigated. Here, we present the results on the recovery of stream macroinvertebrates in two Alpine streams after a supra-seasonal drought. As water resumed, a total of ten sampling sessions were carried out and temporal patterns in diversity, density and taxonomic composition of benthic communities as well as in the percentage of functional feeding groups were investigated. We found that the resistance of invertebrate communities in Alpine streams is generally low: drought significantly reduced the diversity and density of macroinvertebrates. Conversely, our results suggest that the passive dispersal by drift from the upstream river sections seems the main mechanism that promotes the post-drought recovery. Nevertheless, this resilience ability appears to be stream-specific and influenced by intrinsic stream characteristics, including the flow permanence and distance from the nearest upstream perennial reach. This work sheds light on the impacts of climatic and human-induced droughts on benthic invertebrate communities and assumes a primary importance to predict their future composition in relation to the intensification of flow intermittency in Alpine areas under the current global change scenario.

Keywords: benthic invertebrates, Alpine streams, water scarcity, recolonization, biodiversity, resilience

1. Introduction

Climate change is currently one of the most relevant challenges for habitat and species conservation worldwide because the raising in air temperature and alterations in the precipitation regimes are responsible for the habitat loss and fragmentation, changes in species phenology and enhanced rates of biodiversity loss (Dawson, Jackson, House, Prentice, & Mace, 2011; Mantyka-pringle, Martin, & Rhodes, 2012). The increased frequency and magnitude of hydrological extremes, such as floods and droughts, are among the main consequences of these phenomena for lotic ecosystems (Beniston, 2012; Heino, Virkkala, & Toivonen, 2009; Middelkoop et al., 2001; Ledger, & Milner, 2015; Whitehead, Wilby, Battarbee, Kernan, & Wade, 2009; Wu, & Johnson, 2019).

Alpine streams are expected to be extremely sensitive to the effects of droughts because the Alps are one of the most impacted areas by climate change and, at the same time, water abstraction is an increasing pressure (Fenoglio, Bo, Cucco, Mercalli, & Malacarne, 2010; Gorbach, Shoda, Burky, & Benbow, 2014; McKay, & King, 2006; López-Rodríguez, Márquez Muñoz, Ripoll-Martín, & Tierno de Figueroa, 2019). Under similar conditions, drought occurrence represents a major threat for stream macroinvertebrates, as documented by some authors (Bonada, Doledec, & Statzner, 2007; Calapez, Elias, Almeida, & Feio, 2014; Doretto et al., 2018b; Durance, & Ormerod, 2007; Fenoglio, Bo, Cucco, & Malacarne, 2007; Ledger, Brown, Edwards, Milner, & Woodward, 2013; Piano et al. 2019a; Pinna et al., 2016; Smith, McCormick, Covich, & Golladay, 2017; Storey, 2016).

A growing attention is paid by river ecologists on the resistance and resilience mechanisms of benthic organisms to face the drying phase (Chester, & Robson, 2011; Fritz, & Dodds, 2004; Robson, Chester, & Austin, 2011; Aspin et al., 2019). However, the resistance ability of aquatic invertebrates to drought in Alpine streams is generally considered limited, compared to the aquatic biota of other geographical regions, such as the Mediterranean area, where the drying phase is a natural part of the annual flow regime (Leigh et al., 2016; Tierno de Figueroa, López-Rodríguez, Fenoglio, Sánchez-Castillo, & Fochetti, 2013). Benthic communities in Alpine streams, therefore, are generally considered more resilient than resistant (Doretto et al., 2018b), but scientific evidence on this is still limited.

In a recent publication, Van Looy et al. (2019) developed a general framework to explain the resilience of aquatic communities to disturbance in streams, including droughts, based on the relative and combined role of three main drivers: resources competition and/or facilitation, recruitment and refugia. Firstly, the access or limitation to food resources affect the response of aquatic communities in terms of trophic and biotic relationships, acting as the major driver of the

1
2
3 77 post-disturbance recovery especially where the energetic inputs are pulsed- or patchy-distributed
4
5 78 (Richardson & Sato, 2015). In this context, ameliorative effects of large amount of organic matter
6
7 79 on macroinvertebrate communities have been reported also in relation to other physical
8
9 80 disturbances, such as siltation (Doretto et al., 2017). Second, recruitment (i.e. the gain of individuals
10
11 81 by the dispersal from adjacent habitat sources) is expected to play a primary contribution in highly
12
13 82 connected river networks, resulting in a faster post-drought recovery (Flower, 2004; Ledger &
14
15 83 Hildrew, 2001). Finally, habitat heterogeneity promotes the presence of in-stream refugia (mainly
16
17 84 pools and the hyporheic zone) that can be exploited by the benthic taxa, according to their
18
19 85 ecological traits, to survive under drying conditions (Boulton 2003; Chester & Robson, 2011;
20
21 86 Fenoglio, Bo, & Bosi, 2006; Otermin, Basaguren, & Pozo, 2002; Verdonschot, Oosten-Siedlecka,
22
23 87 Braak, & Verdonschot, 2015; Wood, Boulton, Little, & Stubbington, 2010).
24
25 88 In this study we monitored the post-drought recovery of macroinvertebrate communities in two
26
27 89 Alpine streams affected by a supra-seasonal drought (Lake, 2003) and discussed the results in
28
29 90 relation to the contribution of food resources, recruitment and refugia. In particular, temporal
30
31 91 patterns in composition and diversity of benthic invertebrate communities were evaluated during
32
33 92 the water resumption (hereafter rewetting phase) and compared to the upstream permanent
34
35 93 reaches. Our hypotheses were that: i) the invertebrate recruitment, especially in terms of
36
37 94 recolonization by drift from the upstream sections would be the main mechanism of resilience in
38
39 95 our streams, while ii) the resource availability would have a minor role. As the supra-seasonal
40
41 96 drought was one of the most prolonged ever reported for the Italian Alps, lasting for more than 5
42
43 97 months, we expected a negligible effect of the in-stream refugia (i.e. pools and hyporheic zone). In
44
45 98 addition, we postulated that the recovery process would be affected by stream-specific
46
47 99 characteristics, especially in relation to hydrology stability and flow persistence.
48
49 100

47 101 **2. Materials and methods**

49 102 *2.1 Area of study*

51 103 The sampling area is located in the Cottian Italian Alps (Northwestern Italy), where we examined
52
53 104 the post-drought recovery in two lotic systems, namely the Po and Pellice rivers, which originate at
54
55 105 2,022 and 2,387 m.a.s.l. respectively. The former is the longest Italian watercourse: it runs for 652
56
57 106 Km until the Adriatic Sea with a drainage basin of approximatively 71,000 Km². The latter is the
58
59 107 principal tributary of the Po river within the Alpine area and runs for 55 Km (drainage area: 974 Km²)
60

before its confluence (Fig. 1). They represent two good case studies as they have good water and biological qualities but with stretches recently affected by drought.

On each river, two sampling sites were selected: a perennial stretch (P), with permanent flow throughout the year, and an intermittent stretch (I) experiencing recurrent drought events since 2011 (ARPA, 2013; Piano et al., 2019a; 2019b), due to the joint effect of climate change and consequent water abstraction to fulfill human needs. At this scale of investigation, the Po and Pellice are 5-order rivers (Strahler, 1957), with the substrate dominated by coarse mineral elements and a pluvio-nival hydrological regime. Also, the land use is very similar between these two streams: more than 90% is represented by natural areas, while agricultural and urbanized areas on average account for 8% and 1.5% (Table 1).

2.2 Data collection

In 2017, Northwestern Italy experienced the most severe summer droughts ever reported. In the lower sections of the Po and Pellice rivers, including the intermittent sites here considered, surface water ceased in July and August 2017 respectively (Falasco, Piano, Doretto, Fenoglio, & Bona, 2018) and resumed only in January 2018, after conspicuous rainfalls (see Supplementary Materials Fig. SM1). Although a marked reduction of the river discharge, the upstream perennial sites were characterized by the continued permanence of running surface water during this period.

As water resumed, a total of ten sampling dates were carried out in the intermittent sites to monitor the post-drought recovery of benthic communities, covering a 3-month period (Table 2). To better describe the first phases of the recolonization process, samples were initially collected every 3 days, while at the end of the sampling period benthic invertebrates were sampled every two weeks. Moreover, since we expected that the upstream perennial stretches acted as sources of organisms during the recolonization process, macroinvertebrates were sampled also in the perennial sites on two selected occasions, namely on 19th January and 22th March 2018. These samplings allow to obtain an overview of macroinvertebrate communities, at the beginning and at the end of the studied period.

On each sampling occasion, dissolved oxygen concentration (mgL^{-1}), oxygen saturation (%), pH, water temperature ($^{\circ}\text{C}$) and electrical conductivity (μScm^{-1}) were measured with a multiparametric probe (Hydrolab mod. Quanta). Water depth (cm) and water velocity (ms^{-1}) were measured for each sample using a flowmeter (Hydro-bios Kiel). Moreover, the composition of the substrate within the area delimited by the Surber sampler was visually estimated. Based on the Wentworth's grain size

classification (1922), the percentages of boulders (>256 mm), cobbles (256-64 mm), gravel (64-2 mm) and fine sediment (<2 mm) were estimated by the same operator. Macroinvertebrates were collected using a Surber sampler (0.05 m², 250 µm mesh-size) and three samples were taken on each sampling occasion, with the only exception represented by 23th February, when no samples were collected in the intermittent site of the Po river because the stream bed was completely dry (Table 2). Samples were preserved in 70% ethanol and returned in laboratory for the sorting under a stereo-microscope. Specimens were counted and systematically identified to genus (Ephemeroptera and Plecoptera) or family level using the taxonomic keys for the Italian macroinvertebrate fauna (Campaoli, Ghetti, Minelli, & Ruffo, 1994; 1999), and also classified into functional feeding groups (FFGs: collector-gatherers, filterers, predators, scrapers and shredders; Merritt, Cummins, & Berg, 2008).

2.3 Statistical analyses

Significant differences in the environmental parameters between the perennial and intermittent sampling sites (i.e. Pellice I vs Pellice P and Po I vs Po P) as well as between rivers (i.e. Pellice and Po) over the monitored period were visualized by means of Principal Component Analysis (PCA) and tested with Permutational Analysis of Variance (PERMANOVA). The water temperature, pH, electrical conductivity, dissolved oxygen and the mean value of the water velocity, depth and percentage of the four substrate classes was calculated for each sampling site on each date and included in this analysis. To meet the assumptions of normality, percentage data were square-root(arcsin) transformed prior performing the PCA, which was run using the “prcomp” function in the basic package of R. The “adonis” function in the *vegan* R package (Oksanen et al. 2015) was used, instead, to perform the PERMANOVA analysis, for which the Euclidean distance was applied. Changes in the taxonomic composition of benthic communities between sampling occasions and sites were initially visualized by means of a Non-metric Multidimensional Scaling (NMDS). This multivariate analysis was performed using the function “metaMDS” in the *vegan* R package (Oksanen et al. 2015). Surber samples were used as separate replicates: raw data about the abundance of macroinvertebrates were square-root transformed and then a Bray-Curtis dissimilarity index was applied. PERMANOVA was run to test for significant differences in relation to the “time” (as days from the water return) and “site” (Pellice P, Pellice I, Po P and Po I) factors. Generalized Additive Models (GAMs) were used to assess the non-linear response of the community metrics over the time, expressed in terms of days from the water return. Prior to perform the

statistical models, data exploration was carried out according to Zuur, Ieno & Elphick (2010) and outliers were removed. Four taxonomical metrics were considered: the total taxa richness, total density of macroinvertebrate (number of individuals m^{-2}) as well as EPT (Ephemeroptera, Plecoptera and Trichoptera) richness and density. In addition, the percentage of each functional feeding group and the ratio between scrapers and total collectors were also taken into account. The latter parameter has been proposed as an ecosystem indicator for the prevalence of autotrophy (i.e. grazing) or heterotrophy (i.e. detritus chain) in rivers (Cummins, Merritt, & Andrade, 2005). Samples collected in the perennial sites were not included in the regression models, but the mean value of each metric was calculated to better interpret the observed patterns.

All the GAMs were carried out using the “gam” function in the *mgcv* R package (Wood & Wood, 2015): a Poisson distribution was used for count data, while the negative binomial distribution was alternatively used in case of overdispersion. The binomial distribution was instead applied for the percentage variables. All the analyses were performed with the statistical software R (R Development Core Team, 2018).

3. Results

3.1 Environmental parameters

The first and the second axes of PCA accounted for 25.6% and 22.5% respectively of the variance associated to the environmental parameters, for a cumulative percentage equal to 48.1% (Fig. 2). The first axis (PC1) was positively correlated with the electrical conductivity and negatively correlated with the pH and water velocity. By contrast, the second axis (PC2) was positively correlated with the percentage of cobbles and the dissolved oxygen, while it was negatively correlated with the percentage of sand, water depth and water temperature.

In general, samples from the Pellice river were mainly oriented in the top-left part of the plot and showed a less pronounced dispersion, while samples from the Po river were oriented in the bottom-right part of the graph and showed a higher dispersion. Nevertheless, PERMANOVA did not show significant differences in the environmental parameters among sampling sites ($P = 0.184$) and rivers ($P = 0.056$), despite the p-value in this latter case was close to the significant threshold.

3.2 Macroinvertebrates

A total of 12,570 macroinvertebrates were collected, belonging to 38 different taxa (Supplementary Materials: Table SM1). Plecoptera, Ephemeroptera and Diptera were the orders with the highest

number of taxa (8), followed by Trichoptera (6), Coleoptera and Oligochaeta (2), Odonata, Tricladida, Crustacea and Nematomorpha (1). In general, *Baetis* sp., *Rhithrogena* sp., Hydropsychidae, Chironomiidae and Simuliidae were the dominant taxa in the perennial and intermittent sites of both rivers. The average number of taxa per sample was 10, while the mean number of individuals per sample was 182.

Results of the NMDS and PERMANOVA analyses showed a significant effect of the factors “time” ($F_{9,45} = 3.311$; $P < 0.001$) and “site” ($F_{3,45} = 7.302$; $P < 0.001$) on the composition of macroinvertebrate communities (Fig. 3). Invertebrate samples collected in the intermittent site of the Pellice river (Pellice I) showed a similar taxonomical composition, as they clustered together in the central part of the plot. Moreover, a partial overlap with the composition of the upstream perennial site (Pellice P) was observed. On the contrary, macroinvertebrate communities in the intermittent site of the Po river (Po I) showed the highest dispersal indicating a significant variation in the taxonomic composition over the time. Samples from this site were mostly oriented in the left-side of the plot and did not overlap with samples collected in the upstream permanent site (Po P).

When looking at the temporal variation of the diversity and density of macroinvertebrates during the rewetting phase, we found significant differences between the two rivers (Table 3). Total richness in the Pellice river significantly increased over the time, from 8 to 14 taxa (Fig. 4a, Table 3), despite it was lower than that of the upstream perennial site (18 taxa). Conversely, the total richness in the Po river slightly increased within the first 20 days of rewetting and then it markedly dropped (Fig. 4a). The average number of taxa recorded in the intermittent and permanent sites of the Po river at the end of the study were 5 and 20 respectively.

Similar results were obtained for the EPT richness, which showed opposite trends in the two rivers (Fig. 4b, Table 3). The number of EPT taxa significantly increased since the water resumption and completely approached the same value of the upstream permanent site (10 taxa). By contrast, EPT richness progressively decreased in the Po river and at the end of the sampling period was quite lower than that in the upstream permanent site (11 taxa).

Significant temporal variations in the density of macroinvertebrates were also observed in both rivers (Table 3). In the Pellice river the total density of macroinvertebrates significantly increased over time, from 2,000 to approximately 5,000 individuals m^{-2} after 73 days of rewetting (Fig. 4c). However, it was still lower than the average density recorded in the perennial section (7,700 individuals m^{-2}). Conversely, total density of macroinvertebrates in the Po river peaked around 25 days after the water resumption but then it collapsed (Fig. 4c). The numerical gap with the upstream

site (6,400 individuals m^{-2}) was high, despite the increment on the last sampling occasion. As EPT taxa were numerically dominant in this study, the temporal variation of EPT density closely resembled that observed for the total density, especially in the Po river (Fig. 4d, Table 3). The EPT density in the Pellice river, instead, showed a sharp increase after 20 days from the water resumption and then stabilized around a value of 3,500 individuals m^{-2} , that was comparable to the average EPT density in the perennial site (4,290 individuals m^{-2}).

With exception of shredders, percentages of functional feeding groups significantly varied over the time (Table 3). On average, collector-gatherers were the most abundant group in the Pellice river (34%) followed by filterers (32%), despite these two groups showed some fluctuations (Fig. 5a). Also, the percentage of scrapers was high (29%) and relatively constant over the time (Fig. 5a), while a general increase was observed for predators during the rewetting phase but, on average, they accounted for less than 4%.

Benthic communities in the Po river were almost exclusively dominated by collector-gatherers (50%) and scrapers (40%): the former were more abundant on the first and last sampling occasions respectively, while the latter were numerically abundant on the intermediate sampling occasions (Fig. 5b). The percentage of filterers was generally low (8%), despite this functional group peaked after 31 and 45 days from the water resumption (Fig. 5b). Predators were recorded only on few sampling occasions in the Po river and no significant trends were observed for this group (Fig. 5b, Table 3). Most representative taxa of each functional feeding group are listed in Table SM1 (Supplementary Materials).

Changes in the ratio between the scrapers and total collectors (i.e. shredders, collector-gatherers and filterers) were observed only in the Po river, where this indicator rapidly increased during the initial stages of the rewetting phase, peaked around 20 days, and then it decreased at the end of the study (Fig. 5c, Table 3).

4. Discussion

In a review on the response of riverine communities to disturbance, Death (2010) pointed out that, in general, benthic communities recover rapidly because they are more resilient rather than resistant. In this study we monitored the post-drought recovery of macroinvertebrate communities after a supra-seasonal drought in two Alpine streams and our findings corroborate this statement. Drought significantly reduced the diversity and density of invertebrate communities, especially regarding the most sensitive invertebrates, like EPT taxa, and confirmed our hypothesis for which

the resistance of Alpine macroinvertebrates to this disturbance is quite scarce, as demonstrated previously (Doretto et al., 2017; Fenoglio et al., 2007; Herbst, Cooper, Medhurst, Wiseman, & Hunsaker, 2019; Piano et al., 2019a). Moreover, this limited resistance could be explained by the negligible contribution provided by in-stream refugia because of the drought intensity and length. Unlike other studies, where pools and the hyporheic zone have been recognized to be primary drivers of the post-drought recovery of benthic organisms (Vander Vorste, Malard, & Datry, 2016; Verdonschot et al., 2015), the prolonged drying conditions here observed probably nullified the suitability of such refugia. Indeed, pools disappeared in our intermittent sites and also the survival of macroinvertebrates in the moist interstitial spaces appears unlikely under similar circumstances. To confirm this, data acquired by a piezometer showed that, in the intermittent site of the Po river, water was 2.5 m below the ground level for the majority of the time from July to December 2017 (unpublished data, see Supplementary Materials Fig. SM2). Our results showed that the passive recolonization by drift from the upstream section was probably the main factor facilitating the recovery of macroinvertebrates in Alpine streams, according to the results of other authors (Doretto et al., 2018; Flower, 2004).

However, marked differences were found among the two examined lotic systems, thus supporting the role of recruitment in macroinvertebrate community resilience to exceptional droughts. In the Pellice river we observed a progressive and significant increase in all the diversity metrics since the water resumption and multivariate analysis indicated a partial overlap in the community composition of permanent and intermittent sites. As water resumed in this river, no relevant changes in flow and environmental conditions were observed among sampling occasions. This aspect, combined with the shorter distance from the upstream nearest permanent site, probably explains the recovery dynamics here documented, as pointed out by other authors (Bogan, Boersma, & Lytle, 2015; Fritz & Doods, 2004). On the contrary, the rewetting process in the Po river was strongly influenced by the precipitation amount (Supplementary Materials Fig. SM1): after a steady raise in flow, the riverbed shrank over the time and dried completely around 45 days from the water resumption with flowing water that re-established only on the last sampling occasions. As a consequence, richness and density of macroinvertebrates generally peaked within the first 20 days and then collapsed, while even after 73 days the taxonomical composition of the intermittent and permanent sites was still different. In addition, also the greater distance between these two sampling stations probably explains why an appreciable recovery was not reached in this lotic ecosystem.

Van Looy et al. (2018) indicated also that the resource competition/facilitation plays an important role on the resilience after a disturbance of riverine communities. Although we did not assess directly the food availability and biotic interactions, temporal changes in the percentages of FFGs and ratio between scrapers and total collectors were examined. FFGs have been widely invoked to indirectly infer riverine ecosystem attributes and their use in biomonitoring is currently growing (Cummins et al., 2005; Doretto, Piano, Bona & Fenoglio, 2018; Merritt, Fenoglio & Cummins, 2017). Temporal patterns for FFGs were found for both rivers but, interestingly, significant variations in the ratio between scrapers and total collectors were observed only in the Po river. This ratio was here used as an indicator of the prevalence of autotrophy or heterotrophy, and our results suggest that probably the availability and quality of periphyton and organic matter were not influential factors in the Pellice river (Falasco et al., in preparation), while they affected, at least partially, the recolonization process in the Po river.

To conclude, this work stresses the importance of the recolonization by drift as the main mechanism for the post-drought recovery of macroinvertebrates in Alpine streams. This is in accordance with conceptual framework proposed by Van Looy et al. (2018), for which the recruitment from adjacent habitat sources is usually the main drivers of community resilience in connected river network. As the intermittent and permanent sites in this study were located few kilometers aside, we assume that this condition applied to our results. However, we also demonstrated that river-specific attributes, such as local climate conditions, hydrology and the distance from the nearest upstream perennial site can strongly influence the recovery process. Given the predicted increment in the frequency and magnitude of anthropogenic and climate droughts in the mountain areas, the results of this study offer important information for the management and conservation of Alpine streams and their biota.

Data Availability Statement (DAS)

The data that support the findings of this study are available from the corresponding author upon reasonable request.

References

Agenzia Regionale per la Protezione dell'Ambiente (ARPA) (2013) Idrologia in Piemonte nel 2012. Regione Piemonte, pp 23. <http://www.arpa.piemonte.it>

- Aspin, T. W., Khamis, K., Matthews, T. J., Milner, A. M., O'callaghan, M. J., Trimmer, M., ... & Ledger, M. E. (2019). Extreme drought pushes stream invertebrate communities over functional thresholds. *Global change biology*, 25(1), 230-244. <https://doi.org/10.1111/gcb.14495>
- Beniston, M. (2012). Impacts of climatic change on water and associated economic activities in the Swiss Alps. *Journal of Hydrology*, 412, 291-296. <https://doi.org/10.1016/j.jhydrol.2010.06.046>
- Bogan, M. T., Boersma, K. S., & Lytle, D. A. (2015). Resistance and resilience of invertebrate communities to seasonal and suprasedonal drought in arid-land headwater streams. *Freshwater Biology*, 60(12), 2547-2558. <https://doi.org/10.1111/fwb.12522>
- Bonada, N., Doledec, S., & Statzner, B. (2007). Taxonomic and biological trait differences of stream macroinvertebrate communities between mediterranean and temperate regions: implications for future climatic scenarios. *Global Change Biology*, 13(8), 1658-1671. <https://doi.org/10.1111/j.1365-2486.2007.01375.x>
- Boulton, A. J. (2003). Parallels and contrasts in the effects of drought on stream macroinvertebrate assemblages. *Freshwater Biology*, 48(7), 1173-1185. <https://doi.org/10.1046/j.1365-2427.2003.01084.x>
- Calapez, A. R., Elias, C. L., Almeida, S. F., & Feio, M. J. (2014). Extreme drought effects and recovery patterns in the benthic communities of temperate streams. *Limnetica*, 33(2), 281-296. <https://doi.org/10.23818/limn.33.22>
- Campaioli, S., Ghetti, P. F., Minelli, A., & Ruffo, S. (1994). Manuale per il riconoscimento dei macroinvertebrati delle acque dolci italiane (Vol. I). Trento: Provincia Autonoma di Trento.
- Campaioli, S., Ghetti, P. F., Minelli, A., & Ruffo, S. (1999). Manuale per il riconoscimento dei macroinvertebrati delle acque dolci italiane (Vol. II). Trento: Provincia Autonoma di Trento.
- Chester, E. T., & Robson, B. J. (2011). Drought refuges, spatial scale and recolonisation by invertebrates in non-perennial streams. *Freshwater Biology*, 56(10), 2094-2104. <https://doi.org/10.1111/j.1365-2427.2011.02644.x>
- Cummins, K. W., Merritt, R. W., & Andrade, P. C. (2005). The use of invertebrate functional groups to characterize ecosystem attributes in selected streams and rivers in south Brazil. *Studies on Neotropical Fauna and Environment*, 40(1), 69-89. <https://doi.org/10.1080/01650520400025720>
- Dawson, T. P., Jackson, S. T., House, J. I., Prentice, I. C., & Mace, G. M. (2011). Beyond predictions: biodiversity conservation in a changing climate. *Science*, 332(6025), 53-58. <https://doi.org/10.1126/science.1200303>
- Death, R. G. (2010). Disturbance and riverine benthic communities: what has it contributed to general ecological theory? *River Research and Applications*, 26(1), 15-25. <https://doi.org/10.1002/rra.1302>
- Doretto, A., Bona, F., Piano, E., Zanin, I., Eandi, A. C., & Fenoglio, S. (2017). Trophic availability buffers the detrimental effects of clogging in an alpine stream. *Science of the Total Environment*, 592, 503-511. <https://doi.org/10.1016/j.scitotenv.2017.03.108>

- Doretto, A., Piano, E., Bona, F., & Fenoglio, S. (2018a). How to assess the impact of fine sediments on the macroinvertebrate communities of alpine streams? A selection of the best metrics. *Ecological Indicators*, 84, 60-69. <https://doi.org/10.1016/j.ecolind.2017.08.041>
- Doretto, A., Piano, E., Falasco, E., Fenoglio, S., Bruno, M. C., & Bona, F. (2018b). Investigating the role of refuges and drift on the resilience of macroinvertebrate communities to drying conditions: An experiment in artificial streams. *River Research and Applications*, 34(7), 777-785. <https://doi.org/10.1002/rra.3294>
- Durance, I., & Ormerod, S. J. (2007). Climate change effects on upland stream macroinvertebrates over a 25-year period. *Global Change Biology*, 13(5), 942-957. <https://doi.org/10.1111/j.1365-2486.2007.01340.x>
- Falasco, E., Piano, E., Doretto, A., Fenoglio, S., & Bona, F. (2018). Lenticification in Alpine rivers: patterns of diatom assemblages and functional traits. *Aquatic sciences*, 80(4), 36. <https://doi.org/10.1007/s00027-018-0587-y>
- Fenoglio, S., Bo, T., & Bosi, G. (2006). Deep interstitial habitat as a refuge for *Agabus paludosus* (Fabricius)(Coleoptera: Dytiscidae) during summer droughts. *The Coleopterists Bulletin*, 60(1), 37-42. <https://doi.org/10.1649/842.1>
- Fenoglio, S., Bo, T., Cucco, M., & Malacarne, G. (2007). Response of benthic invertebrate assemblages to varying drought conditions in the Po river (NW Italy). *Italian Journal of Zoology*, 74(2), 191-201. <https://doi.org/10.1080/11250000701286696>
- Fenoglio, S., Bo, T., Cucco, M., Mercalli, L., & Malacarne, G. (2010). Effects of global climate change on freshwater biota: A review with special emphasis on the Italian situation. *Italian Journal of Zoology*, 77(4), 374-383. <https://doi.org/10.1080/11250000903176497>
- Fowler, R. T. (2004). The Recovery of Benthic Invertebrate Communities Following Dewatering in Two Braided Rivers. *Hydrobiologia*, 523(1), 17-28. <https://doi.org/10.1023/B:HYDR.0000033077.13139.7f>
- Fritz, K. M., & Dodds, W. K. (2004). Resistance and resilience of macroinvertebrate assemblages to drying and flood in a tallgrass prairie stream system. *Hydrobiologia*, 527(1), 99-112. <https://doi.org/10.1023/B:HYDR.0000043188.53497.9b>
- Gorbach, K. R., Shoda, M. E., Burky, A. J., & Benbow, M. E. (2014). Benthic community responses to water removal in tropical mountain streams. *River research and applications*, 30(6), 791-803. <https://doi.org/10.1002/rra.2679>
- Heino, J., Virkkala, R., & Toivonen, H. (2009) Climate change and freshwater biodiversity: detected patterns, future trends and adaptations in northern regions. *Biological Reviews*, 84(1), 39-54. <https://doi.org/10.1111/j.1469-185X.2008.00060.x>
- Herbst, D. B., Cooper, S. D., Medhurst, R. B., Wiseman, S. W., & Hunsaker, C. T. (2019). Drought ecohydrology alters the structure and function of benthic invertebrate communities in mountain streams. *Freshwater Biology*, 1-17. <https://doi.org/10.1111/fwb.13270>

- 1
2
3 405 Lake, P. S. (2003). Ecological effects of perturbation by drought in flowing waters. *Freshwater*
4 406 *biology*, 48(7), 1161-1172. <https://doi.org/10.1046/j.1365-2427.2003.01086.x>
- 6 407 Ledger, M. E., & Hildrew, A. G. (2001). Recolonization by the benthos of an acid stream following a
8 408 drought. *Archiv für Hydrobiologie*, 1-17. <https://doi.org/10.1127/archiv-hydrobiol/152/2001/1>
- 10 409 Ledger, M. E., Brown, L. E., Edwards, F. K. F. K., Milner, A. M., & Woodward, G. (2013). Drought alters
11 410 the structure and functioning of complex food webs. *Nature Climate Change*, 3(3), 223-227.
12 411 <https://doi.org/10.1038/nclimate1684>
- 14 412 Ledger, M. E., & Milner, A. M. (2015). Extreme events in running waters. *Freshwater Biology*, 60(12),
15 413 2455-2460. <https://doi.org/10.1111/fwb.12673>.
- 18 414 Leigh, C., Bonada, N., Boulton, A. J., Hugueny, B., Larned, S. T., Vander Vorste, R., & Datry, T. (2016).
19 415 Invertebrate assemblage responses and the dual roles of resistance and resilience to drying in
20 416 intermittent rivers. *Aquatic Sciences*, 78(2), 291-301. <https://doi.org/10.1007/s00027-015-0427-2>.
- 22 417 López-Rodríguez, M. J., Muñoz, C. M., Ripoll-Martín, E., & de Figueroa, J. M. T. (2019). Effect of shifts
23 418 in habitats and flow regime associated to water diversion for agriculture on the macroinvertebrate
24 419 community of a small watershed. *Aquatic Ecology*, 1-13. [https://doi.org/10.1007/s10452-019-](https://doi.org/10.1007/s10452-019-09703-6)
26 420 [09703-6](https://doi.org/10.1007/s10452-019-09703-6)
- 28 421 Mantyka-pringle, C. S., Martin, T. G., & Rhodes, J. R. (2012). Interactions between climate and
29 422 habitat loss effects on biodiversity: a systematic review and meta-analysis. *Global Change*
30 423 *Biology*, 18(4), 1239-1252. <https://doi.org/10.1111/j.1365-2486.2011.02593.x>
- 33 424 McKay, S. F., & King, A. J. (2006). Potential ecological effects of water extraction in small,
34 425 unregulated streams. *River Research and Applications*, 22(9), 1023-1037.
35 426 <https://doi.org/10.1002/rra.958>
- 37 427 Merritt, R.W., Cummins, K.W. and Berg, M.B. (2008) An Introduction to Aquatic Insects of North
38 428 America. 4th Edition, Kendall Hunt Publishers, Dubuque.
- 41 429 Merritt, R. W., Fenoglio, S., & Cummins, K. W. (2017). Promoting a functional macroinvertebrate
42 430 approach in the biomonitoring of Italian lotic systems. *Journal of Limnology*, 76(s1), 5-8.
43 431 <https://doi.org/10.4081/jlimnol.2016.1502>
- 45 432 Middelkoop, H., Daamen, K., Gellens, D., Grabs, W., Kwadijk, J. C., Lang, H., ... & Wilke, K. (2001).
46 433 Impact of climate change on hydrological regimes and water resources management in the Rhine
47 434 basin. *Climatic Change*, 49(1), 105-128. <https://doi.org/10.1023/A:1010784727448>
- 50 435 Oksanen, J., Blanchet, F. G., Friendly, M., Kindt, R., Legendre, P., McGlinn, D., Minchin, P. R., O'Hara,
51 436 R.B., Simpson, G. L., Solymos, P., Stevens, M. H. H., Szoecs, E., Wagner, H. (2015). Vegan: community
52 437 ecology package. R Package Version 2.2-1.
- 54 438 Otermin, A., Basaguren, A., & Pozo, J. (2002). Re-colonization by the macroinvertebrate community
55 439 after a drought period in a first-order stream (Agüera Basin, Northern Spain). *Limnetica*, 21(1-2),
56 440 117-128.

- Piano, E., Doretto, A., Falasco, E., Fenoglio, S., Gruppuso, L., Nizzoli, D., ... & Bona, F. (2019a). If Alpine streams run dry: the drought memory of benthic communities. *Aquatic Sciences*, 81(2), 32. <https://doi.org/10.1007/s00027-019-0629-0>
- Piano, E., Doretto, A., Falasco, E., Gruppuso, L., Fenoglio, S., & Bona, F. (2019b). The role of recurrent dewatering events in shaping ecological niches of scrapers in intermittent Alpine streams. *Hydrobiologia*, 1-13. <https://doi.org/10.1007/s10750-019-04021-2>
- Pinna, M., Marini, G., Cristiano, G., Mazzotta, L., Vignini, P., Cicolani, B., & Di Sabatino, A. (2016). Influence of aperiodic summer droughts on leaf litter breakdown and macroinvertebrate assemblages: testing the drying memory in a Central Apennines River (Aterno River, Italy). *Hydrobiologia*, 782(1), 111-126.
- R Development Core Team. (2018). R: a language and environment for statistical computing. Vienna: R Foundation for Statistical Computing.
- Richardson, J. S., & Sato, T. (2015). Resource subsidy flows across freshwater–terrestrial boundaries and influence on processes linking adjacent ecosystems. *Ecohydrology*, 8(3), 406-415. <https://doi.org/10.1002/eco.1488>
- Robson, B. J., Chester, E. T., & Austin, C. M. (2011). Why life history information matters: drought refuges and macroinvertebrate persistence in non-perennial streams subject to a drier climate. *Marine and Freshwater Research*, 62(7), 801-810. <https://doi.org/10.1071/MF10062>
- Smith, C. R., McCormick, P. V., Covich, A. P., & Golladay, S. W. (2017). Comparison of macroinvertebrate assemblages across a gradient of flow permanence in an agricultural watershed. *River Research and Applications*, 33(9), 1428-1438. <https://doi.org/10.1002/rra.3211>
- Storey, R. (2016). Macroinvertebrate community responses to duration, intensity and timing of annual dry events in intermittent forested and pasture streams. *Aquatic Sciences*, 78(2), 395-414. <https://doi.org/10.1007/s00027-015-0443-2>
- Strahler, A. N. (1957). Quantitative analysis of watershed geomorphology. *Eos - Transactions American Geophysical Union*, 38: 913-920. <https://doi.org/10.1029/TR038i006p00913>
- Tierno de Figueroa, J. M., López-Rodríguez, M. J., Fenoglio, S., Sánchez-Castillo, P., & Fochetti, R. (2013). Freshwater biodiversity in the rivers of the Mediterranean Basin. *Hydrobiologia*, 719, 137-186. <https://doi.org/10.1007/s10750-012-1281-z>
- Vander Vorste, R., Malard, F., & Datry, T. (2016). Is drift the primary process promoting the resilience of river invertebrate communities? A manipulative field experiment in an intermittent alluvial river. *Freshwater Biology*, 61(8), 1276-1292. <https://doi.org/10.1111/fwb.12658>
- Van Looy, K., Tonkin, J. D., Flourey, M., Leigh, C., Soininen, J., Larsen, S., ... & Datry, T. (2019). The three Rs of river ecosystem resilience: Resources, recruitment, and refugia. *River Research and Applications*, 35(2), 107-120. <https://doi.org/10.1002/rra.3396>
- Verdonschot, R., Oosten-Siedlecka, A. M., Braak, C. J., & Verdonschot, P. F. (2015). Macroinvertebrate survival during cessation of flow and streambed drying in a lowland stream. *Freshwater Biology*, 60(2), 282-296. <https://doi.org/10.1111/fwb.12479>

1
2
3 479 Wentworth, C. K. (1922). A scale of grade and class terms for clastic sediments. *The journal of*
4 480 *geology*, 30(5), 377-392.
5
6 481 Whitehead, P. G., Wilby, R. L., Battarbee, R. W., Kernan, M., & Wade, A. J. (2009). A review of the
7 482 potential impacts of climate change on surface water quality. *Hydrological Sciences Journal*, 54(1),
8 483 101-123. <https://doi.org/10.1623/hysj.54.1.101>
9
10 484 Wood, P. J., Boulton, A. J., Little, S., & Stubbington, R. (2010). Is the hyporheic zone a refugium for
11 485 aquatic macroinvertebrates during severe low flow conditions? *Fundamental and Applied*
12 486 *Limnology/Archiv für Hydrobiologie*, 176(4), 377-390. [https://doi.org/10.1127/1863-](https://doi.org/10.1127/1863-9135/2010/0176-0377)
13 487 [9135/2010/0176-0377](https://doi.org/10.1127/1863-9135/2010/0176-0377).
14
15 488 Wood, S., & Wood, M. S. (2015). Package ‘mgcv’. R package version, 1-7.
16
17 489 Wu, H., & Johnson, B. R. 2019. Climate change will both exacerbate and attenuate urbanization
18 490 impacts on streamflow regimes in southern Willamette Valley, Oregon. *River Research and*
19 491 *Applications*. <https://doi.org/10.1002/rra.3454>
20
21 492 Zuur, A. F., Ieno, E. N., & Elphick, C. S. (2010). A protocol for data exploration to avoid common
22 493 statistical problems. *Methods in ecology and evolution*, 1(1), 3-14. [https://doi.org/10.1111/j.2041-](https://doi.org/10.1111/j.2041-210X.2009.00001.x)
23 494 [210X.2009.00001.x](https://doi.org/10.1111/j.2041-210X.2009.00001.x)
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Tables

Table 1. Geographical information of the sampling sites.

River	Site	Coordinates	% Natural areas	% Agricultural areas	% Urbanized areas	Elevation (m.a.s.l.)	Distance between stations (Km)
Po	Perennial	367119E; 4945951N	93	6	1	474	5.5
	Intermittent	372959E; 4943103N	89	10	1	246	
Pellice	Perennial	364293E; 4963123N	92	6	2	422	3.1
	Intermittent	366638N; 4964043E	92	6	2	378	

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Table 2. Scheme of the sampling activity. Date = sampling date, Days = days from the water return in the intermittent sites.

Date	Days	Pellice sites		Po sites	
		Perennial	Intermittent	Perennial	Intermittent
12 th January 2018	3	-	Sampling	-	Sampling
16 th January 2018	7	-	Sampling	-	Sampling
19 th January 2018	10	Sampling	Sampling	Sampling	Sampling
23 th January 2018	14	-	Sampling	-	Sampling
30 th January 2018	21	-	Sampling	-	Sampling
1 st February 2018	23	-	Sampling	-	Sampling
9 th February 2018	31	-	Sampling	-	Sampling
23 th February 2018	45	-	Sampling	-	Dry
9 th March 2018	59	-	Sampling	-	Sampling
22 th March 2018	73	Sampling	Sampling	Sampling	Sampling

Table 3. Statistics of the GAMs for the macroinvertebrate community. Int = intercept, SE = standard error, z = z-value, t = t-value, River = studied rivers (i.e. Pellice, Po), χ^2 = Chi-square, F = F-value, P = p-value. Significant values are in bold.

Metric	Int	SE	z	River	χ^2	P
Taxa richness	2.214	0.046	48.370	Pellice	3.895	0.048
				Po	5.208	0.129
Total density	7.661	0.062	123.000	Pellice	10.520	0.020
				Po	111.730	<0.001
EPT richness	1.791	0.057	31.650	Pellice	3.827	0.050
				Po	3.319	0.238
EPT density	7.171	0.062	115.000	Pellice	21.700	<0.001
				Po	121.500	<0.001
% Collector-gatherers	-0.735	0.035	-21.010	Pellice	347.500	<0.001
				Po	152.200	<0.001
% Filterers	-0.637	0.036	-17.500	Pellice	549.700	<0.001
				Po	261.000	<0.001
% Predators	-4.064	0.119	-34.150	Pellice	48.078	<0.001
				Po	4.839	0.188
% Scrapers	-1.076	0.036	-30.010	Pellice	61.030	<0.001
				Po	215.950	<0.001
% Shredders	-4.335	0.121	-35.940	Pellice	5.986	0.055
				Po	2.214	0.137
Metric	Int	SE	t	River	F	P
Scrapers/Total collectors	0.605	0.061	9.905	Pellice	0.406	0.527
				Po	4.193	0.013

1
2
3 509 **Figure captions**
4
5 510 Fig. 1. Sampling sites and elevational range of the sub-basins of the Pellice and Po rivers.
6
7 511
8
9 512 Fig 2. PCA ordination plot. Labels indicate: river (Pe = Pellice, Po = Po), type of site (P = Permanent,
10 513 I = Intermittent) and sampling occasion expressed as days from the water return. Ellipses represent
11 514 standard deviations around the centroids of sampling sites of the two rivers (solid line = Pellice river,
12 515 dashed line = Po river).
13
14
15 516
16
17 517 Fig 3. NMDS ordination plot. Symbols represent the type of site (I = intermittent, P = perennial) for
18 518 each river. Colors represent the sampling occasions, indicated as number of days since the water
19 519 return. Ellipses represent standard deviations around the centroids of sampling sites of the two
20 520 rivers (solid line = Pellice river, dashed line = Po river).
21
22
23 521
24
25 522 Fig 4. Generalized Additive Models (GAMs) for (a) total taxa richness, (b) EPT richness, (c) total
26 523 density of macroinvertebrates and (d) EPT density. Black lines represent the predicted values of the
27 524 models, while the dashed lines represent 95% confidence interval.
28
29
30 525
31
32 526 Fig 5. Bars indicate the percentage of functional feeding groups in the (a) Pellice and (b) Po rivers on
33 527 each sampling occasion, expressed as days from the water return. GAMs for the ratio between
34 528 scrapers and total collectors during the rewetting phase (c): black lines represent the predicted
35 529 values of the models, while the dashed lines represent 95% confidence interval.
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Fig. 1. Sampling sites and elevational range of the sub-basins of the Pellice and Po rivers.

296x209mm (300 x 300 DPI)

Fig 2. PCA ordination plot. Labels indicate: river (Pe = Pellice, Po = Po), type of site (P = Permanent, I = Intermittent) and sampling occasion expressed as days from the water return. Ellipses represent standard deviations around the centroids of sampling sites of the two rivers (solid line = Pellice river, dashed line = Po river).

169x169mm (300 x 300 DPI)

Fig 3. NMDS ordination plot. Symbols represent the type of site (I = intermittent, P = perennial) for each river. Colors represent the sampling occasions, indicated as number of days since the water return. Ellipses represent standard deviations around the centroids of sampling sites of the two rivers (solid line = Pellice river, dashed line = Po river).

169x109mm (300 x 300 DPI)

Fig 4. Generalized Additive Models (GAMs) for (a) total taxa richness, (b) EPT richness, (c) total density of macroinvertebrates and (d) EPT density. Black lines represent the predicted values of the models, while the dashed lines represent 95% confidence interval.

169x114mm (300 x 300 DPI)

Fig 5. Bars indicate the percentage of functional feeding groups in the (a) Pellice and (b) Po rivers on each sampling occasion, expressed as days from the water return. GAMs for the ratio between scrapers and total collectors during the rewetting phase (c): black lines represent the predicted values of the models, while the dashed lines represent 95% confidence interval.

169x129mm (300 x 300 DPI)