AperTO - Archivio Istituzionale Open Access dell'Università di Torino

Gastrointestinal helminths of wolves *Canis lupus* Linnaeus, 1758 in Piedmont, north-western Italy

This is the author's manuscript

Original Citation:

Availability:

This version is available <http://hdl.handle.net/2318/1725561> since 2020-01-28T12:35:46Z

Published version:

DOI:10.1017/S0022149X19000841

Terms of use:

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)

Gastrointestinal helminths of wolves (*Canis lupus* Linnaeus, 1758) in Piedmont, north-western Italy

M.R.P. de Macedo^{1,*}, S. Zanet², S. Bruno², A. Tolosano³, F. Marucco⁴, L. Rossiz, G. Muller¹ and E. Ferroglia²

¹Laboratory of Wildlife Parasitology, University of Pelotas, Avenida Eliseu Maciel s/n 96010-610, Capão do Leão, Brazil; ²Department of Veterinary Sciences, University of Turin, 10095 Grugliasco (TO), Italy; ³Provincia di Torino, Corpo di vigilanza faunistico-ambientale, Torino, Italy and ⁴Project Lupo Interreg, Regione Piemonte, Parco Naturale Alpi Marittime, Valdieri, Italy

Abstract

Free-ranging grey wolves (*Canis lupus*), which are presently recolonizing Italy, can be parasitized by a diversity of helminths, but have rarely been subject to studies of their parasites. Therefore, this study aims to determine the prevalence of gastrointestinal helminths of road-killed grey wolves from the Piedmont region of Italy. Forty-two wolves were collected and examined for the presence of helminths. We recorded 12 helminth species: nine Nematoda and three Cestoda. The nematodes were: *Ancylostoma caninum* (7.1%), *Capillaria* sp. (2.4%), *Molineus* sp. (2.4%), *Pterygodermatites affinis* (11.9%), *Physaloptera sibirica* (9.5%), *Toxocara canis* (9.5%), *Toxascaris leonina* (2.4%) and *Uncinaria stenocephala* (26.2%); the cestodes were: *Dipylidium caninum* (4.8%), *Mesocostoides* sp. (4.8%) and *Taenia multiceps* (76.2%). *Physaloptera sibirica* had the highest mean intensity and *T. multiceps* had the highest prevalence. Based on age and sex, no differences in the intensity or prevalence of helminth species were found among the hosts. *Molineus* sp. was recorded for the first time in wolves from the Palearctic region; *P. affinis* and *P. sibirica* are respectively reported for the first time in wolves from Europe and Italy.

Introduction

The grey wolf is widely distributed in North America and Eurasia, including Italy, where the wolf population declined until the 1970s, when it was estimated to comprise only 100 individuals. From the end of the 1970s to today, the population increased to approximately 1600–1900 individuals, but fluctuations occur every year due to natural and human factors (Mattioli et al., 2018). Wolf populations have expanded through a unidirectional migration from the Apennines to the Western Italian and French Alps (Fabbri et al., 2007). *Canis lupus* Linnaeus, 1758 is regarded as a protected species (Council Directive 92/43/EEC on the Conservation of Natural Habitats and of Wild Fauna and Flora). The Piedmont Region carries out wolf re-colonization throughout the Italian Alps, by the development of a monitoring and research program on the natural increase of wolves; the program aims to collect the technical data necessary to protect the species and confirm its impact on economic activities (Canavese & Marucco, 2016). Within the regional Piedmont territory, wolves are concentrated mainly in the Turin and Cuneo provinces, which are alpine sectors, but certain packs maintain transboundary territories that cross over from the Italian to the French Alps (Marucco et al., 2013). Information on helminth infections in carnivores, which occupy the highest positions in food chains, is still incomplete. However, such information is essential because parasites have roles in the stabilization of food-web dynamics by regulating host (e.g. specific parasites) abundance or by connections among a variety of host species (e.g. heteroxenous parasites). Diseases are recognized as significant issues for wild carnivore conservation, particularly where there is potential for outbreaks of common pathogens transmitted between domestic and wild canids (Alexander et al., 2010).

Wolves share several of their parasite species with other canids, including helminths of medical and veterinary importance (Craig & Craig, 2005). Nevertheless, there is little information regarding the health status or parasitic fauna of wolves in Italy, and specific data regarding the epidemiological situation of parasites in Italian wolves are scarce. Reports concerning *Echinococcus granulosus* (Batsch, 1786) Rudolphi, 1805 and *Trichinella* Railliet, 1895 were published in the 1980s and 2000s (Arru et al., 1986; Guberti et al., 1993, 2004). More recently, new research concerning *Thelazia callipaeda* Railliet & Henry, 1910 (Otranto et al., 2007) and *Angiostrongylus vasorum* (Baillet, 1866) Kamensky, 1905 (Eleni et al., 2014) have been published. Guberti et al. (1993) reported 12 species of intestinal helminths, the most complete

<https://doi.org/10.1017/S0022149X19000841>

Downloaded from <https://www.cambridge.org/core>. Università di Torino, on 28 Jan 2020 at 10:46:31, subject to the Cambridge Core terms of use, available at <https://www.cambridge.org/core/terms>.

study concerning wolves in Italy. Therefore, knowledge about wolf helminths can contribute to analysing specific risks and to planning appropriate methods for intervention, primarily in the Italian Alps where no details regarding helminth fauna have been published.

This study aims to report helminth species infecting the

gastrointestinal tract of free-ranging wolves in the north-western Italian Alps, Piedmont Region.

Materials and methods

Forty-two ($n = 42$) road-killed grey wolves were collected between November 2003 and December 2014 in the Piedmont Region (fig. 1). The carcasses were maintained at 4°C until post-mortem examination and necropsy. The 28 males and 14 females were sorted into two age classes: juveniles (not exceeding their second spring) and adults. In the course of necropsy, the gastrointestinal tract was removed, labelled and frozen at -20°C. Before parasitological examination, the packages were defrosted overnight at room temperature.

The helminths were collected as described by Amato & Amato (2010). The stomach and small and large intestines were individually opened and washed with tap water for helminth collection and identification. The mucosa was scraped exactly to remove the contents entirely. Using a stereomicroscope, the contents of each organ were examined in aliquots of 50 ml in Petri dishes. Nematodes were cleared in lactophenol and mounted between a slide and coverslip for identification (Amato & Amato, 2010). Nematodes were identified using keys and descriptions (Anderson et al., 2009; Gibbons, 2010). Cestoda were compressed and fixed in alcohol-formalin-acetic acid (AFA) solution (alcohol 70%GL 92%, formalin 5%, acetic acid 3%), and stained with Langeron's Carmine. The specimens were mounted in Canada balsam and identified using standard taxonomic keys (Esch & Self, 1965; Khalil et al., 1994; Loos-Frank, 2000; Padgett et al., 2005; Hrcakova et al., 2011).

The parameters for prevalence, mean abundance and mean intensity were calculated according to Bush et al. (1997). Fisher's exact test was used to test prevalence data against age, sex and season (winter or summer). Calculations were performed using the R software package (www.r-project.org), and statistical significance was inferred if the specific P-value of the test was ≤ 0.05 .

Results and discussion

The wolves examined included 28 males (17 adults, 11 juveniles) and 14 females (eight adults, six juveniles). The mean weight of the carcasses for the females was 24.9 kg (standard deviation (SD) 5.06; range 17.50–31.80 kg), for the males 27.41 kg (SD 8; range 15–37 kg), for the adults 32.54 kg (SD 3.85; range 25–37 kg) and for the juveniles 20.74 kg (SD 3.85; range 15–25.25 kg). Thirty-six of the 42 wolves examined contained helminths. The total prevalence of infection was 85.7%. Of the recorded parasites, there were nine species of nematode and three cestode species (table 1). Co-infections involving two or more helminth species per animal were found in 50% of the cases that presented two or more nematodes and one cestode (e.g. *Physaloptera sibirica* Petrov & Gorbunov, 1931, *Uncinaria stenocephala* Railliet, 1884 and *Taenia multiceps* Leske, 1780); two nematodes (e.g. *Toxocara canis* (Werner, 1782) Stiles, 1905 and *U. stenocephala*), two cestodes (e.g. *Mesocostoides* sp. Vaillant, 1863 and *T. multiceps*), one nematode and two cestodes (e.g. *U. stenocephala*, *Dipylidium caninum* (Linnaeus, 1758) Leuckart, 1863 and *T. multiceps*), one nematode and one cestode (e.g. *T. canis* and *T. multiceps*); and three nematodes (e.g. *Pterygodermatites affinis* Jägerskiöld, 1904, *P. sibirica* and *U. stenocephala*). In general, the most prevalent helminths were *T. multiceps* (76.2%), followed by nematodes *U. stenocephala* (26.2%) and *P. affinis* (11.9%); whereas, *P. sibirica* demonstrated the highest mean intensity (43.2%), followed by *Mesocostoides* sp. (9.5%) and *P. affinis* (7.2%). *Taenia multiceps*, *P. sibirica* and *U. stenocephala* were, respectively, the parasites with the highest mean abundance. Our data presented higher prevalences of the same parasite species in adults compared to juveniles and in females compared to males (table 2), but the mean intensity was higher in females

and juveniles, although not statistically significant ($P > 0.05$). On the other hand, due to the small proportion of positive animals for each helminth species, and the low number of parasites in the sample group, it was not possible to explore the effects of sex, age and season against mean intensity for each parasite species. Further, no significant differences in any helminth species were found relative to sex, age or season in terms of their prevalence ($P > 0.05$).

In accordance with other studies in Europe, *U. stenocephala* is among the most prevalent and frequently reported nematode species. Shimalov & Shimalov (2000) reported a prevalence of 15.4%. However, Popiolek et al. (2007) and Bagrale et al. (2009) reported prevalences of 37% and 41.2%, respectively. Our data are in agreement with previous investigations, yet *U. stenocephala* was more prevalent than *Ancylostoma caninum* (Ercolani, 1859) Hall, 1913. From a literature review based on coprological exams and necropsy findings, both nematode species are found in wolves in several areas of Europe and North America. Differences in prevalence between the two species (*U. stenocephala*, 26.2% vs. *A. caninum*, 7.1%) are comparable to those prevalences reported in similar studies. Until the present study, there had been no reports of *P. affinis* parasitizing wolves in Europe or North America. In this case, it was found in 11.9% of samples. The prevalence of infection is within the range of values reported for red foxes in Italy, varying from 8% (Capelli et al., 2003) to 19.4% (Cerbo et al., 2008). We suggest that in the area studied, and more generally in the Alps (where the population of wolves is expanding), suitable ecological conditions exist for *P. affinis* to utilize the wolf as a new host successfully. *Pterygodermatites affinis* depends on the frequency of ingestion by carnivores of intermediate (insect) or paratenic (small mammals, reptiles and birds) hosts. The stomachs of some wolves contained parts of small mammals, reptiles and arthropods, but primarily comprised bones of cervids. Similarly, *P. sibirica* has been reported parasitizing carnivores and mustelid species such as *Martes martes* (Segovia et al., 2007) in the Iberian Peninsula, and *Vulpes vulpes* and *Meles meles* in the Western Italian Alps (Ferroglia et al., 2009), but not as yet wolves. We did find *P. sibirica* in wolves from high-altitude areas with colder climates. This finding is notable because *P. sibirica* constitutes a new component for helminth fauna in Italian wolves.

Molineus Cameron, 1923 is a common roundworm in carnivores. For example, *Molineus patens* Dujardin, 1845 and *Molineus legerae* Durette-Desset & Pesson, 1987 were found in mustelids and canids (Manfredi et al., 2003; Sato & Suzuki, 2006; Segovia et al., 2007; Popiolek et al., 2009), but in wolves, it is the first report of this genus. The new data contribute to the knowledge of the geographical distribution and variety of hosts for this parasite.

The ascarid worms *Toxascaris leonine* (Linstow, 1902) Leiper, 1907 and *T. canis* presented low prevalence in our samples. The

2 M.R.P. de Macedo et al.

<https://doi.org/10.1017/S0022149X19000841>

Downloaded from <https://www.cambridge.org/core>. Università di Torino, on 28 Jan 2020 at 10:46:31, subject to the Cambridge Core terms of use, available at <https://www.cambridge.org/core/terms>.

prevalence of *T. leonine* (2.4%) is similar to the prevalence determined for Poland (Kloch et al., 2005; Popiolek et al., 2007).

Toxocara canis is one of the most common parasites of both wild and domestic canids, and it is a recognized worm in both Italian canids and wolves (Guberti et al., 1993). Thus, our data confirm the persistence of *T. canis* infecting wolves in Italy.

Fig. 1. Map showing locations of where the 42 wolves used in the current study were collected from in Piedmont, north-western Italy, between November 2003 and December 2014.

Table 1. Summary statistics of gastrointestinal helminths isolated from 42 wolves of Piedmont, by post-mortem examination.

Parasite species Number of animals infected_a Minimum/maximum per animal Mean intensity_b

Nematoda

Ancylostoma caninum 3 (7.1%; 0–11) 2–2 2_c

Capillaria sp. 1 (2.4%; 0–7) 1 1_d

Molineus sp. 1 (2.4%; 0–7) 1 1_d

Pterygodermatites affinis 5 (11.9%; 2–22) 1–19 7.2 (1.6–14.6)
Physaloptera sibirica 4 (9.5%; 0–19) 3–138 43.2 (3.5–110)
Toxocara canis 4 (9.5%; 0–19) 1–6 2.7 (1.2–5)
Toxascaris leonina 1 (2.4%; 0–7) 5–5 5_d
Uncinaria stenocephala 11 (26.2%; 12–40) 1–13 3.4 (1.7–6.7)

Cestoda

Dipylidium caninum 2 (4.8%; 0–11) 2–4 3(2–3)
Mesocestoides sp. 2 (4.8%; 2–7) 5–14 9.5 (5–95)
Taenia multiceps 32 (76.2%; 63–90) 1–18 6.1 (4.8–7.9)

^aPrevalence and 95% confidence limits in parentheses.

^bConfidence limits in parentheses.

^cIntensity is constant, confidence limits were not calculated.

^dThere is only one infected host in the sample, confidence limits cannot be calculated.

Journal of Helminthology 3

<https://doi.org/10.1017/S0022149X19000841>

Downloaded from <https://www.cambridge.org/core>. Università di Torino, on 28 Jan 2020 at 10:46:31, subject to the Cambridge Core terms of use, available at <https://www.cambridge.org/core/terms>.

Table 2. Prevalence and mean intensity (MI) of helminth infections in wolves, according to sex and age.

Species of helminth

Females Males Juveniles Adults

n = 14 n = 28 n = 17 n = 25

% MI % MI % MI % MI

Nematoda

Ancylostoma caninum 7.1 (0.4–31.7) 2_a 7.1 (1.3–22.9)_b 2 5.9 (0.3–28.7) 2_a 8 (1.4–25.6) 2_{a,b}
Capillaria sp. 0 0 3.6 (0.2–17.5)_a 1 0 0 4 (0.2–19.6) 1_a
Molineus sp. 7.1 (0.4–31.7) 1_a 0 (0–11.9) 0 5.9 (0.3–28.7) 1_a 0 0
Pterygodermatites affinis 21.4 (6.1–50) 5.3(2–8.7) 7.1 (1.3–22.9) 10 (1–10) 17.6 (5–41.7) 11 (2–16.7) 8 (1.4–25.6) 1.5 (1–1.5)
Physaloptera sibirica 7.1 (0.4–31.7) 138_a 10.7 (3–28.2) 11.7 (3–20) 17.6 (5–41.7) 56.7 (4–101) 4 (0.2–19.6) 3_a
Toxocara canis 0 0 14.3 (5–31.9) 2.7 (1.2–5) 17.6 (5–41.7) 3.3 (2–4.7) 4 (0.2–19.6) 1_a
Toxascaris leonina 7.1 (0.4–31.7) 5_a 0 0 0 0 0
Uncinaria stenocephala 21.4 (6.1–50) 4 (1–6.7) 28.6 (14.2–48.2) 3.2 (1.7–7.5) 23.5 (16.6–59.4) 2.8 (1.3–6.5) 20 (8.2–39.8) 4.2 (1.6–8.8)

Cestoda

Dipylidium caninum 7.1 (0.4–31.7) 2_a 3.6 (0.2–17.5)_a 4_a 0 0 8 (1.4–25.6) 3 (2–3)
Mesocestoides sp. 7.1 (0.4–31.7) 5_a 3.6 (0.2–17.5)_a 14_a 0 0 8 (1.4–25.6) 9.5 (5–9.5)
Taenia multiceps 71.4 (42.6–89.6) 7.3 (4.3–10.9) 78.6 (59–90.2) 5.6 (4.4–7.5) 58.8 (35–80.4) 5.9 (4.1–10.6) 88 (69.7–96.7) 6.2 (4.6–8)

^aThere is only one infected host in the sample, confidence limits cannot be calculated.

^bIntensity is constant, confidence limits were not calculated.

4 M.R.P. de Macedo et al.

<https://doi.org/10.1017/S0022149X19000841>

Downloaded from <https://www.cambridge.org/core>. Università di Torino, on 28 Jan 2020 at 10:46:31, subject to the Cambridge Core terms of use, available at <https://www.cambridge.org/core/terms>.

Three species of cestodes were found in this investigation:

T. multiceps (P = 76.2%), *D. caninum* and *Mesocestoides* sp., all three with prevalence P = 4.8%. The prevalence reported for *T. multiceps*, which frequently infects a variety of canids worldwide, was higher than the prevalence determined in previous studies carried out in Europe (Segovia et al., 2003; Moks et al., 2006), particularly compared to those values reported in Italy by Guberti et al. (1993). We attribute this occurrence to recent conservation efforts to promote both wolves and their prey. The preservation of prey and predator may have led to parasite life-cycle stabilization and subsequent host–parasite relationship consolidation.

The lower prevalence of *Mesocestoides* sp. and *D. caninum* is consistent with data on the species reported by Guberti et al. (1993) – 8.7% in Italy; Shimalov & Shimalov (2000) – 7.7% in Belarus; and Bagrade et al. (2009) – 5.9% in Latvia. The low prevalence of *D. caninum* reported in the present study is consistent with the findings of Guberti et al. (1993) in Italy (2%), and of Shimalov & Shimalov (2000) in Belarus (15.4%). Concerning *Mesocestoides* sp. infection data observed during the investigation, we can correlate parasitism with the dietary preferences of wolves. The infective phase of *Mesocestoides* occurs in several small vertebrates (amphibians, reptiles and small mammals); however, the stomach contents of the examined wolves predominantly comprised deer bones. The *Mesocestoides* species could not be identified because not all of the specimens were complete.

Most of the helminth species occurring in wolves are known to be parasites of both domestic and wild animals, but it is difficult to draw conclusions about the wolves' epidemiological role because the populations are small, and they have an extensive home range. As the populations grow and disperse to new areas, there may be exchanges of parasites between wolves and other carnivore species. The parasite diversity recorded in wolves from the study area was smaller compared to results reported for other study locations,

such as Belarus, where 24 species of helminths were recorded (Shimalov & Shimalov, 2000), but the mean of species is similar to other European areas and Italy, where between four and 14 species were recorded (Guberti et al., 1993; Kloch et al., 2005; Moks et al., 2006; Popiolek et al., 2007; Fiocchi et al., 2016; Bindke et al., 2017). Nevertheless, we have confirmed the stability of parasitic relationships between wolves and helminths over the last few decades. We have also increased knowledge concerning the geographic distribution of *P. affinis*, as well as reporting wolves as hosts to both *P. sibirica* and *Molineus* sp.

Helminths such as *T. canis* and *D. caninum* are commonly found in both domestic and wild carnivores, and preventive measures, treatment and epidemiology are well described in the literature.

Physaloptera spp. is a well-known agent of gastritis, vomiting and weight loss in dogs, but little is known about its pathogenic role in wildlife. Thus, we emphasize that further studies are required to evaluate the pathogenicity of *P. sibirica* in wild animals and its effects on host life-history traits.

Financial support. This work was supported by the Coordination of Improvement of Higher Education Personnel (CAPES), Ministry of Education, Brazil (process number 99999.003555/2014-03).

Conflicts of interest. None.

References

- Alexander KA, McNutt JW, Briggs MB, Standers PE, Funston P, Hemson G, Keet D and van Vuuren M (2010) Multi-host pathogens and carnivore management in southern Africa. *Comparative Immunology, Microbiology and Infectious Diseases* 33, 249–265.
- Amato J and Amato S (2010) Técnicas gerais para coleta e preparação de helmintos endoparasitos de aves. pp. 369–393 in Von Matter S, Straube F, Accordi I, Piacentini V, Cândido JJ (Eds) *Ornitologia e conservação: ciência aplicada, técnicas de pesquisa e levantamento*. Vol. 1. Rio de Janeiro, Technical Books.
- Anderson RC, Chabaud AG and Willmott S (2009) Keys to the nematode parasites of vertebrates: archival Volume. Oxfordshire, CABI Publishing. 443 pp.
- Arru E, Garippa G and Fico R (1986) Sulla presenza di *Echinococcus granulosus* nella volpe (*Vulpes vulpes*) e nel lupo (*Canis lupus*). *Atti della Società Italiana delle Scienze Veterinarie* 42, 1089–1092.
- Bagrade G, Kirjusina M, Vismanis K and Ozolins J (2009) Helminth parasites of the wolf *Canis lupus* from Latvia. *Journal of Helminthology* 83, 63–68.
- Bindke JD, Springer A, Boer M and Strube C (2017) Helminth fauna in captive European gray wolves (*Canis lupus lupus*) in Germany. *Frontiers in Veterinary Science* 4, 228.
- Bush AO, Lafferty KD, Lotz JM and Shostak AW (1997) Parasitology meets ecology on its own terms: Margolis et al. revisited. *Journal of Parasitology* 83, 575–583.
- Canavese G and Marucco F (2016) The Project LIFE WolfAlps: new knowledge for coordinated conservation actions. *Proceedings II Conference LIFE WolfAlps – The Wolf Population in the Alps: Status and Management*, Cuneo, 22 January 2016, Project LIFE 12 NAT/IT/00080 WOLFALPS127.
- Capelli G, Stancampiano L, Magi M, Poglayen G and Guberti V (2003) Diversità delle comunità parassitarie intestinali in tre popolazioni di volpi. *Journal of Mountain Ecology* 7, 199–205.
- Cerbo A, Manfredi M, Trevisiol K, Bregoli M, Ferrari N, Pirinesi F and Bazzoli S (2008) Intestinal helminth communities of the red fox (*Vulpes vulpes* L.) in the Italian Alps. *Acta Parasitologica* 53, 302–311.
- Craig HL and Craig PS (2005) Helminth parasites of wolves (*Canis lupus*): a species list and an analysis of published prevalence studies in Nearctic and Palearctic populations. *Journal of Helminthology* 79, 95–103.
- Eleni C, De Liberato C, Azam D, Morgan ER and Traversa D (2014) *Angiostrongylus vasorum* in wolves in Italy. *International Journal for Parasitology: Parasites and Wildlife* 3, 12–14.
- Esch GW and Self JT (1965) A critical study of the taxonomy of *Taenia pisiformis* Bloch, 1780; *Multiceps multiceps* (Leske, 1780); and *Hydatigera taeniaeformis* Batsch, 1786. *Journal of Parasitology* 51, 932–937.
- Fabbri E, Miquel C, Lucchini V, et al. (2007) From the Apennines to the Alps: colonization genetics of the naturally expanding Italian wolf (*Canis lupus*) population. *Molecular Ecology* 16, 1661–1671.
- Ferroglio E, Ragagli C and Trisciuglio A (2009) *Physaloptera sibirica* in foxes and badgers from the Western Alps (Italy). *Veterinary Parasitology* 163, 164–166.
- Fiocchi A, Gustinelli A, Gelmini L, Rugna G, Renzi M, Fontana MC and Poglayen G (2016) Helminth parasites of the red fox *Vulpes vulpes* (L., 1758) and the wolf *Canis lupus italicus* Altobello, 1921 in

- Emilia-Romagna, Italy. *Italian Journal of Zoology* 83, 503–513.
- Gibbons LM (2010) Keys to the nematode parasite of vertebrates: supplementary Volume. Oxfordshire, CABI Publishing. 416 pp.
- Guberti V, Stancampiano L and Francisci F (1993) Intestinal helminth parasite community in wolves (*Canis lupus*) in Italy. *Parassitologia* 35, 59–65.
- Guberti V, Bolognini M, Lanfranchi P and Battelli G (2004) *Echinococcus granulosus* in the wolf in Italy. *Parassitologia* 46, 425–427.
- Hrckova G, Miterpakova M, O'Connor A, Snabel V and Olson PD (2011) Molecular and morphological circumscription of *Mesocestoides* tapeworms from red foxes (*Vulpes vulpes*) in central Europe. *Parasitology* 138, 638–647.
- Khalil LF, Jones A and Bray RA (1994) Keys to the cestode parasites of vertebrates. Oxford, Oxford University Press. 768 pp.
- Kloch A, Bednarska M and Bajer A (2005) Intestinal macro- and microparasites of wolves (*Canis lupus* L.) from North-Eastern Poland recovered by coprological study. *Annals of Agricultural and Environmental Medicine* 12, 237–245.
- Loos-Frank B (2000) An up-date of Verster's (1969) 'Taxonomic revision of the genus *Taenia* Linnaeus' (Cestoda) in table format. *Systematic Parasitology* 45, 155–183.
- Journal of Helminthology 5**
<https://doi.org/10.1017/S0022149X19000841>
 Downloaded from <https://www.cambridge.org/core>. Università di Torino, on 28 Jan 2020 at 10:46:31, subject to the Cambridge Core terms of use, available at <https://www.cambridge.org/core/terms>.
- Manfredi MT, Giacometti A, Fraquelli C and Piccolo G (2003) Studio della popolazione elmintica in volpi (*Vulpes vulpes*) del Trentino Alto-Adige. *Journal of Mountain Ecology* 7, 261–263.
- Marucco F, Marboutin E, Duchamp C, Avanzinelli E, Potocnik H and Rauer G (2013) Wolf Alpine Group. Alpine Convention Group. Cogne, WISO Platform, 17 p.
- Mattioli L, Canu A, Passilongo D, Scandura M and Apollonio M (2018) Estimation of pack density in grey wolf (*Canis lupus*) by applying spatially explicit capture-recapture models to camera trap data supported by genetic monitoring. *Frontiers in Zoology* 15, 38–38.
- Moks E, Jogisalu I, Saarma U, Talvik H, Jarvis T and Valdmann H (2006) Helminthologic survey of the wolf (*Canis lupus*) in Estonia, with an emphasis on *Echinococcus granulosus*. *Journal of Wildlife Diseases* 42, 359–365.
- Otranto D, Cantacessi C, Mallia E and Lia RP (2007) First report of *Thelazia callipaeda* (Spirurida, Thelaziidae) in wolves in Italy. *Journal of Wildlife Diseases* 43, 508–511.
- Padgett KA, Nadler SA, Munson L, Sacks B and Boyce WM (2005) Systematics of *Mesocestoides* (Cestoda: Mesocestoididae): evaluation of molecular and morphological variation among isolates. *Journal of Parasitology* 91, 1435–1443.
- Popiolek M, Szczesna J, Nowaka S and Myslajeka RW (2007) Helminth infections in faecal samples of wolves *Canis lupus* L. from the western Beskidy Mountains in southern Poland. *Journal of Helminthology* 81, 339–344.
- Popiolek M, Jarnecki H and Luczynski T (2009) The first record of *Molineus patens* (Dujardin, 1845) (Nematoda, Molineidae) in the ermine (*Mustela erminea* L.) in Poland. *Wiadomości Parazytologiczne* 55, 433–435.
- Sato H and Suzuki K (2006) Gastrointestinal helminths of feral raccoons (*Procyon lotor*) in Wakayama Prefecture, Japan. *Journal of Veterinary Medical Science* 68, 311–318.
- Segovia JM, Guerrero R, Torres J, Miquel J and Feliu C (2003) Ecological analyses of the intestinal helminth communities of the wolf, *Canis lupus*, in Spain. *Folia Parasitologica* 50, 231–236.
- Segovia J-M, Torres J, Miquel J, Sospedra E, Guerrero R and Feliu C (2007) Analysis of helminth communities of the pine marten, *Martes martes*, in Spain: mainland and insular data. *Acta Parasitologica* 52, 156–164.
- Shimalov VV and Shimalov VT (2000) Helminth fauna of the wolf (*Canis lupus* Linnaeus, 1758) in Belorussian Polesie. *Parasitology Research* 86, 163–164.
- 6 M.R.P. de Macedo et al.**
<https://doi.org/10.1017/S0022149X19000841>
 Downloaded from <https://www.cambridge.org/core>. Università di Torino, on 28 Jan 2020 at 10:46:31, subject to the Cambridge Core terms of use, available at <https://www.cambridge.org/core/terms>.