AperTO - Archivio Istituzionale Open Access dell'Università di Torino

Clostridium cellulovorans metabolism of cellulose as studied by comparative proteomic approach

This is a pre print version of the following article:

Original Citation:

Availability:

This version is available <http://hdl.handle.net/2318/1728141> since 2020-02-18T16:06:20Z

Published version:

DOI:10.1016/j.jprot.2020.103667

Terms of use:

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)

Manuscript Number:

Title: Clostridium cellulovorans metabolism of cellulose as studied by comparative proteomic approach

Article Type: Full Length Article

Section/Category: Original Article

Keywords: ATP; acetate; ethanol; Alcohol dehydrogenase; pyruvate phosphate dikinase; glucose

Corresponding Author: Dr. Roberto Mazzoli, Ph.D.

Corresponding Author's Institution: University of Torino

First Author: Giulia Usai

Order of Authors: Giulia Usai; Simona Cirrincione; Angela Re; Marcello Manfredi; Andrea Pagnani; Enrica Pessione; Roberto Mazzoli, Ph.D.

Abstract: Clostridium cellulovorans is among the most promising candidates for consolidated bioprocessing (CBP) of cellulosic biomass to liquid biofuels (ethanol, butanol). C. cellulovorans can metabolize all the main plant polysaccharides and its main catabolite is butyrate. This makes this strain a potential butanol producer since most reactions of butyrate and butanol biosynthesis from acetyl-CoA are common. Recent studies demonstrated that introduction of a single heterologous alcohol/aldehyde dehydrogenase diverts the branching-point intermediate, i.e. butyryl-CoA, towards butanol production in this strain. Despite C. cellulovorans potential for CBP of plant biomass, engineering its metabolic pathways towards industrial utilization requires better understanding of its metabolism. The present study aimed at improving comprehension of cellulose metabolism in C. cellulovorans by comparing growth kinetics, substrate consumption/product accumulation and whole-cell soluble proteome (data available via ProteomeXchange, identifier PXD015487) with those of the same strain grown on a soluble carbohydrate, glucose, as the main carbon source. Modulations of the central carbon metabolism in response to different growth substrate were detected, including regulation of glycolytic enzymes, fermentation pathways and nitrogen assimilation possibly affecting the redox balance. Higher energy expenditure seems to occur in cellulose-grown C. cellulovorans, which induces up-regulation of ATP synthetic pathways, e.g. acetate production and ATP synthase.

Suggested Reviewers: Daniel G. Olson PhD
Research assistant professor, Thayer School of Engineering, Dartmouth college, Hanover, NH 03755, USA
Daniel.G.Olson@dartmouth.edu
Expertise in the metabolism and metabolic engineering of anaerobic cellulolytic bacteria

Adam M. Guss Ph.D.

Assistant Professor, Biosciences Division, Oak Ridge National Laboratory,
One Bethel Valley Road, Oak Ridge, TN 37831, USA
gussam@ornl.gov
Expertise in the metabolism and metabolic engineering of anaerobic
cellulolytic bacteria

David B. Levin
Professor, Department of Biosystems Engineering, University of Manitoba,
Winnipeg R3T 2N2, MB, Canada.
david.levin@umanitoba.ca
Expertise in proteomics and metabolism of anaerobic cellulolytic bacteria

Shang-Tian Yang
Professor, Department of Chemical and Biomolecular Engineering, The Ohio
State University, 151 West Woodruff Avenue, Columbus, OH, 43210, USA
yang.15@osu.edu
Expertise in metabolic engineering of Clostridium cellulovorans

Vincent J.J. Martin
Professor, Centre for Structural and Functional Genomics, Concordia
University, 7141 Sherbrooke Street West, Montréal, QC H4B 1R6, Canada
vincent.martin@concordia.ca
Expertise in proteomics and metabolic engineering aimed at biorefinery of
cellulosic biomass

Ariane Bize
Research director, UR HBAN, Irstea, Antony, France
ariane.bize@irstea.fr
Expertise in proteomics of anaerobic cellulolytic bacteria

Opposed Reviewers:

Cover letter

Dear Editor,

please find attached the manuscript entitled “*Clostridium cellulovorans* metabolism of cellulose as studied by comparative proteomic approach”. *C. cellulovorans* is among the most attractive candidates for direct fermentation of lignocellulosic biomass to industrially relevant chemicals, particularly, biofuels. Recently, metabolic engineering has enabled butanol production in this strain, which is a milestone in one-step production of this biofuel from plant biomass. However, currently scarce information on the central carbon metabolism of *C. cellulovorans* hampers further implementation of metabolic engineering strategies towards application of this strain in industrial processes. The present study aimed at improving understanding of cellulose metabolism in *C. cellulovorans*. Growth kinetics, substrate consumption, catabolite accumulation and whole-cell soluble proteome of cellulose-grown cells were compared with those of glucose-grown cells. This analysis showed specific modulations of the central carbon metabolism in response to changes in the growth substrate, including modifications in the redox and energy balance. We think that the results of the present study will help better understanding of *C. cellulovorans* physiology. Furthermore, these data could be useful to determine key genes and possible metabolic bottlenecks to be addressed towards improved metabolic engineering of *C. cellulovorans*.

We hope that the subject and of the present manuscript will be of interest for publication in Journal of Proteomics. We look forward to receive your answer and we send you our best regards.

Sincerely yours,

Roberto Mazzoli

Significance

C. cellulovorans can metabolize all the main plant polysaccharides (cellulose, hemicelluloses and pectins) and, unlike other well established cellulolytic microorganisms, can produce butyrate. *C. cellulovorans* is therefore among the most attractive candidates for direct fermentation of lignocellulose to high-value chemicals and, especially, n-butanol, i.e. one of the most promising liquid biofuels for the future. Recent studies aimed at engineering n-butanol production in *C. cellulovorans* represent milestones towards production of biofuels through one-step fermentation of lignocellulose but also indicated that more detailed understanding of the *C. cellulovorans* central carbon metabolism is essential to refine metabolic engineering strategies towards improved n-butanol production in this strain. The present study helped identifying key genes associated with specific catabolic reactions and indicated modulations of central carbon metabolism (including redox and energy balance) associated with cellulose consumption. This information will be useful to determine key enzymes and possible metabolic bottlenecks to be addressed towards improved metabolic engineering of this strain.

Highlights

- Whole-cell soluble proteome of cellulose- and glucose-grown *C. cellulovorans*
- Cellulose-grown cells produce higher amount of acetate and lower amount of ethanol
- Modulation of glycolysis, fermentative pathways, nitrogen assimilation was detected
- Cellulose induces up-regulation of ATP biosynthetic pathways
- Cellulose-grown cells show lower intracellular ATP content

1 ***Clostridium cellulovorans* metabolism of cellulose as studied by**
2 **comparative proteomic approach**

3 Giulia Usai¹, Simona Cirrincione¹, Angela Re², Marcello Manfredi³, Andrea Pagnani⁴, Enrica
4 Pessione¹, Roberto Mazzoli^{1,#}

5

6 ¹Structural and Functional Biochemistry. Laboratory of Proteomics and Metabolic Engineering of
7 Prokaryotes. Department of Life Sciences and Systems Biology, University of Turin. Via Accademia
8 Albertina 13, 10123 Torino, Italy.

9 ²Centre for Sustainable Future Technologies, Fondazione Istituto Italiano di Tecnologia, Environment
10 Park - Parco Scientifico Tecnologico per l'Ambiente Via Livorno 60, 10144, Torino, Italy

11 ³Center for Translational Research on Autoimmune and Allergic Diseases, Università del Piemonte
12 Orientale, Novara, Italy; Department of Translational Medicine, Università del Piemonte Orientale,
13 Novara, Italy.

14 ⁴DISAT, Politecnico di Torino, Corso Duca degli Abruzzi 24, 10129 Torino, Italy. Italian Institute for
15 Genomic Medicine (IIGM), Via Nizza 52, Torino, Italy. Istituto Nazionale di Fisica Nucleare (INFN),
16 Sezione di Torino, Via Pietro Giuria 1, 10125 Torino, Italy.

17

18

19 [#]Corresponding author: Roberto Mazzoli,

20 Department of Life Sciences and Systems Biology. University of Turin. Via Accademia Albertina 13.
21 10123 Torino. Italy.

22 Tel. +39 011 6704644

23 Fax +39 011 6704508

24 E-mail: roberto.mazzoli@unito.it

25

26 **Key words:** ATP, acetate, ethanol, Alcohol dehydrogenase, pyruvate phosphate dikinase, glucose

27

28 **Running title:** Cellulose metabolism in *C. cellulovorans*

29 **Abbreviations:**

ABC ATP binding cassette

ACAT Acetyl-CoA acetyltransferase

ACK Acetate kinase

ADH Alcohol dehydrogenase

BCD Butyryl-CoA dehydrogenase

BUK Butyrate kinase

CAZY Carbohydrate active enzyme

CBP Consolidated bioprocessing

CoA Coenzyme A

COGs	Cluster of Orthologous Genes
DSMZ	German Collection of Microorganisms and Cell Cultures
DTT	Dithiothreitol
ECH	Enoyl-CoA hydratase
F1,6BP	Fructose 1,6-bisphosphate
F6P	Fructose-6-phosphate
FDR	False discovery rate
GDH	Glutamate dehydrogenase
GlnS	Glutamine synthase
GluS	Glutamate synthase
HBD	Hydroxybutyryl-CoA dehydrogenase
MDH	Malate dehydrogenase
ME	Malic enzyme
PA	Pyruvic acid

PEP	Phosphoenolpyruvate
PEPC	Phosphoenolpyruvate carboxylase
PEPCK	Phosphoenolpyruvate carboxykinase
PEPS	Phosphoenolpyruvate synthetase
PFL	Pyruvate formate lyase
PFOR	Pyruvate ferredoxin oxidoreductase
PK	Pyruvate kinase
PP_i	Pyrophosphate
PPDK	Pyruvate phosphate dikinase
PTA	Phosphate acetyltransferase
PTB	Phosphate butyryltransferase
PTS	Phosphotransferase system
SWATH-MS	Sequential Window Acquisition of All Theoretical Mass Spectra
TCA	Tricarboxylic acid

31 **Abstract**

32 *Clostridium cellulovorans* is among the most promising candidates for consolidated bioprocessing
33 (CBP) of cellulosic biomass to liquid biofuels (ethanol, butanol). *C. cellulovorans* can metabolize all
34 the main plant polysaccharides and its main catabolite is butyrate. This makes this strain a potential
35 butanol producer since most reactions of butyrate and butanol biosynthesis from acetyl-CoA are
36 common. Recent studies demonstrated that introduction of a single heterologous alcohol/aldehyde
37 dehydrogenase diverts the branching-point intermediate, i.e. butyryl-CoA, towards butanol production
38 in this strain. Despite *C. cellulovorans* potential for CBP of plant biomass, engineering its metabolic
39 pathways towards industrial utilization requires better understanding of its metabolism. The present
40 study aimed at improving comprehension of cellulose metabolism in *C. cellulovorans* by comparing
41 growth kinetics, substrate consumption/product accumulation and whole-cell soluble proteome (data
42 available via ProteomeXchange, identifier PXD015487) with those of the same strain grown on a
43 soluble carbohydrate, glucose, as the main carbon source. Modulations of the central carbon
44 metabolism in response to different growth substrate were detected, including regulation of glycolytic
45 enzymes, fermentation pathways and nitrogen assimilation possibly affecting the redox balance. Higher
46 energy expenditure seems to occur in cellulose-grown *C. cellulovorans*, which induces up-regulation of
47 ATP synthetic pathways, e.g. acetate production and ATP synthase.

48

49 **Significance**

50 *C. cellulovorans* can metabolize all the main plant polysaccharides (cellulose, hemicelluloses and
51 pectins) and, unlike other well established cellulolytic microorganisms, can produce butyrate. *C.*
52 *cellulovorans* is therefore among the most attractive candidates for direct fermentation of lignocellulose
53 to high-value chemicals and, especially, n-butanol, i.e. one of the most promising liquid biofuels for the
54 future. Recent studies aimed at engineering n-butanol production in *C. cellulovorans* represent
55 milestones towards production of biofuels through one-step fermentation of lignocellulose but also

56 indicated that more detailed understanding of the *C. cellulovorans* central carbon metabolism is
57 essential to refine metabolic engineering strategies towards improved n-butanol production in this
58 strain. The present study helped identifying key genes associated with specific catabolic reactions and
59 indicated modulations of central carbon metabolism (including redox and energy balance) associated
60 with cellulose consumption. This information will be useful to determine key enzymes and possible
61 metabolic bottlenecks to be addressed towards improved metabolic engineering of this strain.

62

63

64 **Introduction**

65 Current awareness of the effects that fossil fuel exploitation brings about on global warming and
66 climate changes has prompted search for alternative energy sources with lower environmental footprint
67 [1]. Bio-based processes, such as biorefineries, have been especially promoted because of their
68 potential significant benefits to environmental, economic and societal sustainability issues [2].
69 Lignocellulosic biomass is the most abundant raw material on the Earth, hence, it is among the most
70 promising feedstock for the so-called 2nd generation biorefineries. Unlike 1st generation biorefineries,
71 these processes rely on non-food biomass, such as agricultural/land by-products (e.g. cereal straw,
72 forest residues), municipal or industrial wastes (e.g. paper mill sludge) [3]. However, lignocellulose is
73 very recalcitrant to biodegradation and its bioconversion currently requires physical/chemical or
74 enzymatic pre-treatment to improve its accessibility to enzymes and fermenting microorganisms [4,5].
75 Moreover, multiple bioreactors in series are necessary for enzyme production and/or polysaccharide
76 hydrolysis and/or fermentation of soluble sugar(s) [2]. Development of consolidated bioprocessing
77 (CBP) of lignocellulose, that is single-pot fermentation to high-value chemicals, could significantly
78 reduce current costs of lignocellulose fermentation [6,7]. Since natural microorganisms isolated so far
79 do not possess the biochemical features enabling CBP, metabolic engineering has been employed to
80 develop improved microbial strains [7–10]. Candidates for CBP include both cellulolytic
81 microorganisms such as *Clostridium thermocellum* and *Clostridium cellulolyticum* [9] and high-value
82 compound producing yeasts and bacteria (e.g. *Saccharomyces cerevisiae*, lactic acid bacteria) [7,8].

83 *C. cellulovorans* is a strict anaerobic, mesophilic bacterium [11] among the most interesting
84 candidates for CBP of plant biomass. *C. cellulovorans* shows some advantageous metabolic features
85 with respect to other well established cellulolytic microorganisms such as *C. thermocellum*, *C.*
86 *cellulolyticum* or *Thermoanaerobacterium saccharolyticum*: i) it ferments a larger panel of substrates
87 that include all the main plant polysaccharides, namely cellulose, hemicelluloses and pectins [12,13];
88 ii) it produces butyryl-CoA, which is a key intermediate for n-butanol production [11]. n-butanol is

89 considered the most promising liquid biofuel for future use in transportation [14]. With respect to
90 ethanol, butanol has higher combustion energy and can be used in pure form in engines (while ethanol
91 must be blended with gasoline) [15].

92 However, so far, research on *C. cellulovorans* has been mainly focused on its enzyme system
93 for depolymerizing plant polysaccharides. The latter features dozens of carbohydrate active enzymes
94 (CAZys) including both glycosyl hydrolases, carbohydrate esterases and polysaccharide lyases [16].
95 Remarkably, the number of CAZys encoded by *C. cellulovorans* genome is 37% higher than that found
96 in *C. thermocellum* (one of the most efficient cellulolytic microorganisms isolated so far) [8]. Several
97 of these enzymes, that is those containing a dockerin domain, are physically associated to form huge
98 protein complexes called cellulosomes, which are tethered to the *C. cellulovorans* cell surface [17].
99 Several studies have shown that the expression of *C. cellulovorans* cellulosomal and non-cellulosomal
100 CAZYs is modulated by the available growth substrate(s) [18–21].

101 Recently, gene tools for the manipulation of *C. cellulovorans* have been developed [22–24].
102 This led to improvement of production of liquid biofuels, namely ethanol and n-butanol, in *C.*
103 *cellulovorans* by metabolic engineering [25,26]. These studies have been milestones towards n-butanol
104 production by CBP of plant biomass, using a single microorganism. However, the n-butanol titer
105 obtained by these investigations (i.e. 3.47 g/L) is insufficient for industrial application and further
106 optimization of these strains is necessary [25,27]. Improvements of gene systems for *C. cellulovorans*
107 manipulation are desirable to increase transformation efficiency and success rate of gene modification
108 attempts [24,27]. Moreover, detailed understanding of *C. cellulovorans* central carbon metabolism is
109 essential to identify key genes and possible metabolic bottlenecks that should be addressed by
110 metabolic engineering strategies. As far as we know, only two previous studies have focused on
111 intracellular proteins involved in the metabolism of cellulose or other plant polysaccharides in *C.*
112 *cellulovorans* [12,13] which seem insufficient for the aforementioned purposes.

113 In the present study, a comparative approach was applied to *C. cellulovorans* cells grown with
114 either crystalline cellulose (i.e. avicel) or a soluble carbohydrate, namely glucose, as the main carbon
115 source. Previous studies on other cellulolytic clostridia have shown that overall metabolism on
116 cellulose (and other polysaccharides) may highly differ from that observed on soluble mono- /di-
117 saccharides, because of different kinetics and energetics of complex versus simple carbohydrate
118 metabolism [2]. To gauge the global effects caused by cellulose utilization, we analyzed *C.*
119 *cellulovorans* growth kinetics and substrate consumption/metabolite accumulation in conjunction with
120 label-free quantitative proteomics.

121 **Materials and Methods**

122 GROWTH CONDITIONS

123 *Clostridium cellulovorans* was grown anaerobically in the DSMZ medium 320
124 (https://www.dsmz.de/microorganisms/medium/pdf/DSMZ_Medium320.pdf), with some
125 modifications. Trypticase peptone and rumen fluid were not supplemented because not required for
126 growth [11]. Na₂CO₃ was replaced by PIPES [piperazine-N,N'-bis(2-ethanesulfonic acid)] (15.12 g/l)
127 as pH buffering agent [11]. Media were purged with pure N₂ (instead of 80:20 N₂-CO₂ mixture) and
128 sterilized by autoclaving (20 min 121°C). Either 5 g/l D-glucose or 10 g/l avicel® PH-101
129 microcrystalline cellulose (50 µm particle size; Sigma-Aldrich Inc., St. Louis, MO, USA) were
130 supplemented as the main carbon source. Inocula were grown in glucose-supplemented medium until
131 exponential growth phase and then transferred into 500 ml butyl-stoppered bottles containing glucose-
132 or avicel-supplemented medium. Cultures were incubated at 37°C without agitation. For each growth
133 condition three independent cultures were performed.

134

135 ESTIMATION OF BACTERIAL GROWTH

136 Microbial growth was monitored at regular time intervals though estimation of total protein content.
137 For protein extraction, 15 ml of culture was collected by centrifugation (4000 xg, 4°C, 20 min) and
138 washed twice with 0.9% (w/v) NaCl. The cell pellet was re-suspended in 1 ml of 0.2 M NaOH and
139 incubated 10 min at 100°C. Protein samples were quantified by using the Bradford reagent (Sigma-
140 Aldrich Inc., St. Louis, MO, USA) following the manufacturer's instruction. Bovine serum albumin
141 was used as the standard.

142

143 SUBSTRATE CONSUMPTION DETERMINATION

144 Glucose concentration in cell free supernatants was measured hourly using the glucose oxidase-
145 peroxidase kit (K-GLUC, gopod format, Megazyme International, Bray, Ireland), following the
146 manufacturer's instructions.

147 To determinate cellulose consumption, the cellulose concentration was measured from cell pellet every
148 three days. Briefly, after two washing steps with 0.9% (w/v) NaCl, the pellet was re-suspended in 67%
149 (v/v) H₂SO₄ and incubated for one hour under stirring at room temperature, to promote cellulose
150 hydrolysis. The total carbohydrate quantification was performed with the phenol-sulfuric acid method
151 [28], using glucose for the standard curve.

152

153 END-PRODUCT AND INTRACELLULAR ATP QUANTIFICATION

154 Accumulation of acetic acid, lactic acid, formic acid and butyric acid in the growth medium was
155 quantified by high-pressure liquid chromatography (HPLC; Agilent Technologies 1200 series),
156 equipped with an Aminex HPX-87H column (Bio-Rad, Hercules, CA, USA), a Micro-guard (cation H;
157 Bio-Rad, Hercules, CA, USA) and a UV-Vis detector set at 210 nm. The mobile phase was 5 mM
158 H₂SO₄ at a flux of 0.5 ml/min and a temperature of 50°C.

159 Ethanol in the growth medium was quantified by the Ethanol assay kit (K-EtOH, Megazyme
160 International, Bray, Ireland), following the manufacturer's instructions.

161 For quantification of total intracellular ATP content, 1 ml of culture was collected and centrifuged at
162 10000 xg for 10 min. Pellets were resuspended in 200 µl of PBS, lysed with CellTiter-Glo® One
163 Solution (Promega Corporation, Madison, WI, USA), and ATP content was measured by following the
164 manufacturer's instructions.

165

166 PROTEOMIC ANALYSIS

167 *Cytosolic protein extraction*

168 Biomass samples were harvested (4000 xg, 4°C, 20 min) 5 hours and 7 days after inoculum from
169 glucose- and avicel-supplemented cultures, respectively, and washed twice with 0.9% (w/v) NaCl.
170 Protein were extracted according to Munir *et al.*, 2015 [29]. Briefly, the pellets were resuspended in 6
171 ml of 2% (v/v) SDT-lysis buffer (2 mM Tris-HCl, 0.4% (w/v) SDS, pH 7.6), with 100 mM DTT. The
172 samples were incubated 10 min at 95°C and centrifuged (4000 xg, 20 min). The supernatant was
173 centrifuged again (10 min, 10000 xg) to discard unlysed cells and cell debris. Proteins were
174 precipitated from supernatants by methanol-chloroform method [30]. The protein pellets were
175 resuspended in 25 mM NH₄HCO₃ with 0.1% (w/v) SDS and the protein concentration was measured by
176 the 2-D Quant kit (GE Healthcare, Chicago, IL, USA), following the manufacturer's instructions.

177 *In-solution protein digestion*

178 Prior to SWATH-MS (Sequential Window Acquisition of all Theoretical fragment ion spectra Mass
179 Spectrometry) [31,32] analysis, proteins were digested with trypsin. Briefly, the samples were prepared
180 to have 100 µg of proteins in 25 µl of 100 mM NH₄HCO₃. The proteins were reduced by adding 2.5 µl
181 of 200 mM DTT and incubating them at 90°C for 20 min, and alkylated with 10 µl of 200 mM
182 iodoacetamide for 1 h at room temperature in the dark. Excess of iodoacetamide was finally removed
183 by 200 mM DTT. After dilution with 300 µl of water and 100 µl of NH₄HCO₃ to raise the pH to 7.5-
184 8.0, 5 µg of trypsin (Sequence Grade, Promega Corporation, Madison, WI, USA) was added and

185 digestion was performed overnight at 37 °C. Trypsin activity was stopped by adding 2 µl of neat formic
186 acid and digests were dried by Speed Vacuum [33]. The samples were desalted on the Discovery®
187 DSC-18 solid phase extraction (SPE) 96-well Plate (25 mg/well) (Sigma-Aldrich Inc., St. Louis, MO,
188 USA) and then analyzed as previously described [34].

189 *SWATH-MS analysis*

190 LC–MS/MS analyses were performed using a micro-LC Eksigent Technologies (Dublin, OH, USA)
191 system with a stationary phase of a Halo Fused C18 column (0.5 × 100 mm, 2.7 µm; Eksigent
192 Technologies, Dublin, OH, USA). The mobile phase was a mixture of 0.1% (v/v) formic acid in water
193 (A) and 0.1% (v/v) formic acid in acetonitrile (B), eluting at a flow-rate of 15.0 µl/min at an increasing
194 concentration of solvent B from 2% to 40% in 30 min. The LC system was interfaced with a 5600+
195 TripleTOF system (SCIEX, Concord, Canada). Samples used to generate the SWATH-MS spectral
196 library were subjected to the traditional data-dependent acquisition (DDA) and to cyclic data
197 independent analysis (DIA) of the mass spectra, using a 25-Da window as reported elsewhere [35]. The
198 MS data were acquired with Analyst TF 1.7 (SCIEX, Concord, Canada). Three instrumental replicates
199 for each sample were subjected to the DIA analysis [36].

200 *Protein data search*

201 The MS files were searched using Protein Pilot v. 4.2 (SCIEX, Concord, Canada) with the following
202 parameters: cysteine alkylation, digestion by trypsin, no special factors and False Discovery Rate at
203 1%; The files were search also with Mascot v. 2.4 (Matrix Science Inc., Boston, USA) using trypsin as
204 enzyme, with 2 missed cleavages and a search tolerance of 50 ppm was specified for the peptide mass
205 tolerance, and 0.1 Da for the MS/MS tolerance, charges of the peptides to search for were set to 2 +, 3
206 + and 4 +, and the search was set on monoisotopic mass. The instrument was set to ESI-QUAD-TOF
207 and the following modifications were specified for the search: carbamidomethyl cysteines as fixed
208 modification and oxidized methionine as variable modification. The UniProt/Swiss-Prot reviewed

209 database containing *C. cellulovorans* proteins (NCBI_Clostridium_cellulovorans743B, version
210 30102017, 4278 sequence entries) was used.

211 *Protein quantification*

212 The label-free quantification was performed by integrating the extracted ion chromatogram of all the
213 unique ions for a given peptide using PeakView 2.0 and MarkerView 1.2. (Sciex, Concord, ON,
214 Canada). SwathXtend was employed to build an integrated assay library, built with the DDA
215 acquisitions, using a protein FDR threshold of 1 %. Six peptides per protein and six transitions per
216 peptide were extracted from the SWATH files. Shared peptides were excluded as well as peptides with
217 modifications. Peptides with FDR lower than 1 % were exported in MarkerView for the t-test. The up-
218 and down-regulated proteins were selected using p-value < 0.05 and fold change > 1.5.

219 The mass spectrometry proteomics data have been deposited to the ProteomeXchange Consortium via
220 the PRIDE [37] partner repository (<https://www.ebi.ac.uk/pride/archive/>) with the dataset identifier
221 PXD015487.

222 *Protein classification and statistical analysis*

223 For each primary annotations of proteins, files were downloaded from the National Center for
224 Biotechnology Information resource
225 ([ftp://ftp.ncbi.nlm.nih.gov/genomes/refseq/bacteria/Clostridium cellulovorans/](ftp://ftp.ncbi.nlm.nih.gov/genomes/refseq/bacteria/Clostridium_cellulovorans/)), the PATRIC Bacterial
226 Bioinformatics Resource Center (<https://www.patricbrc.org>) and the eggNOG database of orthologous
227 groups and functional annotation (<http://eggnogdb.embl.de/#/app/home>). The eggNOG resource
228 provides Clusters of Orthologous Groups (COGs) of proteins
229 (<https://www.ncbi.nlm.nih.gov/pubmed/9381173>), which represent a framework for functional protein
230 classification on the basis of accurately deciphered evolutionary relationships
231 (<https://www.ncbi.nlm.nih.gov/pubmed/26582926>). Protein-level annotations overall *C. cellulovorans*
232 proteins are detailed in the **Supplementary File 1**. Fisher's exact test and fold enrichment were used to

233 identify significantly overrepresented COG categories in the up- and down-regulated proteins
234 compared to the overall quantified proteins. Multiple testing adjustment for p-values derived from the
235 Fisher's exact test was carried out using the Benjamini-Hochberg method.
236 For predicting both signal peptides and subcellular localization of the proteins involved in
237 lignocellulose depolymerization Signal-P 4.1 (cut-off > 0.45) [38]
238 (<http://www.cbs.dtu.dk/services/SignalP-4.1/>) and PSORTb v.3.0 [39] (<https://www.psorth.org/psorth/>)
239 were used, respectively.

240

241 **Results and discussion**

242 **GROWTH CHARACTERISTICS AND FERMENTATION PRODUCTS**

243 *C. cellulovorans* growth and substrate consumption in cultures with 5 g/l D-glucose and 10 g/l
244 avicel (microcrystalline cellulose) are shown in **Fig. 1A, B**. Glucose supported around 16-fold higher
245 growth rate ($\mu = 0.39 \text{ h}^{-1}$) with respect to cellulose ($\mu = 0.025 \text{ h}^{-1}$). These observations essentially
246 confirm previous reports on *C. cellulovorans* [13,22]. The supplied substrates were only partially
247 consumed, that is 3.1 g/l of cellulose (in 28 days) and 1.4 g/l of glucose (in 9 hours) were consumed in
248 avicel- and glucose-supplemented cultures, respectively. Therefore, factors other than carbon substrate
249 depletion (*e.g.* metabolite accumulation, pH) determined growth arrest. Actually, complete
250 consumption of glucose (about 10 g/l in about 4 days) and higher consumption of cellulose (about 7 g/l
251 in 10 days) by *C. cellulovorans* was previously reported by maintaining the pH of the culture between
252 6.0 and 7.0 [22] while no pH regulation was used in the present study.

253 All the main fermentation products, that is acetic acid, butyric acid, ethanol, formic acid and
254 lactic acid (**Fig. 1C, Supplementary Figure 1**) were produced at higher yield in glucose-grown
255 cultures. The sum of their final concentrations indicates that about 98% of the carbon derived from
256 glucose was converted into these products. However, only 27% of the cellulose consumed by avicel-

grown *C. cellulovorans* was converted into these catabolites. Since the final bacterial biomass was similar in the two growth conditions studied, the rest of consumed cellulose should have taken other metabolic fates. Other cellulolytic bacteria, such as *Ruminiclostridium cellulolyticum* (previously *Clostridium cellulolyticum*) and *C. thermocellum*, may accumulate glycogen and/or exopolysaccharides and/or extracellular cellodextrins and/or amino acids [40,41]. This has never been reported in *C. cellulovorans*, but it could not be excluded, because of the scarce number of available studies on the metabolism of this strain. Further investigations necessary to test this hypothesis were beyond the scope of the present study. Butyric, formic and lactic acid were accumulated in similar amounts in the two growth conditions tested (**Fig. 1C**). Butyric and formic acid were the most abundant end-products while low amounts of lactic acid were detected. However, cellulose- and glucose-grown *C. cellulovorans* produced different amounts of acetate and ethanol (**Fig. 1C**). More in detail, cellulose-grown cultures accumulated higher amounts of acetate (t-test, p-value = 0.01) and lower amounts of ethanol (t-test, p-value = 0.02). Previous investigations reported no ethanol accumulation by glucose- or cellulose-grown *C. cellulovorans* [22,25]. However, growth media with significantly different composition and different pH regulation of bacterial cultures in these studies may justify the different catabolite profiles observed. Changes in metabolite production profiles between different growth conditions have been reported for other cellulolytic bacteria. In *Clostridium termitidis* CT1112, reduction in ethanol (and formate) production and increase in acetate accumulation was observed when it was grown on cellulose instead of a soluble sugar (*i.e.* cellobiose) [42]. In *R. cellulolyticum*, the carbon flux partition between the main catabolic products (*i.e.* acetate, ethanol and lactate) is greatly affected by pH and entering carbon flows, with higher acetate production from substrates which are more slowly metabolized [43]. Lower acetate production and higher ethanol accumulation as the growth rate increases has also been observed in *C. thermocellum* [44]. The results obtained in the present study are therefore consistent with previous observations on other cellulolytic clostridia. Interestingly, this study shows that cellulose promotes a pathway, *i.e.* acetate production, involved in

282 ATP synthesis, while it down-regulates ethanol production which is involved in NAD(P)H-
283 consumption in *C. cellulovorans*.

284

285 **COMPARATIVE PROTEOMIC STUDY**

286 *Overall findings*

287 The present study was focused on soluble whole-cell extracts in order to identify proteins specifically
288 associated with glucose or crystalline cellulose metabolism. Most previous proteomic studies on
289 cellulose metabolism by *C. cellulovorans* were interested in extracellular proteins [19–21,42]. Only
290 very recently, whole-cell proteomes of *C. cellulovorans* grown on either glucose- or cellulose-
291 supplemented medium were compared for a total of 1016 identified proteins in both conditions [13].
292 Samples for the present investigation were harvested in the late exponential phase (**Fig. 1A, B**).
293 Quantitative proteomic analysis was performed by the data independent acquisition-based Sequential
294 Window Acquisition of All Theoretical Mass Spectra (SWATH-MS) approach [31,32] combining deep
295 proteome coverage capabilities with quantitative consistency and accuracy. In this study, 621 proteins
296 were quantified (**Supplementary File 1**) corresponding to about 15 % of the *C. cellulovorans*
297 annotated proteins [16]. To analyze the distribution of their biological functions, the quantified proteins
298 were annotated by means of the Cluster of Orthologous Genes (COGs) categories
299 (<http://eggnogdb.embl.de/#/app/home>). The large majority (522 proteins, 84 %) of quantified proteins
300 were associated with at least a known function grasped by a COG category whereas the remaining 16
301 % consisted of proteins with unknown function (**Fig. 2A**). Of the total quantified proteins, 319 were
302 found as differentially expressed when comparing the two growth conditions, that is they were at least
303 1.5-fold more abundant ($p\text{-value} < 0.05$) in one growth condition compared to the other.
304 Most (258 proteins, 81 %) differentially expressed proteins were classified into the variety of COG
305 categories shown in **Fig. 2A** while the remaining 19% are still functionally uncharacterized. This
306 suggests that proteins among those with yet unknown biological function could be involved in

cellulose/glucose metabolism thus confirming previous reports [19]. We noticed that 124 proteins corresponding to 39 % of the differentially expressed proteins were annotated as having a transport function. Actually, differentially expressed proteins were not fairly distributed across the COG categories (**Fig. 2B**). The proteins involved in cell duplication and protein translation were the largest COG categories shared by the proteins resulting down-regulated in cellulose-grown cells. This latter observation is likely related to the higher growth rate shown by glucose-grown cells. The proteins involved in carbohydrate transport and metabolism were instead the largest COG category in up-regulated proteins in cellulose-grown cultures. The latter observation was corroborated by the analysis of the functional enrichment with respect of all quantified proteins (Fisher's exact test, FDR < 0.05) whereby overexpressed proteins in cellulose-grown cultures were functionally enriched in the category of metabolism and transport of carbohydrates (**Fig. 2C**) and especially (74% of them) in plant polysaccharide depolymerization. In the following sections, differentially expressed proteins will be thoroughly discussed according to their functional classification.

Proteins involved in plant polysaccharide depolymerization

Many proteins that are more abundant in avicel-grown cultures are involved in plant polysaccharide depolymerization (**Table 1**). They include 24 out of 57 cellulosomal subunits encoded by the *C. cellulovorans* genome [45]. Cellulosomal subunits represent 24 out of 39 (62%) over-expressed proteins belonging to the class of carbohydrate transport and metabolism. Exoglucanase S (ExgS, Clocel_2823), *i.e.* a cellulase, Mannanase A (ManA, Clocel_2818), *i.e.* a hemicellulase, and CbpA (Clocel_2824), *i.e.* the main scaffolding protein of the *C. cellulovorans* cellulosome, showed the largest differential expression, since they were 44-fold, 36-fold and 33-fold more abundant in cellulose-grown bacteria, respectively. These proteins, together with additional up-regulated proteins identified in the current study (*i.e.*, EngH, Clocel_2822; EngK, Clocel_2821; EngL, Clocel_2819; EngM, Clocel_2816; HbpA, Clocel_2820), are encoded by a large cellulosomal gene cluster on the *C. cellulovorans* chromosome [45]. Endoglucanase Z (EngZ, Clocel_2741) and Endoglucanase E

333 (Clocel_2576) were also among the most strongly up-regulated (19- and 17-fold, respectively) proteins,
334 consistently with their key role in hydrolysis of crystalline cellulose [46] and for the function and
335 architecture of the *C. cellulovorans* cellulosome [47,48], respectively.

336 Over-expressed non-cellulosomal proteins include three cellulases (EngD, Clocel_3242; EngO,
337 Clocel_1478; Clocel_2606) and two hemicellulases (β -mannanase, Clocel_1134; β -xylosidase,
338 Clocel_2595).

339 The present analysis showed that hemicellulases were expressed in cellulose-grown cells and
340 the same cellulases and hemicellulases up-regulated in cellulose-grown cells were detected also in
341 glucose-grown cells, although at a lower abundance. These data agree with most results obtained by
342 previous transcriptomic/proteomic studies on *C. cellulovorans* which reported that several cellulosome
343 components (e.g. CbpA, EngE, EngL, EngY, ExgS, ManA, and the products of genes Clocel_0619,
344 Clocel_2575, Clocel_2576, Clocel_2607, Clocel_4119), including hemicellulases, are constitutively
345 biosynthesized [19,49]. On the other hand, overexpression of some of these proteins (e.g. CbpA, EngB,
346 EngE, and EngZ) is induced by crystalline cellulose [46,49].

347

348 ***Proteins involved in protein secretion***

349 Three components of the Sec protein-translocation complex (SecF, Clocel_2074; SecD, Clocel_2075;
350 SecY, Clocel_3713) were shown to be more abundant (1.9-, 1.8- and 1.7-fold, respectively) in
351 cellulose-grown cells. This finding could be related to increased need to secrete CAZYS involved in
352 cellulose depolymerization. Almost no information is available on mechanisms of secretion of
353 cellulases in native cellulolytic microorganisms. For most (91 %) cellulases reported in UniProtKB
354 (<https://www.uniprot.org/help/uniprotkb>) no signal peptide is annotated [50,51]. Among cellulases
355 from Gram-positive bacteria, 10 % contained twin-arginine translocation (Tat)-like signal peptides,
356 while 11 % featured amino acid patterns ascribable to the Sec secretory system [50,51]. The main
357 components of the Sec system are a protein-conducting channel SecYEG, and an ATP-dependent
358 motor protein SecA [52]. The auxiliary SecDF membrane protein-complex seems to enhance

translocation efficiency in a proton motive force-powered manner [52]. A signal peptide was predicted at the N-terminus of all the over-expressed proteins involved in cellulose/hemicellulose depolymerization identified in this study by Signal-P 4.1 (<http://www.cbs.dtu.dk/services/SignalP-4.1/>), except for Clocel_2595. More in detail, the N-terminus of 72 % of these proteins shows the typical structure of Sec-type signal peptides, that is a positively charged N-terminal (N-region), a hydrophobic core (H-region) and a negative charged C-region with alanine-rich cleavage site [53]. The variability occurring at the C-terminus of the cleavage site, with 67% of proteins displaying a VXA motif instead of the most typical AXA motif of the Gram-positive bacteria (**Fig. 3**) [51], may indicate that a different consensus sequence is used by *C. cellulovorans* for secreting many of its cellulases. Future investigations will be necessary to confirm this hypothesis and the possible role of the Sec machinery proteins (SecD, Clocel_2074; SecF, Clocel_2075; SecY, Clocel_3713) that were over-expressed in this study in cellulase secretion in *C. cellulovorans*. Understanding cellulase secretion in native cellulolytic microorganisms is crucial for improving comprehension of plant polysaccharide degradation and its exploitation in biotechnological applications such as the construction of recombinant cellulolytic microorganisms, which are frequently hampered by inefficient expression of heterologous cellulases [7].

375

376 *Substrate uptake*

It is generally considered that most cellulose is extracellularly converted to different length cellodextrins (rather than glucose) by anaerobic cellulolytic bacteria [40]. Cellodextrins are then transported into the cytoplasm and depolymerized mainly through phosphorolysis [40]. Seven proteins related to ATP binding cassette-type (ABC) transporters (Clocel_0040; Clocel_1357; Clocel_4050; Clocel_3636; Clocel_3461; Clocel_3460; Clocel_3857) were more abundant in cellulose-grown cells. Protein sequence alignments by the protein BLAST(<https://www.uniprot.org/blast/>) highlighted sequence identity ranging from 54 % to 86 % with proteins involved in the transport of a variety of

384 substrates in other clostridia. In addition, four soluble-binding proteins (Clocel_0038; Clocel_3201;
385 Clocel_0435; Clocel_1358), that is proteins which present soluble substrates to the transport channel,
386 were overexpressed in cellulose-grown cultures [54]. Amino acid sequence analysis of these proteins
387 strongly indicated that some of them could be involved in the uptake of mono-/oligo-saccharides
388 derived from extracellular cellulose depolymerization. More in detail Clocel_0040 (5-fold over-
389 expressed in cellulose-grown cells) featured 64 % amino acid sequence identity with a carbohydrate
390 ABC transporter membrane protein from *Clostridium* sp. DMS 8431 (SAMN04487886_10195).
391 Similarly, the ABC transporter encoded by Clocel_3857 (1.6-fold more abundant in cellulose-grown
392 cells) has 72 % sequence identity with the sugar ABC transporter ATP-binding protein of *Clostridium*
393 *pasteurianum* (C1I91_02605). However, the proteins encoded by Clocel_1357, Clocel_3460 and
394 Clocel_3461 are likely involved in other functions since they show higher sequence identity (50-70 %)
395 with Clostridial peptide- and sodium-transport systems. The protein products of Clocel_4050 and
396 Clocel_3636 lacked conserved residues required for the propagation of feature annotation.
397 Furthermore, a modulation of the expression of sugar phosphotransferase systems (PTS) was observed.
398 PTS are multicomponent transporters that couple sugar transport with phosphorylation using
399 phosphoenolpyruvate (PEP) as energy source [55]. Our analysis showed that some PTS components,
400 namely the phosphotransferase system lactose/cellobiose-specific IIB subunit (Clocel_2881) and the
401 permease IIC component (Clocel_2880), were overexpressed while other subunits, such as the
402 phosphoryl carrier protein (Clocel_2058) and the phosphoenolpyruvate-protein phosphotransferase
403 (Clocel_3686), were down-regulated in cellulose-grown cultures. A cytoplasmic cellodextrin
404 phosphorylase (Clocel_2717) was also identified in this study, but its expression level did not
405 significantly change between the two growth conditions considered.

406

407 ***Central carbon metabolism***

408 *Glycolysis*

409 *C. cellulovorans* metabolizes glucose through the Embden-Meyerhof-Parnas pathway [16] (**Fig. 4A**).
 410 Our analysis revealed that the expression of several *C. cellulovorans* glycolytic enzymes was affected
 411 by carbon source change, albeit at different extent (**Fig. 4A, B**). Phosphofructokinases generally control
 412 glycolysis and are allosterically activated by ADP and inhibited by fructose-6-P (F6P) and ATP in
 413 Clostridia [13,56]. The *C. cellulovorans* genome encodes two ATP-dependent 6-phosphofructokinases
 414 (namely Clocel_2901 and Clocel_0388), *i.e.* enzymes that catalyze the phosphorylation of F6P to
 415 fructose 1,6-bisphosphate (F1,6BP) by consuming ATP. The present study revealed that Clocel_0388
 416 was not differentially expressed, whereas Clocel_2901 was 5-fold more abundant in cellulose-grown
 417 cells. Overexpression of Clocel_2901 in cellulose-grown *C. cellulovorans* (fold change, FC, comprised
 418 between 4 and 16) was also reported by another recent study [13]. *C. cellulovorans* genome also
 419 encodes a pyrophosphate (PP_i)-fructose 6-phosphate 1-phosphotransferase (Clocel_1603) (that
 420 catalyzes the phosphorylation of F6P using PP_i) but this enzyme was not found as differentially
 421 expressed in the current study. This observation could indicate a preferential ATP-dependent
 422 conversion of F6P in the cells grown on cellulose. Furthermore, four glycolytic enzymes were present
 423 in lower abundance in cellulose-grown cells: triosephosphate isomerase (Clocel_0721, FC = 0.03),
 424 glyceraldehyde-3-phosphate dehydrogenase (Clocel_0719, FC = 0.23), glucose-6-phosphate isomerase
 425 (Clocel_1364, FC = 0.25), and phosphoglycerate kinase (Clocel_0720, FC = 0.6). In general, these
 426 results do not seem to differ from those previously reported by Aburaya et al. [13], since most of these
 427 enzymes were slightly down-regulated in cellulose- versus glucose grown *C. cellulovorans* (with the
 428 only exception of the late stationary phase). Glyceraldehyde-3-phosphate dehydrogenase is among the
 429 most important glycolytic enzymes since it catalyzes NADH production through oxidation of
 430 glyceraldehyde-3-phosphate and has previously been identified as a probable bottleneck in glycolysis
 431 [57] and towards alcoholic biofuel production [58]. Glyceraldehyde-3-phosphate dehydrogenase up-
 432 regulation in glucose-grown *C. cellulovorans* agrees with previous studies on *C. termitidis* [42] and *C.*
 433 *thermocellum* [59], that reported that the same enzyme was one of the most abundantly expressed
 434 proteins during growth on cellobiose, a soluble β -glucose disaccharide.

435

436 *Pyruvate metabolism*

437 Interconversion between pyruvate (PA) and phosphoenolpyruvate (PEP) is most frequently mediated
438 by the antagonistic enzymatic couple consisting of ADP-dependent pyruvate kinase (PK) (catabolic
439 role) and phosphoenolpyruvate synthetase (PEPS) (anabolic role). In *C. cellulovorans* PK is encoded
440 by Clocel_0389 whereas no PEPS is annotated. However, *C. cellulovorans* genome encodes a pyruvate
441 phosphate dikinase (PPDK, Clocel_1454) that may catalyze the reversible production of PA, ATP and
442 P_i from PEP, AMP and PP_i [60]. In some organisms such as *Acetobacter xylinum*, *Propionibacterium*
443 *shermanii* and *Microbispora rosea* [61–63], PPDK seems to exert its activity in the anabolic (PEP)
444 direction [64,65], whereas it fulfils a catabolic role in others, such as *Clostridium symbiosum* [66]. In
445 *Thermoproteus tenax*, the combined action of PPDK, PK and PEPS was shown to control the
446 interconversion between PEP and PA [67]. In *C. thermocellum*, which does not possess PK, PPDK can
447 substantially support the production of PA from PEP [68]. In the present study, PK did not show any
448 differential expression between the two growth conditions, while PPDK showed 11-fold higher
449 abundance in cellulose-grown cells. Consistently, a 3.5-fold up-regulation was observed also for a
450 PPDK putative regulatory protein (Clocel_4349) [69]. Up-regulation of PPDK (Cter_0809) in
451 cellulose-grown cells was reported also in *Clostridium termitidis* CT1112 [42]. Previous
452 characterization of PK and PPDK from different microorganisms has shown that their activity is
453 generally allosterically regulated. More in details, PPDK is regulated by the ATP/AMP ratio in
454 *Trypanosoma brucei* and *Acetobacter xylinum* [70,71] whereas PK activity by intracellular
455 concentration of ADP, AMP and phosphate sugars [72–74]. The present study points at PPDK as a
456 possible key enzyme regulating carbon flux during cellulose metabolism by *C. cellulovorans*. Further
457 investigations on this enzyme, such as understanding potential regulation by allosteric effectors, appear
458 essential to better understand its role in *C. cellulovorans* metabolism.

459 In some bacteria, malic enzyme (ME), malate dehydrogenase (MDH) and phosphoenolpyruvate
460 carboxykinase (PEPCK) are involved in an alternative three-step pathway converting PEP to pyruvate

that is called malate shunt [75] (**Fig. 4A**). In *C. thermocellum*, the malate shunt was proposed as a strategy to transfer electrons from NADH to NADP⁺, thus supplying most of the NADPH necessary for biosynthetic routes [68]. A putative ME (Clocel_0393) was down-expressed, *i.e.* 3-fold less abundant, in cellulose-grown *C. cellulovorans*. We were therefore interested to understand if a malate shunt could be present in *C. cellulovorans* also. In its genome, two lactate/malate dehydrogenase are annotated (Clocel_1533; Clocel_2700). The protein product of Clocel_1533 shares an amino acid sequence identity of 40% with *C. thermocellum* MDH (Cthe_0345) which includes key residues in the active site [76]. Although a confirmation by enzyme activity assay is necessary, these findings strongly suggest that Clocel_1533 encodes a MDH. No PEP carboxykinase is annotated in the *C. cellulovorans* genome, but PEP carboxylase (PEPC, Clocel_1149), that catalyzes the conversion of PEP to oxaloacetate [77], could functionally replace it. PEPC was identified in the present study, although in similar amounts in the two growth conditions. Hence, a malate shunt could be hypothesized in *C. cellulovorans*, especially to supply NADPH for biosynthetic reactions in fast growth conditions such as in glucose-supplemented cultures. Further investigations by enzyme activity assay of the products of Clocel_1533 and Clocel_1149 will be necessary to confirm this hypothesis. Since previous studies reported that ME is allosterically inhibited by PP_i [78] and PPDK activity controls the intracellular PP_i concentration, a possible regulatory interconnection between the two pathways could exist.

478

479 ***Tricarboxylic acid cycle and nitrogen assimilation***

C. cellulovorans can partially operate the tricarboxylic acid (TCA) cycle in a reductive manner [77]. Clostridia use TCA cycle mainly to produce intermediates for biosynthetic routes and regulate the redox balance inside the cells. According to the present study, biosynthetic levels of aconitase (Clocel_1405), citrate synthase (Clocel_3688), isocitrate dehydrogenase (Clocel_2469) and fumarate hydratase (Clocel_0392) were not affected by the carbon sources used. However, glutamate dehydrogenase (GDH, Clocel_1284) was 10-fold more abundant in glucose-grown bacteria. Furthermore, glutamine synthetase (GlnS, Clocel_3873) and glutamate synthase (GluS, Clocel_2665)

487 were found as 3- and 1.5-fold more abundant in glucose- versus cellulose-grown *C. cellulovorans*,
488 respectively. All these proteins are typically involved in nitrogen assimilation and synthesis of
489 components of cell biomass in bacteria. GDH catalyzes the reversible NAD(P)H-dependent reductive
490 amination of 2-ketoglutarate to glutamate [79]; GlnS aminates glutamate to glutamine in ATP-
491 dependent manner; Glus catalyzes the reversible transfer of an amino group from glutamine to 2-
492 ketoglutarate with the consumption of NAD(P)H (or reduced ferredoxin) and the production of 2
493 glutamate molecules and NAD(P)⁺ (or oxidized ferredoxin) [41]. In addition, intracellular
494 accumulation of glutamate and secretion of amino acids (up to 15-17 % of the total substrate
495 consumed) have been reported in other cellulolytic clostridia (e.g. *C. thermocellum* and *R.*
496 *cellulolyticum*) [41,80]. However, overexpression of GDH in *C. thermocellum* was reported for cells
497 growing on cellulose with respect to cultures growing on a soluble carbohydrate (*i.e.* cellobiose) [81].
498 Higher abundance of these enzymes in glucose-grown *C. cellulovorans* could be related with higher
499 growth rates measured in this condition. In addition, activities of these proteins affect the redox balance
500 of the cell. A recent study on *C. thermocellum* demonstrated that deletion of the gene *glnA* encoding its
501 main glutamine synthetase significantly decreases amino acid secretion and increases ethanol yield
502 likely by increasing NADH availability in the cell [41]. Finally, GDH, GlnS and Glus have been
503 referred as involved in pH homeostasis in *C. thermocellum* [80]. However, no significant difference in
504 extracellular pH between glucose- and cellulose grown *C. cellulovorans* cultures has been detected in
505 the present study (data not shown).

506

507 ***End-product synthesis pathways***

508 The main products of *C. cellulovorans* metabolism are H₂, CO₂, acetic acid, lactic acid, butyric acid
509 and ethanol [11]. The proteins encoded by three out of the four hydrogenase genes annotated in the *C.*
510 *cellulovorans* genome [82] were not identified in this study while the Fe-only hydrogenase
511 (Clocel_4097) resulted 1.5-fold more abundant in glucose-grown cells. Therefore, the protein encoded
512 by Clocel_4097 may be the main responsible for H₂ production in this strain. Hydrogen production is

the fastest way to dispose of excess of reduced cofactors generated through carbohydrate metabolism [83]. Up-regulation of Clocel_4097 in glucose-grown cells, appears as a strategy to maximize glycolytic turnover in this condition. This Fe-only hydrogenase probably receives electrons from the oxidative activity of pyruvate ferredoxin oxidoreductase PFOR (Clocel_1684), which converts pyruvate to acetyl-CoA, yielding CO₂ and reduced ferredoxin (**Fig. 4A**) Consistently, PFOR was found 1.6-fold more abundant in glucose-grown cells. According to gene annotation, *C. cellulovorans* genome encodes another PFOR (Clocel_2840) and an indole pyruvate oxidoreductase subunit (Clocel_4184), that were not detected in the present study. These data therefore point at Clocel_1684 as the main PFOR in *C. cellulovorans*.

Another enzyme that was more abundant (2-fold) in glucose-grown cells is pyruvate formate lyase (PFL, Clocel_1811). This enzyme catalyzes pyruvate conversion to formate and acetyl-CoA production from pyruvate (**Fig. 4A**). Consistently, a pyruvate-formate lyase activating enzyme (Clocel_1812) [84] was also found as 5-fold more abundant in the same growth condition.

C. cellulovorans genome encodes seven alcohol dehydrogenases (ADHs), four of which were overexpressed in cellulose-grown bacteria: Clocel_2402 (2.5-fold), Clocel_1990 (8-fold), Clocel_3817 (1.6-fold), Clocel_1140 (2.2-fold) (**Supplementary File 1**). The presence of multiple ADHs is common in solvent-producing clostridia and other microorganisms [85,86]. ADHs can differ for their substrate and coenzyme specificity, but the physiological significance of multiple ADHs in the same strain has not always been elucidated. Clocel_2402, Clocel_3817 and Clocel_1140 encode bifunctional aldehyde/alcohol dehydrogenases, possibly involved in 2-step reduction of acetyl-CoA to acetaldehyde and finally to ethanol with consumption of two NADH (**Fig. 4A**). The amino acid sequence of ADH encoded by Clocel_1990 shows 89% identity with glycerol dehydrogenases from other Clostridia, *i.e.* enzymes that catalyze the oxidation of glycerol [87]. However, owing to the scarce substrate specificity generally shown by ADHs, it is difficult to definitely assess their role in ethanol or other alcohol production without enzyme activity assays. Up-regulation of these ADHs in cellulose-grown *C. cellulovorans* is not reflected by increased ethanol production since a lower amount of ethanol is

539 accumulated in this growth condition (**Fig. 1C**). Apart from possible involvement of these up-regulated
 540 enzymes in other alcohol synthesizing pathways, a reasonable explanation for this observation is that
 541 ADHs may be reversible enzymes and also catalyze alcohol oxidation. Most studies on characterization
 542 of ADH from clostridia focused on their substrate and coenzyme specificity but few papers reported
 543 clostridial ADHs able to oxidize alcohols, although their activity was mainly in the aldehyde-reducing
 544 direction [88,89]. Previous studies showed that relative protein abundances do not always correlate
 545 with end-product distribution profiles. In particular, higher abundance of ADHs with respect to lactate
 546 dehydrogenase was observed in some growth conditions in *Thermoanaerobacter*
 547 *thermohydrosulfuricus* WC1, however, lactate was always the major end-product of this strain [90].
 548 The present study confirms that protein abundance is not the only player determining carbon flux
 549 distribution in cells, which is also influenced by other parameters such as catabolic bottlenecks,
 550 allosteric regulation or cofactor availability.

551 Phosphate acetyltransferase (PTA, Clocel_1891), which catalyzes the conversion of acetyl-CoA to
 552 acetyl-phosphate, was found up-regulated (1.9-fold) in cellulose-grown cells whereas acetate kinase
 553 (ACK, Clocel_1892), which catalyzes the subsequent conversion of acetyl-phosphate to acetate and
 554 ATP did not result as differentially expressed in the present study (**Fig. 4B**). Genes encoding PTA and
 555 ACK form an operon in *C. cellulovorans*, so, our results indicate that post-transcriptional events may
 556 differentially regulate the expression of these genes. Changes in proteomic levels of PTA are consistent
 557 with increased production of acetate by cellulose-grown *C. cellulovorans* (**Fig. 1C**). These data also
 558 suggest that the level of PTA activity could be a bottleneck for acetate production in *C. cellulovorans*,
 559 while ACK could have a higher specific activity.

560 Butyric acid derives from acetyl-CoA through a multi-step pathway as shown in **Fig. 4A**. Some of the
 561 enzymes involved in this pathway are also implicated in other metabolic processes, such as fatty acid
 562 and amino acid metabolism. Maybe for this reason, some of them, that is acetyl-CoA acetyltransferase
 563 (ACAT, Clocel_3058), hydroxybutyryl-CoA dehydrogenase (HBD, Clocel_2972), enoyl-CoA
 564 hydratase (ECH, Clocel_2976) and phosphate butyryltransferase (PTB, Clocel_3675), were down-

565 regulated while butyryl-CoA dehydrogenase (BCD, Clocel_2975) and butyrate kinase (BUK,
566 Clocel_3674) were up-regulated in cellulose-grown cells. Actually, no significant changes in butyrate
567 amounts accumulated by cellulose- or glucose-grown *C. cellulovorans* was observed in this study.

568

569 ENERGY PRODUCTION

570 Three components of the F₀F₁-ATP synthase, i.e. ATP synthase F₁ sector subunit alpha (Clocel_3051)
571 and subunit beta (Clocel_3049) and F₀ portion subunit c (Clocel_3054), were 2-fold more abundant in
572 cellulose-grown *C. cellulovorans*. ATP synthase F₀ subunit b (Clocel_3053) and F₁ gamma chain
573 (Clocel_3050) did not show any differential expression between the two growth conditions. The
574 genes encoding these proteins together with Clocel_3048, Clocel_3052 and Clocel_3055 form an
575 operon which codes for the entire ATP synthase complex [16]. Some F₀F₁-ATP synthases of strict
576 anaerobes mainly hydrolyze ATP to pump H⁺ out of the cell, such as in *C. thermocellum* where this
577 function has been related to pH homeostasis [80,91,92]. However, no significant difference in
578 extracellular pH between glucose- and cellulose-grown *C. cellulovorans* cultures has been detected in
579 the present study (data not shown). Other strict anaerobic bacteria have ATP synthases which mainly
580 function towards ATP synthesis by using either Na⁺, such as *Propionigenium modestum* [93] and
581 *Acetobacterium woodii* [94], or H⁺, e.g. *Moorella thermoacetica* [95], transmembrane gradient as
582 energy source. In anaerobes, Na⁺/H⁺ gradient can be generated by several mechanisms such as
583 anaerobic electron chains, H₂ oxidation [91] and membrane bound proton-translocating
584 pyrophosphatases [59]. It can therefore be hypothesized that up-regulation of F₀F₁-ATP synthase could
585 provide additional ATP to that generated by substrate-level phosphorylation in *C. cellulovorans* (**Fig.**
586 **4A**).

587

588 ATP QUANTIFICATION

Metabolite and proteomic analyses indicated an up-regulation of acetate biosynthetic pathway in cellulose-grown *C. cellulovorans*. Acetate, as well as butyrate, production is involved in ATP synthesis through substrate-level phosphorylation. Furthermore, over-expression of some F₀F₁ ATP synthase subunits was detected in cellulose-grown *C. cellulovorans*. These observations suggest possible differences in ATP-content of glucose- or cellulose-grown cells. In most growth phases, the ATP content of cellulose-grown bacteria ranged between 0 and 3 ng/mg, that is much lower (about a 100 times) than that measured in glucose-grown cells (**Fig. 5**). Interestingly, ATP content of cellulose-grown cultures at the time of inoculum is of the same order of magnitude of that observed in glucose-grown cells. It is worth reminding that both glucose- and cellulose-grown cultures were inoculated with glucose-grown bacteria, hence, just after inoculum, *C. cellulovorans* cells likely still show most characteristics associated with glucose metabolism. Our findings seem to contradict previous studies claiming bioenergetic benefits for anaerobic bacteria grown on cellulose instead of simple carbohydrates [40,96]. Increased acetate production and up-regulation of ATP synthase subunits by cellulose-grown *C. cellulovorans* can be interpreted as a mean to improve ATP cell supply in conditions requiring high ATP consumption (e.g. for cellulase biosynthesis and secretion) [97].

604

605 **Conclusions**

The present comparative analysis indicated that the metabolism of cellulose-grown *C. cellulovorans* significantly differs from that of the same strain grown on simple carbohydrates, i.e. glucose, thus confirming previous reports on other cellulolytic microorganisms [40]. Apart from modulation of the expression of cellulosomal and non-cellulosomal enzymes directly involved in cellulose depolymerization, the present investigation revealed that other modifications of the metabolic network are induced by the growth substrate. The up-regulation of some subunits of the Sec-machinery in cellulose-grown *C. cellulovorans* point at them as interesting candidates possibly involved in cellulase secretion in this strain. This could be a starting point for filling the gaps in understanding the

614 mechanisms of cellulase secretion in cellulolytic microorganisms. The most original findings brought
615 by the present study concern modifications of the central metabolism and fermentative pathways. It is
616 worth noting that previous studies did not report any significant modification in fermentation product
617 profile [22] or any major change in the expression of enzymes involved in the central metabolism
618 between glucose- and cellulose-grown *C. cellulovorans* [13]. This is most probably caused by different
619 composition [22] and/or different pH regulation of the growth medium [13]. In some cases, the present
620 study helped to identify the key genes associated with specific catabolic reactions or pathways among
621 multiple paralogs with the same annotation in *C. cellulovorans* based on their expression levels (e.g.
622 H₂ase, PFOR). In other cases (e.g. ADHs, MDH), our results clearly indicated that further studies (e.g.
623 substrate and cofactor specificity, catalytic activity regulation) are essential to understand the
624 physiological role of certain gene products. The main findings of the present investigation can be
625 summarized as follows:

- 626 • Proteomic data indicate that a re-distribution of the central carbon flow occurs between glucose-
627 and cellulose-grown *C. cellulovorans*, through a modulation of the biosynthesis of different key
628 enzymes that include ATP-dependent 6-phosphofructokinase (Clocel_2901), PPKK
629 (Clocel_1454), GDH (Clocel_1284), PTA (Clocel_1891), PFL (Clocel_1811) and different
630 ADHs.
- 631 • A macroscopic consequence of this re-arranged metabolic network is that cellulose promotes
632 acetate accumulation, while glucose induces higher ethanol production. Based on previous
633 reports, this could depend on the different substrate used, or the different growth rate/carbon
634 flux supported by these substrates [42–44].
- 635 • Cellulose-grown cells have significantly lower ATP content, possibly related to higher energy
636 expenditure for cellulase biosynthesis and secretion. Up-regulation of acetate pathway and ATP
637 synthase subunits may help cope with this high energy demanding condition. Therefore, these
638 observations seem to contradict previous studies claiming bioenergetic benefits of anaerobic
639 bacteria growing on cellulose instead of simple carbohydrates [40,96].

640 In conclusion, this study pointed out some aspects of glucose and cellulose metabolism in *C.*
641 *cellulovorans*, which could be useful for better understanding the physiology of this strain and also
642 towards engineering of its metabolic pathways for application in processes for biorefining plant
643 biomass.

644

645 **Acknowledgments**

646 This study was financially supported by “Progetti di Ricerca Ateneo/Compagnia di San Paolo - anno
647 2014” program (Torino_call2014_L2_138 grant). AP acknowledges financial support from Marie
648 Sklodowska-Curie, grant agreement No. 734439 (INFERNET). We would like to thank Chiara Gandini
649 for her contribution to the first steps of this study.

650

651 **Conflict of interest**

652 The authors declare no conflict of interest

653

654 **Author contributions**

655 RM designed research and supervised experiments. GU, SC, AR and MM performed experiments and
656 analyzed proteomic data. All the Authors contributed in discussing experimental data and writing the
657 manuscript.

658

659 **References**

660 [1] IPCC, T.F. Stocker, D. Qin, G.-K. Plattner, M.M.B. Tignor, S.K. Allen, J. Boschung, A. Nauels, Y. Xia,

- 661 V. Bex, P.M. Midgley, Climate Change 2013 - The Physical Science Basis, 2013. doi:10.1038/446727a.
- 662 [2] R. Mazzoli, Development of Microorganisms for Cellulose-Biofuel Consolidated Bioprocessings:
663 Metabolic Engineers' Tricks, Comput. Struct. Biotechnol. J. 3 (2012) e201210007. doi:10.5936/csbj.201210007.
- 664 [3] R.E.H. Sims, W. Mabee, J.N. Saddler, M. Taylor, An overview of second generation biofuel
665 technologies, Bioresour. Technol. 101 (2010) 1570–1580. doi:10.1016/j.biortech.2009.11.046.
- 666 [4] A. Procentese, E. Johnson, V. Orr, A. Garruto Campanile, J.A. Wood, A. Marzocchella, L. Rehmann,
667 Deep eutectic solvent pretreatment and subsequent saccharification of corncob, Bioresour. Technol. 192 (2015)
668 31–36. doi:10.1016/j.biortech.2015.05.053.
- 669 [5] S. Tabasso, D. Carnaroglio, E. Calcio Gaudino, G. Cravotto, Microwave, ultrasound and ball mill
670 procedures for bio-waste valorisation, Green Chem. 17 (2015) 684–693. doi:10.1039/C4GC01545B.
- 671 [6] L.R. Lynd, X. Liang, M.J. Biddy, A. Allee, H. Cai, T. Foust, M.E. Himmel, M.S. Laser, M. Wang, C.E.
672 Wyman, Cellulosic ethanol: status and innovation, Curr. Opin. Biotechnol. 45 (2017) 202–211.
673 doi:https://doi.org/10.1016/j.copbio.2017.03.008.
- 674 [7] L. Tarraran, R. Mazzoli, Alternative strategies for lignocellulose fermentation through lactic acid
675 bacteria: the state of the art and perspectives, FEMS Microbiol. Lett. 365 (2018). doi:10.1093/femsle/fny126.
- 676 [8] H. Liu, J. Sun, J.-S. Chang, P. Shukla, Engineering microbes for direct fermentation of cellulose to
677 bioethanol, Crit. Rev. Biotechnol. 38 (2018) 1089–1105. doi:10.1080/07388551.2018.1452891.
- 678 [9] C. Cheng, T. Bao, S.-T. Yang, Engineering *Clostridium* for improved solvent production: recent
679 progress and perspective, Appl. Microbiol. Biotechnol. 103 (2019) 5549–5566. doi:10.1007/s00253-019-09916-
680 7.
- 681 [10] F. Xin, W. Dong, W. Zhang, J. Ma, M. Jiang, Biobutanol Production from Crystalline Cellulose through
682 Consolidated Bioprocessing, Trends Biotechnol. 37 (2019) 167–180. doi:10.1016/j.tibtech.2018.08.007.
- 683 [11] R. Sleat, R.A. Mah, R. Robinson, Isolation and characterization of an anaerobic, cellulolytic bacterium,
684 *Clostridium cellulosovorans* sp. nov., Appl. Environ. Microbiol. 48 (1984) 88–93. doi:10.1099/00207713-32-1-87.
- 685 [12] S. Aburaya, K. Esaka, H. Morisaka, K. Kuroda, M. Ueda, Elucidation of the recognition mechanisms for

686 hemicellulose and pectin in *Clostridium cellulovorans* using intracellular quantitative proteome analysis, *AMB*
687 *Express.* 5 (2015). doi:10.1186/s13568-015-0115-6.

688 [13] S. Aburaya, W. Aoki, K. Kuroda, H. Minakuchi, M. Ueda, Temporal proteome dynamics of *Clostridium*
689 *cellulovorans* cultured with major plant cell wall polysaccharides, *BMC Microbiol.* 19 (2019) 118.
690 doi:10.1186/s12866-019-1480-0.

691 [14] N.H. Al Makishah, Bioenergy: Microbial Biofuel Production Advancement, *Int. J. Pharm. Res. Sci.* 6(3)
692 (2017) 93–106.

693 [15] W. Kaminski, E. Tomczak, A. Gorak, Biobutanol - Production and Purification Methods, *Ecol. Chem.*
694 *Eng.* 18 (2011) 31–37.

695 [16] Y. Tamaru, H. Miyake, K. Kuroda, A. Nakanishi, Y. Kawade, K. Yamamoto, M. Uemura, Y. Fujita,
696 R.H. Doi, M. Ueda, Genome sequence of the cellulosome-producing mesophilic organism *Clostridium*
697 *cellulovorans* 743B, *J. Bacteriol.* 192 (2010) 901–902. doi:10.1128/JB.01450-09.

698 [17] Y. Tamaru, H. Miyake, K. Kuroda, A. Nakanishi, C. Matsushima, R.H. Doi, M. Ueda, Comparison of
699 the mesophilic cellulosome-producing *Clostridium cellulovorans* genome with other cellulosome-related
700 clostridial genomes, *Microb. Biotechnol.* 4 (2011) 64–73. doi:10.1111/j.1751-7915.2010.00210.x.

701 [18] W. Cho, S.D. Jeon, H.J. Shim, R.H. Doi, S.O. Han, Cellulosomic profiling produced by *Clostridium*
702 *cellulovorans* during growth on different carbon sources explored by the cohesin marker, *J. Biotechnol.* 145
703 (2010) 233–239. doi:10.1016/j.jbiotec.2009.11.020.

704 [19] H. Morisaka, K. Matsui, Y. Tatsukami, K. Kuroda, H. Miyake, Y. Tamaru, M. Ueda, Profile of native
705 cellulosomal proteins of *Clostridium cellulovorans* adapted to various carbon sources, *AMB Express.* 2 (2012)
706 37. doi:10.1186/2191-0855-2-37.

707 [20] K. Matsui, J. Bae, K. Esaka, H. Morisaka, K. Kuroda, M. Ueda, Exoproteome profiles of *Clostridium*
708 *cellulovorans* grown on various carbon sources, *Appl. Environ. Microbiol.* 79 (2013) 6576–6584.
709 doi:10.1128/AEM.02137-13.

710 [21] K. Esaka, S. Aburaya, H. Morisaka, K. Kuroda, M. Ueda, Exoproteome analysis of *Clostridium*

711 *cellulovorans* in natural soft-biomass degradation, AMB Express. 5 (2015) 2. doi:10.1186/s13568-014-0089-9.

712 [22] X. Yang, M. Xu, S.T. Yang, Metabolic and process engineering of *Clostridium cellulovorans* for biofuel
713 production from cellulose, Metab. Eng. 32 (2015) 39–48. doi:10.1016/j.ymben.2015.09.001.

714 [23] X. Yang, M. Xu, S.T. Yang, Restriction modification system analysis and development of in vivo
715 methylation for the transformation of *Clostridium cellulovorans*, Appl. Microbiol. Biotechnol. 100 (2016) 2289–
716 2299. doi:10.1007/s00253-015-7141-9.

717 [24] Z. Wen, N.P. Minton, Y. Zhang, Q. Li, J. Liu, Y. Jiang, S. Yang, Enhanced solvent production by
718 metabolic engineering of a twin-clostridial consortium, Metab. Eng. 39 (2017) 38–48.
719 doi:10.1016/j.ymben.2016.10.013.

720 [25] T. Bao, J. Zhao, J. Li, X. Liu, S.-T. Yang, n-Butanol and ethanol production from cellulose by
721 *Clostridium cellulovorans* overexpressing heterologous aldehyde/alcohol dehydrogenases, Bioresour. Technol.
722 285 (2019) 121316. doi:https://doi.org/10.1016/j.biortech.2019.121316.

723 [26] Z. Wen, R. Ledesma-Amaro, J. Lin, Y. Jiang, S. Yangd, Improved n-butanol production from
724 *Clostridium cellulovorans* by integrated metabolic and evolutionary engineering, Appl. Environ. Microbiol. 85
725 (2019). doi:10.1128/AEM.02560-18.

726 [27] T. Bao, J. Zhao, Q. Zhang, S.-T. Yang, Development of a shuttle plasmid without host restriction sites
727 for efficient transformation and heterologous gene expression in *Clostridium cellulovorans*, Appl. Microbiol.
728 Biotechnol. 103 (2019) 5391–5400. doi:10.1007/s00253-019-09882-0.

729 [28] M. Dubois, K.A. Gilles, J.K. Hamilton, P.A. Rebers, F. Smith, Colorimetric Method for Determination
730 of Sugars and Related Substances, Anal. Chem. 28 (1956) 350–356. doi:10.1021/ac60111a017.

731 [29] R.I. Munir, V. Spicer, D. Shamshurin, O. V. Krokhn, J. Wilkins, U. Ramachandran, R. Sparling, D.B.
732 Levin, Quantitative proteomic analysis of the cellulolytic system of *Clostridium termitidis* CT1112 reveals
733 distinct protein expression profiles upon growth on α -cellulose and cellobiose, J. Proteomics. 125 (2015) 41–53.
734 doi:10.1016/j.jprot.2015.04.026.

735 [30] D. Wessel, U.I. Flügge, A method for the quantitative recovery of protein in dilute solution in the

736 presence of detergents and lipids, *Anal. Biochem.* 138 (1984) 141–143. doi:<https://doi.org/10.1016/0003->
737 2697(84)90782-6.

738 [31] B.C. Collins, C.L. Hunter, Y. Liu, B. Schilling, G. Rosenberger, S.L. Bader, D.W. Chan, B.W. Gibson,
739 A.C. Gingras, J.M. Held, M. Hirayama-Kurogi, G. Hou, C. Krisp, B. Larsen, L. Lin, S. Liu, M.P. Molloy, R.L.
740 Moritz, S. Ohtsuki, R. Schlapbach, N. Selevsek, S.N. Thomas, S.C. Tzeng, H. Zhang, R. Aebersold, Multi-
741 laboratory assessment of reproducibility, qualitative and quantitative performance of SWATH-mass
742 spectrometry, *Nat. Commun.* 8 (2017) 1–11. doi:10.1038/s41467-017-00249-5.

743 [32] C. Ludwig, L. Gillet, G. Rosenberger, S. Amon, B.C. Collins, R. Aebersold, Data-independent
744 acquisition-based SWATH-MS for quantitative proteomics: a tutorial, *Mol. Syst. Biol.* 14 (2018) e8126.
745 doi:10.15252/msb.20178126.

746 [33] E. Dalla Pozza, M. Manfredi, J. Brandi, A. Buzzi, E. Conte, R. Pacchiana, D. Cecconi, E. Marengo, M.
747 Donadelli, Trichostatin A alters cytoskeleton and energy metabolism of pancreatic adenocarcinoma cells: An in
748 depth proteomic study, *J. Cell. Biochem.* 119 (2018) 2696–2707. doi:10.1002/jcb.26436.

749 [34] S. Martinotti, M. Patrone, M. Manfredi, F. Gosetti, M. Pedrazzi, E. Marengo, E. Ranzato, HMGB1
750 Osteo-Modulatory Action on Osteosarcoma SaOS-2 Cell Line: An Integrated Study From Biochemical and -
751 Omics Approaches, *J. Cell. Biochem.* 117 (2016) 2559–2569. doi:10.1002/jcb.25549.

752 [35] L.D. Carbonare, M. Manfredi, G. Caviglia, E. Conte, E. Robotti, E. Marengo, S. Cheri, F. Zamboni, D.
753 Gabbiani, M. Deiana, D. Cecconi, F. Schena, M. Mottes, M.T. Valenti, Can half-marathon affect overall health?
754 The yin-yang of sport, *J. Proteomics.* 170 (2018) 80–87. doi:<https://doi.org/10.1016/j.jprot.2017.09.004>.

755 [36] P. Albanese, M. Manfredi, A. Meneghesso, E. Marengo, G. Saracco, J. Barber, T. Morosinotto, C.
756 Pagliano, Dynamic reorganization of photosystem II supercomplexes in response to variations in light
757 intensities, *Biochim. Biophys. Acta - Bioenerg.* 1857 (2016) 1651–1660. doi:10.1016/j.bbabi.2016.06.011.

758 [37] Y. Perez-Riverol, A. Csordas, J. Bai, M. Bernal-Llinares, S. Hewapathirana, D.J. Kundu, A. Inuganti, J.
759 Griss, G. Mayer, M. Eisenacher, E. Pérez, J. Uszkoreit, J. Pfeuffer, T. Sachsenberg, Ş. Yilmaz, S. Tiwary, J.
760 Cox, E. Audain, M. Walzer, A.F. Jarnuczak, T. Ternent, A. Brazma, J.A. Vizcaíno, The PRIDE database and
761 related tools and resources in 2019: improving support for quantification data, *Nucleic Acids Res.* 47 (2018)

762 D442–D450. doi:10.1093/nar/gky1106.

763 [38] H. Nielsen, Predicting Secretory Proteins with SignalP, in: D. Kihara (Ed.), Protein Funct. Predict.
 764 Methods Protoc., Springer New York, New York, NY, 2017: pp. 59–73. doi:10.1007/978-1-4939-7015-5_6.

765 [39] N.Y. Yu, J.R. Wagner, M.R. Laird, G. Melli, S. Rey, R. Lo, P. Dao, S.C. Sahinalp, M. Ester, L.J. Foster,
 766 F.S.L. Brinkman, PSORTb 3.0: improved protein subcellular localization prediction with refined localization
 767 subcategories and predictive capabilities for all prokaryotes, Bioinformatics. 26 (2010) 1608–1615.
 768 doi:10.1093/bioinformatics/btq249.

769 [40] M. Desvaux, Unravelling carbon metabolism in anaerobic cellulolytic bacteria, Biotechnol. Prog. 22
 770 (2006) 1229–1238. doi:10.1021/bp060016e.

771 [41] T. Rydzak, D. Garcia, D.M. Stevenson, M. Sladek, D.M. Klingeman, E.K. Holwerda, D. Amador-
 772 Noguez, S.D. Brown, A.M. Guss, Deletion of Type I glutamine synthetase deregulates nitrogen metabolism and
 773 increases ethanol production in *Clostridium thermocellum*, Metab. Eng. 41 (2017) 182–191.
 774 doi:https://doi.org/10.1016/j.ymben.2017.04.002.

775 [42] R.I. Munir, V. Spicer, O. V. Krokhin, D. Shamshurin, X. Zhang, M. Tallefer, W. Blunt, N. Cicek, R.
 776 Sparling, D.B. Levin, Transcriptomic and proteomic analyses of core metabolism in *Clostridium termitidis*
 777 CT1112 during growth on α -cellulose, xylan, cellobiose and xylose, BMC Microbiol. 16 (2016) 1–21.
 778 doi:10.1186/s12866-016-0711-x.

779 [43] N. Badalato, A. Guillot, V. Sabarly, M. Dubois, N. Pourette, B. Pontoire, P. Robert, A. Bridier, V.
 780 Monnet, D.Z. Sousa, S. Durand, L. Mazéas, A. Buléon, T. Bouchez, G. Mortha, A. Bize, Whole Proteome
 781 Analyses on *Ruminiclostridium cellulolyticum* Show a Modulation of the Cellulolysis Machinery in Response to
 782 Cellulosic Materials with Subtle Differences in Chemical and Structural Properties, PLoS One. 12 (2017)
 783 e0170524. doi:10.1371/journal.pone.0170524.

784 [44] A. Riederer, T.E. Takasuka, S. Makino, D.M. Stevenson, Y. V Bukhman, N.L. Elsen, B.G. Fox, Global
 785 Gene Expression Patterns in *Clostridium thermocellum* as Determined by Microarray Analysis of Chemostat
 786 Cultures on Cellulose or Cellobiose, Appl. Environ. Microbiol. 77 (2011) 1243 LP-1253.
 787 doi:10.1128/AEM.02008-10.

788 [45] R.H. Doi, Y. Tamaru, The *Clostridium cellulovorans* cellulosome: an enzyme complex with plant cell
789 wall degrading activity., Chem. Rec. 1 (2001) 24–32. doi:10.1002/1528-0691(2001)1:1<24::AID-
790 TCR5>3.0.CO;2-W [pii].

791 [46] S.D. Jeon, K.O. Yu, S.W. Kim, S.O. Han, The processive endoglucanase EngZ is active in crystalline
792 cellulose degradation as a cellulosomal subunit of *Clostridium cellulovorans*, N. Biotechnol. 29 (2012) 365–371.
793 doi:10.1016/j.nbt.2011.06.008.

794 [47] K. Murashima, A. Kosugi, R.H. Doi, Synergistic Effects on Crystalline Cellulose Degradation between
795 Cellulosomal Cellulases from, Society. 184 (2002) 5088–5095. doi:10.1128/JB.184.18.5088.

796 [48] R. Kozaki, H. Miyake, Enzymatic and molecular characterization of an endoglucanase E from
797 *Clostridium cellulovorans* 743B, J. Biosci. Bioeng. (2019). doi:10.1016/j.jbiosc.2019.03.013.

798 [49] S.O. Han, H.Y. Cho, H. Yukawa, M. Inui, R.H. Doi, Regulation of expression of cellulosomes and
799 noncellulosomal (hemi)cellulolytic enzymes in *Clostridium cellulovorans* during growth on different carbon
800 sources, J. Bacteriol. 186 (2004) 4218–4227. doi:10.1128/JB.186.13.4218-4227.2004.

801 [50] S. Yan, G. Wu, Secretory pathway of cellulase: A mini-review, Biotechnol. Biofuels. 6 (2013) 1–12.
802 doi:10.1186/1754-6834-6-177.

803 [51] S. Yan, G. Wu, Signal peptide of cellulase, Appl. Microbiol. Biotechnol. 98 (2014) 5329–5362.
804 doi:10.1007/s00253-014-5742-3.

805 [52] A. Tsirigotaki, J. De Geyter, N. Šoštaric', A. Economou, S. Karamanou, Protein export through the
806 bacterial Sec pathway, Nat. Rev. Microbiol. 15 (2016) 21. doi: 10.1038/nrmicro.2016.161.

807 [53] P. Natale, T. Brüser, A.J.M. Driessen, Sec- and Tat-mediated protein secretion across the bacterial
808 cytoplasmic membrane-Distinct translocases and mechanisms, Biochim. Biophys. Acta - Biomembr. 1778
809 (2008) 1735–1756. doi:10.1016/j.bbamem.2007.07.015.

810 [54] R. Tam, M.H. Saier, Structural, functional, and evolutionary relationships among extracellular solute-
811 binding receptors of bacteria, Microbiol. Rev. 57 (1993) 320–46. PMC372912.

812 [55] J.-M. Jeckelmann, B. Erni, Carbohydrate Transport by Group Translocation: The Bacterial

813 Phosphoenolpyruvate: Sugar Phosphotransferase System BT - Bacterial Cell Walls and Membranes, in: A.
814 Kuhn (Ed.), Springer International Publishing, Cham, 2019: pp. 223–274. doi:10.1007/978-3-030-18768-2_8.

815 [56] K. Uyeda, S. Kurooka, Crystallization and properties of phosphofructokinase from *Clostridium*
816 *pasteurianum*, J. Biol. Chem. 245 (1970) 3315–3324.

817 [57] S. Payot, E. Guedon, C. Cailliez, E. Gelhaye, H. Petitdemange, Metabolism of cellobiose by *Clostridium*
818 *cellulolyticum* growing in continuous culture: Evidence for decreased NADH reoxidation as a factor limiting
819 growth, Microbiology. 144 (1998) 375–384. doi:10.1099/00221287-144-2-375.

820 [58] L. Tian, S.J. Perot, D. Stevenson, T. Jacobson, A.A. Lanahan, D.A. Noguez, D.G. Olson, L.R. Lynd,
821 Biotechnology for Biofuels Metabolome analysis reveals a role for glyceraldehyde 3- phosphate dehydrogenase
822 in the inhibition of *C. thermocellum* by ethanol, Biotechnol. Biofuels. (2017) 1–11. doi:10.1186/s13068-017-
823 0961-3.

824 [59] T. Rydzak, P.D. McQueen, O. V. Krokhn, V. Spicer, P. Ezzati, R.C. Dwivedi, D. Shamshurin, D.B.
825 Levin, J.A. Wilkins, R. Sparling, Proteomic analysis of *Clostridium thermocellum* core metabolism: Relative
826 protein expression profiles and growth phase-dependent changes in protein expression, BMC Microbiol. 12
827 (2012) 1–18. doi:10.1186/1471-2180-12-214.

828 [60] H.J. Evans, H.G. Wood, The mechanism of the pyruvate, phosphate dikinase reaction, Proc. Natl. Acad.
829 Sci. U. S. A. 61 (1968) 1448–1453. doi:10.1073/pnas.61.4.1448.

830 [61] M. Benziman, N. Eisen, A. Palgi, Properties and physiological role of the pep-synthase of *A. xylinum*,
831 FEBS Lett. 3 (1969) 156–159. doi:10.1016/0014-5793(69)80123-7.

832 [62] H.G. Wood, H.J. Evans, Purification and properties of pyruvate phosphate dikinase from propionic acid
833 bacteria, Biochemistry. 10 (1971) 721–729. doi:10.1021/bi00781a001.

834 [63] N. Eisaki, H. Tatsumi, S. Murakami, T. Horiuchi, Pyruvate phosphate dikinase from a thermophilic
835 actinomyces *Microbispora rosea* subsp. aerata: Purification, characterization and molecular cloning of the gene,
836 Biochim. Biophys. Acta - Protein Struct. Mol. Enzymol. 1431 (1999) 363–373. doi:10.1016/S0167-
837 4838(99)00057-6.

838 [64] M. Østerås, B.T. Driscoll, T.M. Finan, Increased pyruvate orthophosphate dikinase activity results in an
839 alternative gluconeogenic pathway in *Rhizobium (Sinorhizobium) meliloti*, *Microbiology*. 143 (1997) 1639–
840 1648. doi:10.1099/00221287-143-5-1639.

841 [65] D. Rodriguez-Contreras, N. Hamilton, Gluconeogenesis in *Leishmania mexicana*: contribution of
842 glycerol kinase, phosphoenolpyruvate carboxykinase, and pyruvate phosphate dikinase., *J. Biol. Chem.* 289
843 (2014) 32989–33000. doi:10.1074/jbc.M114.569434.

844 [66] R.E. Reeves, A New Enzyme with Function of Pyruvate Kinase, *J. Biol. Chem.* 243 (1968) 3202–3204.
845 <http://www.jbc.org/content/243/11/3202.short>.

846 [67] B. Tjaden, A. Plagens, C. Dörr, B. Siebers, R. Hensel, Phosphoenolpyruvate synthetase and pyruvate,
847 phosphate dikinase of *Thermoproteus tenax*: Key pieces in the puzzle of archaeal carbohydrate metabolism, *Mol.*
848 *Microbiol.* 60 (2006) 287–298. doi:10.1111/j.1365-2958.2006.05098.x.

849 [68] D.G. Olson, M. Hörl, T. Fuhrer, J. Cui, J. Zhou, M.I. Maloney, D. Amador-Noguez, L. Tian, U. Sauer,
850 L.R. Lynd, Glycolysis without pyruvate kinase in *Clostridium thermocellum*, *Metab. Eng.* 39 (2017) 169–180.
851 doi:10.1016/j.ymben.2016.11.011.

852 [69] C.J. Chastain, L.M. Baird, M.T. Walker, C.C. Bergman, G.T. Novbatova, C.S. Mamani-Quispe, J.N.
853 Burnell, Maize leaf PPK regulatory protein isoform-2 is specific to bundle sheath chloroplasts and
854 paradoxically lacks a Pi-dependent PPK activation activity, *J. Exp. Bot.* 69 (2018) 1171–1181.
855 doi:10.1093/jxb/erx471.

856 [70] K. Deramchia, P. Morand, M. Biran, Y. Millerioux, M. Mazet, M. Wargnies, J.M. Franconi, F.
857 Bringaud, Contribution of pyruvate phosphate dikinase in the maintenance of the glycosomal ATP/ADP balance
858 in the *Trypanosoma brucei* procyclic form, *J. Biol. Chem.* 289 (2014) 17365–17378.
859 doi:10.1074/jbc.M114.567230.

860 [71] M. Benziman, Pyruvate-Phosphate Dikinase and the Control Gluconeogenesis in *Acetobacter xylinum*, *J.*
861 *Biol. Chem.* 246 (1971) 57–61.

862 [72] L.B. Collins, T.D. Thomas, Pyruvate Kinase of *Streptococcus lactis*, *J. Bacteriol.* 120 (1974) 52 LP-58.
863 <http://jb.asm.org/content/120/1/52.abstract>.

- 864 [73] N. Veith, A. Feldman-Salit, V. Cojocaru, S. Henrich, U. Kummer, R.C. Wade, Organism-Adapted
 865 Specificity of the Allosteric Regulation of Pyruvate Kinase in Lactic Acid Bacteria, PLOS Comput. Biol. 9
 866 (2013) 1–15. doi:10.1371/journal.pcbi.1003159.
- 867 [74] M. Enriqueta Muñoz, E. Ponce, Pyruvate kinase: current status of regulatory and functional properties,
 868 Comp. Biochem. Physiol. Part B Biochem. Mol. Biol. 135 (2003) 197–218. doi:https://doi.org/10.1016/S1096-
 869 4959(03)00081-2.
- 870 [75] S. Spaans, R. Weusthuis, J. Van Der Oost, S. Kengen, NADPH-generating systems in bacteria and
 871 archaea, Front. Microbiol. 6 (2015) 742. doi:10.3389/fmicb.2015.00742.
- 872 [76] M. Taillefer, T. Rydzak, D.B. Levin, I.J. Oresnik, R. Sparling, Reassessment of the
 873 transhydrogenase/malate shunt pathway in *Clostridium thermocellum* ATCC 27405 through kinetic
 874 characterization of malic enzyme and malate dehydrogenase, Appl. Environ. Microbiol. 81 (2015) 2423–2432.
 875 doi:10.1128/AEM.03360-14.
- 876 [77] M. Shinohara, H. Sakuragi, H. Morisaka, H. Miyake, Y. Tamaru, E. Fukusaki, K. Kuroda, M. Ueda,
 877 Fixation of CO₂ in *Clostridium cellulovorans* analyzed by ¹³C-isotopomer-based target metabolomics, AMB
 878 Express. 3 (2013) 1–6. doi:10.1186/2191-0855-3-61.
- 879 [78] M. Ljunggren, K. Willquist, G. Zacchi, E.W. van Niel, A kinetic model for quantitative evaluation of the
 880 effect of hydrogen and osmolarity on hydrogen production by *Caldicellulosiruptor saccharolyticus*, Biotechnol.
 881 Biofuels. 4 (2011) 31. doi:10.1186/1754-6834-4-31.
- 882 [79] K. Shimizu, Metabolic Regulation of a Bacterial Cell System with Emphasis on *Escherichia coli*
 883 Metabolism, ISRN Biochem. 2013 (2013) 1–47. doi:10.1155/2013/645983.
- 884 [80] J.M. Whitham, J.W. Moon, M. Rodriguez, N.L. Engle, D.M. Klingeman, T. Rydzak, M.M. Abel, T.J.
 885 Tschaplinski, A.M. Guss, S.D. Brown, *Clostridium thermocellum* LL1210 pH homeostasis mechanisms
 886 informed by transcriptomics and metabolomics, Biotechnol. Biofuels. 11 (2018) 1–13. doi:10.1186/s13068-018-
 887 1095-y.
- 888 [81] E. Burton, V.J.J. Martin, Proteomic analysis of *Clostridium thermocellum* atcc 27405 reveals the
 889 upregulation of an alternative transhydrogenase-malate pathway and nitrogen assimilation in cells grown on

cellulose, Can. J. Microbiol. 58 (2012) 1378–1388. doi:10.1139/cjm-2012-0412.

[82] M. Calusinska, T. Happe, B. Joris, A. Wilmotte, The surprising diversity of clostridial hydrogenases: A comparative genomic perspective, Microbiology. 156 (2010) 1575–1588. doi:10.1099/mic.0.032771-0.

[83] P.M. Vignais, B. Billoud, J. Meyer, Classification and phylogeny of hydrogenases1, FEMS Microbiol. Rev. 25 (2001) 455–501. doi:10.1111/j.1574-6976.2001.tb00587.x.

[84] A. Becker, W. Kabsch, X-ray structure of pyruvate formate-lyase in complex with pyruvate and CoA. How the enzyme uses the Cys-418 thiyl radical for pyruvate cleavage, J. Biol. Chem. 277 (2002) 40036–40042. doi:10.1074/jbc.M205821200.

[85] J. Lo, T. Zheng, S. Hon, D.G. Olson, L.R. Lynd, The Bifunctional Alcohol and Aldehyde Dehydrogenase Gene, Is Necessary for Ethanol Production in *Clostridium thermocellum* and *Thermoanaerobacterium saccharolyticum*, J. Bacteriol. 197 (2015) 1386 LP-1393. doi:10.1128/JB.02450-14.

[86] Z. Dai, H. Dong, Y. Zhang, Y. Li, Elucidating the contributions of multiple aldehyde/alcohol dehydrogenases to butanol and ethanol production in *Clostridium acetobutylicum*, Sci. Rep. 6 (2016) 28189. <https://doi.org/10.1038/srep28189>.

[87] F. Luers, M. Seyfried, R. Daniel, G. Gottschalk, Glycerol conversion to 1,3-propanediol by *Clostridium pasteurianum*: cloning and expression of the gene encoding 1,3-propanediol dehydrogenase, FEMS Microbiol. Lett. 154 (1997) 337–345. doi:10.1111/j.1574-6968.1997.tb12665.x.

[88] R.W. Welch, F.B. Rudolph, E.T. Papoutsakis, Purification and characterization of the NADH-dependent butanol dehydrogenase from *Clostridium acetobutylicum* (ATCC 824), Arch. Biochem. Biophys. 273 (1989) 309–318. doi: 10.1016/0003-9861(89)90489-X.

[89] A.A. Ismaiel, C.X. Zhu, G.D. Colby, J.S. Chen, Purification and characterization of a primary-secondary alcohol dehydrogenase from two strains of *Clostridium beijerinckii*, J. Bacteriol. 175 (1993) 5097 LP-5105. doi:10.1128/jb.175.16.5097-5105.1993.

[90] T.J. Verbeke, V. Spicer, O. V Krokhnin, X. Zhang, J.J. Schellenberg, B. Fristensky, J.A. Wilkins, D.B.

915 Levin, R. Sparling, *Thermoanaerobacter thermohydrosulfuricus* WC1 Shows Protein Complement Stability
 916 during Fermentation of Key Lignocellulose-Derived Substrates, *Appl. Environ. Microbiol.* 80 (2014) 1602 LP-
 917 1615. doi:10.1128/AEM.03555-13.

918 [91] K. Ozawa, T. Meikari, K. Motohashi, M. Yoshida, H. Akutsu, Evidence for the presence of an F-type
 919 ATP synthase involved in sulfate respiration in *Desulfovibrio vulgaris*, *J. Bacteriol.* 182 (2000) 2200–2206.
 920 doi:10.1128/JB.182.8.2200-2206.2000.

921 [92] P. Lund, A. Tramonti, D. De Biase, Coping with low pH: molecular strategies in neutralophilic bacteria,
 922 *FEMS Microbiol. Rev.* 38 (2014) 1091–1125. doi:10.1111/1574-6976.12076.

923 [93] P. Dimroth, Primary sodium ion translocating enzymes, *Biochim. Biophys. Acta - Bioenerg.* 1318
 924 (1997) 11–51. doi: 10.1016/S0005-2728(96)00127-2.

925 [94] S. Rahlfs, S. Aufurth, H. Schägger, V. Müller, Identification of Subunits a , b and c 1 from
 926 *Acetobacterium woodii* Na⁺ + -F₁F₀-ATPase, *J. Biol. Chem.* 275 (2000) 33297–33301.
 927 doi:10.1074/jbc.m005134200.

928 [95] J. Hugenholtz, L.G. Ljungdahl, Metabolism and energy generation in homoacetogenic clostridia, *FEMS*
 929 *Microbiol. Rev.* 7 (1990) 383–389. doi:10.1111/j.1574-6968.1990.tb04941.x.

930 [96] Y.-H.P. Zhang, L.R. Lynd, Determination of the Number-Average Degree of Polymerization of
 931 Cellodextrins and Cellulose with Application to Enzymatic Hydrolysis, *Biomacromolecules.* 6 (2005) 1510–
 932 1515. doi:10.1021/bm049235j.

933 [97] L. Girbal, P. Soucaille, Regulation of *Clostridium acetobutylicum* metabolism as revealed by mixed-
 934 substrate steady-state continuous cultures: role of NADH/NAD ratio and ATP pool., *J. Bacteriol.* 176 (1994)
 935 6433–6438. doi:10.1128/jb.176.21.6433-6438.1994.

936

Figure captions

938

939 **Figure 1.** *C. cellulovorans* growth kinetics in a medium containing 5 g/l glucose (A) or 10 g/l cellulose
940 (B) as the main carbon source. Growth curves are represented by solid lines, while dashed lines
941 represent substrate concentration. Bacterial biomass was measured by total cell protein determination.
942 Syringe symbols indicate the cell sampling points for proteomic analysis, i.e. 5 hours and 7 days after
943 inoculum for glucose- and cellulose-supplemented media, respectively. End-product final concentration
944 (g/l) measured in glucose- (purple) or cellulose- (green) supplemented *C. cellulovorans* cultures (C).
945 Data are displayed as mean $\pm$ SD across triplicate cultures. Asterisks indicate statistically significant
946 differences (t-test, p-value ≤ 0.05) between the two growth conditions.

947

948 **Figure 2.** Functional categorization of quantified proteins. (A) Cluster of Orthologous Genes (COG)
949 functional classification of the quantified proteins along with the differentially expressed proteins
950 (DEP) resulting from the comparison between *C. cellulovorans* cultures grown on cellulose or glucose
951 as the main carbon source. (B) COG-based functional classification of the proteins resulting up- or
952 down-regulated when comparing *C. cellulovorans* cultures grown on cellulose or glucose as the main
953 carbon source. (C) Functional enrichment analysis of up- and down-regulated proteins according to
954 COG annotation. The dashed line corresponds to a fold enrichment equal to 1.5. Statistically significant
955 over-representation of a COG category in the up- or down-regulated proteins (Fisher's exact test, FDR
956 < 0.05) is labelled with an asterisk.

957

958 **Figure 3.** Consensus sequence of N-terminal signal peptide of the *C. cellulovorans* proteins involved in
959 plant polysaccharide depolymerization that were up-regulated in cellulose-grown cultures.

960

961 **Figure 4.** (A) Scheme representing the metabolic changes observed in both *C. cellulovorans* central
962 carbon metabolism and end-product synthetic pathways, based on protein differential expression. (B)
963 Function, gene locus and fold change of the proteins involved in the central carbon metabolism and
964 end-product synthesis in *C. cellulovorans*. Up- and down-regulated proteins are displayed in green and
965 purple, respectively. Proteins that are not differentially expressed or not quantified in the present study
966 are indicated in grey. GPI, glucose 6-phosphate isomerase; PFK, phosphofructokinase; ALD, aldolase;
967 TPI, triosephosphate isomerase; GAPDH, glyceraldehyde-3-phosphate dehydrogenase; PGK,

968 phosphoglycerate kinase; PMG, phosphoglycerate mutase; ENO, enolase; PK, pyruvate kinase; PPK, pyruvate
969 pyruvate phosphate dikinase; PEPC, phosphoenolpyruvate carboxylase; PFOR, pyruvate ferredoxin
970 oxidoreductase; PFL, pyruvate formate lyase; ME, malic enzyme; MDH, malate dehydrogenase. Fd-
971 H2ase, ferredoxin-hydrogenase; ADH, bifunctional aldehyde/alcohol dehydrogenase; PTA, phosphate
972 acetyltransferase; ACK, acetate kinase; ACAT, acetyl-CoA acetyltransferase; HBD, hydroxybutyryl-
973 CoA dehydrogenase; ECH, enoyl-CoA hydratase; BCD, butyryl-CoA dehydrogenase; PTB, phosphate
974 butyryltransferase; BUK, butyrate kinase.

975

976 **Figure 5.** Intracellular ATP concentration measured in glucose- (A) and cellulose-grown *C.*
977 *cellulovorans* (B). Intracellular ATP concentration is expressed as Log₂ ng of ATP per mg of proteins
978 extracted from cell biomass. Data are displayed as mean ± SD of triplicate cultures.

979

980 **Supplementary File 1.** List of the quantified proteins in *C. cellulovorans* grown on avicel and glucose.
981 The fold-change is given by the ratio between the average of the protein abundances (based on
982 SWATH-MS data acquisition) for the three replicates of *C. cellulovorans* grown on avicel and the
983 average of the protein abundances (based on SWATH-MS data acquisition) of the three replicates in *C.*
984 *cellulovorans* grown on glucose. The proteins with fold-change > 1.5 and p-value > 0.05 are considered
985 as up-regulated (green), while the proteins with fold-change < 0.67 and p-value > 0.05 are considered
986 down-regulated (purple).

987

988 **Supplementary Figure 1.** Kinetics of accumulation of end-product measured in glucose- (A) or
989 cellulose- (B) supplemented *C. cellulovorans* cultures. Data are displayed as mean ± SD across
990 triplicate cultures.

Table 1. Up-regulated proteins involved in plant polysaccharide depolymerization. Protein function, gene locus, CAZy category, fold change, signal peptide prediction (Signal-P 4.1, D-value > 0.5) and cellular localization prediction (PsortB v3.0) are shown. GH, glycoside hydrolase; CBM, carbohydrate binding module; PL, pectate lyase; SLH, S-layer homology; NA, not annotated.

Function	Gene locus	CAZy	FC	Signal-P	Psortb
<i>Cellulosomal proteins</i>					
Exoglucanase S	Clocel_2823	GH48	43.62	0.88	Extracell
Mannanase A	Clocel_2818	GH5	35.86	0.78	Unknown
Cellulose-binding protein A	Clocel_2824	CBM3, SLH, HBD	33.49	0.89	Extracell
Endoglucanase	Clocel_2576	GH9, CBM3	19.41	0.74	Extracell
Endoglucanase Z	Clocel_2741	GH9, CBM3	18.64	0.55	Extracell
Dockerin type 1	Clocel_3193	NA	17.38	0.57	Extracell
Endoglucanase E	Clocel_3359	GH5, CBM65, SLH	16.93	0.82	Unknown
Endoglucanase L	Clocel_2819	GH9	14.49	0.80	Membrane
β-xylanase	Clocel_2900	GH10, CBM22	11.29	0.80	Extracell
Endoglucanase H	Clocel_2822	GH9, CBM3	10.97	0.57	Extracell
Mannanase	Clocel_4119	GH26, CBM35	9.83	0.73	Unknown
Endoglucanase	Clocel_0983	GH5	8.48	0.85	Unknown
Mannanase	Clocel_2575	GH26, CBM35	7.81	0.61	Unknown
Mannanase	Clocel_2607	GH26, CBM35	6.83	0.84	Extracell
Endoglucanase Y	Clocel_1624	GH9, CBM30	6.71	0.75	Extracell
Endoglucanase	Clocel_2600	GH5, CBM32	5.97	0.78	Extracell
Endoglucanase M	Clocel_2816	GH9, CBM4	5.94	0.54	Extracell
Endoglucanase B	Clocel_1150	GH5	5.03	0.68	Membrane
Endoglucanase	Clocel_3111	GH5	5.00	0.61	Extracell
Hydrophobic protein A	Clocel_2820	SLH, HBD	4.97	0.66	Extracell
Endoglucanase	Clocel_0619	GH5	4.69	0.86	Extracell
Endoglucanase	Clocel_0930	GH9, CBM3	4.50	0.78	Extracell
Endoglucanase K	Clocel_2821	GH9, CBM4	4.38	0.80	Extracell
Pectate lyase A	Clocel_1623	PL1, PL9	4.35	0.57	Extracell
<i>Non-cellulosomal proteins</i>					
Endoglucanase O	Clocel_1478	GH9, CBM4	6.06	0.60	Extracell
B-mannanase	Clocel_1134	GH26, CBM23	4.22	0.66	Cell wall
Endoglucanase D	Clocel_3242	GH5, CBM2	3.35	0.59	Membrane
β-xylosidase	Clocel_2595	GH43	3.16	-	Unknown
Endoglucanase	Clocel_2606	GH5, CBM46	2.64	0.64	Cell wall

Figure 1

[Click here to download Figure: Figure 1.pptx](#)

Figure 1

Figure 2
[Click here to download Figure: Figure 2.pptx](#)

Figure 2

Figure 3

[Click here to download Figure: Figure 3.pptx](#)

Figure 3

[Click here to download Figure: Figure 4.pptx](#)

Figure 4

A

B

Figure 5

[Click here to download Figure: Figure 5.pptx](#)

Figure 5

Supplementary figure 1

[Click here to download Supplementary material: Supplementary Figure 1.pptx](#)

Supplementary file 1

[Click here to download Supplementary material: Supplementary File 1.xlsx](#)

***Conflict of Interest**

[Click here to download Conflict of Interest: coi_disclosure RM.pdf](#)