



# AperTO - Archivio Istituzionale Open Access dell'Università di Torino

# Pre- and postnatal exposure to glyphosate-based herbicide causes behavioral and cognitive impairments in adult mice: evidence of cortical ad hippocampal dysfunction

# This is the author's manuscript

Original Citation:

Availability:

This version is available http://hdl.handle.net/2318/1731182 since 2020-04-29T23:49:41Z

Published version:

DOI:10.1007/s00204-020-02677-7

Terms of use:

**Open Access** 

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)

- Pre- and postnatal exposure to glyphosate-based herbicide causes behavioral and cognitive impairments
   in adult mice: evidence of cortical and hippocampal dysfunction
- Yassine Ait bali<sup>1,2</sup>, Saadia Ba-M'hamed<sup>1</sup>, Giovanna Gambarotta<sup>3</sup>, Marco Sassoè-Pognetto<sup>2,4</sup>, Maurizio
   Giustetto<sup>2,4</sup>, Mohamed Bennis<sup>1\*</sup>
- <sup>1</sup>Laboratory of Pharmacology, Neurobiology and Behavior (URAC-37), Cadi Ayyad University, Marrakech,
   Morocco.
- 8 <sup>2</sup>Department of Neuroscience, University of Turin, Turin, Italy.
- 9 <sup>3</sup>Department of Clinical and Biological Sciences, University of Turin, Turin, Italy
- 10 <sup>4</sup>National Institute of Neuroscience-Italy, Turin, Italy
- 11

# 12 \*Corresponding author:

- 13 Mohamed BENNIS:
- 14 Lab of Pharmacology, Neurobiology and Behavior (URAC-37),
- 15 Faculty of Sciences Semlalia, Cadi Ayyad University,
- 16 Bd. Prince My Abdallah, BP. 2390, 40000,
- 17 Marrakech, Morocco
- 18
- 19 Phone: 212 670099315
- 20 FAX: 212 524 436769
- 21 e-mail: mbennis@uca.ma
- 22

# 23 Acknowledgments:

24 The authors are grateful to Prof I. Elatiri for technical assistance and to Prof A. Buffo for fruitful discussions

25 about astrocytes and microglia analysis. Y. A-B. was a recipient of a fellowship from the Italian Ministry of

- 26 Foreign Affairs and International Cooperation. We are grateful to Fondo di Beneficienza Intesa Sanpaolo (Italy)
- 27 for financial support.

# 28 Abstract

29 Glyphosate-based herbicides (GBH) are the most widely used pesticides worldwide. Despite considerable 30 progress in describing the neurotoxic potential of GBH, the harmful effects on brain cytoarchitecture and 31 behavior are still unclear. Here, we addressed the developmental impact of GBH by exposing female mice to 250 32 or 500 mg/kg doses of GBH during both pregnancy and lactation and then examined the downstream effects at 33 the behavioral, neurochemical and molecular levels. We show that pre- and neonatal exposure to GBH impairs 34 fertility and reproduction parameters as well as maternal behavior of exposed mothers. In offspring, GBH was 35 responsible for a global delay in innate reflexes and a deficit in motor development. At the adult age, exposed 36 animals showed a decrease of locomotor activity, sociability, learning and short and long-term memory 37 associated with alterations of cholinergic and dopaminergic systems. Furthermore, GBH activated microglia and 38 astrocytes, sign of neuroinflammation event in the medial prefrontal cortex and hippocampus. At the molecular 39 level, a down-regulation of brain-derived neurotrophic factor (BDNF) expression and an up-regulation of 40 tyrosine related kinase receptor (TrkB), NR1 subunit of NMDA receptor as well as tumor necrosis factor a 41 (TNF $\alpha$ ) were found in the brain of GBH-exposed mice. 42 The present work demonstrates that GBH induces numerous behavioral and cognitive abnormalities closely 43 associated with significant histological, neurochemical and molecular impairments. It also raises fundamental

44 concerns about the ability of current safety testing to assess risks of pesticide exposure during developmental

- 45 periods of central nervous system.
- 46 Key words: Glyphosate, behavior, cognition, acetylcholinesterase, neuroinflammation, BDNF signaling.

#### 47 Introduction

48 The American National Academy of Sciences estimated that 3% of brain developmental disorders, such as 49 autism spectrum disorders and learning disabilities, may be directly linked to exposure to environmental 50 chemicals (National Research Council 2000). Existing evidence supports the idea that lifelong susceptibility to 51 anxiety disorders can be partly determined by environmental factors during early development (Gross and Hen 52 2004). Furthermore, it is important to note that early perturbations of brain development may lead to 53 neurobehavioral disturbances expressed either in childhood or with delayed onset in adulthood (Olney, 2002). In 54 line with this, the potential of in utero or early postnatal pesticide exposure to affect brain development has been 55 shown and an emerging literature provides evidence for neurobehavioral consequences resulting from early 56 exposure to organophosphates (OPs) pesticides (for review, see Heyer and Meredith 2017).

57 Glyphosate (Gly) (N-[phosphonomethyl]glycine; CAS registry number 1071-83-6), the active ingredient present 58 in Roundup® (Monsanto Company, St. Louis, MO), is the most heavily OP herbicide used worldwide (Powles et 59 al. 1997). The herbicidal action of Gly is due to inhibition of a key plant enzyme (5-enolpyruvyl shikimate-3-60 phosphate synthase) (Franz et al. 1997). Since this enzyme is not present in vertebrates, it has long been assumed 61 that Gly does not affect non-target species. In addition, experimental data showing that the blood Gly 62 concentrations peaked at around 1-2 h post ingestion and that Gly is poorly metabolized and generally 63 eliminated in the urine and feces (Chan and Mahler 1992; Brewster et al. 1991) supported the safety hypothesis 64 of Gly. However, over the last few years, several concerns have been raised by the scientific community and 65 regulatory agencies regarding the potential adverse effects of Gly and its adjuvants on the environment and 66 human health (Solomon et al. 2007). Evidence of exposure to Gly has been revealed by its detection in urine 67 samples of people living in farm and non-farm household (Conrad et al. 2017) and in maternal and umbilical 68 cord serum of pregnant women in Thailand (Kongtip et al. 2017). Clinical reports of intoxication with 69 commercial formulations of Gly described negative effects on the nervous system, including Parkinsonism 70 (Barbosa et al. 2001; Wang et al. 2011), anxiety and short-term memory impairments (Nishiyori et al. 2014). 71 Moreover, an epidemiological study highlighted a strong correlation between the increasing application of Gly in 72 agriculture and the occurrence of several neurological diseases, including autism, dementia and anxiety disorder 73 at different ages (Seneff et al. 2015). Concomitantly, experimental studies also revealed neurotoxic effects of 74 glyphosate-based herbicide (GBH). Indeed, oral administration of GBH to pregnant rats alters the activity of 75 brain enzymes in both mothers and offspring (Daruich et al. 2001). These data suggest that GBH could impact 76 maternal behavior and subsequently offspring's sociability given the crucial role of early maternal care on 77 offspring's abilities (Branchi et al. 2013). However, there have been no systematic studies evaluating the effects 78 of GBH on maternal care and social behavior of their offspring. At the behavioral level, it has been shown that 79 early exposure to GBH results later in life in depressive-like behavior (Cattani et al. 2017) and contradictory 80 effects on the anxiety-related behavior (Gallegos et al. 2016; Baier et al. 2017). Recently, we found that GBH 81 induces cognitive alterations in young male mice, supporting the idea that early exposure to GBH can interfere 82 with brain structures involved in learning and memory leading to cognitive deficits (Bali et al. 2019). However, 83 the effect of prenatal and neonatal exposure to GBH on learning and memory is still unexplored. In this context, 84 the present study was conducted using a multifaceted behavioral battery to assess the profile of gestational and 85 lactational GBH effects in a mouse model. Behavioral assays covering neonatal age and adulthood were selected 86 to measure a range of early reflex development as well as locomotor, affective, sociability and cognitive

- 87 functions. Mechanistic, cytoarchitectural, neurochemical and molecular mechanisms underlying of GBH-
- 88 induced neurobehavioral deficits were evaluated in this follow-up study, with the goal to provide benchmark
- 89 data for GBH risk assessment in the brain.

# 90 Materials and Methods

# 91 Pesticide

- 92 Roundup herbicide (glyphosate concentration: 360g/l in the form of glyphosate isopropylamine salt 486 g/l), was
- 93 used in the liquid commercial form supplied by Monsanto Company (St. Louis, MO, USA). The molecular
- 94 formula is C<sub>6</sub>H<sub>17</sub>N<sub>2</sub>O<sub>5</sub>P (molecular weight: 228.183 g/mol; melting point: 200 °C; density: 1.218 g/cm3)
- 95 Animals
- 96 Male and female Swiss mice (3-months-old) were obtained from the animal husbandry of the Faculty of
- 97 Sciences, Cadi Ayyad University, Marrakech, Morocco. The animals were housed in Plexiglas cages (30 x 15 x
- 98 12 cm) under standard conditions of temperature  $(22 \pm 2^{\circ}C)$  and photoperiod (12h / 12h) with food and water *ad*
- 99 libitum. All procedures were approved by the Council Committee of Research Laboratories of the Faculty of
- 100 Sciences, Cadi Ayyad University (Marrakech, Morocco) and conducted in accordance with European Council
- 101 Directive: EU2010/63. All efforts were made to minimize animal suffering.

# 102 Doses and protocol of exposure

- Females were mated with breeding males (2 females for 1 male) over a day, and were examined on the following day by vaginal plug inspection to assess successful mating. If judged copulated, the female was removed from the cage of the male and housed individually. This was considered as the day 0 of gestation (G0).
- 106 Female exposure to GBH through oral gavage occurred daily from G0 to postnatal day 21 (P21) (Fig. 1). Three
- 107 experimental groups were formed, each one including a minimum of six female mice: a group exposed to a
- 108 lower dose (250 mg/kg) of GBH, a group exposed to a higher dose (500 mg/kg) of GBH, and a control group
- 109 which received vehicle (tap water). These doses were selected based on the no-observed adverse effect level
- 110 (NOAEL) indications (i.e. 500 mg/kg/day) (EPA, 1993). The GBH doses used in the present study were higher
- 111 than the GBH levels to which the population is normally exposed (Solomon, 2016). However, as in other
- 112 toxicological studies, exposure to relatively high doses was used in order to demonstrate a plausible drug-
- 113 action (see for example Ford et al. 2017).
- 114 Gestation outcomes, maternal behavior and body weight of pups
- 115 In order to detect any signs of poisoning, all pregnant mice were observed daily from the first administration day
- 116 (G0) until parturition. In addition, several parameters of maternal behavior, fertility and reproduction were
- 117 evaluated according to Ema et al. (2007).

#### 118 Motor and sensory development assessment

- The behavioral testing (negative geotaxis, righting reflex, cliff avoidance and rotarod tests) was performed as described by Ait bali et al. (2016). Animals (n = 10 for each group: control, 250 mg/kg and 500 mg/kg; two males from each litter) were tested during morning sessions starting at 9 a.m. The tests for sensorimotor
- 122 development assessed during the same day were separated by an interval of 30 min.

# 123 Adult behavior

- 124 From P60, behavioral tests were performed in order to assess locomotor activity (open field, OF), levels of
- 125 anxiety (OF and elevated-plus maze, EPM), social interaction (three-chambered sociability test, TCS), working
- 126 memory (Y-maze), recognition memory (novel object recognition test, NOR), and learning and emotional
- 127 memory performances (passive avoidance test, PA). The behavior of a total number of 10 mice for each group
- 128 was evaluated between 9 a.m. and 13 p.m and recorded with Ethovision XT Noldus 8.5 video tracking program

129 (Noldus Information Technology b.v., Wageningen, The Netherlands), connected to a video camera (JVC,

130 Japan).

# 131 Open field

132 This test was performed to assess the general locomotor activity (Walsh and Cummins 1976). Activity

- 133 monitoring was conducted in a square shaped, white arena, measuring  $50 \times 50 \times 50$  cm. Mice were placed
- individually into the arena and monitored for 20 min. The assessed parameters were the total distance travelled,
- 135 the velocity and the time spent in the center.

# 136 Elevated-plus maze

- 137 The elevated plus maze is a widely accepted paradigm used to assess anxiety-like behavior in rodents (Pellow et 138 al. 1985). The elevated plus-maze included two opposing open arms (OA) ( $50 \times 5$  cm) and two closed arms (CA) 139 ( $50 \times 5 \times 15$  cm) joining at a square central area ( $5 \times 5$  cm) to form a plus sign. The entire apparatus was
- 140 elevated to a height of 45 cm above the floor. Each mouse was tested within a 5-min test session. At the
- beginning, each mouse was placed individually in the central area facing one of the open arms and allowed to
- 142 freely explore the maze. The time spent in the OA and CA as well as the number of entries into each arm were
- 143 quantified. An anxiety index (1- [([open arm time/total time] + [open arm entries/total number of entries])/2])
- 144 was determined according to Cohen et al. (2013).

# 145 Three-chambered sociability test

146 The social interaction test was run in a three-chambered arena made of clear glass. Retractable doorways, built 147 into the two dividing walls, controlled access to the side chambers. Each of the two outside chambers had an 148 inverted empty wire cup, one housing a male "stimulus" mouse age-matched to the "test" mouse, and the other 149 with a plastic object ("plastic mouse"). The test session began with a 5 min habituation session with the test 150 mouse free to explore the entire arena. This mouse was then briefly confined to the center chamber while the 151 plastic object was placed in the cup on one side and an adult male mouse on the other side. The "stimulus" 152 mouse and the "plastic mouse" sides were alternated, left and right, between tests. Once the stimuli were in 153 position, the two side doors were simultaneously raised and the test mouse could access all three chambers for 5 154 min. Automatic monitoring recorded and scored the time spent in contact with each wire cup as well as the 155 number of visits. The apparatus was cleaned between tests using a 70% ethanol/ water solution.

#### 156 *Y-maze*

Y-maze was used in order to assess working memory performances (Hughes, 2004). This 3 arms apparatus was made of brown wood ( $60 \text{ cm} \times 15 \text{ cm} \times 30 \text{ cm}$ ), positioned at equal angles of  $120^{\circ}$ . Mice were placed at the end of one arm and allowed to freely explore the maze over an 8 min session. The series of arm entries were recorded, and alternation was defined as a triplet of explored arms. Alternation was considered as successful when the 3 arms were different. The percentage of spontaneous alternation was calculated according to the following equation: % alternation = [(number of alternations) / (total arm entries - 2)] x 100 (Chen et al. 2016).

#### 163 Novel object recognition

164 This test was used to evaluate recognition memory. It was based on the natural preference of mice for a new 165 object with respect to a familiar one. The apparatus consisted of an OF made in Plexiglas (50 x 50 x 50 cm)

- 166 containing two identical or different objects according to the phase of the test. The objects to be discriminated
- 167 were three plastic objects. The task procedure consists of three phases: habituation, training, and retention phase,

- 168 according to the protocol described by Bevins and Besheer (2006) for a one trial nonmatching to sample learning
- 169 procedure. The habituation phase with the apparatus was conducted for 10 min without the presence of the
- 170 objects. The next day, during the training session, two identical objects were placed in the back corner of the
- 171 box. The experimental mouse was then placed midway at the front of the box and the total time spent exploring
- 172 the two objects was recorded for 5 min. During the retention session (1 h after), one of the two identical objects
- 173 was replaced by a new one and the mouse was allowed to explore the different objects for 10 min. The time
- 174 spent next to each of the two objects (the familiar one and the novel one) was recorded and discrimination
- 175 between them was calculated using a discrimination index [DI = (novel object exploration time/ total exploration
- 176 time of the two objects) (familiar object exploration time/ total exploration time of two objects) x 100].

# 177 Passive avoidance

178 The test is based on the association formed between an aversive stimulus (a foot shock) and a specific 179 environmental context. The apparatus consisted of a two-compartment (light-dark) box. The light compartment 180  $(10 \times 13 \times 15 \text{ cm})$  was illuminated while the dark one  $(10 \times 13 \times 15 \text{ cm})$  was equipped with energized grid floor, 181 separated by a guillotine door. The entrance of animals to the dark box was punished by an electric foot shock 182 (0.2 mA for 1 s duration). 24 h before the training session, the mice were allowed to explore freely the apparatus 183 for 3 min (habituation). On the training day, each mouse was placed in the center of the light compartment facing 184 away from the guillotine door. After 10 s of adaptation the guillotine door was opened exposing the dark 185 compartment. When the mouse entered the dark box with all four paws, the guillotine door was closed, and the 186 foot shock was delivered. On the test day, the mouse was returned to the illuminated compartment and the 187 procedure was repeated except that no shock was delivered. The test session was carried out 2 h (short-term 188 memory) or 24 h (long-term memory) after the training. Each time, the latency to enter the dark compartment 189 was recorded. Mice whose latency on the training session exceeded 60 s were excluded from the experiment in 190 order to minimize the deviation of baseline data. If the animal did not enter the dark compartment during the test 191 within 300 s, the trial was fstopped and the final score was established as 300 s. After each session, the apparatus 192 was cleaned using 70% ethanol.

# 193 Determination of acetylcholinesterase enzyme activity

- After the behavioral analyses, mice were killed by decapitation, their brains immediately removed from the skull and the PFC and hippocampi were dissected for biochemical analyses. Acetylcholinesterase (AChE) activity was determined in tissue homogenates as described previously (Bali et al. 2019). Briefly, acetylthiocholine (ASCh) was used as a substrate of AChE with Thiocholine (SCh) as a reaction product. The activity of AChE was determined according to the colorimetric method of Ellman et al. (1961) based on the reaction of SCh with DTNB (5,5'-Dithiobis (2-nitrobenzoic acid)), which gives a yellow compound (TNB: S Thio-2-*N* nitro-benzoate) absorbing light at 412 nm. The absorbance of the TNB measured with a spectrophotometer is proportional to the
- 201 enzymatic activity of AChE. The specific activity is calculated and presented as a percentage:

$$SA = \frac{\Delta OD / \min \times 1000}{\varepsilon \times [weight \ of brain \ tissue]}$$

- 203  $\Delta OD/min$ : Variation of OD per min;
- 204 [weight of brain tissue of each mouse (g);
- 205  $\varepsilon$ : Molar extinction coefficient of TNB at 412 nm with  $\varepsilon TNB = 13.6 \times 10^{-6} M^{-1} cm$ .
- 206 Tissue sampling and immunofluorescence

207 Upon conclusion of behavioral testing, control and treated mice were anesthetized with an intraperitoneal 208 injection of urethane 40% (1 g/kg, from Sigma–Aldrich, France) and transcardially perfused with saline solution 209 (0.9%), followed by ice-cold 4% formaldehyde in phosphate buffered saline (PBS; 0.1 M). The brains were then 210 removed, post-fixed in the same fixative for 12 h and cryoprotected overnight in 30% sucrose. They were then 211 cut on a freezing cryostat (Leica Microsystems, Germany) into 30 µm frontal sections. The sections containing 212 the substantia nigra pars compacta (SNc), the ventral tegmental area (VTA) and the striatum were used for 213 tyrosine hydroxylase (TH) immunofluorescence, while sections containing PFC and dorsal hippocampus stocked 214 for GFAP and Iba-1 immunofluorescence. The regions of interest were determined according to stereotaxic atlas 215 of Paxinos and Franklin (2001). Sections were kept in PBS containing 0.05% Triton X-100 and 10% normal 216 donkey serum (NDS) for 1 h. Thereafter, they were incubated with the appropriate primary antibodies (mouse 217 monoclonal anti-TH 1 :1000, Immunostar cat. 22941; rabbit polyclonal anti-GFAP; 1:500, Abcam, Ab7260; 218 rabbit polyclonal anti-Iba-1; 1 :1000; Wako, cat. 019-19741) diluted in PBS with 3% NDS and 0.05 Triton X-219 100 with gentle stirring at 25°C. The following day, the sections were rinsed and incubated for 2h with the 220 adequate secondary antibodies (1:1000; Jackson ImmunoResearch, West Grove, PA, USA) at room temperature 221 for 2h. After PBS rinsing, the sections were mounted on gelatin-coated glass slides and coverslipped with Dako 222 fluorescence mounting medium (Dako Italia, Milan, Italy).

#### 223 Image analysis

224 The TH-immunostained sections were used to assess the number of dopaminergic cells in the SNc and VTA 225 (Bregma: - 6.30 mm) and dopaminergic fibers in the striatum (Bregma: 0.48 mm), while the GFAP and Iba-1 226 immunostained sections were used to assess reactive astocytes and microglia, respectively, in the PFC (Bregma: 227 3.20 mm) and dorsal hippocampus (Bregma: - 3.80 mm). Three mice per group were used for these analyses. For 228 each brain region, three representative sections (0.5 µm Z-step size) from anterior to posterior were acquired 229 with a laser scanning confocal microscope (LSM5 Pascal; Zeiss, DE, Germany) using either a 20x objective (for 230  $TH^+$ , GFAP<sup>+</sup> and Iba-1<sup>+</sup> cells count) or a 40x objective (for measuring the density of  $TH^+$  innervation) with the 231 pinhole was set at 1 Airy unit. TH<sup>+</sup>, GFAP<sup>+</sup>, and Iba-1<sup>+</sup> cells were manually counted using the point tool in 232 ImageJ software (Image processing and analysis in Java, NIH, USA). The ROI Manager tool in Image-J software 233 was employed to quantify integral optical density of TH, GFAP and Iba-1 expression. All analyses were carried

234 out by an operator blinded to the experimental groups.

# 235 RNA isolation, cDNA preparation, and quantitative real-time PCR

236 Twenty four hours following the completion of behavioral and cognitive tests, RNA was extracted from PFC and 237 hippocampus tissues from control and GBH 500 mg/kg exposed mice (n = 3 each). Isolation of the total RNA 238 was carried out using Trizol (Sigma-Aldrich, St. Louis, MO, USA) according to a previously described protocol 239 (Rio et al. 2010). RNA level was quantified by measuring absorbance at 260 and 280 nm. The final RNA 240 concentration and purity was determined using Nanodrop (Thermo Fisher Scientific, Waltham, MA, USA), 241 obtaining absorbance 260/280 ratios between 2 and 2.2 in all the samples. Retrotranscription of 1 µg total RNA 242 was carried out in a 25 µl reaction volume containing: 1x RT-Buffer, 0.1 µg/µl bovine serum albumin (BSA), 243 0.05% Triton, 1 mM dNTPs, 7.5 µM Random Hexamer Primers, 40 U RIBOlock, and 200 U RevertAid® 244 Reverse Transcriptase (all RT ingredients were provided by Thermo Scientific). The reaction was performed 10 245 min at 25°C, 90 min at 42°C, 15 min at 70°C. Quantitative real-time PCR (qRT-PCR) was carried out using an 246 ABI Prism 7300 (Applied Biosystems) detection system. Analyses were performed in technical duplicate and 247 biological triplicate. Data from qRT-PCR experiments were analyzed using the delta cycle threshold ( $\Delta Ct$ ) 248 method. The expression levels were normalized to reference gene: TBP (TATA box Binding Protein). For each 249 tissue (PFC, hippocampus), the  $\Delta$ Ct average of control samples was used as calibrator. Primers were designed 250 using Annhyb software (http://www.bioinformatics.org/annhyb/) and were synthesized by Invitrogen. Primer 251 sequences are reported in supplementary table 1.

# 252 Statistical analysis

253 Fertility and reproduction parameters as well as maternal behavior were analyzed by one-way ANOVA, while

- body weight gain and early sensorimotor endpoint results were analyzed using the repeated measure two-way
- ANOVA (GBH dose and age), followed by a Holm-Sidak's *post hoc* test for multiple comparisons. The dataset
- 256 of behavioral tests in adult mice, enzyme activity results and histological assays were compared between
- 257 different groups (treated and control) and analyzed using one-way ANOVA, followed by a Holm-Sidak's post
- 258 hoc for multiple comparisons. The biomolecular data were analyzed with t-test. The results are presented as
- 259 mean  $\pm$  S.E.M, and a value of p < 0. 05 was considered statistically significant. All statistical analyses were
- 260 carried out using the software SigmaPlot 11.0 for Windows and all graphs were generated with Prism 7.0 for
- 261 Windows (GraphPad software).

#### 262 **Results**

263 Gestation outcome and maternal behavior following GBH exposure

264 Administration of GBH to pregnant females affected fertility and reproduction parameters. Indeed, the fertility

265 rate and the gestational index were lower in treated groups compared to control. Similarly, both the number of

266 litters and the total number of mice per litter were significantly lower in the GBH exposed groups. Treatment 267 with GBH 500 mg/kg, but not with the lower dosage, also affected retrieving and nesting index (table 2). In

268 contrast, the statistical analysis did not reveal a significant difference in the gestation length between treated and

269 control groups ( $F_{(2.18)} = 1.08, p > 0.05$ ) (table 1).

270	Table 1	Reproductive	findings	in mice	given	GBH	during	pregnancy	and	lactation
-----	---------	--------------	----------	---------	-------	-----	--------	-----------	-----	-----------

	G	lyphosate-based herbicide	
—	0 (Control)	250 mg/kg	500 mg/kg
No. of females copulated	8	10	24
No. of pregnant females	7	6	6
Fecundity Index (%)	87	60	25
No. of death during pregnancy	0	0	1
Gestation length (days)	$19.8\pm0.37$	$20 \pm 0.54$	$19.2\pm0.2$
No. of females with live born	7	6	5
Gestation index (%)	100	100	83
No. of females with totally litter loss	0	0	1
No. of litters	7	6	5
Total no. of pups born	56	40	35
No. of pups born alive	51	34	33
No. of dead pups	5	6	2
Delivery index (%)	100	100	100
Lactation index (%)	88.2	79.4	69.6
Nest building index (%)	100	100	80
Retrieving index (%)	100	100	80
% of males	37	45	40

Fecundity Index (%) = (no. of pregnant females/ no. of females copulated) x 100.

Values are given as the mean  $\pm$  S.D.

Gestation index (%) = (no. of females with live pups born / no. of pregnant females) x 100.

Delivery index (%) = (no. of females delivering / no. of pregnant females) x 100.

Lactation index (%) = (no. of living offspring on day 21 / no. of offspring born alive) x 100.

Nesting index = (no. of females building theirs nests for a maximum duration of 30 min/ no. of females delivering) x100.

271 272 273 274 275 276 277 278 Retrieving index = (no. of females retrieving over pups for a maximum duration of 30 min/ no. of females delivering) x 100.

% of males = (no. of males / no. of pups) x 100.

#### 279 GBH decreased body weight of pregnant females and their offspring

280 The two-way repeated measures ANOVA showed a significant effect of treatment ( $F_{(2,19)} = 11.80, p < 0.01$ ) and

281 period of treatment ( $F_{(2,19)} = 211.62$ , p < 0.001) on dam weights. However, the interaction between the two

282 factors was not significant ( $F_{(2,19)} = 0.95$ , p > 0.05). The *post-hoc* comparisons showed that body weight gain was

- 283 significantly reduced following GBH treatment at both 250 and 500 mg/kg with respect to the control (p < 0.05)
- 284 (table 2). Likewise, our results indicated a significant effect of treatment and age on pup's body weight ( $F_{(2,19)} =$
- 285 11.80, p < 0.01;  $F_{(2,19)} = 11.80$ , p < 0.01, respectively) as well as for the interaction between the two factors
- 286  $(F_{(2,19)} = 11.80, p < 0.01)$ . Furthermore, Holm-sidak comparisons revealed a significant reduction of body weight
- 287 gain in offspring delivered from both GBH exposed dams groups p < 0.05) (table 2).

	Chu	hasata hasad har	biaida			Post-	hoc		
	Слур	nosate-based ner	biciue	250 vs a	control	500 vs c	control	250 vs	s 500
	0 (Control)	250 mg/kg	500 mg/kg	t	р	t	р	t	р
Pregnancy									
G1	$0.78\pm0.34$	$-1.22 \pm 0.48$	$-3.66 \pm 0.14$	3.59	ns	7.98	***	1.94	ns
G7	$5.24\pm0.55$	$\textbf{-2.20}\pm0.66$	$0.24\pm0.49$	5.46	*	8.98	***	1.56	ns
G15	$10.38\pm0.67$	$8.12\pm0.55$	$5.18\pm0.56$	4.06	ns	9.34	***	2.34	ns
G19	$16.34 \pm 0.65$	$14.16\pm0.49$	$10.48\pm0.78$	3.91	ns	10.53	***	2.93	**
Lactation									
L1	$9.22\pm0.95$	$5.04 \pm 1.19$	$3.80\pm0.76$	4.34	ns	5.63	*	0.98	ns
L7	$10.00\pm0.93$	$6.76 \pm 1.28$	$5.52 \pm 1.01$	3.37	ns	4.66	ns	0.33	ns
L15	$11.06\pm0.66$	$7.54 \pm 1.31$	$6.30\pm0.83$	3.66	ns	4.95	*	2.80	ns
L19	$11.44\pm0.83$	$8.44 \pm 1.01$	$7.20\pm0.51$	3.12	ns	4.41	ns	0.33	ns
Litter									
P1	$1.31\pm0.02$	$1.15\pm0.07$	$1.05\pm0.88$	1.02	ns	1.65	ns	0.51	ns
P 7	$2.51\pm0.03$	$2.08\pm0.14$	$1.91\pm0.11$	2.68	ns	3.79	ns	0.90	ns
P 15	$5.47\pm0.23$	$4.06\pm0.22$	$3.17\pm0.13$	8.92	***	14.52	***	4. 59	***
P 21	$7.56\pm0.18$	$6.57\pm0.24$	$6.43\pm0.20$	6.23	**	7.10	***	0.71	ns

#### 288 Table 2 GBH affects body weight gain of mice mothers and their offspring

289

G gestation day; L lactation day; P postnatal day; \*p < 0.05; \*\*p < 0.01; \*\*\*p < 0.001; ns (no significant) p > 0.05

# 290 GBH delayed developmental skills of pre- and postnatally exposed offspring

**Righting reflex:** We investigated whether pre- and post-natal exposure to GBH caused atypical sensorimotor skills development in offspring. Two-way repeated measure ANOVA showed a significant main effect of treatment ( $F_{(2,23)} = 13.83$ , p < 0.001) and the expected effect of age, indicating a reduction in righting time as the mice developed ( $F_{(2,23)} = 309.10$ , p < 0.001). Notably, GBH treatment delayed the development of the righting reflex as shown by a significant interaction between the two factors ( $F_{(2,23)} = 8.38$ , p < 0.001). *Post-hoc* analysis revealed that mice treated with 250 mg/kg or 500 mg/kg GBH had slower righting reflexes time than the controls only at P5 (250 vs control mg/kg: q = 7.45, p < 0.001; 500 mg/kg vs control: q = 10.32, p < 0.001) (Fig. 2a).

298Negative geotaxis:The two-way ANOVA revealed a significant effect of treatment ( $F_{(2,23)} = 83.84$ , p < 0.001)299and the expected maturation effect reflected by the significant effect of age ( $F_{(2,23)} = 59.44$ , p < 0.001). The300interaction was also significant ( $F_{(2,23)} = 23.41$ , p < 0.001). Post-hoc comparisons confirmed that mice treated301with 500 mg/kg had higher negative geotaxis time compared to the controls at P5, 7 and 9 (q = 19.27, p < 0.001;302q = 10.45, p < 0.001 and q = 5.97, p < 0.01, respectively) (Fig. 2b).

303 *Cliff avoidance:* A main effect of treatment was found ( $F_{(2,23)} = 4.05$ , p < 0.05), as well as the expected effect of 304 age ( $F_{(2,23)} = 23.05$ , p < 0.001). However, the interaction was not significant ( $F_{(2,23)} = 2.40$ , p > 0.05). *Post-hoc* 305 comparisons confirmed that mice treated with 500 mg/kg of GBH had slower cliff avoidance time than controls 306 only at P 5 (q = 4.84, p < 0.05) (Fig. 2c).

307 *Traction test:* One-way ANOVA analysis showed a significant difference between treated and control groups

308 ( $F_{(2,23)} = 21.38$ , p < 0.001). Post-hoc analysis revealed that mice treated with 250 mg/kg or 500 mg/kg GBH

309 showed shorter fall down latencies than the controls at P10 (250 mg/kg vs control: q = 2.41, p < 0.05; 500 mg/kg

310 vs control: q = 6.47, p < 0.001). The difference between the two doses of GBH was also statistically significant 311 (250 mg/kg vs 500 mg/kg; q = 4.06, p < 0.01) (Fig. 2d).

312 *Rotarod test:* The two-way ANOVA analysis revealed that the motor coordination was significantly affected by

313 the treatment ( $F_{(2,23)} = 13.61$ , p < 0.001), as well as by the age of the animals ( $F_{(2,23)} = 64.04$ , p < 0.001).

- 314 Likewise, the interaction was significant ( $F_{(2,23)} = 4.76$ , p < 0.001). Multiple comparisons confirmed that both
- 315 GBH-treated groups had lower fall down latency than the control group on P23 (250 mg/kg vs control: q = 4.95,
- 316 p < 0.05; 500 mg/kg vs control: q = 8.01, p < 0.001) and P24 (250 mg/kg vs control: q = 4.83, p < 0.05; 500
- 317 mg/kg vs control: q = 5.42, p < 0.01 (Fig. 2e).

# 318 Effects of GBH on offspring lasting into adulthood

- 319 In order to investigate whether GBH exposure during prenatal and postnatal developmental might translate into
- 320 long-lasting consequences on adult behavior, we tested the behavioral repertoire of adults' offspring by assaying
- 321 locomotor activity, anxiety-like phenotype, social interaction as well as different forms of memory.

#### 322 Open field test

323 As indicators of locomotor activity and anxiety-like levels, we recorded the total distance traveled over the open 324 field as well as the velocity and the time spent in the central zone of the maze (anxiety index). One-way ANOVA 325 analysis of the total distance traveled and the percentage of the time spent in the central zone revealed significant 326 differences between treated and control groups ( $F_{(2,17)} = 3.76$ , p < 0.05;  $F_{(2,17)} = 50.67$ , p < 0.001, respectively). 327 However, no difference was found between groups for the velocity ( $F_{(2,17)} = 3.76$ , p > 0.05) (Fig. 3c), even 328 though treated mice seemed slower than controls. Multiple comparisons confirmed that the group of 500 mg/kg 329 exhibited significant decrease of distance traveled compared to the control group (t = 2.69, p < 0.05) (Fig. 3b). 330 Similarly, post hoc analysis revealed that both GBH-treated groups spent significantly less time in the center of 331 the open field compared to the control (250 mg/kg vs control: t = 6.57, p < 0.001; 500 mg/kg vs control: t = 9.88, 332 p < 0.001). In addition, the same analysis revealed a significant difference between treated groups (t = 3.30,  $p < 10^{-10}$ 

333 0.01) (Fig. 3d).

#### 334 Elevated-plus maze

The open field result was confirmed by EPM data. Indeed, One-way ANOVA analysis of the ratio of time spent in the OA and the anxiety index showed a significant difference between treated and control group ( $F_{(2,17)} =$ 72.38, p < 0.001;  $F_{(2,17)} = 18.17$ , p < 0.001, respectively). However, the analysis of the ratio of entries number into the OA did not showed any statistical difference between groups ( $F_{(2,17)} = 1.05$ , p > 0.05) (Fig. 3g). Multiples comparisons confirmed that the ratio of time spent in the OA was significantly lower in treated groups compared to the control group (250 mg/kg *vs* control: t = 9.29; p < 0.001; 500 mg/kg *vs* control: t = 11.26; p <0.001) (Fig. 3f). Moreover, the anxiety index was significantly higher in treated groups (250 mg/kg *vs* control: t

342 = 4.36; p < 0.001; 500 mg/kg vs control: t = 5.78; p < 0.001) (Fig. 3h).

#### 343 Three-chambered sociability

344 This test was used to investigate the voluntary social interaction of animals. The data analysis revealed a 345 significant interaction between the wire cup (holding mouse vs object) and treatment ( $F_{(2,17)} = 73.53$ , p < 0.001). 346 However, there was no significant main effect of both treatment and wire cup ( $F_{(2,17)} = 2.00, p > 0.05$ ;  $F_{(2,17)} =$ 347 1.10, p > 0.05, respectively). Post-hoc comparisons revealed that mice treated either with 250 or 500 mg/kg of 348 GBH spent less time with the wire cup holding another conspecific (250 mg/kg vs control: t = 7.42, p < 0.001; 349 500 mg/kg vs control: t = 9.42, p < 0.001; 250 mg/kg vs 500 mg/kg: t = 1.99, p < 0.05) and more time with the 350 wire cup holding the object (250 mg/kg vs control: t = 4.89, p < 0.001; 500 mg/kg vs control: t = 7.06, p < 0.001; 500 mg/kg vs control: t = 7.06, p < 0.001; 500 mg/kg vs control: t = 7.06, p < 0.001; 500 mg/kg vs control: t = 7.06, p < 0.001; 500 mg/kg vs control: t = 7.06, p < 0.001; 500 mg/kg vs control: t = 7.06, p < 0.001; 500 mg/kg vs control: t = 7.06, p < 0.001; 500 mg/kg vs control: t = 7.06, p < 0.001; 500 mg/kg vs control: t = 7.06, p < 0.001; 500 mg/kg vs control: t = 7.06, p < 0.001; 500 mg/kg vs control: t = 7.06, p < 0.001; 500 mg/kg vs control: t = 7.06, p < 0.001; 500 mg/kg vs control: t = 7.06, p < 0.001; 500 mg/kg vs control: t = 7.06, p < 0.001; 500 mg/kg vs control: t = 7.06, p < 0.001; 500 mg/kg vs control: t = 7.06, p < 0.001; 500 mg/kg vs control: t = 7.06, p < 0.001; 500 mg/kg vs control: t = 7.06, p < 0.001; 500 mg/kg vs control: t = 7.06, p < 0.001; 500 mg/kg vs control: t = 7.06, p < 0.001; 500 mg/kg vs control: t = 7.06, p < 0.001; 500 mg/kg vs control: t = 7.06, p < 0.001; 500 mg/kg vs control: t = 7.06, p < 0.001; 500 mg/kg vs control: t = 7.06, p < 0.001; 500 mg/kg vs control: t = 7.06, p < 0.001; 500 mg/kg vs control: t = 7.06, p < 0.001; 500 mg/kg vs control: t = 7.06, p < 0.001; 500 mg/kg vs control: t = 7.06, p < 0.001; 500 mg/kg vs control: t = 7.06, p < 0.001; 500 mg/kg vs control: t = 7.000; 500 mg/kg vs control: t = 7.000 mg/kg vs con 351 0.001; 250 mg/kg vs 500 mg/kg; t = 2.16, p < 0.05) compared to the controls (Fig. 4b). We also found a 352 significant effect of the wire cup factor on the visit number as well as its interaction with the treatment factor 353  $(F_{(2,17)} = 26.25, p < 0.001; F_{(2,17)} = 17.89, p < 0.001,$  respectively). Multiple comparisons confirmed that only 354 mice exposed to 250 mg/kg showed less visit number of wire cup holding another conspecific (t = 4.89,  $p < 10^{-10}$ 355 0.001) while both GBH-exposed groups showed higher visit number of the wire cup holding the object compared 356 to the controls (250 mg/kg vs control: t = 4.39, p < 0.001; 500 mg/kg vs control: t = 6.32, p < 0.001) (Fig. 4c).

### 357 *Passive avoidance*

358 We further examined mice by step-through avoidance learning, an experimental paradigm used for assessing 359 learning, short and long-term memory. The change in the latency to enter into the dark compartment was 360 compared among the three treatment groups and was found to be affected by both treatment and time of test 361  $(F_{(2,17)} = 13.96, p < 0.01; F_{(2,17)} = 31.00, p < 0.01$ , respectively). The analysis revealed also a significant 362 interaction between the two factors ( $F_{(2,17)} = 7.46$ , p < 0.05). Post-hoc analysis indicated that only the group 363 exposed to 500 mg/kg of GBH showed a significant decrease of latency after 2h (500 mg/kg vs control: t = 6.32, 364 p < 0.001) while both treated groups exhibited a significant decrease of latency 24 h of after the electrical shock 365 administration (250 mg/kg vs control: t = 4.39, p < 0.001; 500 mg/kg vs control: t = 6.32, p < 0.001) (Fig. 4d).

- 366 *Y-maze*
- The effect of GBH on working memory was evaluated by Y-maze task. One-way ANOVA analysis showed a significant difference in spontaneous activity between treated and control groups ( $F_{(2,17)} = 11.50$ , p < 0.001). *Post-hoc* comparisons confirmed that the spontaneous alternation in GBH-treated mice was lower than that in
- 370 control mice (250 mg/kg vs control: t = 2.14, p < 0.05; 500 mg/kg vs control: t = 4.78, p < 0.001) and this effect
- 371 was significantly more pronounced in 500 mg/kg than in 250 mg/kg (t = 2.64, p < 0.001) (Fig. 4e).

### 372 Novel object recognition

This test was used to assess the potential effects of GBH on recognition memory. One-way ANOVA analysis showed a significant difference in both the ratio of time spent beside the new object and the discrimination index  $(F_{(2,17)} = 37.70, p < 0.001)$ . Multiple comparisons revealed that GBH-exposed mice spent less time exploring the novel object and had low discrimination index compared to the controls (250 mg/kg *vs* control: t= 5.43, *p* < 0.001; 500 mg/kg *vs* control: t = 8.58, *p* < 0.001). In addition, the same analysis revealed a significant difference between treated groups (t = 3.15, *p* < 0.01) (Fig. 4f-g).

# 379 Biochemical, histological and molecular changes within brain of GBH-exposed mice

# 380 GBH effects on AChE activity

- 381 Because the cholinergic system is closely associated with anxious and cognitive functions (Mineur et al. 2013;
- 382 Coyle et al. 1983), we sought to assess the impact of GBH exposure on AChE in the supernatants of specific
- brain areas homogenates. ANOVA analysis showed significant differences between groups in PFC and whole
- brain ( $F_{(2,2)}$ = 12.78; p < 0.05;  $F_{(2,2)}$  = 13.93; p < 0.05, respectively), while no statistical difference was found in the hippocampus ( $F_{(2,2)}$  = 5.79; p > 0.05) (Fig. 5c). *Post-hoc* comparisons revealed that only the group exposed
- 386 to 500 mg/kg of GBH showed a significant decrease of AChE activity in the whole brain (500 mg/kg vs control:
- t = 5.24, p < 0.05 (Fig. 5a), while the activity of this enzyme was significantly decreased in PFC for both doses
- 388 250 and 500 mg/kg doses (250 mg/kg vs control: t = 4.85, p < 0.05; 500 mg/kg vs control: t = 3. 65, p < 0.05)
- 389 (Fig. 5b).

# 390 GBH effect on dopaminergic system

391 A growing number of data correlate movement control abnormalities with dopaminergic systems dysfunction 392 (Rodríguez et al. 2013; Gallo et al. 2015). To examine the impact of GBH exposure on dopaminergic neurons, 393 we evaluated the expression of tyrosine hydroxylase (TH), a key enzyme involved in dopamine synthesis. 394 Indeed, the quantitative analysis of TH-immunolabeled cells number in the SNc, in the VTA and in the striatum 395 indicated a significant difference between control and treated groups ( $F_{(2,17)} = 17.89, p < 0.001$ ;  $F_{(2,17)} = 14.78, p$ 396 < 0.01; F<sub>(2,17)</sub> = 8.41, p < 0.05, respectively). Post-hoc comparisons confirmed a significant reduction in the 397 number of TH positive cell bodies (TH<sup>+</sup>) in the SNc (250 mg/kg vs control: 3.66, p < 0.05; 500 mg/kg vs control: 398 t = 5. 92, p < 0.01) (Fig. 6b) and in the VTA (250 mg/kg vs control: 4.47, p < 0.01; 500 mg/kg vs control: t = 399 4.91, p < 0.01) of GBH-exposed animals compared to the controls (Fig. 6c). Moreover, the same analysis 400 showed that 500 mg/kg treated group showed a significant decrease in the integral optical density of the TH<sup>+</sup> 401 fibers immunofluorescence in the striatum compared to the control and 250 mg/kg groups (500 mg/kg vs control: 402 t = 4.07, p < 0.01; 250 mg/kg vs 500 mg/kg: t = 2.46, p < 0.05) (Fig. 6d).

# 403 **GBH** causes neuroinflammation

404 Clear evidence showed that organophosphate OPs intoxication is associated with inflammatory responses as 405 revealed the activation of both astrocytes and microglial cells (Banks and Lin 2012). In this perspective, we 406 evaluated neuroinflammation by assessing the expression of both GFAP and Iba-1 proteins in astrocytes and 407 microglial cells, respectively, in different regions of the PFC and the dorsal hippocampus. Interestingly, the 408 quantitative analysis of GFAP and Iba-1 immunofluorescence showed a significant difference among groups. 409 The treated groups showed a significant increase in the number, integral optical density and area occupied by 410 GFAP<sup>+</sup> and Iba-1<sup>+</sup> cells in both PFC and hippocampus (Fig. 7-8 and supplementary table 2-3).

#### 411 *GBH* affects the expression of genes associated with neuroinflammation, synaptic plasticity and cell survival

- 412 Since one major hallmarks of the inflammatory response is the release of cytokines and chemokines from
- 413 activated glial cells, we measured the mRNA expression of tumor necrosis factor alpha (TNFα) by quantitative
- 414 real time PCR (qRT-PCR). The statistical analysis showed significant increased levels of TNFα mRNA
- 415 expression in the hippocampus of exposed mice compared to controls (t = 5.26, p < 0.05) while no statistical
- 416 difference was found in the PFC (t = 0.41, p > 0.05) (Fig. 9a).
- 417 It is well reported that the activation of NMDA glutamate receptors (NMDARs) is crucial for synaptic plasticity
- 418 underlying both short- and long-term memory storage, and that the disruption of these receptors function can
- 419 cause profound cognitive deficits (Lewis 1997; Malenka and Nicoll 1999). Moreover, the interaction between
- 420 proinflammatory cytokines and the glutamatergic neurotransmission has been previously described (Fogal and
- 421 Hewett 2008). Indeed, excessive activation of NMDA receptors leads to elevated calcium influx into cells, which
- 422 perturbs mitochondria and increases generation of reactive oxygen species, inflammation leading ultimately to
- 423 neurodegeneration. To test whether GBH could also impact glutamatergic signaling, we assessed the expression
- 424 level of genes encoding for different NMDA receptors subunits. Our results showed that GBH induced
- 425 significant increase in the mRNA expression of NR1 subunit in the PFC (t = 4.92, p < 0.01) (Fig. 9b) while no
- 426 significant differences could be observed for both NR2A and NR2B subunits in either the PFC or the
- 427 hippocampus (p > 0.05) (Fig. 9c-d).
- 428 The role of various memory associated neurotrophins such as brain derived neurotrophic factor (BDNF), has
- 429 also been suggested in memory dysfunction (Alonso et al. 2005). Based on our data related to cognitive deficits,
- 430 we assessed the impact of GBH exposure on the expression of brain-derived neurotrophic factor (BDNF) and its
- 431 receptor tyrosine regulated kinase B (TrkB) within the PFC and hippocampus. The analysis showed a
- 432 statistically significant decreasing effect of GBH on the expression of BDNF (t = 3.76, p < 0.05) and a
- 433 significant increasing effect on the expression of its receptor TrkB (t = 3.08, p < 0.05) in the cortices of treated
- 434 mice (Fig. 9 e-f). However, no statistical significant differences were found in the hippocampi (t = 1.65, p > 1.65
- 435 0.05; t = 0.05, p > 0.05, respectively) (figure 9 e-f).

#### 436 **Discussion**

- In the present study we characterized the long-term effects of GBH exposure, which may interfere with braindevelopment and lead to permanent abnormalities. Thus, we developed an experimental murine model of chronic
- 439 pre- and postnatal exposure to GBH and examined the downstream effects at the behavioral, histological and
- 440 molecular levels in pups and adult mice.

# 441 *GBH affects reproduction parameters*

442 Our results showed that continuous exposure to GBH during pregnancy and lactation affected fertility and 443 reproduction in mouse females. Furthermore, GBH exposure exerted negative effects also on maternal behavior. 444 Reproduction and fertility toxicity observed in this work are in agreement with the results published in a report 445 of the World Health Organization (WHO, 1994). Indeed, exposure of 3500 mg/kg of Gly in rats from day 6 to 19 446 of pregnancy caused increased maternal mortality, augmented incidence of early resorptions, and decreased 447 number of implantations and viable fetuses (WHO, 1994). Although the precise reprotoxicity mechanism is vet 448 to be clarified, it has been previously reported that Gly targets two crucial steps of steroidogenesis in mammals: 449 at the first rate-limiting level of mitochondrial cholesterol transport (Walsh et al. 2000), and at the last 450 irreversible conversion of sexual androgens into estrogens, via a direct action on the aromatase enzyme 451 (Richard et al. 2005).

# 452 GBH delayed sensorimotor development

453 So far, it has been shown that GBH is able to cross the placental barrier which could possibly alter the 454 developmental process of the fetus (Mose et al. 2008). More recently, the report published by the non-455 governmental american organization "Moms Across America" suggested that detection of relatively high levels 456 of Gly in breast milk in three out of 10 women raised a bioaccumulation problem (Honeycutt and Rowlands 457 2014) and therefore posed potential risks for breastfed offspring. Our results support these hypotheses: early 458 after birth, we found that GBH-exposed offspring displayed delayed developmental reflexes likely related to 459 abnormal maturity of sensorimotor, vestibular and/or proprioceptive functions (Secher et al. 2006; Santillán et al. 460 2010). In the present work, we also highlighted that offspring prenatally exposed to GBH show a significantly 461 shorter latency to fall from the rotarod than controls which could be to alterations in the cerebellar function, 462 since this center is heavily involved in the rotarod performance (Hamm et al. 1994). The reduction in the motor 463 coordination could also be due to a failure in neuromuscular maturity as mentioned by Perez-Reyes et al. (1998). 464 This suggestion is further reinforced by our results obtained through the suspension test showing reduced muscle 465 strength in exposed animals. In sum, the pattern of developmental deficits emerged from this study is in 466 agreement with previous human and animal studies showing similar reflexes defects (Engel et al. 2011; 467 Laugeray et al. 2014; Lan et al. 2017).

# 468 GBH reduces locomotor activity and affects dopaminergic system

469 Normal development of neonatal reflexes can be considered as an index of brain maturation, and late acquisition 470 of these milestones represent a predictive factor of other behavioral changes in adulthood. In line with this idea, 471 our results show that adult progeny from GBH treated groups presented locomotor hypoactivity which is in 472 agreement with Gallegos et al. (2016), showing that rats exposed to 200 mg/kg of GBH during pregnancy and 473 lactation are hypoactive, and with our previous observations in juvenile mice (Ait bali et al. 2017). Albeit 474 locmotor hypoactivity was previously reported following prenatal exposure to GBH (Gallegos et al. (2016), the 475 neuronal basis still unclear. Thus, to generate a mechanistic understanding whereby GBH produces locomotor 476 hypoactivity, we assessed the outcomes of GBH exposure on dopaminergic system. The nigrostriatal pathway, 477 crucial in movement control, has been shown to be vulnerable to herbicides (Thiruchelvam et al. 2000; 478 Rodríguez et al. 2013). Considering the idea that reduction in locomotor activity is positively correlated with loss 479 of dopaminergic neurons in the SNc (Bano et al. 2014, Gallo et al. 2015), one of the most intriguing findings of 480 our work is that GBH-exposed mice show a robust decrease in the number of dopaminergic neurons, both in the 481 SNc and VTA, and fibers, in the striatum. It should be noted that since pesticide is used in commercial 482 formulation which combine an active ingredient with adjuvants, the toxicity exerted by GBH cannot therefore be 483 exclusively due to the active ingredient but either to the toxicity of adjuvants or to the possible synergy between 484 Gly and the other formulation ingredients (El-Shenawy, 2009, Mesnage et al., 2013). The decrease of 485 dopaminergic cells we observed may be due to apoptotic events triggered by elevated oxidative stress observed 486 after Gly exposure (Astiz et al. 2009). Our results are in agreement with Hernandez-Plata et al. (2015) showing 487 that intraperitoneal exposure to 150 mg/kg of Gly for 2 weeks produces a decrease of DA level in the striatum of 488 adult rats that is associated with hypoactivity. Interestingly, human reports support the central effects of GBH on 489 basal ganglia circuits following intoxication with commercial formulations of Gly, described as alterations in the 490 GP and SNc closely related to Parkinsonian syndrome (Barbosa et al. 2001). Finally, although the hypoactivity 491 induced by Gly administration is similar to that observed after administration of a DA antagonist (Hernandez-492 Plata et al. 2015), Gly can affect other neurotransmitter systems involved in motor control (Martinez et al. 2018). 493 Quantitative analysis of serotonin, dopamine and norepinephrine levels confirmed a dramatic loss of these 494 neurotransmitters mainly in the striatum, PFC and hippocampus of rats exposed to 35, 75, 150 or 800 mg/kg of 495 Gly (Martinez et al. 2018), strongly indicating that further studies are needed to decipher the contribution of 496 these systems to GBH-produced hypoactivity.

### 497 GBH increase anxiety level and reduces social behavior

498 Our data clearly showed that early exposure to GBH increases anxiety levels, which is consistent with another 499 study (Baier et al. 2017) in which Gly was administered to adult mice by intranasal irrigation. In contrast, 500 Gallegos et al. (2016) reported that rats exposed to 100 or 200 mg/kg of GBH (Glifloglex®) during pregnancy 501 and lactation experienced low levels of anxiety in adulthood. The difference observed between our results and 502 those of Gallegos et al. (2016) could be explained by the diversity of the commercial formula of Gly 503 administered in the two studies - i.e.: different chemical composition in terms of adjuvants Indeed, it has been 504 pointed out that the herbicidal activity of Gly is potentiated by the presence of adjuvants (Mitchell et al. 1987), 505 thus supporting their potential role in in the differential effects of Gly on anxiety levels.

506 It is well established that neuroinflammation expressed by overproduction of proinflammatory cytokines and 507 excessive activation of glia plays an important role in the etiology of many psychiatric disorders including 508 anxiety (Holloway-Erickson et al. 2012). Accordingly, the neuroinflammatory condition that we report in GBH-509 treated mice, associated with increased anxiety levels, supports this idea. The PFC is strongly involved in the 510 expression of behavioral and autonomic responses to emotionally relevant stimuli, and imaging studies highlight 511 abnormalities in the structure and function of this region in patients with mood disorder (Kennedy et al. 2001; 512 Drevets et al. 2008). We hypothesized that GBH-induced neuroinflammation of this cortical area could lead to 513 cellular disorganization within the PFC since we already observed that GBH exposure induced loss of 514 serotoninergic fibers within this structure in young mice (Ait bali et al. 2017). Nevertheless, the outcomes of 515 neuroinflammation within the PFC was not investigated in the present work and future studies will be necessary

516 to completely understand the mechanism whereby GBH-induced neuroinflammation produces functional 517 alteration of the PFC.

518 Our results show that perinatal exposure to GBH has a strong effect on the social skills of adult mice, as reflected 519 by significant alterations in the three-chamber test, an analysis conventionally used to test sociability also in 520 murine models of autism (Silverman et al. 2010). Despite that mice are a social species, engaging in high levels 521 of social interactions and sexual/parenting behaviors (Arakawa et al. 2008), perinatally GBH-exposed mice are 522 unable to distinguish a social partner from a new object in adulthood, indicating that Gly effects on social 523 behavior are long-lasting. Our results are in agreement with those of Laugeray et al. (2014) who found that 524 exposure to another OP, glufosinate, during pregnancy and suckling periods reduced sociability in mice. There is 525 evidence that the early social environment (e.g.: maternal care and interactions) has a profound impact on the 526 offspring ability to develop normal social skills (Branchi et al. 2013). Our results showing that GBH 527 significantly alters maternal weight gain, nest building capacity, and retrieving behavior suggest that atypical 528 sociability observed in GBN-exposed mice could be, at least in part, due to physiological or behavioral deficits 529 induced in the mothers. However, the cellular origins and circuit mechanisms contributing to these phenotypes 530 have yet to be identified, a future goal that will shed light into the pathogenic mechanisms that underlie social 531 interaction impairment after GBH exposure.

# 532 GBH induces cognitive and cholinergic system impairment and triggers neuroinflammation

533 The current study showed that GBH affects recognition, working and contextual memory. These results are 534 similar to those recently published by our laboratory in juvenile mice exposed to the same doses of GBH (Bali et 535 al. 2019), and support those of Gallegos et al. (2018) revealing a significant impairment of recognition memory 536 in adult rat prenatally exposed to 100 or 200 mg/kg of GBH. These results are in agreement with clinical reports 537 showing that populations accidentally exposed to the Gly have developed short and long-term memory 538 impairments accompanied with hippocampal lesion (Barbosa et al. 2001, Nishiyori et al. 2014). Human and 539 animal studies have implicated the central cholinergic system as an important regulator of cognitive functions 540 such as learning and memory. Indeed, brain cholinergic hypofunction associated with memory deficits was 541 previously observed in patients with Alzheimer's disease (Coyle et al. 1983). Similarly, learning and cognitive 542 defects paralleled with cholinergic alteration were recorded in different animal models (for review, see Voorhees 543 et al. 2017). In agreement with these studies, our results showed that GBH triggers cholinergic crisis expressed 544 by significant decrease of AChE activity in the PFC and the whole brain. Although Gly is a weak inhibitor of the 545 AChE activity (Larsen et al., 2016), the results of the present work corroborate the previous data from our 546 laboratory showing that GBH reduced AchE activity in young mice (Bali et al. 2019). These findings support the 547 previous evidence showing that the commercial formulation is more toxic than Gly (Folmar et al., 1979; Richard 548 et al., 2005). It is well known that in addition to its involvement in adult neurotransmission by its catalytic 549 functions (hydrolysis of acetylcholine), AChE also plays a morphogenic role during the development of the 550 nervous system (Grisaru et al. 1999). Therefore, inhibition of AChE activity observed following GBH exposure 551 could interfere with the morphogenic role of this enzyme and would be the main mechanism inherent to the 552 neurobehavioral alterations induced by these compounds. Furthermore, the cognitive impairments associated 553 with the AChE inhibition following exposure to GBH could be also explained by over-stimulation and persistent 554 activation of both muscarinic and nicotinic receptors due to the accumulation of acetylcholine in the synaptic 555 cleft (Scheffel et al. 2018). This finding might offer a partial explanation for the complaints on memory loss

- reported by workers chronically exposed to OPs compounds even though the neurotoxicity of Gly or GBH may
- 557 not be entirely due to disturbances of the cholinergic system.
- 558 It is recognized that the co-operation of the PFC and the hippocampus is vital for fundamental cognitive 559 functions and that disconnection or damage to either one of these two brain regions induces impaired cognitive 560 behaviors (Yoon et al. 2008). Furthermore, evidence linking inflammation to neurodegeneration and cognitive 561 defects (Dziedzic, 2006) suggests the possibility that induction of inflammation by chronic exposure to OPs may 562 be mechanistically related to deficits in cognitive ability. Based on these, our morphological analysis showed an 563 increase in the GFAP expression in the PFC and hippocampus of adult progeny from GBH treated groups. 564 Likewise, GBH elevated the expression of Iba-1 in PFC and hippocampus indicating that microglia, as the most 565 sensitive immune cells in the brain, was activated after GBH exposure. In addition, our study demonstrated a 566 significant increase in the expression of proinflammatory cytokines TNF $\alpha$  in the PFC and hippocampus after 567 GBH exposure. These results indicate that repeated exposure to GBH can trigger neuroinflammation response in 568 PFC and hippocampus of exposed mice. In agreement with our results, several previous studies elaborated that 569 exposure to OPs induced glial cells activation as well as cytokine and chemokine elevation contributing to 570 neuronal damage in the hippocampus and piriform cortex. Indeed, soman (an OP nerve agent) intoxication 571 upregulates the GFAP expression and activates microglia in many regions of the brain, including the 572 hippocampus (Angoa-Perez et al. 2010). Similarly, chronic administration of the chlorpyrifos increases GFAP 573 expression in rat hippocampus (Lim et al. 2011).

# 574 GBH increase the expression of glutamatergic receptor and affects cell survival pathway

575 There is evidence that neuroinflammation exacerbates neuronal damage due to excitotoxicity via interactions 576 between proinflammatory cytokines and glutamatergic pathways (Fogal and Hewett 2008). Indeed, excessive 577 activation of NMDA receptors leads to elevated calcium influx into cells, which perturbs mitochondria and 578 increases generation of reactive oxygen species ultimately leading to cells death. Our biomolecular analysis 579 corroborates these findings showing that, in addition to neuroinflammation event, GBH increases the transcript 580 level of NR1 subunit of NMDA receptors in the PFC. Then, it would seem that GBH could facilitates and 581 enhances NMDA activation either through: 1) increase of NMDA receptors expression, 2) excessive release of 582 glutamate by activated astrocytes or 3) reduction of glutamate uptake and metabolism within glial cells, 583 associated with an increased release of this neurotransmitter in the synaptic cleft (Cattani et al. 2014). Taken 584 together, our data present additional mechanistic bases to explain excitotoxic condition associated with cognitive 585 impairments observed following GBH exposure.

- 586 In the PFC and hippocampus, signaling through the BDNF pathway plays an important role in the survival, 587 maintenance and growth and promotes neuronal plasticity and neurogenesis, processes that have been described 588 as potential cellular mechanisms for learning and memory (Sakata et al. 2013). Our results showed that mice 589 exposed to GBH display decreased BDNF levels in PFC, which is coupled with impaired cognitive impairments. 590 In the brain, BDNF binds to TrkB receptor causing its phosphorylation and subsequent activation of the 591 intracellular signaling cascade to promote synaptic plasticity (Cunha et al. 2010). Our study demonstrated that 592 the GBH increased TrkB transcript in the PFC. Therefore, the increased TrkB mRNA level we observed in the 593 PFC after exposure to GBH may reflect a compensatory mechanism to balance reduction of BDNF expression.
- 594 In agreement with our results, Jain et al. (2013) demonstrated that chronic exposure to Trizophos, another OP,

- 595 significantly reduced the mRNA expression and protein levels of BDNF in rat hippocampi correlated with 596 learning and memory deficits (Jain et al. 2013).
- 597 In summary, our results show that GBH has pervasive harmful effects when administered during the highly
- 598 sensitive pre- and postnatal periods. Our results indicate that GBH exposure induces multiple behavioral
- box abnormalities involving motor, emotional, social and cognitive functions and targets CNS integrity, affecting
- 600 cholinergic, dopaminergic and glutamatergic systems as well as neuroinflammation and cellular stress induction.
   601 These results strongly shed light on additional (noncholinergic) mechanisms mediating neurological injury
- 602 induced by this OP pesticide. All data presented about the behavioral changes as well as brain abnormalities
- 603 induced by GBH pave the way for further analyses: brain microarray analysis is needed to gain a better
- 604 understanding regarding the molecular and cellular mechanisms involved in the neurodevelopmental effects of
- 605 GBH. Such analysis will, with no doubt, contribute to our knowledge of the constellations of genes involved in
- 606 pathological processes mediated by pre- and postnatal exposure to GBH.

# References

Ait Bali Y, Ba-Mhamed S, Bennis M (2017) Behavioral and immunohistochemical study of the effects of subchronic and chronic exposure to glyphosate in mice. Frontiers in behavioral neuroscience 11: 146

Ait-Bali Y, Ba-M'hamed S, Bennis M (2016) Prenatal Paraquat exposure induces neurobehavioral and cognitive changes in mice offspring. Environmental toxicology and pharmacology 48: 53–62.

Alonso M, Bekinschtein P, Cammarota M, Vianna MRM, Izquierdo I, Medina JH (2005) Endogenous BDNF is required for long-term memory formation in the rat parietal cortex. Learn Mem 12: 504–510

Angoa-Pérez M, Kreipke CW, Thomas DM, Van Shura KE, Lyman M, McDonough JH, Kuhn DM (2010) Soman increases neuronal COX-2 levels: Possible link between seizures and protracted neuronal damage. Neurotoxicology 31 (6): 738–746

Arakawa H, Blanchard DC, Arakawa K, Dunlap C, Blanchard RJ (2008) Scent marking behavior as an odorant communication in mice. Neuroscience and Biobehavioral Reviews 32 (7): 1236–1248

Astiz M, Alaniz MJTd, Marra CA (2009) Effect of pesticides on cell survival in liver and brain rat tissues. Ecotoxicology and Environmental Safety 72 (7): 2025–2032

Baier CJ, Gallegos CE, Raisman-Vozari R, Minetti A (2017) Behavioral impairments following repeated intranasal glyphosate-based herbicide administration in mice. Neurotoxicology and Teratology 64: 63–72

Bali YA, Kaikai NE, Ba-M'hamed S, Bennis M (2019) Learning and memory impairments associated to acetylcholinesterase inhibition and oxidative stress following glyphosate based-herbicide exposure in mice. Toxicology 415: 18-25

Banks CN, Lein PJ (2012) A review of experimental evidence linking neurotoxic organophosphorus compounds and inflammation. Neurotoxicology 33 (3): 575–584

Bano F, Ahmed A, Parveen T, Haider S (2014) Anxiolytic and hyperlocomotive effects of aqueous extract of Nigella sativa L. seeds in rats Pak. J Pharm Sci 27: 1547-1552

Barbosa ER, Leiros da Costa MD, Bacheschi LA, Scaff M, Leite CC (2001) Parkinsonism after glycinederivate exposure. Movement Disorders: Official Journal of the Movement Disorder Society 16: 565–568

Bevins RA, Besheer, J (2006) Object recognition in rats and mice: a one-trialnon-matching-to sample learning task to study recognition memory. Nat Protoc 1: 1306–1311

Branchi I, Curley JP, D'andrea I, Cirulli F, Champagne FA, Alleva E (2013) Early interactions with mother and peers independently build adult social skills and shape BDNF and oxytocin receptor brain levels. Psychoneuroendocrinology 38: 522–532

Cattani D, Cesconetto PA, Tavares MK, Parisotto EB, De Oliveira PA, Rieg CEH, Zamoner, A (2017) Developmental exposure to glyphosate-based herbicide and depressive-like behavior in adult offspring: Implication of glutamate excitotoxicity and oxidative stress. Toxicology 387: 67–80

656 Cattani D, de Liz Oliveira Cavalli VL, Heinz Rieg CE, Domingues JT, Dal-Cim T, Tasca C.I, Zamoner A
657 (2014) Mechanisms underlying the neurotoxicity induced by glyphosate-based herbicide in immature rat
658 hippocampus: Involvement of glutamate excitotoxicity. Toxicology 320: 34–45

660 Chen Z, Huang C, Ding W (2016) Z-Guggulsterone improves the scopolamine-induced memory
661 impairments through enhancement of the BDNF signal in C57BL/6J mice. Neurochemical research 41 (12):
662 3322–3332

664 Cohen H, Matar MA, Joseph Z (2013) Animal Models of Post-traumatic Stress Disorder. Current Protocols
665 in Neuroscience. John Wiley and Sons (chapter 9, unit 9.45)

667 668 669	Conrad A, Schroter-Kermani C, Hoppe HW, Ruther M, Pieper S, Kolossa-Gehring M (2017) Glyphosate in German adults - time trend (2001 to 2015) of human exposure to a widely used herbicide. Int J Hyg Environ Health 220: 8–16
670 671 672	Coyle JT, Price DL, Delong MR (1983) Alzheimer's disease: a disorder of cortical cholinergic innervation. Science 219 (4589): 1184–1190
673 674 675	Cunha. (2010). A simple role for BDNF in learning and memory? Frontiers in Molecular Neuroscience 3: 1
676 677 678	Daruich J, Zirulnik F, Gimenez MS (2001) Effect of the herbicide glyphosate on enzymatic activity in pregnant rats and their fetuses. Environmental research 85(3): 226-231
679 680 681	Drevets, WC (2001) Neuroimaging and neuropathological studies of depression: implications for the cognitive-emotional features of mood disorders. Current Opinions in Neurobiology 11: 240–249
682 683	Dziedzic T (2006) Systemic Inflammatory Markers and Risk of Dementia. American Journal of Alzheimer's Disease and Other Dementias, 21 (4): 258–262
685 686 687	Ellman GL, Courtney KD, Andres V, Featherstone RM (1961) A new and rapid colorimetric determination of acetylcholinesterase activity. Biochem. Pharmacol. 7 (2): 88IN191-9095
688 689	EMA, Makoto, HARA, Hiroaki, Matsumoto, Mariko (2008) Evaluation of developmental neurotoxicity of polysorbate 80 in rats. Reproductive Toxicology 25: 89-99
690 691 692 693	Engel SM, Wetmur J, Chen J, Zhu C, Barr DB, Canfield RL, Wolff MS (2011) Prenatal exposure to organophosphates, paraoxonase 1, and cognitive development in childhood. Environmental health perspectives 119(8): 1182-1188
694 695 696 697 698	EPA (1993). Registration Eligibility Decision (RED) Glyphosate. Office of Prevention, Pesticides and Toxic Substances. EPA-738-F-93-011. Washington, DC. Fogal B, Hewett SJ (2008) Interleukin-1β: a bridge between inflammation and excitotoxicity? Journal of Neurochemistry 106 (1): 1–23
699 700 701 702	Ford B, Bateman LA, Gutierrez-Palominos L, Park R, Nomura DK (2017) Mapping proteome-wide targets of glyphosate in mice. Cell Chem Biol 24: 133–140
702 703 704	Fox WM (1965). Reflex and behavioural development of the mouse. Anim. Behav.13: 234 24
705 706 707	Franz JE, Mao MK, Sikorski JA (1997) Glyphosate: A Unique Global Herbicide (ACS Monograph No. 189). American Chemical Society, Washington, DC
708 709 710 711	Gallegos CE, Baier CJ, Bartos M, Bras C, Domínguez S, Mónaco N, Minetti A (2018) Perinatal Glyphosate- Based Herbicide Exposure in Rats Alters Brain Antioxidant Status, Glutamate and Acetylcholine Metabolism and Affects Recognition Memory. Neurotoxicity Research 34 (3): 363–374
711 712 713 714 715	Gallegos CE, Bartos M, Bras C, Gumilar F, Antonelli MC, Minetti A (2016) Exposure to a glyphosate- based herbicide during pregnancy and lactation induces neurobehavioral alterations in rat offspring. Neurotoxicology 53: 20–28
715 716 717 718 710	Gallo EF, Salling MC, Feng B, Moron JA, Harrison NL, Javitch JA (2015). Upregulation of dopamine D2 receptors in the nucleus accumbens indirect pathway increases locomotion but does not reduce alcohol consumption. Neuropsychopharmacology 40: 1609–1618
719 720 721 722	Grisaru D, Sternfeld M, Eldor A, Glick D, Soreq H (1999) Structural roles of acetylcholinesterase variants in biology and pathology. European Journal of Biochemistry 264 (3): 672–686
723 724	Gross C, Hen R (2004) The developmental origins of anxiety. Nat. Rev. Neurosci 5: 545–552

Hernández-Plata I, Giordano M, Díaz-Muñoz M, Rodríguez VM (2015) The herbicide glyphosate causes
behavioral changes and alterations in dopaminergic markers in male Sprague-Dawley
rat. Neurotoxicology 46: 79–91

- Heyer DB, Meredith RM (2017) Environmental toxicology: Sensitive periods of development and neurodevelopmental disorders. Neurotoxicology 58: 23-41
- Holloway-Erickson CM, McReynolds JR, McIntyre CK (2012) Memory-enhancing intra basolateral amygdala infusions of clenbuterol increase Arc and CaMKIIalpha protein expression in the rostral anterior cingulate cortex. Front Behav Neurosci 6: 17
- Honeycutt Z, Rowlands H (2014) Glyphosate testing report: Findings in American mothers' breast milk, urine and water. Unpublished report, dated 7 April 2014, available from the websites of —Moms Across Americal and —Sustainable Pulse
  - Hughes, RN (2004) The value of spontaneous alternation behavior (SAB) as a test of retention in pharmacological investigations of memory. Neuroscience and Biobehavioral Reviews 28 (5): 497–505
  - Iezhitsa IN, Spasov AA, Bugaeva LI (2001) Effects of bromantan on offspring maturation and development of reflexes. Neurotoxicology and Teratology 23 (2): 213–222
  - Jain S, Banerjee BD, Ahmed RS, Arora VK, Mediratta PK (2013) Possible role of oxidative stress and brain derived neurotrophic factor in triazophos induced cognitive impairment in rats. Neurochemical research 38(10): 2136–2147
  - Jiraungkoorskul W, Upatham ES, Kruatrachue M, Sahaphong S, Vichasri-Grams S, Pokethitiyook P (2003) Biochemical and histopathological effects of glyphosate herbicide on Nile tilapia (Oreochromis niloticus). Environmental Toxicology: An International Journal 18(4): 260–267
    - Kennedy SH, Evans KR, Kruger S, Mayberg HS, Meyer JH, McCann S (2001) Changes in regional brain glucose metabolism measured with positron emission tomography after paroxetine treatment of major depression. American Journal of Psychiatry 158: 899–905
    - Kongtip P, Nankongnab N, Phupancharoensuk R, Palarach C, Sujirarat D, Sangprasert S, Woskie SR (2017) Glyphosate and Paraquat in Maternal and Fetal Serums in Thai Women. Journal of Agromedicine 22: 282–
    - Lan A, Kalimian M, Amram B, Kofman O (2017) Prenatal chlorpyrifos leads to autism-like deficits in C57Bl6/J mice. Environmental Health 16(1): 43
    - Laugeray A, Herzine A, Perche O, Hébert, B, Aguillon-Naury M, Richard O, Mortaud S. (2014) Pre- and Postnatal Exposure to Low Dose Glufosinate Ammonium Induces Autism-Like Phenotypes in Mice. Frontiers in Behavioral Neuroscience 8: 1–14
    - Lewis DA (1997) Development of the prefrontal cortex during adolescence: insights into vulnerable neural circuits in schizophrenia. Neuropsychopharmacology 16: 385–398
    - Lim KL, Tay A, Nadarajah VD, Mitra NK (2011) The effect of consequent exposure of stress and dermal application of low doses of chlorpyrifos on the expression of glial fibrillary acidic protein in the hippocampus of adult mice. Journal of Occupational Medicine and Toxicology 6 (1): 4
  - Malenka RC, Nicoll RA (1999) Long-term potentiationa decade of progress? Science 285: 1870–1874
  - Martínez MA, Ares I, Rodríguez JL, Martínez M, Martínez-Larrañaga MR, Anadón A (2018) Neurotransmitter changes in rat brain regions following glyphosate exposure. Environmental Research 161: 212–219
- Meyer OA, Tilson HA, Byrd WC, Riley MT (1979) A method for the routine assessment of foreand hindlimb grip strength of rats and mice. Neurobehav Toxicol 1(Fall (3)): 233–236

785	Mineur VS, Obayemi A, Wigestrand MB, Fote GM, Calarco CA, Li AM, Picciotto MB (2013) Cholinergic
786	signaling in the hippocampus regulates social stress resilience and anxiety- and depression-like behavior.
787	Proc Natl Acad Sci 110: 3573–3578
788	
789	Mitchell DG, Chapman PM, Long TJ (1987) Acute toxicity of Roundup and Rodeo herbicides to rainbow
790	trout, chinook, and coho salmon. Bulletin of Environmental Contamination and Toxicology 39 (6): 1028-
791	1035
792	
793	Mose T, Kjaerstad MB, Mathiesen L, Nielsen JB, Edelfors S, Knudsen LE (2008) Placental Passage of
794	Benzoic acid, Caffeine, and Glyphosate in an Ex Vivo Human Perfusion System. Journal of Toxicology and
795	Environmental Health, Part A 71 (15): 984–991
796	
/9/	National Research Council (2000) Scientific Frontiers in Developmental Toxicology and Risk Assessment.
/98	Washington, DC: National Academy Press
799	Nichiyari V. Nichida M. Shiada K. Suda S. Kata S. (2014) Unilateral hippogeneral inferation according to with
800	an attempted suicide: A case report Journal of Medical Case Penerts 8 (1): 25
802	an attempted sucide. A case report. Journal of Medical Case Reports 8 (1). 25
803	Olney IW (2002) New insights and new issues in developmental neurotoxicology. Neurotoxicology 23:
804	659–668
805	
806	Paxinos G. Franklin KBJ (2001) The Mouse Brain in Stereotaxic Coordinates, 2ndedition, Academic Press,
807	San Diego, California, USA, pp. 216
808	
809	Pellow S, Chopin P, File SE, Briley M (1985) Validation of open: closed arm entries in an elevated plus-
810	maze as a measure of anxiety in the rat. Journal of neuroscience methods 14(3): 149-167
811	
812	Perez-Reyes E, Cribbs LL, Daud A, Lacerda AE, Barclay J, Williamson MP, Lee JH (1998) Molecular
813	characterization of a neuronal low-voltage-activated T-type calcium channel. Nature 391 (6670): 896–900
814	
815	Pettersson M, Ekelund NG (2006) Effects of the herbicides Roundup and Avans on Euglena
810 917	gracilis. Archives of environmental contamination and toxicology 50(2): 1/5-181
818	Powles SP. Preston C. Pryon IP. Jutsum AP (1007) Herbigide registence: impact and management. Adv.
810	Agron 58: 57, 93
820	Agion 58. 57-75
821	Ramirez RL Spear LP (2010) Ontogeny of ethanol-induced motor impairment following acute ethanol-
822	Assessment via the negative geotaxis reflex in adolescent and adult rats. Pharmacology Biochemistry and
823	Behavior 95(2): 242–248
824	
825	Richard S, Moslemi S, Sipahutar H, Benachour N, Seralini GE (2005) Differential effects of glyphosate and
826	roundup on human placental cells and aromatase. Environmental health perspectives 113(6): 716-720
827	
828	Rio DC, Ares M, Hannon GJ, Nilsen TW (2010) Purification of RNA using TRIzol (TRI reagent). Cold
829	Spring Harbor Protocols (6): pdb-prot5439
830	
831	Rodier PM (1995) Developing brain as a target of toxicity. Environmental Health Perspectives 103 (6): 73–
0 <i>32</i> 022	/6
837	Podríguaz VM Limán Dochago III Mandoza Traio MS Conzélaz Callardo A Harnándaz Plata L
835	Giordano M (2013) Repeated exposure to the herbicide atrazine alters locomotor activity and the
836	nigrostriatal donaminergic system of the albino rat Neurotoxicology 34: 82–94
837	ingrosultuur dopuninergie system of the dromo fut. Predrotoxicology 54: 02-94
838	Sakata K. Martinowich K. Woo NH. Schloesser RJ. Jimenez DV. Ji Y. Lu B (2013) Role of activity-
839	dependent BDNF expression in hippocampal-prefrontal cortical regulation of behavioral
840	perseverance. Proceedings of the National Academy of Sciences 110(37): 15103–15108
841	
842	Santillán ME, Vincenti LM, Martini AC, Fiol de Cuneo M, Ruiz RD, Mangeaud A, Stutz G (2010)
843	Developmental and neurobehavioral effects of perinatal exposure to diets with different $\omega$ -6: $\omega$ -3 ratios in
844	mice. Nutrition 26(4): 423–431.

Scheffel C, Niessen KV, Rappenglück S, Wanner KT, Thiermann H, Worek F, Seeger T (2018)
Counteracting desensitization of human α7-nicotinic acetylcholine receptors with bispyridinium compounds as an approach against organophosphorus poisoning. Toxicol Lett 293: 149–150

Secher T, Novitskaia V, Berezin V, Bock E, Glenthoj B, Klementiev BA (2006) Neural cell adhesion
 molecule-derived fibroblast Growth factor receptor agonist, the FGL peptide, promotes early postnatal
 sensorymotor development and enhances social memory retention. Neurosciences 141: 1289–1299

- Seneff S, Swanson N, Li C (2015) Aluminum and glyphosate can synergistically induce pineal gland pathology: connection to gut dysbiosis and neurological disease. J Agric Sci 6: 42–701
- Silverman JL, Yang M, Lord C, Crawley JN (2010) Behavioural phenotyping assays for mouse models of
  autism. Nature Reviews Neuroscience 11 (7): 490–502
  - Solomon KR (2016) Glyphosate in the general population and in applicators: a critical review of studies on exposures. Critical reviews in toxicology 46: 21–27.
- Solomon KR, Anadón A, Carrasquilla G, Cerdeira AL, Marshall J, Sanin LH (2007) Coca and poppy
  eradication in Colombia: environmental and human health assessment of aerially applied glyphosate. Rev
  Environ Contam Toxicol 190: 43–125
- Thiruchelvam, M, Richfield EK, Baggs RB, Tank AW, Cory-Slechta DA (2000) The nigrostriatal dopaminergic system as a preferential target of repeated exposures to combined paraquat and maneb:
  implications for Parkinson's disease. Journal of Neuroscience 20 (24): 9207–9214
  - Voorhees JR, Rohlman DS, Lein PJ, Pieper AA (2017) Neurotoxicity in preclinical models of occupational exposure to organophosphorus compounds. Frontiers in Neuroscience 10: 590
    - Walsh RN, Cummins RA (1976) The open-field test: a critical review. Psychological bulletin 83 (3): 482

Walsh LP, McCormick C, Martin C, Stocco DM (2000) Roundup inhibits steroidogenesis by disrupting steroidogenic acute regulatory (StAR) protein expression. Environmental health perspectives 108(8): 769-776

- Wang G, Fan XN, Tan YY, Cheng Q, Chen SDi (2011) Parkinsonism after chronic occupational exposure to glyphosate. Parkinsonism and Related Disorders 17: 486–487
- World Health Organisation (WHO), United Nations Environment Programme, the International Labour
  Organisation. (1994). Glyphosate. Environmental Health Criteria #159. Geneva, Switzerland: World Health
  Organization
- Yoon T, Okada J, Jung MW, Kim JJ (2008) Prefrontal cortex and hippocampus subserve different components of working memory in rats. Learn Mem 15: 97–105

# 890 Figures

891 892



893

894 Fig. 2 Pre-and postnatal GBH exposure resulted in neurodevelopmental endpoints changes. a Righting reflex test. b Negative 895 geotaxis test. c Cliff avoidance test. d Traction test. e Rotarod test. Results are presented as mean  $\pm$  SEM. \*p < 0.05; \*\*p <

896 0. 01; \*\*\*p < 0.001. The "\*" refers to 250 mg/kg or 500 mg/kg vs control groups comparison



897 898

**Fig. 3** Pre- and postnatal GBH exposure resulted in behavioral alterations in the offspring during adulthood. **a** Recording of the trajectory of in the OF test. **b-d** Effect of GBH on locomotor activity and anxiety-like phenotype in the open field test. **e** Recording of the trajectory of mice in the EPM test. **f-h** Effect of GBH on anxiety-like phenotype in EPM test. Results are presented as mean  $\pm$  SEM. \*p < 0.05; \*\*\*p < 0.001; #p < 0.05; ##p < 0.01. The "\*" refers to 250 mg/kg or 500 mg/kg vs control groups comparison and the "#" refers to the 250 mg/kg vs 500 mg/kg groups comparison. **OA** open arm; **CA** closed arm



**Fig. 4** Pre- and postnatal GBH exposure resulted in sociability and cognitive alterations in the offspring during adulthood. **a** Recording of the trajectory in the TCS test. **b**-**c** Effect of GBH on social interaction in the TCS test. **d** Effect of GBH on short and long-term memory in the PA test. **e** Effect of GBH on working memory in the Y-maze test. **f**-**g** Effect of GBH exposures on recognition memory in the NOR test. Results are presented as mean  $\pm$  SEM. \*p < 0.05; \*\*p < 0.01; \*\*\*p < 0.001; \*\*p <





Fig. 5 Pre- and postnatal GBH exposure resulted in AChE inhibition. **a** whole brain. **b** PFC. **c** hippocampus. Results are presented as mean  $\pm$  SEM. \*p < 0.05. The "\*" refers to 250 mg/kg or 500 mg/kg vs control group comparison



914 915

Fig. 6 Pre- and postnatal GBH exposure resulted in dopaminergic circuit defects. a Photomicrographs of mice brain cross 916 sections showing the tyrosine hydroxylase (TH)-immunoreactive neurons. b Count of TH positive cells in the SNc and c in 917 the VTA. **d** The density of TH Immunoreactivity in the striatum. Results are presented as mean  $\pm$  SEM. \*p < 0.05; \*\*p < 0.05; \*p < 0.05; \*p918 0.01;  ${}^{\#}p < 0.05$ . The "\*" refers to 250 mg/kg or 500 mg/kg vs control groups comparison and the "#" refers to the 250 mg/kg 919 vs 500 mg/kg groups comparison. SNc substantia nigra pars compacta; VTA ventral tegmental area



920 921 922 Fig. 7 Pre- and postnatal GBH exposure resulted in reactive astrocytes. a and k Micrographs showing the expression of GFAP by immunofluorescence in the PFC and hippocampus. b-d Count of GFAP positive cells in the PFC and l-n in the 923 hippocampus. e-g The integral optical density of GFAP positive cells in the PFC and o-q in the hippocampus. h-j Area 924 occupied by GFAP positive cells in the PFC and r-t in the hippocampus. Results are presented as mean  $\pm$  SEM. \*p < 0.05; 925 \*\*p < 0.01; \*\*\*p < 0.001; ##p < 0.01. The "\*" refers to 250 mg/kg or 500 mg/kg vs control groups comparison and the "#" 926 refers to the 250 mg/kg vs 500 mg/kg groups comparison. ACCx anterior cingulate cortex; PrLCx prelimbic cortex; ILCx 927 infralimbic cortex; DG dentate gyrus; CA1 Ammon's horn 1; CA3 Ammon's horn 3



928 929 930 Fig. 8 Pre- and postnatal GBH exposure resulted in reactive microglia. a and k Micrographs showing the expression of Iba-1 by immunofluorescence in the PFC and hippocampus. b-d Count of Iba-1 positive cells in the PFC and I-n in the 931 hippocampus. e-g The integral optical density of Iba-1 positive cells in the PFC and o-q in the hippocampus. h-j Area 932 occupied by Iba-1 positive cells in the PFC and r-t in the hippocampus. Results are presented as mean  $\pm$  SEM. \*p < 0.05; \*\*p933 < 0.01; \*\*\*p < 0.001; p < 0.05; p < 0.05; p < 0.01. The "\*" refers to 250 mg/kg or 500 mg/kg vs control groups comparison and the 934 "#" refers to the 250 mg/kg vs 500 mg/kg groups comparison. ACCx anterior cingulate cortex; PrLCx prelimbic cortex; 935 ILCx infralimbic cortex; DG dentate gyrus; CA1 Ammon's horn 1; CA3 Ammon's horn 3



936PFCHippocampusPFCHippocampusPFCHippocampus937Fig. 9 Pre- and postnatal GBH exposure resulted in genes expression in the PFC and hippocampus of adult progeny. a qRT-938PCR analysis of TNFα transcript. b-d qRT-PCR analysis of NMDA receptor subunits transcripts. e qRT-PCR analysis of939BDNF and f its receptor TrkB transcripts. For each tissue (PFC, Hippocampus), the gene expression is shown relatively to its940control samples. Results are presented as mean ± SEM. \*p < 0.05; \*\*p < 0.01. The "\*" refers to 250 mg/kg or 500 mg/kg vs</td>941control groups comparison

# 944 Supplementary data

	11	•	
945	Table 1	rimers for	quantitative real time PCR analysis

Gene	N° Genbank	Forward primer (5'-3')	Reverse primer (5'-3')	(Size)
BDNF	NM_001285416.1	GCGTGTGTGACAGTATTAGCGAGTG	CAGTTGGCCTTTGGATACCGGG	116
TrkB	NM_001282961.1	GAAAAACAGCAACCTGCGGCAC	GAACGGATTACCCGTCAGGATCAGG	115
NR1	NM_001177657.2	CGTCCTGGGGCTGACTACCC	GCTGGACTGGTGGGAGTAGGG	97
NR2A	NM_008170.2	GACCCACTGACTGAGACCTGCG	CCCCTTGCAGCACTTCTTCACATTC	108
NR2B	NM_008171.3	GAACAAGGAGAGGAAGTGGGAGAGG	CAGTCTCAGGACACATTCGAGGCC	95
ΤΝΓα	NM_013693.3	CTCAGCCTCTTCTCATTCCTGCTTG	GGCCATTTGGGAACTTCTCATCCC	106
TBP	NM_013684.3	GATCAAACCCAGAATTGTTCTCC	GGGGTAGATGTTTTCAAATGCTTC	106

# **Table 2** Statistical analysis of GFAP data

		A NI				Pe	ost-hoc		
		AIN	JVA	250 1	vs control	<b>500</b> 1	vs control	250 v	vs 500
		<b>F</b> <sub>(2, 2)</sub>	р	t	р	t	р	t	р
	Number of GFAP <sup>+</sup> cells								
	ACCx	18.66	**	4.00	**	6.00	***	2.00	Ns
	PrLCx	7.58	*	3.86	**	2.37	Ns	1.48	Ns
	ILCx	22.58	**	3.18	*	6.71	***	3.53	*
Ų	Integral optical density								
PF	ACCx	7.39	*	2.27	Ns	3.82	**	1.55	Ns
	PrLCx	0.78	Ns						
	ILCx	6.27	*	2.36	Ns	3.44	*	1.08	Ns
	Occupied surface								
	ACCx	5.58	*	2.63	*	3.09	*	0.46	Ns
	PrLCx	4.98	Ns						
	ILCx	9.38	*	4.02	**	3.39	*	0.63	Ns
	Number of GFAP <sup>+</sup> cells								
	DG	7.38	*	3.16	*	3.47	*	0.30	Ns
	CA1	41.56	***	6.86	***	8.62	***	1.76	Ns
SI	CA3	72.11	***	7.98	***	11.7	***	3.78	**
ā						6			
an	Integral optical density	10.04		• • • •				4 60	
ÿ	DG	10.96	*	3.01	*	4.61	**	1.60	Ns
ğ	CAI	2.81	Ns						
	CA3	8.76	*	2.37	Ns	4.17	**	1.79	Ns
	Occupied surface								
	DG	5.27	*	2.76	*	2.86	*	0.10	Ns
	CA1	1.44	Ns						
	CA3	4.04	Ns						

949 \*p < 0.05; \*\*p < 0.01; \*\*\*p < 0.001; ns: no significant (p > 0.05).

# **Table 3** Statistical analysis of Iba-1 data

			7	Post hoc					
		ANU	AIOVA		250 vs control		control	250 vs 500	
		<b>F</b> <sub>(2, 2)</sub>	р	t	р	t	р	t	р
	Number of Iba-1 <sup>+</sup> cells								
Ç	ACCx	6.24	*	1.96	Ns	3.52	*	1.56	Ns
	PrLCx	7.55	*	3.10	*	3.58	*	0.47	Ns
	ILCx	20.57	**	3.09	*	6.44	***	3.34	*
	Integral optical density								
H	ACCx	8.41	*	2.58	*	4.05	**	1.47	Ns
	PrLCx	2.28	Ns						
	ILCx	9.67	*	2.11	Ns	4.39	**	2.50	Ns
	Occupied surface								
	ACCx	5.76	*	2.96	*	2.91	*	0.05	Ns
	PrLCx	1.10	Ns						
	ILCx	5.91	*	2.54	Ns	3.20	*	0.93	Ns
	Number of Iba-1 <sup>+</sup> cells								
	DG	13.52	**	4.36	**	4.63	**	0.26	Ns
	CA1	5.89	*	1.83	Ns	3.42	*	1.59	Ns
sn	CA3	2.55	Ns						
du	Integral optical density								
an	DG	6.19	*	2.33	Ns	3.44	*	1.10	Ns
ĕ	CA1	6.70	Ns	2.97	*	3.33	*	0.36	Ns
dd	CA3	5.43	*	2.13	Ns	3.24	*	1.11	Ns
H	Occupied surface								
—	DG	8.80	*	1.93	Ns	4.19	**	2.25	Ns
	CA1	2.33	Ns						
	CA3	15.65	**	1.51	Ns	5.42	Ns	3.90	**

p < 0.05; \*\*p < 0.01; \*\*\*p < 0.001; ns: no significant (p > 0.05).