

AperTO - Archivio Istituzionale Open Access dell'Università di Torino

Effects of 3-hydroxy-3-methylglutaryl coenzyme A reductase inhibitors on ageing: Molecular mechanisms

This is the author's manuscript

Original Citation:

Availability:

This version is available <http://hdl.handle.net/2318/1739573> since 2020-05-21T15:37:04Z

Published version:

DOI:10.1016/j.arr.2020.101024

Terms of use:

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)

Effects of 3-hydroxy-3-methylglutaryl coenzyme A reductase inhibitors on ageing: molecular mechanisms

Abstract

Human ageing is determined by degenerative alterations and processes with different manifestations such as gradual organ dysfunction, tissue function loss, increased population of aged (senescent) cells, incapability of maintaining homeostasis, reduced repair capacity, which collectively lead to an increased risk of diseases and death. The inhibitors of HMG-CoA reductase statins are the most widely used lipid-lowering agents which can reduce cardiovascular morbidity and mortality. Accumulating evidence has documented several pleiotropic effects of statins in addition to their lipid-lowering properties. Recently, several studies have highlighted that statins may have the potential to delay the ageing process and inhibit the onset of senescence. In this review, we focused on the statin anti-ageing mechanisms and effects on cardiovascular and non-cardiovascular diseases.

Keywords: RhoA; klotho; Sirtuin-1; senescence-associated secretory phenotype; telomerase

1. Introduction

Human ageing, the process of becoming older after reaching sexual maturity, is due to a complex interplay of degenerative alterations and processes, with different manifestations such as gradual organ dysfunction or tissue function loss, increment of the aged (senescent) cell population, incapability of maintaining homeostasis, reduced repair potency, and increased risk of pathological changes, disease and mortality (Alichniewicz et al., 2012; Marchand et al., 2011).

Inhibitors of 3-hydroxy-3-methylglutaryl coenzyme A (HMG-CoA) reductase known as statins are the most widely used lipid-lowering agents that can reduce cardiovascular morbidity and mortality (Grundy, 1988; Werner et al., 2002). In addition to lipid-reduction properties, statins have shown pleiotropic beneficial properties, such as anti-inflammatory, anti-oxidant benefits, antiproliferative and many other cholesterol-independent effects, thus probably regulating many critical processes involved in vascular biology (Bahrami et al., 2018; Ferretti et al., 2015; Marrone et al., 2013; Owens, 2012; Sahebkar et al., 2015; Serban et al., 2015). Recently, several studies have suggested that statins may have the potential of delaying ageing and inhibiting the onset of senescence. In this review, we have focused on the anti-ageing mechanisms and effects of statins in cardiovascular and non-cardiovascular diseases.

2. Cellular ageing or senescence

Ageing is defined as a reduction in the capability of maintaining homeostasis in several tissues.

In biology, senescence refers to deteriorative processes and irreversible growth arrest that follow development and maturation. Cellular senescence known also as cellular ageing is the hallmark of ageing, i.e. the reduction of the regenerative capacity of cells. In this regard,

cellular senescence seems to be detrimental because it consists in a deficit of tissue renewal and functionality (Campisi and di Fagagna, 2007).

Genetic background of ageing

The main evolutionary conservative pathways known to influence organism longevity are the insulin/insulin-like growth factor (IGF) signaling (IIS) (Kenyon et al., 1993; Tatar et al., 2001), the serine/threonine kinase mechanistic target of rapamycin complex (mTORC) (Kaeberlein et al., 2005; Powers et al., 2006), and the protein kinase A (PKA) pathway (Enns et al., 2009). IIS is a master inducer of cell proliferation and an inhibitor of apoptosis. *In-vivo* experiments have revealed that mutations that interfere with IGF-I/IIS activity are associated with an extended lifespan through several mechanisms such as decrement of insulin concentrations, increased insulin sensitivity, change in carbohydrate/lipid metabolism, reduction in oxidative stress (OS) and in the production of Reactive Oxygen Species (ROS), enhanced resistance to stress, slowdown of the age-associated onset of diseases (Bartke, 2005). Centenarian females when compared to controls showed overrepresentation of heterozygous mutations in the IGF1 receptor (IGF1R) gene, which is related to high IGF1 serum concentrations and lower IGFIR activity (Suh et al., 2008). Therefore, genetic variations in the human IGF1R that lead to alteration of IIS are associated with an increase in human longevity, supporting a role of this axis in the determination of lifespan extension (Suh et al., 2008). Moreover, in laboratory animals, mutations of growth hormone (GH)/GH receptors or GH resistance cause secondary inhibition of circulating IGF1 and insulin levels, a strong extension of lifespan and multiple manifestations of delayed/slower ageing (Bartke, 2011). Possible explanations for this mechanism could be elevated stress resistance and change in insulin/mTOR signaling and metabolic regulations (Bartke, 2011). A major intracellular target of IIS is mTORC, which is activated in response to abundant nutrient

supplies and growth factor signals. It has been shown that suppression of the mTOR pathway through genetic interventions or pharmacological inhibitors is related to longevity among invertebrates and human species (Harrison et al., 2009). PKA is a tetramer, composed of two regulatory (R) and two catalytic (C) subunits, which exist in different isoforms. Functional activation of regulatory subunits, RI and RII, takes place in response to cAMP. cAMP binds to the R subunits, releasing the C subunits that are free to connect to downstream molecules and phosphorylate them (Niswender et al., 2002). Mutant mice for PKA regulatory subunit II β (PRKARII β) showed extended lifespan and were resistant to age-associated diseases (Enns et al., 2009). Forkhead box O (FOXO) proteins are a group of conserved transcription factors (TFs) that are essential elements upstream of multiple critical cellular pathways (Carter and Brunet, 2007). FOXO transcriptional targets are implicated in the regulation of cell cycle, cell death, oxidative stress resistance, hematopoiesis, metabolism, cell proliferation, life span and tumor suppression (Accili and Arden, 2004). FOXOs are downstream of IIS, while IIS lowers function/expression of FOXOs via activation of phosphoinositide-3-kinase (PI3K)/AKT (PKB). FOXO3 may be crucial in ageing and increasing lifespan since polymorphisms in this gene are closely associated with longevity in numerous human cohorts (Morris et al., 2015).

Inflammatory pathways in ageing

Continuous low-grade inflammatory processes known as “Inflammageing” are a main feature of ageing and possibly causes to some age-related metabolite, genes and pathways. Mild inflammation is closely related to multiple ageing phenotypes such as alterations in body composition, energy balance, metabolic homeostasis, stress tolerance, immune senescence, and neuronal well-being (Biagi et al., 2010; Franceschi and Campisi, 2014). So, targeting inflammation is one of the promising anti-ageing approaches in the near future. Local

expression of inflammatory cytokines e.g. tumor necrosis factor-alpha (TNF- α), interleukin (IL) 1 and IL-6 could activate both local and systemic inflammation, which are known risk factors for age-dependent diseases. Actually, chronic over-expression of pro-inflammatory mediators underlying ageing due to the age-related redox imbalance activates multiple pro-inflammatory pathways, such as NF- κ B signaling (Chung et al., 2009). In addition, the induction of NF- κ B is a main activator of gene expression of inflammatory proteins, i.e. cytokines, chemokines, growth factors, interleukins and adhesion molecules, all contributing to the spread of the systemic inflammatory process (Makarov, 2000).

Cardiac and vascular ageing

Cardiovascular ageing consists in functional and structural modifications of the cardiovascular system leading to cardiac hypertrophy, myocardial fibrosis, decreased ventricular compliance, elevated vascular stiffness and risk of diastolic heart failure (Fowler et al., 2007; Stepan et al., 2012). Among several factors, peroxisome proliferator-activated receptors (PPARs), a class of **TFs** belonging to the nuclear hormone receptor family, have been recognized as relevant in the heart ageing process (Guellich et al., 2007; Poynter and Daynes, 1998).

Vascular ageing is closely related to the senescence of vascular cells (Minamino et al., 2004), that is the senescence of endothelial cells (ECs) and vascular smooth muscle cells, which are implied in the pathology of vascular dysfunction and atherogenesis (Minamino and Komuro, 2007; Orlandi et al., 2006). Nitric oxide (NO) also plays a key role in the control of vascular cell senescence (Förstermann and Sessa, 2011; Minamino and Komuro, 2007). NO is synthesized from L-arginine by NO synthases (NOSs), which exist in different isoforms: endothelial NOS (eNOS), inducible NOS (iNOS), and neuronal NOS (nNOS) (Förstermann and Sessa, 2011). Remarkable signs of vascular ageing are down-regulation of eNOS,

interruption of eNOS activation, enhanced oxidative stress-induced NO inactivation and finally low NO bioavailability. The integrity and functionality of the endothelial monolayer is fundamental shield from the onset of atherosclerosis (Lusis, 2000). Endothelial progenitor cells (EPCs) might be involved in endothelial integrity maintenance, substitution of apoptotic ECs and neovascularization of ischemic tissue (Asahara et al., 1997; Kawamoto et al., 2001). Furthermore, administration of EPCs in patients with ischemic heart disease promotes neovascularization, and, subsequently, improvement of cardiac function and reduction of myocardial ischemia (Kawamoto et al., 2001).

Telomeres and ageing

Mammalian telomeres are maintained through telomerase that adds polynucleotides repeats at the ends of eukaryotic linear chromosomal DNA (de Lange, 2005). Telomeres are made up of several thousand non-coding, short double-stranded tandem repeats of a guanidine-rich DNA sequence (TTAGGG). The telomere ends in a single-stranded DNA (ssDNA) sequence, known as the G-rich strand overhang, added by telomerase and coated by capping proteins extended over 2–15 kb. The G-overhang folds back onto itself and invade the telomeric ends, making the two higher terminal loop structures, large duplex lariat structure (T-loop) and forming a single-stranded displacement loop (D-loop) (Doksani et al., 2013). Although their function is not fully understood, they possibly contribute to successful DNA replication and chromosomal stability and facilitate the creation of a lariat-like structure to protect the exposed ends of DNA from the genome damage such as degradation, telomere uncapping, unwanted repair, end-to-end fusion, and inappropriate recombination (de Lange, 2005; Greider, 1991). Shelterin or telosome complex, a very dynamic structure, is characterized by six essential proteins including TRF1 (telomeric repeat binding factor 1), TRF2 (TRF1-interaction nuclear factor 2), POT1 (protection of telomeres 1), TIN2 (TRF1 and TRF2

interacting nuclear protein 2 (TIN2), TPP1 (Tripeptidyl-peptidase 1), Rap1 (repressor/activator protein 1), and constitutes a shield cap at the 3'-end of chromosome. TRF1 and TRF2 interact with the double-stranded hexanucleotide telomeric repeats (TTAGGG) and binds to TIN2 and RAP1, respectively. POT1 binds specifically to ssDNA telomeric portion and links to TRF1 and TRF2 via a binding partner, TPP1, forming a complex with TIN2 (Maciejowski and de Lange, 2017).

Telomerase, a ribonucleoprotein (RNP) complex, is a family member of the reverse transcriptases (RTs) and is composed of a telomerase reverse transcriptase (hTERT) protein, a template-containing RNA component (TERC) and dyskerin (Jiang et al., 2018; Musgrove et al., 2018). Since DNA polymerase is not able to complete the replication of the 3'-end of eukaryotic DNA, telomeres become shorter at each round of chromosomal division (Verdun and Karlseder, 2007). Telomerase complex synthesizes and appends multi tandem repeats DNA sequences at the 3'-chromosomal ends (5'→3') throughout successive cell replication cycles and is responsible for compensating the attrition of the DNA-ends and maintaining telomere length (TL) (Blackburn, 1991; Klegarth and Eisenberg, 2018). Telomerase is not adequate to counteract telomere erosion determined by cell division, therefore telomeres shorten with age (Martínez and Blasco, 2017). The first report of this phenomenon was proposed by Leonard Hayflick and co-researchers who found that isolated human embryo-derived fibroblasts have limited proliferative potential with every subsequent cell division (Hayflick, 1965). This biological clock (also known as Hayflick limit) that is the progressive loss of telomeres after each mitosis, is at present defined as end-replication problem or replicative senescence (Olovnikov, 1973). TL shortening determines a continual DNA damage reaction, which results in replicative cellular senescence and eventually massive cell death (Saliques et al., 2010). In almost all cells, senescence and consequent cell death takes place when the mean TL reaches a critical threshold (Allsopp and Harley, 1995). Therefore,

mean TL provides a potential indicator of cellular biological age, where shorter TL represents increased biological age. However, telomeres can remain genetically stable and intact if the telomere renewal machinery, telomerase, remains entirely functional. Dysfunctional telomere leads to genome fragmentation as a result of the removal of “shelterin protection”(Sfeir and De Lange, 2012). Partial or full loss of telomerase associates with inability of tissues to regenerate and, consequently progressive tissue damage and ultimately reduction of life span (García-Cao et al., 2006; Mitchell et al., 1999).

Cellular senescence is an irreversible growth arrest by which cells cease to replicate; it takes place in somatic cells and restricts their proliferative life span. Cellular senescence is conventionally divided into the following two major forms: intrinsic telomere-dependent (replicative senescence) and extrinsic telomere-independent known as stress-induced premature senescence (SIPS) (Itahana et al., 2004; Wlaschek et al., 2003). When telomeres shorten, the cell goes into replicative senescence, which determines enormous alterations in the cell-cycle gene expression profile resulting in reduced proliferation and ultimately apoptosis (Satyanarayana et al., 2003). SIPS is induced via a variety of external stresses such as OS, oncogenic Ras activation, DNA damage/mutation, mitochondrial injury and chemotherapeutic regimens and radiation. These stresses cause the activation of the premature senescence process independent of TL (Blazkova et al., 2010; Collins, 2000; Jin et al., 2017; Mirzayans et al., 2012). The physiological phenotype of a senescent cell, either reached via an intrinsic or extrinsic path, has been named as the senescence-associated secretory phenotype (SASP) or complex senescence-messaging secretome (Young and Narita, 2009).

Salient features of senescent cells include altered cell size and large flat smoothed shape, emergence of senescence markers such as the senescence-associated β -galactosidase (SA- β -gal)(Dimri et al., 1995), senescence associated heterochromatin foci (SAHF) (Kosar et al.,

2011), lipofuscin accumulation (Georgakopoulou et al., 2013), DNA damage response (DDR) signaling (di Fagagna, 2008), and expression of decoy death receptor 2 (DCR2), embryonic chondrocyte-expressed 1 (DEC1) (Collado et al., 2005), and p15^{INK4b} (Kim and Sharpless, 2006). SA- β -gal, a lysosomal hydrolysis, has an optimal pH 4.0 in young or immortal cells (Lee et al., 2006). However, this activity is increased in senescent cells because of the increments of lysosomal content and hydrolysis degree at suboptimal pH (pH=6) displayed by senescent cells (Lee et al., 2006). In the cellular nucleus during senescence, the epigenetic alterations are related to a global change in heterochromatin via the arrangement of facultative domains heterochromatin termed SAHF (Collado et al., 2005; Lee et al., 2006; Narita et al., 2003). DEC1, as a TF contributes to cell growth, proliferation, differentiation, cell death, and senescence (Boudjelal et al., 1997; Guillaumond et al., 2008; Shen et al., 1997). Indeed, senescent cells are able to secrete different factors for instance growth factors, cytokines, chemokines, some microRNAs (miRNAs) and proteases (Acosta et al., 2013; Collado et al., 2007).

Accumulation of different senescence markers in tissues of aged mammals indicate that senescent cells take part in pathologies (Chkhotua et al., 2003; Dimri et al., 1995; Liu et al., 2009). In addition, development and evolution of age-related diseases could be attributed to the reduction of the regenerative actions of stem cells because of increasing age (Sharpless and DePinho, 2007). Senescent cells can mount up with age along with age-related pathologies, like in osteoarthritis (Martin and Buckwalter, 2002) and atherosclerotic plaques (Gorenne et al., 2006), and can have an effect on the normal function of the tissues, leading to an accelerated degeneration.

Statin drugs

Inhibitors of 3-hydroxy-3-methylglutaryl coenzyme A (HMG-CoA) reductase (or statins) represent a family of cholesterol-lowering drugs and are widely used in medical practice for patients at increased risk of atherosclerotic cardiovascular disease (Armitage, 2007; Unit, 2005). Statins influence an essential stage of cholesterol biosynthesis, the “mevalonate pathway”. Through inhibiting HMG-CoA reductase and preventing L-mevalonic acid synthesis, statins reduce the generation of subsequent isoprenoid elements of this pathway, i.e. farnesyl pyrophosphate (FPP) as well as geranylgeranylpyrophosphate (GGPP) (Istvan and Deisenhofer, 2001). These intermediates are crucial lipid attachments for post-translational modification of important proteins such as small GTPases of the Ras, Rap, Rab and Rho families (Figure 1) (Van Aelst and D’Souza-Schorey, 1997; Zhang and Casey, 1996). After isoprenylation, RhoA activates its downstream target Rho-kinase (ROCK) and afterwards induces several cellular effects (Yang et al., 2010). FPP and GGPP are essential for Ras farnesylation and Rho geranylgeranylation, respectively (Iguchi et al., 2009). By blocking the mevalonate pathway, statins prevent the biosynthesis of isoprenoid intermediates thus restraining small GTPases isoprenylation that support multiple lipid-independent or “pleiotropic” activities of statins (Laufs and Liao, 1998). Another cholesterol-independent activity of statins is the over-expression and activation of endothelial nitric oxide synthase (eNOS). The inhibition of RhoA geranylgeranylation by statins promotes eNOS expression and NO production so that statins alleviate endothelial dysfunction (Laufs et al., 1998). GGPP, but not FPP, can counteract the eNOS overexpression-induced by statins, indicating an important role of RhoA as a negative regulator of eNOS (Laufs et al., 1998).

Statins are usually prescribed as therapeutic agents as either free acid forms or lactone forms. The prodrug lactone form requires metabolization to the acid form in the liver in order to be activated and exert pharmacological activities (Schachter, 2005). Prodrug lactone and active acid forms vary regarding to their pharmacological properties and water solubility, half-life,

and potency. The lactones [i.e. lovastatin (LOV) and simvastatin (SMV)] are lipophilic whereas the acids are hydrophilic [i.e. mevastatin (MEV), pravastatin (PRV), atorvastatin (ATV), fluvastatin (FLV), and rosuvastatin (RSV)] (Corsini et al., 1999). Hydrophilic statins are predominantly targeted to liver tissue, whilst vascular cells are the main target for lipophilic statins (Schachter, 2005).

Statins reduce blood cholesterol levels and thereby delay the onset of atherosclerosis, lower the risk of cardiovascular diseases (CVDs), particularly ischemic heart disease, strokes, and reduce CVD-associated morbidity and mortality (Mehta et al., 2006; Nakamura et al., 2006). Statins are also potentially therapeutic in hypertension (Strazzullo et al., 2007) and several malignancies (Gronich and Rennert, 2013). Statins decrease the production of ROS and expression of angiotensin II (AngII)-type 1 receptor in vessels determining vasodilation. Shang *et al* claimed that SMV inhibits lipopolysaccharide-induced TNF- α expression in rat cardiomyocytes via suppression of NADPH oxidase activation and following ROS production (Shang et al., 2006). Statins can decrease ROS through activating NF-E2-related factor 2 (Nrf2) via the PI3K/Akt pathway. Nrf2 as a TFs mediate the transcriptional cells response to OS and electrophilic xenobiotics (Chartoumpekis et al., 2010). Furthermore, statins can reduce large artery stiffness and enhance arterial wall compliance (Strazzullo et al., 2007). Moreover, statins have anti-tumor properties through the reduction of p-MEK1/MEK2 protein levels, which are associated to the Ras/Raf/MEK/ERK signaling pathway responsible for cell proliferation. After statin treatment, malignant cells had a proapoptotic alterations in the Bcl-2/Bax protein ratio, and a translation of their potential from high-invasive to low-invasive (Gronich and Rennert, 2013).

Anti-ageing effect of statins

Accumulating evidences indicate that the pleiotropic effects of statins involve restoring endothelial function, stabilization of the atherosclerotic plaques, remodeling of myocardial

interstitial collagen, **modifying** apoptosis in vascular ECs and reducing oxidative stress and vascular inflammation, plus anti-neoplastic, immune-modulative, and reno-protective properties (Choi et al., 2010; Kureishi et al., 2000; Sverdrup et al., 2010). Recently, the anti-ageing effects of statins have attracted the attention of researchers (Table 1).

The *klotho* gene, originally discovered through insertional mutagenesis in mouse, is currently described as an anti-ageing gene (Kuro-o et al., 1997). Homozygous *klotho* gene knock-out mice (*klotho* mice) show reduced ageing-related morbidity. Deregulation of *klotho* is associated with ageing-related manifestations like infertility, arteriosclerosis, soft tissues calcification, neural degeneration, osteoporosis, skin atrophy, pulmonary emphysema and lower lifespan, while high-expression of *klotho* suppresses the ageing process (Kuro-o et al., 1997; Nagai et al., 2003). *Klotho* transcript and protein are mostly found in the kidney and play a key role in contrasting endothelial dysfunction, enhancing **vascular action** (Saito et al., 2000), protecting from AngII-stimulated renal damage and ameliorating kidney function (Mitani et al., 2002). Several studies demonstrated that statins can increase the renal transcription of *klotho* mRNA in a dose-dependent way (Figure 2) (Kuwahara et al., 2008; Yoon et al., 2012). Actually, inhibition of RhoA via statins is necessary for *klotho* amplification. ATV appears to be more potent on RhoA/ROCK pathway inhibition and *klotho* mRNA expression induction when compared to PTV. Controversially, AngII activates RhoA and concurrently hampers *klotho* transcription. However, ATV pretreatment reduces AngII-induced response without any tubular damage. Altogether, these findings reveal that the Rho/ROCK pathway suppression is involved in *klotho* mRNA transcription induction through statins (Narumiya et al., 2004). In another study, the effects of NOS inhibition on *klotho* protein expression as well as the anti-arteriosclerotic effects of statin were explored (Kuwahara et al., 2008). Chronic NOS blockade significantly reduces kidney *klotho* protein expression, whereas administration of ATV or PTV entirely counteracts the NOS-induced

decrement in *klotho* expression in rats (Kuwahara et al., 2008). Besides, statin therapy attenuates the downregulation of renal *klotho* mRNA in cyclosporine A (CsA)-induced nephropathy. CsA prescription promotes phosphorylated FOXO1 (p-FOXO1) expression and reduces p-FOXO3a expression, while statin administration reverses those alterations by up-regulating antioxidant enzymes and inhibiting the transcription of pro-apoptotic proteins (Yoon et al., 2012).

Sabbatini and colleagues examined the effects of ATV on ischemic acute renal failure (ARF) in aged rats with a higher risk of developing ischemia. ATV pretreatment attenuates kidney vasoconstriction in aged rats and restores GFR levels to those of young rats. ATV does not influence the Ras pathway in aged rats but can partially block Rho activation. Low-dose and short-term ATV therapy increases NO availability in old rats, ameliorating renal dynamics and providing a specific tubular protection post-ischemia. Altogether, ATV decreases NO deficiency in old rats via a remarkable impact on eNOS and RhoA proteins. These favorable effects of statins in aged animals may also be mediated by alterations in lipid oxidation and by reduced ROS generation and cell turnover (Sabbatini et al., 2004). The ability of statins in reducing ROS in stressed cells may be due to reduction of oxidized-LDL, down-expression of vascular AngII (AT₁) receptor (Wassmann et al., 2001), suppression of small G-proteins (i.e. Rho) expression (16) and inactivation of the mitogen activated protein kinase (MAPK)-extracellular signal-regulated kinase1/2 (ERK1/2) (Gueler et al., 2002).

It has been suggested that statins may have a potential role in delaying cardiac ageing. The cardioprotective mechanism of statins has been investigated in several reports (Han et al., 2013; Han et al., 2012; Takemoto and Liao, 2001; Thunyakitpisal and Chaisuparat, 2004; Wilson et al., 2005). For instance, during ATV treatment, myocardial lipofuscin amounts, an index of oxidative stress and cardiac ageing status, are remarkably decreased in the aged rats. Moreover, long-term ATV treatment can delay or rather reverse many ageing-associated

pathophysiological alterations, such as cardiac hypertrophy, myocardial apoptosis, and myocardial oxidative stress in a dose-dependent manner. Furthermore, owing to their anti-inflammatory effects, statins are able to delay the heart ageing process (Han et al., 2013).

Statin therapy leads to remarkable reduction of cardiomyocytes diameter, left ventricular wall thickness, collagen condensation, type I/III collagen ratio, heart β -galactosidase levels, heart tissues malondialdehyde activity and to over-activation of superoxide dismutase, catalase and NOS in Wistar rats (Han et al., 2012). Statin administration could reduce expression of several ageing-associated inflammatory cytokines (i.e. IL-1 and TNF- α) on both the transcriptional and translational levels, and enhances the expression of PPARs (Han et al., 2012). Statin may also reduce the expression and secretion of a family of proteolytic enzymes and matrix metalloproteinases (MMPs), thanks to the blockade of mevalonate synthesis (Takemoto and Liao, 2001; Thunyakitpisal and Chaisuparat, 2004; Wilson et al., 2005). In atherogenesis, MMPs may take part in plaque rupture and thereby treatment with statins may relieve the symptoms of cardiac ageing (Davies, 1995; Richardson et al., 1989).

Long-term treatment with ATV lessens the SASP, enhances endothelium relaxation, reduces malondialdehyde levels, increases superoxide dismutase, eNOS, and SIRT1 regulation and translation, and normalizes eNOS/iNOS imbalance in aortas of ageing-rat in comparison with non-treated controls (Gong et al., 2014). These data suggest the efficacy of long-term ATV treatment in improving age-associated endothelial dysfunction, representing a promising approach for atherosclerosis prevention (Figure 2).

Sirtuin-1 (SIRT1; silent information regulator 1), belongs to the Sir2 family proteins whose expression falls with age (Hall et al., 2013). SIRT1 is a NAD⁺-dependent deacetylase that targets both histone and non-histone proteins, like TFs (Zhang and Kraus, 2010) in order to preserve chromatin silencing, genomic stability, cell differentiation, and is a regulator of metabolism and endothelial function (Menghini et al., 2009). SIRT1 can suppress

atherosclerosis, endothelial dysfunction and vascular cell senescence via interaction with and stimulation of eNOS (Potente and Dimmeler, 2008a, b). Recently, accumulating evidences showed that mammalian SIRT1 activity increases longevity in animal models (Guarente, 2013; Imai and Guarente, 2016; Mercken et al., 2014), and statin administration counteracts endothelial senescence in human umbilical vein ECs through SIRT1 increment (Ota et al., 2010).

MicroRNAs (miRs) belong to a family of short non-coding RNAs that control mRNA expression of target genes through the **cleavage** or repression of translation (Amerizadeh et al., 2018; Bahreyni et al., 2019). It has been shown that SIRT1-related miRs, such as *miR-34a*, can target SIRT1 causing endothelial senescence. Ota *et al* studied the expression pattern of SIRT1-related miR profiles, including *miR-9*, *-34a*, *-132*, *-181a*, *-195*, *-199a*, *-199b* and *-204*, SIRT1 in **EPCs** of CAD patients, and the statin influence on these patterns. Among all the analyzed miRs, only *miR-34a* levels were increased and SIRT1 protein levels lower in CAD patients when compared to non-CAD controls. Values of *miR-34a* were slightly inversely correlated with SIRT1 protein amounts. After ATV supplementation, *miR-34a* significantly decreased and SIRT1 increased, while after RSV treatment no change was observed (Ota et al., 2010). Concentrations of other miRs did not differ before and after treatment with ATV or RSV. Conclusively, *miR-34a* may modulate SIRT1 transcription in EPCs and the ATV-induced over-expression of SIRT1 through suppression of *miR-34a* might concur to the beneficial effects of ATV on endothelial function among CAD patients (Ota et al., 2010).

Progeroid syndromes are heritable human diseases characterized by the very early occurrence of signs/symptoms of premature ageing. Hutchinson-Gilford progeria syndrome (HGPS), a rare childhood premature ageing syndrome, caused by a spontaneous point mutation in *lamin A (LMNA)* gene, leads to an aberrant alternative splicing of prelamin A towards the synthesis

of a truncated nuclear protein named progerin (Gordon et al., 2014). Accumulation of farnesylated forms of progerin at the nuclear membrane distorts nuclear architecture and affects chromatin stability determining many severe damages. There is an association between HGPS and progressive TL shortening (Burla et al., 2016; Chojnowski et al., 2015; McCord et al., 2013) that causes DNA damage. Combined treatment with statins and aminobisphosphonates (aBP) effectively blocks farnesylation and geranylgeranylation of prelamin A and progerin and significantly reduces the ageing-related phenotypes of mice mutant for the *Zmpste24*-MMP. Differently from statins, aBP act on the last enzymes of the mevalonate synthetic pathway, blocking FPP synthase and isopentenyl pyrophosphate isomerase (IPP) (Konstantinopoulos and Papavassiliou, 2007). The additive/synergistic effects of statins and aBP on *Zmpste24*^{-/-} ageing-related phenotypes may be due to the sequential actions on different steps of the mevalonate axis, thereby inhibiting both protein farnesylation and geranylgeranylation (Varela et al., 2008).

Statins were found to be able to effectively prevent inflammatory arthritis (Leung et al., 2003). In recent reports, chondrocyte ageing was reported to be linked to the progression of cartilage damages (Yudoh and Karasawa, 2010). Yudoh and co-researchers demonstrated that the OA-associated catabolic factor IL-1 β induces down-expression of cellular function, over-activation of the senescent marker β -galactosidase and reduces the cellular lifetime of chondrocytes *in vitro*. Statin administration in chondrocytes suppresses the IL-1 β -triggered synthesis of cartilage matrix degenerating enzymes (MMP-1 and MMP-13), reduces premature senescence, and promotes the biosynthesis of cartilage matrix proteoglycans. In an animal model of OA, statin therapy remarkably decreased the degradation of articular cartilage, whereas the control group displayed progressive knee joints cartilage destruction over time. Statin may have the great advantage to counteract the catabolic stress-stimulated

chondrocyte dysfunction found in the articular cartilages of the elderly (Yudoh and Karasawa, 2010).

Effect of statins on cellular senescence and Telomere

Cellular senescence interferes with tumor development by inhibiting the proliferation of damaged cells and enforcing oncogenic cells to enter cell cycle arrest, though senescent cells are able to induce malignancy by the pro-inflammatory SASP (Figure 2) (Campisi, 2001). It has been shown that SMV attenuates the SASP and its tumor-inducing potential in senescent human fibroblasts by hindering protein prenylation and inactivation of Rho family GTPases Rac1 and Cdc42. IL-6 is one of the SASP factors that can trigger proliferation of breast tumor cells by activation of the MEK-ERK1/2 pathway, but SMV impedes this SASP-induction proliferation (Liu et al., 2015).

Cellular OS can determine a unique phenotype characterized by telomeres shortening, DNA/protein/lipid peroxidation, and cellular senescence-induced by p38-MAPK activation (Behnia et al., 2016; Behnia et al., 2015). p38MAPK protein kinases are a family of intracellular acute stress response proteins that contribute to inflammation, cell death and senescence (Coulthard et al., 2009). p38MAPK is important for the senescence because of its capability to induce the p53 and pRb/p16 growth arrest pathways (Iwasa et al., 2003). Thus, regulating p38MAPK activation and diminishing premature senescence activation could reduce the incidence of OS-associated disorders. Ayad *et al* reported that both SMV and RSV potentially down-express OS-stimulated p38MAPK activation, premature senescence and SASP, whereas RSV exhibits a promoting effect (Ayad et al., 2018).

Inflammation and OS are two major factors that induce age-associated telomere shortening, suggesting that telomere erosion could be considered a promising marker of OS and inflammation (Babizhayev et al., 2011; Houben et al., 2008). Recent evidences pointed out

that statins can affect cell senescence, suggesting a connection between statins and telomere/telomerase system (Nielsen et al., 2012). The TL of EPCs has poor inverse correlation with oxidative DNA stress (Satoh et al., 2008). It has been demonstrated that EPCs obtained from CAD patients present shorter TL and higher telomere shortening rate and oxidative damage to DNA than those from control subjects (Satoh et al., 2008). Evidence supported that EPC telomere shortening due to the oxidative DNA lesions may contribute to the development of CAD. Other supporting evidence comes from a study which assessed both the association between history of statin treatment and leukocyte telomere length (LTL) and its connection with plausible novel indicators of oxidative DNA damages and ROS-induced inflammation. As telomere shortening is nearly equal in various human tissues, blood circulating mononuclear cells can be used as an easily applicable surrogate tissue for TL assessment. Results demonstrated that patients undergoing statin treatment have longer mean LTL when compared with non-treated patients. Indeed, the expression levels of two potent biomarkers of oxidative damage and inflammation, leukocyte Finkel-Biskis-Jinkins osteosarcoma (FOS) and 8-oxoguanine DNA glycosylase (OGG1), were not different between two groups. This finding showed that statin treatment was correlated with longer LTL. Notably, higher FOS and OGG1 can be considered as a novel pertinent biomarkers of LTL (Saliques et al., 2011). A randomized controlled trial compared the effect of intensive cholesterol-reduction treatment (ATV; 10 mg per day) and moderate cholesterol-reduction treatment (PTV; 10 mg per day) on EPC telomere **biology** (Satoh et al., 2009). Intensive cholesterol-reduction treatment **enhanced EPC frequencies** and limited EPC telomere shortening, but moderate cholesterol-reduction treatment did not affect EPC populations or withhold EPC telomere shortening (Satoh et al., 2009).

Boccardi and co-workers investigated the effect of statin administration on peripheral blood leukocyte telomerase activity (TA), leukocyte telomere length (LTL) instability, and the

relationship with telomere shortening rate through ageing (Boccardi et al., 2013). In this cross-sectional survey, individuals receiving statin showed significantly increased TA, longer LTL and less telomere erosion *versus* the control group, after adjustment for potential confounders including age, sex, smoking, blood pressure and blood levels of glucose, lipids, and inflammatory parameters. LTL reduction was 0.03 Kb per each year of age in statin receiving group, and 0.06 Kb in controls, respectively. Interestingly, a major difference in telomere erosion between the two groups was found in older subjects (65 year) (Boccardi et al., 2013). Several studies demonstrated that low TA and subsequent weaker telomere maintenance potency are connected to a higher risk for CVD, independent of chronological age (Serrano and Andrés, 2004), whereas over-expression of telomerase without net telomere lengthening possibly enhances cellular longevity and genome stability *in vitro* (Zhu et al., 1999). In healthy women, TA rather than TL is negatively related with main CVD risk factors (Sato et al., 2009).

Another study recruited 484 cases at high risk for coronary heart disease (CHD) events and 1058 controls to evaluate the relationship between TL and observed clinical advantage of statin therapy. Results showed that mean LTL declined with age by 9% and 5.9% per decade in controls and cases, respectively. Subjects in the middle and in the lowest tertiles of LTL were at significantly greater risk of developing a CHD event when compared to subjects in the highest tertile (odds ratio [OR] for CHD: 1.5; 95% confidence interval [CI]:1.1–2.0 in the middle tertile; 1.4; 1.1–1.9 in the lowest). In the placebo group, risk of CHD was approximately two-fold in individuals in the lower two tertiles of LTL when compared to individuals in the highest tertile (1.9; 1.3–2.8, $p < 0.001$ and 1.9; 1.3–2.85, $p < 0.001$ in the middle and lowest tertiles, respectively). In patients treated with PRV, the elevated risk due to shorter telomeres, was remarkably mitigated (1.1; 0.7–1.7, $p = 0.58$ and 1.02; 0.7–1.5, $p = 0.94$ in the middle and lowest tertiles, respectively). Mean LTL seems to be a predictor of

CHD events. It seems that patients with higher risk according to mean LTL did greatly benefit from statin treatment in term of CHD prevention (Brouillette et al., 2007).

Low-dose FLV or valsartan or their combination significantly raised TA by 106.9%, 59.5%, and 228.0% respectively when compared to paired controls who received placebo. Remarkably, increased TA from the combination arm was significantly related to the improvement of arterial function as evaluated by flow-mediated dilation as well as inflammation/OS reduction as assessed by C-reactive protein and total antioxidant capacity. The authors recommended the introduction of a combination of low-dose FLV and valsartan as a novel innovative approach for “arterial rejuvenation” (Janić et al., 2016).

In a prospective randomized trial, 100 hypercholesterolemic patients in primary prevention were randomized to receive either ATV (20 mg daily) or placebo for one year. At the trial end, the log-value of TA changed from 0.46 to 0.68 ($p = 0.004$) and from 0.67 to 0.60 ($p = 0.48$) in the ATV and placebo arms, respectively. ATV treatment was the only significant predictor of TA alterations, independently of inflammatory and oxidative biomarkers levels (Strazhesko et al., 2016).

EPCs play a significant role in neo-angiogenesis after ischemia (Zhang et al., 2002). ATV and MEV hamper the onset of EPC senescence in culture in a dose-dependent way. In addition, ATV accelerates proliferation of EPCs, while FPP or GGPP attenuate the anti-ageing effect of ATV. On the other hand, NOS withholding, antioxidants, or Rho kinase inhibitors do not affect ATV effects. ATV regulates several cell cycle genes transcription, for instance inducing the expression of cyclins and reducing expression of p27Kip1, the cyclin dependent kinase–G1 phase inhibitor. Accordingly, statins prevented EPCs senescence independently of NO, ROS, and Rho kinase, but dependently on GGPP. ATV-induced inhibition of EPCs senescence seems to be mediated by the regulation of different cell cycle proteins via the PI3K pathway. The increase of cell cycle related proteins together with a

decrement of the p27 may contribute to cell cycle progression and thereby inhibit the initiation of replicative senescence. Therefore, activation of the PI3K/Akt axis via statins may have several protective consequences on EPCs such as the expansion of EPCs **frequency** and ultimately suppression of apoptosis and senescence (Assmus et al., 2003).

Opposing evidences about anti-ageing effects of statin

Suppression of HMG-CoA reductase leads to a reduced production of cholesterol and other metabolites downstream of mevalonate, a precursor of the coenzyme Q10 (CoQ10, ubiquinone) synthesis (Mabuchi et al., 2007). CoQ10 is a **unique** lipid soluble benzoquinone and a major component of the respiratory chain that is located in the hydrophobic sections of cellular membranes and involved in oxidative phosphorylation for adenosine triphosphate (ATP) biosynthesis (Langsjoen, 1994).

The cellular roles of CoQ10 in humans include transport of electrons from mitochondrial respiratory complexes I and II to complex III, production of superoxide anion radicals via autoxidation ubiquinone, as well as anti-oxidant scavenging of free radicals (Kishimoto et al., 2003; Sun et al., 1992).

The antioxidant activity of **CoQ** is particularly important in inhibiting ceramide-associated apoptosis (Navas et al., 2007), a crucial regulator of longevity **concerning of normal aging** (Martin-Montalvo et al., 2016). Aging seems to contribute in lowering the concentrations of CoQ10 (Littarru and Langsjoen, 2007). A remarkable decrement in the speed of CoQ production has been found during the aging process and aging-related disorders (Allewa et al., 1995; Kalén et al., 1989). It was suggested that aging-associated elevation in mitochondrial OS may be due to CoQ depletion (Miles et al., 2004). On the other hand, Ishii *et al.* demonstrated that CoQ10 extends the life span of wild-type *Caenorhabditis elegans*, the most widely used organism as a model for ageing in longevity researches. In fact, CoQ10

supplementation can significantly increase *C. elegans* longevity possibly by lowering the superoxide anion levels and subsequently decreasing OS in mitochondria (Ishii et al., 2004). However, there are several inconsistencies about the association between CoQ concentrations and the aging process. Mice deficient of one of the alleles of the *COQ7* gene present extended lifespan even though their CoQ concentrations are similar to those of wild-type mice, indicating that factors different from CoQ could be implicated in life time extension (Lapointe and Hekimi, 2008). However, a direct association between lifespan and mitochondrial CoQ contents has been documented in a senescence-accelerated mice model (Tian et al., 2014). Furthermore, CoQ treatment delays senescence (Tian et al., 2014).

Since cholesterol and coQ10 share a similar biosynthetic pathway, a statin-induced decrement in cholesterol may also lead to a block of CoQ10 generation. A recent systematic review and meta-analysis of eight placebo-controlled trials reported a significant reduction in plasma CoQ10 levels after statins therapy (weighted mean difference=−0.44 mol/L, 95%CI: −0.52 to −0.37, $p<0.001$) (Banach et al., 2015b).

Adverse side effects related to statins, known as statin-associated muscle symptoms (SAMS), are mainly attributed to the reduction of CoQ10 in muscle tissue and subsequent mitochondrial dysfunction (Banach et al., 2015a; Deichmann et al., 2010). However, not all studies support the possible mitochondrial dysfunction induced by statins. In humans, Laaksonen and coworkers found no alterations in CoQ10 levels in muscle biopsies from subjects before and after SMV therapy (Laaksonen et al., 1996).

We previously mentioned that longevity is closely related to decrement in circulating insulin concentrations and increased insulin sensitivity. However, whether statins by decreasing insulin sensitivity and secretion triggers the progression of diabetes is an issue of concern. Mechanisms behind the relationship of statins with diabetes mellitus remain unknown. Researches about the effects of statin therapy on insulin sensitivity are contradicting. Some

studies reported detrimental or neutral effects of statins on insulin sensitivity and secretion (Cederberg et al., 2015; Gannagé-Yared et al., 2005; Moutzouri et al., 2011; Puurunen et al., 2013; Szendroedi et al., 2009), whereas others have demonstrated beneficial effects (Naples et al., 2008; Paolisso et al., 2000; Sugiyama et al., 2007). In a systematic review and meta-analysis of clinical trials, PRV and SMV were reported to respectively improve and worsen significantly insulin sensitivity (Baker et al., 2010). It may be possible that lipophilic and hydrophilic statins have various effect on glucose tolerance and thereby differential metabolic effects (Axsom et al., 2013). **Therefore, statins have unique affinities to cell membranes and thus subjects competencies to influence cell functions.** Future researches should focus not only on shedding light on the impact of the different statins on the glycemic control, but also on their association with the ageing processes.

Conclusion

Cellular ageing or senescence is a complex response to different stimuli that lead to a progressive loss of tissue and organ function. Statin therapy improves cell functionality, delays senescence, and suppresses telomere shortening and apoptosis. Since senescence is an irreversible process, slowing its progression through statins could offer new opportunities and targets for preventive approaches. However, the interplay between TL and statin therapy has not been specifically studied in humans; maintaining the shelterin complex stability, a high TA or a beneficial effect on telomere function are the potential effects of statins in humans. Finally, statins display several benefits on cardiac/vascular cells and endothelial function, including inflammation and oxidative stress reduction, which could participate to the pleiotropic lipid-lowering independent effects of statins. Statin administration could be considered as a promising strategy for the treatment of age-related pathologies.

Competing interests: The authors declare no conflict of interests.

Funding: None.

Figure legends:

Figure 1. Effects of statins on the cholesterol biosynthesis pathway. **Abbreviations:** eNOS, endothelial nitric oxide synthases; HMG-CoA, 3-hydroxy-3-methylglutaryl coenzyme A; FPP, farnesyl pyrophosphate; GGPP, geranylgeranylpyrophosphate; ROCK, Rho-kinase; ROS, Reactive Oxygen Species.

Figure 2. Mechanisms by which statins influence cellular ageing and senescence. Statins inhibit telomerase shortening, *miR-34a*, eNOS down-regulation, eNOS inactivation, and OS-stimulated p38MAPK activation; statins induce SIRT1, FoxO3 and klotho mRNA over-expression. Statins also minimize the SASP and its ageing-enhancing consequences in senescent cells. **Abbreviations:** FoxO, Forkhead box O; SASP, senescence-associated secretory phenotype; SA- β -gal, senescence-associated β -galactosidase; SAHF, senescence associated heterochromatin foci; DDR, DNA damage response; DCR2, decoy death receptor 2; DEC1, embryonic chondrocyte-expressed 1.

Table 1. Effect of statin treatments on ageing and longevity.

Condition	Arm	Experiment model	Outcome	Ref.
Ageing	ATV (1 μ mol/l)	-mIMCD3 tubular cell line	-RhoA/ROCK pathway inhibition -klotho mRNA expression induction	(Narumiya et al., 2004)
CsA-nephropathy	PRV	- normal mice - chronic CsA nephropathy-induced mice	-attenuation of the down-regulation of renal klotho mRNA in CsA-nephropathy induced by oxidative stress -reduction of p-FoxO1 expression and increase of p-FoxO3a expression	(Yoon et al., 2012)
ARF	ATV (12 mg/kg/d for 14 d)	-young (3 mo old) rats -old (18 mo old) rats	-block the Rho activation -enhancement of NO availability	(Sabbatini et al., 2004)
Cardiac ageing	ATV (10 or 1mg/kg/day); 4 months -saline	Aged rats	-reversion of the ageing-associated pathophysiological alterations (i.e. cardiac hypertrophy, myocardial apoptosis, myocardial oxidative stress) -decrement of myocardial lipofuscin	(Han et al., 2013)
Cardiac ageing	-AVT (10 or 1 mg/kg/ day); 4 months -saline	-old Wistar rats	-decrement of the transcription and translation of several ageing-associated inflammatory cytokines (i.e. IL-1 and TNF- α) -overexpression of PPARs -inhibition of the expression and secretion MMPs	(Han et al., 2012)
Endothelial dysfunction	-ATV (5 mg/kg/day); 8 months	-Male old Wistar rats	-reduction of the SASP -enhancement of the relaxation of endothelium -decrement of MDA -increase of SOD, eNOS, and SIRT1 regulation -normalization of eNOS/iNOS imbalance	(Gong et al., 2014)
CAD	- ATV(10 mg/day) ; 8 months -RSV (2.5 mg/day); 8 months	-cultured EPCs from 70 patients with CAD and 48 subjects without CAD	-ATV leads to over-expression of SIRT1 through suppression of <i>miR-34a</i>	(Ota et al., 2010)
Premature ageing	-PRV(100 mg/kg/d) -amino-bisphosphonates	-Zmpste24 ^{-/-} mice	-blocking of farnesylation and geranylgeranylation of prelamin A and progerin -reduction of the ageing-related phenotypes	(Varela et al., 2008)
Osteoarthritis	-statin (1.0 or 10.0 μ M)	-human articular cartilage samples from OA patients -STR/OrtCrlj mice	-suppression of the IL-1 β -triggered generation of cartilage matrix degenerating enzymes -increase of the biosynthesis of cartilage matrix proteoglycan -stress-stimulated chondrocyte dysfunction	(Yudoh and Karasawa, 2010)
Breast cancer -induced by senescent cells	-SMV	- HCA2 human fibroblast cells - MCF7 and ZR-75 human breast cancer cells	-minimizing the SASP and its tumor-enhancing consequences in senescent human fibroblasts through hindering protein prenylation and inactivation of Rho family GTPases Rac1 and Cdc42	(Liu et al., 2015)
Premature	-SMV (100 and 200	-fetal membranes	-down-expression of OS-stimulated	(Ayad et

senescence	ng/mL) -RSV (100 and 200 ng/mL) - progesterone (10 ⁻⁶ mol/L)		p38MAPK activation, premature senescence and SASP	al., 2018)
Arteriosclerosis	-ATV (10 mg per day) -PTV (10 mg per day)	-Wistar rats	-ATV enhances EPC frequencies and barricaded EPC telomere shortening	(Sato et al., 2009)
Ageing	-RSV -ATV -SMV	-230 Caucasians	-increment in TA, LTL levels and reduction in telomere erosion	(Boccardi et al., 2013)
CHD	- PRV (40 mg daily); 4.9 years -placebo	-6595 statin-naive men	-mitigation of elevated risk with shorter telomeres	(Brouillette et al., 2007)
Arterial wall rejuvenation	-FLV (10 mg daily); 30 days -valsartan (20 mg daily); 30 days -combination (10 mg daily); 30 days - placebo	-130 middle-aged, apparently healthy	-low-dose FLV or valsartan or combination significantly elevate TA by 106.9%, 59.5%, and 228.0% when compared to controls -improvement of arterial function and reduction of inflammation/oxidative stress	(Janić et al., 2016)
CVD	- ATV (20 mg/day); 12 months -placebo)	-100 hypercholesterolemic patients	-increment in TA	(Strazhesko et al., 2016)
Ischemic tissue	-ATV -MEV	-Mononuclear cells	-acceleration of the proliferation of EPCs -modification of mRNA transcription of several cell cycle genes -prevention of the GGPP-dependent senescence in EPCs	(Boccardi et al., 2013)

ARF, ischemic acute renal failure; ATV, atorvastatin; CAD, coronary artery disease; CHD, coronary heart disease; CsA; cyclosporine; EPCs: Endothelial progenitor cells; FLV, fluvastatin; GGPP: geranylgeranylpyrophosphate; IL, interleukin 1; LTL: leukocyte telomere length; MAPK, mitogen activated protein kinase; MDA, Malonyldialdehyde; MEV, mevastatin; miR, microRNA; MMPs, matrix metalloproteinases; NO, nitric oxide; NOS, nitric oxide synthases; iNOS: inducible NOS; OA: osteoarthritis; OS, oxidative stress; p-FoxO1, phosphorylated Forkhead box O; PRV, pravastatin; ROCK, Rho-kinase; PPARs, peroxisome proliferator-activated receptors; RSV, rosuvastatin; SASP, senescence-associated secretory phenotype; SIRT1, Sirtuin-1 ; SMV, simvastatin ; SOD, superoxide dismutase; TA, telomerase activity; TNF- α , tumor necrosis factor;

References

- Accili, D., Arden, K.C., 2004. FoxOs at the crossroads of cellular metabolism, differentiation, and transformation. *Cell* 117, 421-426.
- Acosta, J.C., Banito, A., Wuestefeld, T., Georgilis, A., Janich, P., Morton, J.P., Athineos, D., Kang, T.-W., Lasitschka, F., Andrusis, M., 2013. A complex secretory program orchestrated by the inflammasome controls paracrine senescence. *Nature Cell Biology* 15, 978-990.
- Alichniewicz, K.K., Brunner, F., Klünemann, H.H., Greenlee, M.W., 2012. Structural and functional neural correlates of visuospatial information processing in normal aging and amnesic mild cognitive impairment. *Neurobiology of Aging* 33, 2782-2797.
- Alleva, R., Tomasetti, M., Battino, M., Curatola, G., Littarru, G., Folkers, K., 1995. The roles of coenzyme Q10 and vitamin E on the peroxidation of human low density lipoprotein subfractions. *Proceedings of the National Academy of Sciences* 92, 9388-9391.
- Allsopp, R.C., Harley, C.B., 1995. Evidence for a critical telomere length in senescent human fibroblasts. *Experimental Cell Research* 219, 130-136.
- Amerizadeh, F., Khazaei, M., Maftouh, M., Mardani, R., Bahrami, A., 2018. miRNA Targeting Angiogenesis as a Potential Therapeutic Approach in the Treatment of Colorectal Cancers. *Current Pharmaceutical Design* 24, 4668-4674.
- Armitage, J., 2007. The safety of statins in clinical practice. *The Lancet* 370, 1781-1790.
- Asahara, T., Murohara, T., Sullivan, A., Silver, M., van der Zee, R., Li, T., Witzenbichler, B., Schatteman, G., Isner, J.M., 1997. Isolation of putative progenitor endothelial cells for angiogenesis. *Science* 275, 964-967.
- Assmus, B., Urbich, C., Aicher, A., Hofmann, W.K., Haendeler, J., Rössig, L., Spyridopoulos, I., Zeiher, A.M., Dimmeler, S., 2003. HMG-CoA reductase inhibitors reduce senescence and increase proliferation of endothelial progenitor cells via regulation of cell cycle regulatory genes. *Circulation Research* 92, 1049-1055.
- Axson, K., Berger, J.S., Schwartzbard, A.Z., 2013. Statins and diabetes: the good, the bad, and the unknown. *Current Atherosclerosis Reports* 15, 299. doi: 10.1007/s11883-012-0299-z
- Ayad, M.T., Taylor, B.D., Menon, R., 2018. Regulation of p38 mitogen-activated kinase-mediated fetal membrane senescence by statins. *American Journal of Reproductive Immunology* 80, e12999. doi: 10.1111/aji.12999
- Babizhayev, M.A., Savel'yeva, E.L., Moskvina, S.N., Yegorov, Y.E., 2011. Telomere length is a biomarker of cumulative oxidative stress, biologic age, and an independent predictor of survival and therapeutic treatment requirement associated with smoking behavior. *American Journal of Therapeutics* 18, e209-e226.
- Bahrami, A., Parsamanesh, N., Atkin, S.L., Banach, M., Sahbekar, A., 2018. Effect of statins on toll-like receptors: a new insight to pleiotropic effects. *Pharmacological Research* 135, 230-238.
- Bahreyni, A., Rezaei, M., Bahrami, A., Khazaei, M., Fiuji, H., Ryzhikov, M., Ferns, G.A., Avan, A., Hassanian, S.M., 2019. Diagnostic, prognostic, and therapeutic potency of microRNA 21 in the pathogenesis of colon cancer, current status and prospective. *Journal of Cellular Physiology* 234, 8075-8081.
- Baker, W.L., Talati, R., White, C.M., Coleman, C.I., 2010. Differing effect of statins on insulin sensitivity in non-diabetics: a systematic review and meta-analysis. *Diabetes Research and Clinical Practice* 87, 98-107.

- Banach, M., Rizzo, M., Toth, P.P., Farnier, M., Davidson, M.H., Al-Rasadi, K., Aronow, W.S., Athyros, V., Djuric, D.M., Ezhov, M.V., 2015a. Statin intolerance—an attempt at a unified definition. Position paper from an International Lipid Expert Panel: This paper is also published in parallel in Archives of Medical Science [Banach M, Rizzo M, Toth PP, et al. Statin intolerance—an attempt at a unified definition. Position paper from an International Lipid Expert Panel. Arch Med Sci 2015; 11 (1): 1–23]. Expert Opinion on Drug Safety 14, 935-955.
- Banach, M., Serban, C., Ursoniu, S., Rysz, J., Muntner, P., Toth, P.P., Jones, S.R., Rizzo, M., Glasser, S.P., Watts, G.F., 2015b. Statin therapy and plasma coenzyme Q10 concentrations—a systematic review and meta-analysis of placebo-controlled trials. Pharmacological Research 99, 329-336.
- Bartke, A., 2005. Minireview: role of the growth hormone/insulin-like growth factor system in mammalian aging. Endocrinology 146, 3718-3723.
- Bartke, A., 2011. Growth hormone, insulin and aging: the benefits of endocrine defects. Experimental Gerontology 46, 108-111.
- Behnia, F., Sheller, S., Menon, R., 2016. Mechanistic differences leading to infectious and sterile inflammation. American Journal of Reproductive Immunology 75, 505-518.
- Behnia, F., Taylor, B.D., Woodson, M., Kacerovsky, M., Hawkins, H., Fortunato, S.J., Saade, G.R., Menon, R., 2015. Chorioamniotic membrane senescence: a signal for parturition? American Journal of Obstetrics and Gynecology 213, 359.e1-16
- Biagi, E., Nylund, L., Candela, M., Ostan, R., Bucci, L., Pini, E., Nikkila, J., Monti, D., Satokari, R., Franceschi, C., 2010. Through ageing, and beyond: gut microbiota and inflammatory status in seniors and centenarians. PloS One 5, e10667.
- Blackburn, E.H., 1991. Structure and function of telomeres. Nature 350, 569-573.
- Blazkova, H., Krejčíková, K., Moudry, P., Frisan, T., Hodny, Z., Bartek, J., 2010. Bacterial intoxication evokes cellular senescence with persistent DNA damage and cytokine signalling. Journal of Cellular and Molecular Medicine 14, 357-367.
- Boccardi, V., Barbieri, M., Rizzo, M.R., Marfella, R., Esposito, A., Marano, L., Paolisso, G., 2013. A new pleiotropic effect of statins in elderly: modulation of telomerase activity. The FASEB Journal 27, 3879-3885.
- Boudjelal, M., Taneja, R., Matsubara, S., Bouillet, P., Dollé, P., Chambon, P., 1997. Overexpression of Stra13, a novel retinoic acid-inducible gene of the basic helix-loop-helix family, inhibits mesodermal and promotes neuronal differentiation of P19 cells. Genes & Development 11, 2052-2065.
- Brouillette, S.W., Moore, J.S., McMahon, A.D., Thompson, J.R., Ford, I., Shepherd, J., Packard, C.J., Samani, N.J., Group, W.o.S.C.P.S., 2007. Telomere length, risk of coronary heart disease, and statin treatment in the West of Scotland Primary Prevention Study: a nested case-control study. The Lancet 369, 107-114.
- Burla, R., La Torre, M., Saggio, I., 2016. Mammalian telomeres and their partnership with lamins. Nucleus 7, 187-202.
- Campisi, J., 2001. Cellular senescence as a tumor-suppressor mechanism. Trends in Cell Biology 11, S27-S31.
- Campisi, J., di Fagagna, F.d.A., 2007. Cellular senescence: when bad things happen to good cells. Nature Reviews in Molecular Cell Biology 8, 729–740.
- Carter, M.E., Brunet, A., 2007. FOXO transcription factors. Current Biology 17, R113-R114.
- Cederberg, H., Stančáková, A., Yaluri, N., Modi, S., Kuusisto, J., Laakso, M., 2015. Increased risk of diabetes with statin treatment is associated with impaired insulin sensitivity

- and insulin secretion: a 6 year follow-up study of the METSIM cohort. *Diabetologia* 58, 1109-1117.
- Chartoumpakis, D., Ziros, P.G., Psyrogiannis, A., Kyriazopoulou, V., Papavassiliou, A.G., Habeos, I.G., 2010. Simvastatin lowers reactive oxygen species level by Nrf2 activation via PI3K/Akt pathway. *Biochemical and Biophysical Research Communications* 396, 463-466.
- Chkhotua, A.B., Gabusi, E., Altimari, A., D'Errico, A., Yakubovich, M., Vienken, J., Stefoni, S., Chieco, P., Yussim, A., Grigioni, W.F., 2003. Increased expression of p16 (INK4a) and p27 (Kip1) cyclin-dependent kinase inhibitor genes in aging human kidney and chronic allograft nephropathy. *American Journal of Kidney Diseases* 41, 1303-1313.
- Choi, E.-Y., Chang, W., Lim, S., Song, B.-W., Cha, M.-J., Kim, H.-J., Choi, E., Jang, Y., Chung, N., Hwang, K.-C., 2010. Rosuvastatin inhibits norepinephrine-induced cardiac hypertrophy via suppression of Gh. *European Journal of Pharmacology* 627, 56-62.
- Chojnowski, A., Ong, P.F., Wong, E.S., Lim, J.S., Mutalif, R.A., Navasankari, R., Dutta, B., Yang, H., Liow, Y.Y., Sze, S.K., 2015. Progerin reduces LAP2 α -telomere association in Hutchinson-Gilford progeria. *Elife* 4, e07759.
- Chung, H.Y., Cesari, M., Anton, S., Marzetti, E., Giovannini, S., Seo, A.Y., Carter, C., Yu, B.P., Leeuwenburgh, C., 2009. Molecular inflammation: underpinnings of aging and age-related diseases. *Ageing Research Reviews* 8, 18-30.
- Collado, M., Blasco, M.A., Serrano, M., 2007. Cellular senescence in cancer and aging. *Cell* 130, 223-233.
- Collado, M., Gil, J., Efeyan, A., Guerra, C., Schuhmacher, A.J., Barradas, M., Benguría, A., Zaballos, A., Flores, J.M., Barbacid, M., 2005. Tumour biology: senescence in premalignant tumours. *Nature* 436, 642-642.
- Collins, K., 2000. Mammalian telomeres and telomerase. *Current Opinion in Cell Biology* 12, 378-383.
- Corsini, A., Bellocchia, S., Baetta, R., Fumagalli, R., Paoletti, R., Bernini, F., 1999. New insights into the pharmacodynamic and pharmacokinetic properties of statins. *Pharmacology & Therapeutics* 84, 413-428.
- Coulthard, L.R., White, D.E., Jones, D.L., McDermott, M.F., Burchill, S.A., 2009. p38MAPK: stress responses from molecular mechanisms to therapeutics. *Trends in Molecular Medicine* 15, 369-379.
- Davies, M., 1995. Acute coronary thrombosis—the role of plaque disruption and its initiation and prevention. *European Heart Journal* 16, 3-7.
- de Lange, T., 2005. Shelterin: the protein complex that shapes and safeguards human telomeres. *Genes Dev* 19, 2100-2110.
- Deichmann, R., Lavie, C., Andrews, S., 2010. Coenzyme q10 and statin-induced mitochondrial dysfunction. *Ochsner Journal* 10, 16-21.
- di Fagagna, F.d.A., 2008. Living on a break: cellular senescence as a DNA-damage response. *Nature Reviews Cancer* 8, 512–522.
- Dimri, G.P., Lee, X., Basile, G., Acosta, M., Scott, G., Roskelley, C., Medrano, E.E., Linskens, M., Rubelj, I., Pereira-Smith, O., 1995. A biomarker that identifies senescent human cells in culture and in aging skin in vivo. *Proceedings of the National Academy of Sciences* 92, 9363-9367.
- Doksani, Y., Wu, J.Y., de Lange, T., Zhuang, X., 2013. Super-resolution fluorescence imaging of telomeres reveals TRF2-dependent T-loop formation. *Cell* 155, 345-356.

- Enns, L.C., Morton, J.F., Treuting, P.R., Emond, M.J., Wolf, N.S., McKnight, G., Rabinovitch, P.S., Ladiges, W.C., 2009. Disruption of protein kinase A in mice enhances healthy aging. *PLoS One* 4, e5963.
- Ferretti, G., Bacchetti, T., Sahebkar, A., 2015. Effect of statin therapy on paraoxonase-1 status: a systematic review and meta-analysis of 25 clinical trials. *Progress in Lipid Research* 60, 50-73.
- Förstermann, U., Sessa, W.C., 2011. Nitric oxide synthases: regulation and function. *European Heart Journal* 33, 829-837.
- Fowler, M.R., Naz, J.R., Graham, M.D., Orchard, C.H., Harrison, S.M., 2007. Age and hypertrophy alter the contribution of sarcoplasmic reticulum and Na⁺/Ca²⁺ exchange to Ca²⁺ removal in rat left ventricular myocytes. *Journal of Molecular and Cellular Cardiology* 42, 582-589.
- Franceschi, C., Campisi, J., 2014. Chronic inflammation (inflammaging) and its potential contribution to age-associated diseases. *Journals of Gerontology Series A: Biomedical Sciences and Medical Sciences* 69, S4-S9.
- Gannagé-Yared, M.-H., Azar, R.R., Amm-Azar, M., Khalifé, S., Germanos-Haddad, M., Neemtallah, R., Halaby, G., 2005. Pravastatin does not affect insulin sensitivity and adipocytokines levels in healthy nondiabetic patients. *Metabolism* 54, 947-951.
- García-Cao, I., García-Cao, M., Tomás-Loba, A., Martín-Caballero, J., Flores, J.M., Klatt, P., Blasco, M.A., Serrano, M., 2006. Increased p53 activity does not accelerate telomere-driven ageing. *EMBO Reports* 7, 546-552.
- Georgakopoulou, E., Tsimaratou, K., Evangelou, K., Fernandez, M.-P., Zoumpourlis, V., Trougakos, I., Kletsas, D., Bartek, J., Serrano, M., Gorgoulis, V., 2013. Specific lipofuscin staining as a novel biomarker to detect replicative and stress-induced senescence. A method applicable in cryo-preserved and archival tissues. *Aging (Albany NY)* 5, 37-50.
- Gong, X., Ma, Y., Ruan, Y., Fu, G., Wu, S., 2014. Long-term atorvastatin improves age-related endothelial dysfunction by ameliorating oxidative stress and normalizing eNOS/iNOS imbalance in rat aorta. *Experimental Gerontology* 52, 9-17.
- Gordon, L.B., Rothman, F.G., López-Otín, C., Misteli, T., 2014. Progeria: a paradigm for translational medicine. *Cell* 156, 400-407.
- Gorenne, I., Kavurma, M., Scott, S., Bennett, M., 2006. Vascular smooth muscle cell senescence in atherosclerosis. *Cardiovascular Research* 72, 9-17.
- Greider, C.W., 1991. Telomeres. *Current Opinion in Cell Biology* 3, 444-451.
- Gronich, N., Rennert, G., 2013. Beyond aspirin—cancer prevention with statins, metformin and bisphosphonates. *Nature Reviews in Clinical Oncology* 10, 625-642.
- Grundy, S.M., 1988. HMG-CoA reductase inhibitors for treatment of hypercholesterolemia. *New England Journal of Medicine* 319, 24-33.
- Guarente, L., 2013. Calorie restriction and sirtuins revisited. *Genes & Development* 27, 2072-2085.
- Gueler, F., Rong, S., Park, J.-K., Fiebeler, A., Menne, J., Elger, M., Mueller, D.N., Hampich, F., Dechend, R., Kunter, U., 2002. Postischemic acute renal failure is reduced by short-term statin treatment in a rat model. *Journal of the American Society of Nephrology* 13, 2288-2298.
- Guellich, A., Damy, T., Lecarpentier, Y., Conti, M., Claes, V., Samuel, J.-L., Quillard, J., Hébert, J.-L., Pineau, T., Coirault, C., 2007. Role of oxidative stress in cardiac dysfunction of

- PPAR α -/- mice. *American Journal of Physiology-Heart and Circulatory Physiology* 293, H93-H102.
- Guillaumond, F., Lacoche, S., Dulong, S., Grechez-Cassiau, A., Filipski, E., Li, X.-M., Lévi, F., Berra, E., Delaunay, F., Teboul, M., 2008. Altered Stra13 and Dec2 circadian gene expression in hypoxic cells. *Biochemical and Biophysical Research Communications* 369, 1184-1189.
- Hall, J.A., Dominy, J.E., Lee, Y., Puigserver, P., 2013. The sirtuin family's role in aging and age-associated pathologies. *The Journal of Clinical Investigation* 123, 973-979.
- Han, L., Li, M., Liu, X., 2013. Effects of long-term atorvastatin treatment on cardiac aging. *Experimental and Therapeutic Medicine* 6, 721-726.
- Han, L., Li, M., Liu, Y., Han, C., Ye, P., 2012. Atorvastatin may delay cardiac aging by upregulating peroxisome proliferator-activated receptors in rats. *Pharmacology* 89, 74-82.
- Harrison, D.E., Strong, R., Sharp, Z.D., Nelson, J.F., Astle, C.M., Flurkey, K., Nadon, N.L., Wilkinson, J.E., Frenkel, K., Carter, C.S., 2009. Rapamycin fed late in life extends lifespan in genetically heterogeneous mice. *Nature* 460, 392-395.
- Hayflick, L., 1965. The limited in vitro lifetime of human diploid cell strains. *Experimental Cell Research* 37, 614-636.
- Houben, J.M., Moonen, H.J., van Schooten, F.J., Hageman, G.J., 2008. Telomere length assessment: biomarker of chronic oxidative stress? *Free Radical Biology and Medicine* 44, 235-246.
- Iguchi, Y., Katsuno, M., Niwa, J.-i., Yamada, S.-i., Sone, J., Waza, M., Adachi, H., Tanaka, F., Nagata, K.-i., Arimura, N., 2009. TDP-43 depletion induces neuronal cell damage through dysregulation of Rho family GTPases. *Journal of Biological Chemistry* 284, 22059-22066.
- Imai, S.-I., Guarente, L., 2016. It takes two to tango: NAD⁺ and sirtuins in aging/longevity control. *NPJ Aging and Mechanisms of Disease* 2, 16017. doi: 10.1038/npjamd.2016.17.
- Ishii, N., Senoo-Matsuda, N., Miyake, K., Yasuda, K., Ishii, T., Hartman, P.S., Furukawa, S., 2004. Coenzyme Q10 can prolong *C. elegans* lifespan by lowering oxidative stress. *Mechanisms of Ageing and Development* 125, 41-46.
- Istvan, E.S., Deisenhofer, J., 2001. Structural mechanism for statin inhibition of HMG-CoA reductase. *Science* 292, 1160-1164.
- Itahana, K., Campisi, J., Dimri, G.P., 2004. Mechanisms of cellular senescence in human and mouse cells. *Biogerontology* 5, 1-10.
- Iwasa, H., Han, J., Ishikawa, F., 2003. Mitogen-activated protein kinase p38 defines the common senescence-signalling pathway. *Genes to Cells* 8, 131-144.
- Janić, M., Lunder, M., Cerkovnik, P., Prosenc Zmrzljak, U., Novaković, S., Šabovič, M., 2016. Low-dose fluvastatin and valsartan rejuvenate the arterial wall through telomerase activity increase in middle-aged men. *Rejuvenation Research* 19, 115-119.
- Jiang, J., Wang, Y., Susac, L., Chan, H., Basu, R., Zhou, Z.H., Feigon, J., 2018. Structure of Telomerase with Telomeric DNA. *Cell* 173, 1179-1190 e13.
- Jin, J., Tao, J., Gu, X., Yu, Z., Wang, R., Zuo, G., Li, Q., Lv, X., Miao, D., 2017. P16 INK4a Deletion Ameliorated Renal Tubulointerstitial Injury in a Stress-induced Premature Senescence Model of Bmi-1 Deficiency. *Scientific Reports* 7, 7502.
- Kaeberlein, M., Powers, R.W., Steffen, K.K., Westman, E.A., Hu, D., Dang, N., Kerr, E.O., Kirkland, K.T., Fields, S., Kennedy, B.K., 2005. Regulation of yeast replicative life span by TOR and Sch9 in response to nutrients. *Science* 310, 1193-1196.

- Kalén, A., Appelkvist, E.L., Dallner, G., 1989. Age-related changes in the lipid compositions of rat and human tissues. *Lipids* 24, 579-584.
- Kawamoto, A., Gwon, H.C., Iwaguro, H., Yamaguchi, J.I., Uchida, S., Masuda, H., Silver, M., Ma, H., Kearney, M., Isner, J.M., Asahara, T., 2001. Therapeutic potential of ex vivo expanded endothelial progenitor cells for myocardial ischemia. *Circulation* 103, 634-637.
- Kenyon, C., Chang, J., Gensch, E., Rudner, A., Tabtiang, R., 1993. A *C. elegans* mutant that lives twice as long as wild type. *Nature* 366, 461-464.
- Kim, W.Y., Sharpless, N.E., 2006. The regulation of INK4/ARF in cancer and aging. *Cell* 127, 265-275.
- Kishimoto, C., Tomioka, N., Nakayama, Y., Miyamoto, M., 2003. Anti-oxidant effects of coenzyme Q10 on experimental viral myocarditis in mice. *Journal of Cardiovascular Pharmacology* 42, 588-592.
- Klegarth, A.R., Eisenberg, D.T.A., 2018. Mammalian chromosome-telomere length dynamics. *Royal Society Open Science* 5, 180492.
- Konstantinopoulos, P.A., Papavassiliou, A.G., 2007. Multilevel modulation of the mevalonate and protein-prenylation circuitries as a novel strategy for anticancer therapy. *Trends in Pharmacological Sciences* 28, 6-13.
- Kosar, M., Bartkova, J., Hubackova, S., Hodny, Z., Lukas, J., Bartek, J., 2011. Senescence-associated heterochromatin foci are dispensable for cellular senescence, occur in a cell type- and insult-dependent manner and follow expression of p16ink4a. *Cell Cycle* 10, 457-468.
- Kureishi, Y., Luo, Z., Shiojima, I., Bialik, A., Fulton, D., Lefer, D.J., Sessa, W.C., Walsh, K., 2000. The HMG-CoA reductase inhibitor simvastatin activates the protein kinase Akt and promotes angiogenesis in normocholesterolemic animals. *Nature Medicine* 6, 1004-1010.
- Kuro-o, M., Matsumura, Y., Aizawa, H., Kawaguchi, H., Suga, T., Utsugi, T., Ohyama, Y., Kurabayashi, M., Kaname, T., Kume, E., 1997. Mutation of the mouse *klotho* gene leads to a syndrome resembling ageing. *Nature* 390, 45-51.
- Kuwahara, N., Sasaki, S., Kobara, M., Nakata, T., Tatsumi, T., Irie, H., Narumiya, H., Hatta, T., Takeda, K., Matsubara, H., 2008. HMG-CoA reductase inhibition improves anti-aging *klotho* protein expression and arteriosclerosis in rats with chronic inhibition of nitric oxide synthesis. *International Journal of Cardiology* 123, 84-90.
- Laaksonen, R., Jokelainen, K., Laakso, J., Sahi, T., Härkönen, M., Tikkanen, M.J., Himberg, J.-J., 1996. The effect of simvastatin treatment on natural antioxidants in low-density lipoproteins and high-energy phosphates and ubiquinone in skeletal muscle. *The American Journal of Cardiology* 77, 851-854.
- Langsjoen, P.H., 1994. Introduction to coenzyme Q10. Texas, University of Texas Medical Branch at Galveston <http://faculty.washington.edu/ely/coenzq10.html> (25.4.2009).
- Lapointe, J., Hekimi, S., 2008. Early mitochondrial dysfunction in long-lived *Mcl1+/-* mice. *Journal of Biological Chemistry* 283, 26217-26227.
- Laufs, U., La Fata, V., Plutzky, J., Liao, J.K., 1998. Upregulation of endothelial nitric oxide synthase by HMG CoA reductase inhibitors. *Circulation* 97, 1129-1135.
- Laufs, U., Liao, J.K., 1998. Post-transcriptional regulation of endothelial nitric oxide synthase mRNA stability by Rho GTPase. *Journal of Biological Chemistry* 273, 24266-24271.

- Lee, B.Y., Han, J.A., Im, J.S., Morrone, A., Johung, K., Goodwin, E.C., Kleijer, W.J., DiMaio, D., Hwang, E.S., 2006. Senescence-associated β -galactosidase is lysosomal β -galactosidase. *Aging Cell* 5, 187-195.
- Leung, B.P., Sattar, N., Crilly, A., Prach, M., McCarey, D.W., Payne, H., Madhok, R., Campbell, C., Gracie, J.A., Liew, F.Y., 2003. A novel anti-inflammatory role for simvastatin in inflammatory arthritis. *The Journal of Immunology* 170, 1524-1530.
- Littarru, G.P., Langsjoen, P., 2007. Coenzyme Q10 and statins: biochemical and clinical implications. *Mitochondrion* 7, S168-S174.
- Liu, S., Uppal, H., Demaria, M., Desprez, P.-Y., Campisi, J., Kapahi, P., 2015. Simvastatin suppresses breast cancer cell proliferation induced by senescent cells. *Scientific Reports* 5, 17895.
- Liu, Y., Sanoff, H.K., Cho, H., Burd, C.E., Torrice, C., Ibrahim, J.G., Thomas, N.E., Sharpless, N.E., 2009. Expression of p16INK4a in peripheral blood T-cells is a biomarker of human aging. *Aging Cell* 8, 439-448.
- Lusis, A.J., 2000. Atherosclerosis. *Nature* 407, 233-241.
- Mabuchi, H., Nohara, A., Kobayashi, J., Kawashiri, M.-a., Katsuda, S., Inazu, A., Koizumi, J., Group, H.L.R., 2007. Effects of CoQ10 supplementation on plasma lipoprotein lipid, CoQ10 and liver and muscle enzyme levels in hypercholesterolemic patients treated with atorvastatin: a randomized double-blind study. *Atherosclerosis* 195, e182-e189.
- Maciejowski, J., de Lange, T., 2017. Telomeres in cancer: tumour suppression and genome instability. *Nature reviews. Molecular Cell Biology* 18, 175-186.
- Makarov, S.S., 2000. NF-kappaB as a therapeutic target in chronic inflammation: recent advances. *Molecular Medicine Today* 6, 441-448.
- Marchand, W.R., Lee, J.N., Suchy, Y., Garn, C., Johnson, S., Wood, N., Chelune, G., 2011. Age-related changes of the functional architecture of the cortico-basal ganglia circuitry during motor task execution. *Neuroimage* 55, 194-203.
- Marrone, G., Russo, L., Rosado, E., Hide, D., García-Cardena, G., García-Pagán, J.C., Bosch, J., Gracia-Sancho, J., 2013. The transcription factor KLF2 mediates hepatic endothelial protection and paracrine endothelial-stellate cell deactivation induced by statins. *Journal of Hepatology* 58, 98-103.
- Martin-Montalvo, A., Sun, Y., Diaz-Ruiz, A., Ali, A., Gutierrez, V., Palacios, H.H., Curtis, J., Siendones, E., Ariza, J., Abulwerdi, G.A., 2016. Cytochrome b 5 reductase and the control of lipid metabolism and healthspan. *NPJ Aging and Mechanisms of Disease* 2, 16006.
- Martin, J.A., Buckwalter, J.A., 2002. Aging, articular cartilage chondrocyte senescence and osteoarthritis. *Biogerontology* 3, 257-264.
- Martínez, P., Blasco, M.A., 2017. Telomere-driven diseases and telomere-targeting therapies. *Journal of Cell Biology* 216, 875-887.
- McCord, R.P., Nazario-Toole, A., Zhang, H., Chines, P.S., Zhan, Y., Erdos, M.R., Collins, F.S., Dekker, J., Cao, K., 2013. Correlated alterations in genome organization, histone methylation, and DNA-lamin A/C interactions in Hutchinson-Gilford progeria syndrome. *Genome Research* 23, 260-269.
- Mehta, J.L., Bursac, Z., Hauer-Jensen, M., Fort, C., Fink, L.M., 2006. Comparison of mortality rates in statin users versus nonstatin users in a United States veteran population. *The American Journal of Cardiology* 98, 923-928.
- Menghini, R., Casagrande, V., Cardellini, M., Martelli, E., Terrinoni, A., Amati, F., Vasa-Nicotera, M., Ippoliti, A., Novelli, G., Melino, G., 2009. MicroRNA 217 modulates

- endothelial cell senescence via silent information regulator 1. *Circulation Research* 120, 1524-1532.
- Mercken, E.M., Hu, J., Krzysik-Walker, S., Wei, M., Li, Y., McBurney, M.W., de Cabo, R., Longo, V.D., 2014. SIRT 1 but not its increased expression is essential for lifespan extension in caloric-restricted mice. *Aging Cell* 13, 193-196.
- Miles, M.V., Horn, P.S., Tang, P.H., Morrison, J.A., Miles, L., DeGrauw, T., Pesce, A.J., 2004. Age-related changes in plasma coenzyme Q10 concentrations and redox state in apparently healthy children and adults. *Clinica Chimica Acta* 347, 139-144.
- Minamino, T., Komuro, I., 2007. Vascular cell senescence: contribution to atherosclerosis. *Circulation Research* 100, 15-26.
- Minamino, T., Miyauchi, H., Yoshida, T., Tateno, K., Kunieda, T., Komuro, I., 2004. Vascular cell senescence and vascular aging. *Journal of Molecular and Cellular Cardiology* 36, 175-183.
- Mirzayans, R., Andrais, B., Hansen, G., Murray, D., 2012. Role of in Replicative Senescence and DNA Damage-Induced Premature Senescence in p53-Deficient Human Cells. *Biochemistry Research International* 2012, 951574.
- Mitani, H., Ishizaka, N., Aizawa, T., Ohno, M., Usui, S.-i., Suzuki, T., Amaki, T., Mori, I., Nakamura, Y., Sato, M., 2002. In vivo klotho gene transfer ameliorates angiotensin II-induced renal damage. *Hypertension* 39, 838-843.
- Mitchell, J.R., Wood, E., Collins, K., 1999. A telomerase component is defective in the human disease dyskeratosis congenita. *Nature* 402, 551-555.
- Morris, B.J., Willcox, D.C., Donlon, T.A., Willcox, B.J., 2015. FOXO3: a major gene for human longevity-a mini-review. *Gerontology* 61, 515-525.
- Moutzouri, E., Liberopoulos, E., Mikhailidis, D., Kostapanos, M., Kei, A., Milionis, H., Elisaf, M., 2011. Comparison of the effects of simvastatin vs. rosuvastatin vs. simvastatin/ezetimibe on parameters of insulin resistance. *International Journal of Clinical Practice* 65, 1141-1148.
- Musgrove, C., Jansson, L.I., Stone, M.D., 2018. New perspectives on telomerase RNA structure and function. *Wiley Interdisciplinary Reviews RNA* 9, 1-15.
- Nagai, T., Yamada, K., Kim, H.-C., Kim, Y.-S., Noda, Y., Imura, A., Nabeshima, Y.-i., Nabeshima, T., 2003. Cognition impairment in the genetic model of aging klotho gene mutant mice: a role of oxidative stress. *The FASEB Journal* 17, 50-52.
- Nakamura, H., Arakawa, K., Itakura, H., Kitabatake, A., Goto, Y., Toyota, T., Nakaya, N., Nishimoto, S., Muranaka, M., Yamamoto, A., 2006. Primary prevention of cardiovascular disease with pravastatin in Japan (MEGA Study): a prospective randomised controlled trial. *The Lancet* 368, 1155-1163.
- Naples, M., Federico, L.M., Xu, E., Nelken, J., Adeli, K., 2008. Effect of rosuvastatin on insulin sensitivity in an animal model of insulin resistance: evidence for statin-induced hepatic insulin sensitization. *Atherosclerosis* 198, 94-103.
- Narita, M., Nuñez, S., Heard, E., Narita, M., Lin, A.W., Hearn, S.A., Spector, D.L., Hannon, G.J., Lowe, S.W., 2003. Rb-mediated heterochromatin formation and silencing of E2F target genes during cellular senescence. *Cell* 113, 703-716.
- Narumiya, H., Sasaki, S., Kuwahara, N., Irie, H., Kusaba, T., Kameyama, H., Tamagaki, K., Hatta, T., Takeda, K., Matsubara, H., 2004. HMG-CoA reductase inhibitors up-regulate anti-aging klotho mRNA via RhoA inactivation in IMCD3 cells. *Cardiovascular Research* 64, 331-336.

- Navas, P., Villalba, J.M., de Cabo, R., 2007. The importance of plasma membrane coenzyme Q in aging and stress responses. *Mitochondrion* 7, S34-S40.
- Nielsen, S.F., Nordestgaard, B.G., Bojesen, S.E., 2012. Statin use and reduced cancer-related mortality. *New England Journal of Medicine* 367, 1792-1802.
- Niswender, C.M., Ishihara, R.W., Judge, L.M., Zhang, C., Shokat, K.M., McKnight, G.S., 2002. Protein engineering of protein kinase A catalytic subunits results in the acquisition of novel inhibitor sensitivity. *Journal of Biological Chemistry* 277, 28916-28922.
- Olovnikov, A.M., 1973. A theory of marginotomy: the incomplete copying of template margin in enzymic synthesis of polynucleotides and biological significance of the phenomenon. *Journal of Theoretical Biology* 41, 181-190.
- Orlandi, A., Bochaton-Piallat, M.-L., Gabbiani, G., Spagnoli, L.G., 2006. Aging, smooth muscle cells and vascular pathobiology: implications for atherosclerosis. *Atherosclerosis* 188, 221-230.
- Ota, H., Eto, M., Kano, M.R., Kahyo, T., Setou, M., Ogawa, S., Iijima, K., Akishita, M., Ouchi, Y., 2010. Induction of endothelial nitric oxide synthase, SIRT1, and catalase by statins inhibits endothelial senescence through the Akt pathway. *Arteriosclerosis, Thrombosis and Vascular Biology* 30, 2205-2211.
- Owens, C.D., 2012. Statins and other agents for vascular inflammation. *Journal of Vascular Surgery* 56, 1799-1806.
- Paolisso, G., Barbagallo, M., Petrella, G., Ragno, E., Barbieri, M., Giordano, M., Varricchio, M., 2000. Effects of simvastatin and atorvastatin administration on insulin resistance and respiratory quotient in aged dyslipidemic non-insulin dependent diabetic patients. *Atherosclerosis* 150, 121-127.
- Potente, M., Dimmeler, S., 2008a. Emerging roles of SIRT1 in vascular endothelial homeostasis. *Cell Cycle* 7, 2117-2122.
- Potente, M., Dimmeler, S., 2008b. NO targets SIRT1: a novel signaling network in endothelial senescence. *Arteriosclerosis Thrombosis and Vascular Biology* 28, 1577-1579.
- Powers, R.W., Kaeberlein, M., Caldwell, S.D., Kennedy, B.K., Fields, S., 2006. Extension of chronological life span in yeast by decreased TOR pathway signaling. *Genes & Development* 20, 174-184.
- Poynter, M.E., Daynes, R.A., 1998. Peroxisome proliferator-activated receptor α activation modulates cellular redox status, represses nuclear factor- κ B signaling, and reduces inflammatory cytokine production in aging. *Journal of Biological Chemistry* 273, 32833-32841.
- Puurunen, J., Piltonen, T., Puukka, K., Ruokonen, A., Savolainen, M.J., Bloigu, R., Morin-Papunen, L., Tapanainen, J.S., 2013. Statin therapy worsens insulin sensitivity in women with polycystic ovary syndrome (PCOS): a prospective, randomized, double-blind, placebo-controlled study. *The Journal of Clinical Endocrinology & Metabolism* 98, 4798-4807.
- Richardson, P.D., Davies, M., Born, G., 1989. Influence of plaque configuration and stress distribution on fissuring of coronary atherosclerotic plaques. *The Lancet* 334, 941-944.
- Sabbatini, M., Pisani, A., Uccello, F., Serio, V., Serù, R., Paternò, R., Cianciaruso, B., Fuiano, G., Andreucci, M., 2004. Atorvastatin improves the course of ischemic acute renal failure in aging rats. *Journal of the American Society of Nephrology* 15, 901-909.
- Sahebkar, A., Kotani, K., Serban, C., Ursoniu, S., Mikhailidis, D.P., Jones, S.R., Ray, K.K., Blaha, M.J., Rysz, J., Toth, P.P., 2015. Statin therapy reduces plasma endothelin-1

- concentrations: A meta-analysis of 15 randomized controlled trials. *Atherosclerosis* 241, 433-442.
- Saito, Y., Nakamura, T., Ohyama, Y., Suzuki, T., Iida, A., Shiraki-Iida, T., Kuro-o, M., Nabeshima, Y.-i., Kurabayashi, M., Nagai, R., 2000. In vivo klotho gene delivery protects against endothelial dysfunction in multiple risk factor syndrome. *Biochemical and Biophysical Research Communications* 276, 767-772.
- Saliques, S., Teyssier, J.-R., Vergely, C., Lorgis, L., Lorin, J., Farnier, M., Donzel, A., Sicard, P., Berchoud, J., Lagrost, A.-C., 2011. Circulating leukocyte telomere length and oxidative stress: a new target for statin therapy. *Atherosclerosis* 219, 753-760.
- Saliques, S., Zeller, M., Lorin, J., Lorgis, L., Teyssier, J.-R., Cottin, Y., Rochette, L., Vergely, C., 2010. Telomere length and cardiovascular disease. *Archives of Cardiovascular Diseases* 103, 454-459.
- Satoh, M., Ishikawa, Y., Takahashi, Y., Itoh, T., Minami, Y., Nakamura, M., 2008. Association between oxidative DNA damage and telomere shortening in circulating endothelial progenitor cells obtained from metabolic syndrome patients with coronary artery disease. *Atherosclerosis* 198, 347-353.
- Satoh, M., Minami, Y., Takahashi, Y., Tabuchi, T., Itoh, T., Nakamura, M., 2009. Effect of intensive lipid-lowering therapy on telomere erosion in endothelial progenitor cells obtained from patients with coronary artery disease. *Clinical Science* 116, 827-835.
- Satyanarayana, A., Wiemann, S., Buer, J., Lauber, J., Dittmar, K., Wüstefeld, T., Blasco, M., Manns, M., Rudolph, K., 2003. Telomere shortening impairs organ regeneration by inhibiting cell cycle re-entry of a subpopulation of cells. *The EMBO Journal* 22, 4003-4013.
- Schachter, M., 2005. Chemical, pharmacokinetic and pharmacodynamic properties of statins: an update. *Fundamental & Clinical Pharmacology* 19, 117-125.
- Sen, P., Shah, P.P., Nativio, R., Berger, S.L., 2016. Epigenetic mechanisms of longevity and aging. *Cell* 166, 822-839.
- Serban, C., Sahebkar, A., Ursoniu, S., Mikhailidis, D.P., Rizzo, M., Lip, G.Y., Hovingh, G.K., Kastelein, J.J., Kalinowski, L., Rysz, J., 2015. A systematic review and meta-analysis of the effect of statins on plasma asymmetric dimethylarginine concentrations. *Scientific Reports* 5, 9902.
- Serrano, A.L., Andrés, V., 2004. Telomeres and cardiovascular disease: does size matter? *Circulation Research* 94, 575-584.
- Sfeir, A., De Lange, T., 2012. Removal of shelterin reveals the telomere end-protection problem. *Science* 336, 593-597.
- Shang, F., Zhao, L., Zheng, Q., Wang, J., Xu, Z., Liang, W., Liu, H., Liu, S., Zhang, L., 2006. Simvastatin inhibits lipopolysaccharide-induced tumor necrosis factor- α expression in neonatal rat cardiomyocytes: The role of reactive oxygen species. *Biochemical and Biophysical Research Communications* 351, 947-952.
- Sharpless, N.E., DePinho, R.A., 2007. How stem cells age and why this makes us grow old. *Nature Reviews Molecular Cell Biology* 8, 703-713.
- Shen, M., Kawamoto, T., Yan, W., Nakamasu, K., Tamagami, M., Koyano, Y., Noshiro, M., Kato, Y., 1997. Molecular characterization of the novel basic helix-loop-helix protein DEC1 expressed in differentiated human embryo chondrocytes. *Biochemical and Biophysical Research Communications* 236, 294-298.
- Steppan, J., Tran, H., Benjo, A.M., Pellakuru, L., Barodka, V., Ryoo, S., Nyhan, S.M., Lussman, C., Gupta, G., White, A.R., 2012. Alagebrium in combination with exercise

- ameliorates age-associated ventricular and vascular stiffness. *Experimental Gerontology* 47, 565-572.
- Strazhesko, I.D., Tkacheva, O.N., Akasheva, D.U., Dudinskaya, E.N., Plokhova, E.V., Pykhtina, V.S., Kruglikova, A.S., Kokshagina, N.V., Sharashkina, N.V., Agaltsov, M.V., 2016. Atorvastatin therapy modulates telomerase activity in patients free of atherosclerotic cardiovascular diseases. *Frontiers in Pharmacology* 7, 347.
- Strazzullo, P., Kerry, S.M., Barbato, A., Versiero, M., D'Elia, L., Cappuccio, F.P., 2007. Do statins reduce blood pressure? A meta-analysis of randomized, controlled trials. *Hypertension* 49, 792-798.
- Sugiyama, S., Fukushima, H., Kugiyama, K., Maruyoshi, H., Kojima, S., Funahashi, T., Sakamoto, T., Horibata, Y., Watanabe, K., Koga, H., 2007. Pravastatin improved glucose metabolism associated with increasing plasma adiponectin in patients with impaired glucose tolerance and coronary artery disease. *Atherosclerosis* 194, e43-e51.
- Suh, Y., Atzmon, G., Cho, M.-O., Hwang, D., Liu, B., Leahy, D.J., Barzilai, N., Cohen, P., 2008. Functionally significant insulin-like growth factor I receptor mutations in centenarians. *Proceedings of the National Academy of Sciences* 105, 3438-3442.
- Sun, I., Sun, E., Crane, F., Morre, D., Lindgren, A., Löw, H., 1992. Requirement for coenzyme Q in plasma membrane electron transport. *Proceedings of the National Academy of Sciences* 89, 11126-11130.
- Sverdrup, F.M., Yates, M.P., Vickery, L.E., Klover, J.A., Song, L.R.-H., Anglin, C.P., Misko, T.P., 2010. Protein geranylgeranylation controls collagenase expression in osteoarthritic cartilage. *Osteoarthritis and Cartilage* 18, 948-955.
- Szendroedi, J., Anderwald, C., Krssak, M., Bayerle-Eder, M., Esterbauer, H., Pfeiler, G., Brehm, A., Nowotny, P., Hofer, A., Waldhäusl, W., 2009. Effects of high-dose simvastatin therapy on glucose metabolism and ectopic lipid deposition in nonobese type 2 diabetic patients. *Diabetes Care* 32, 209-214.
- Takemoto, M., Liao, J.K., 2001. Pleiotropic effects of 3-hydroxy-3-methylglutaryl coenzyme a reductase inhibitors. *Arteriosclerosis, Thrombosis, and Vascular Biology* 21, 1712-1719.
- Tatar, M., Kopelman, A., Epstein, D., Tu, M.-P., Yin, C.-M., Garofalo, R., 2001. A mutant *Drosophila* insulin receptor homolog that extends life-span and impairs neuroendocrine function. *Science* 292, 107-110.
- Thunyakitpisal, P.D., Chaisuparat, R., 2004. Simvastatin, an HMG-CoA reductase inhibitor, reduced the expression of matrix metalloproteinase-9 (Gelatinase B) in osteoblastic cells and HT1080 fibrosarcoma cells. *Journal of Pharmacological Sciences* 94, 403-409.
- Tian, G., Sawashita, J., Kubo, H., Nishio, S.-y., Hashimoto, S., Suzuki, N., Yoshimura, H., Tsuruoka, M., Wang, Y., Liu, Y., 2014. Ubiquinol-10 supplementation activates mitochondria functions to decelerate senescence in senescence-accelerated mice. *Antioxidants & Redox Signaling* 20, 2606-2620.
- Unit, E.S., 2005. Efficacy and safety of cholesterol-lowering treatment: prospective meta-analysis of data from 90 056 participants in 14 randomised trials of statins. *Lancet* 366, 1267-1278.
- Van Aelst, L., D'Souza-Schorey, C., 1997. Rho GTPases and signaling networks. *Genes & Development* 11, 2295-2322.

- Varela, I., Pereira, S., Ugalde, A.P., Navarro, C.L., Suárez, M.F., Cau, P., Cadinanos, J., Osorio, F.G., Foray, N., Cobo, J., 2008. Combined treatment with statins and aminobisphosphonates extends longevity in a mouse model of human premature aging. *Nature Medicine* 14, 767-772.
- Verdun, R.E., Karlseder, J., 2007. Replication and protection of telomeres. *Nature* 447, 924-931.
- Wassmann, S., Laufs, U., Bäumer, A.T., Müller, K., Ahlbory, K., Linz, W., Itter, G., Rösen, R., Böhm, M., Nickenig, G., 2001. HMG-CoA reductase inhibitors improve endothelial dysfunction in normocholesterolemic hypertension via reduced production of reactive oxygen species. *Hypertension* 37, 1450-1457.
- Werner, N., Nickenig, G., Laufs, U., 2002. Pleiotropic effects of HMG-CoA reductase inhibitors. *Basic Research in Cardiology* 97, 105-116.
- Wilson, W., Evans, J., Bell, P., Thompson, M., 2005. HMG-CoA reductase inhibitors (statins) decrease MMP-3 and MMP-9 concentrations in abdominal aortic aneurysms. *European Journal of Vascular and Endovascular Surgery* 30, 259-262.
- Wlaschek, M., Ma, W., Jansen-Dürr, P., Scharffetter-Kochanek, K., 2003. Photoaging as a consequence of natural and therapeutic ultraviolet irradiation—studies on PUVA-induced senescence-like growth arrest of human dermal fibroblasts. *Experimental Gerontology* 38, 1265-1270.
- Yang, G., Caldwell, R.B., Yao, L., Romero, M.J., Toque, H.A., Caldwell, R.W., 2010. The role of RhoA/Rho kinase pathway in endothelial dysfunction. *Journal of Cardiovascular Disease Research* 1, 165-170.
- Yoon, H.E., Lim, S.W., Piao, S.G., Song, J.-H., Kim, J., Yang, C.W., 2012. Statin upregulates the expression of klotho, an anti-aging gene, in experimental cyclosporine nephropathy. *Nephron Experimental Nephrology* 120, e123-e133.
- Young, A.R., Narita, M., 2009. SASP reflects senescence. *EMBO Reports* 10, 228-230.
- Yudoh, K., Karasawa, R., 2010. Statin prevents chondrocyte aging and degeneration of articular cartilage in osteoarthritis (OA). *Aging (Albany NY)* 2, 990-998.
- Zhang, F.L., Casey, P.J., 1996. Protein prenylation: molecular mechanisms and functional consequences. *Annual Review of Biochemistry* 65, 241-269.
- Zhang, T., Kraus, W.L., 2010. SIRT1-dependent regulation of chromatin and transcription: linking NAD⁺ metabolism and signaling to the control of cellular functions. *Biochimica et Biophysica Acta (BBA)-Proteins and Proteomics* 1804, 1666-1675.
- Zhang, Z.G., Zhang, L., Jiang, Q., Chopp, M., 2002. Bone marrow-derived endothelial progenitor cells participate in cerebral neovascularization after focal cerebral ischemia in the adult mouse. *Circulation Research* 90, 284-288.
- Zhu, J., Wang, H., Bishop, J.M., Blackburn, E.H., 1999. Telomerase extends the lifespan of virus-transformed human cells without net telomere lengthening. *Proceedings of the National Academy of Sciences* 96, 3723-3728.