



Efficacy of access flap procedures compared to subgingival debridement in the treatment of periodontitis. A systematic review and meta-analysis

Ignacio Sanz-Sánchez¹ | Eduardo Montero¹ | Filippo Citterio² | Federica Romano² | Ana Molina¹ | Mario Aimetti²

¹Etiology and Therapy of Periodontal Diseases (ETEP) Research Group, University Complutense, Madrid, Spain

²Department of Surgical Sciences, C.I.R. Dental School, University of Turin, Turin, Italy

Correspondence

Ignacio Sanz Sánchez, Facultad de Odontología, Universidad Complutense de Madrid, Plaza Ramón y Cajal s/n. 28040 Madrid, Spain.
Email: ignaciosanz@ucm.es

Funding information

This systematic review was self-funded.

Abstract

Aim: This systematic review aimed to answer the following focused questions: (a) “In patients with periodontitis, how effective are access flaps (AFs) as compared to subgingival debridement in attaining probing depth (PD) reduction?” and (b) “In patients with periodontitis, does the type of AF impact PD reduction?”.

Material and Methods: Randomized clinical trials were searched in three databases. Besides PD, information concerning clinical attachment level (CAL) and other relevant outcomes was also collected. Meta-analyses were performed whenever possible and results were categorized based on the initial PD.

Results: Thirty-six publications were included. AFs resulted in a significantly greater PD reduction in deep pockets (>6 mm or ≥6 mm), as compared to subgingival debridement, in short- ($n = 4$; weighted mean difference [WMD] = 0.67 mm; 95% confidence interval [CI] 0.37,0.97; $p < .001$) and long-term studies ($n = 4$; WMD = 0.39 mm; 95% CI 0.09,0.70; $p = .012$), while in moderately deep pockets (4–6, 5–6 or 4–5 mm) only in short-term studies ($n = 4$; WMD = 0.34; 95% CI 0.21,0.46; $p < .001$). In shallow pockets (1–3 or 1–4 mm), AFs led to greater CAL ($n = 7$; WMD = -0.43 mm; 95% CI -0.56, -0.28; $p < .001$). There was not enough evidence to answer question PICO 2.

Conclusions: AFs resulted in greater PD reduction in the treatment of deep and moderate pockets.

KEYWORDS

clinical attachment level, periodontal debridement, periodontitis, probing depth, systematic review

1 | INTRODUCTION

The role of subgingival biofilm in the initiation and progression of periodontal diseases has been clearly established (Roberts & Darveau, 2015). Thus, adequate individual plaque control practices, and professional elimination of supra- and subgingival plaque and calculus are essential for the successful treatment of periodontitis (Graziani, Karapetsa, Alonso, & Herrera, 2017). In this context,

cause-related periodontal therapy aims to mechanically remove subgingival biofilm and to control inflammation, either with non-surgical or surgical approaches, in order to arrest further attachment loss by reducing probing depth (PD), as sites with $PD \leq 4$ mm are associated with lower risk of disease progression and tooth loss (Matuliene et al., 2008).

Non-surgical periodontal therapy consists on subgingival debridement. Traditionally, this approach has been done by means of

subgingival scaling and root planing (SRP). Although, theoretically, these are well-differentiated procedures, they are performed clinically at the same time (Aimetti, 2014; Graziani et al., 2017; Laleman et al., 2017).

A consistent amount of evidence has indicated that subgingival debridement is effective in reducing bleeding on probing (BOP) and PD, and gaining clinical attachment level (CAL) (Sanz, Alonso, Carasol, Herrera, & Sanz, 2012; Smiley et al., 2015; Van der Weijden & Timmerman, 2002). However, it is technically demanding, and complete calculus removal is difficult to achieve (Jepsen, Deschner, Braun, Schwarz, & Eberhard, 2011). Indeed, in vitro studies have shown that after therapy 3%–80% of instrumented root surfaces harboured some residual calculus (Buchanan & Robertson, 1987; Caffesse, Sweeney, & Smith, 1986; Geisinger, Mealey, Schoolfield, & Mellonig, 2007; Rabbani, Ash, & Caffesse, 1981). More hard deposits are generally detected in deep sites (PD > 5 mm) and at molars (particularly in furcation areas), than at single-rooted teeth (Caffesse et al., 1986; Rateitschak-Pluss, Schwarz, Guggenheim, Duggelin, & Rateitschak, 1992; Waerhaug, 1978), what limits the efficacy of closed subgingival debridement as definitive therapy (Heitz-Mayfield, Trombelli, Heitz, Needleman, & Moles, 2002; Serino, Rosling, Ramberg, Socransky, & Lindhe, 2001).

Access flaps were introduced to improve the efficiency of subgingival debridement by gaining direct access to the root surface, root concavities and furcations in sites with residual pockets irrespective of the pattern of bone resorption. In these conservative surgeries, there is no active removal of alveolar bone and no or minimal resection of soft tissues, as the treatment strategy focuses on enhancing the cleaning of the colonized root surface to achieve shallower pockets, greater CAL gain and lower levels of BOP. Graziani Karapetsa Mardas Leow and Donos (2018) classified conservative surgical procedures into open flap debridement (Kirkland, 1931), minimally resective flaps (e.g. the modified Widman flap [MWF]; Ramfjord & Nissle, 1974) and flaps aimed at conserving interdental soft tissues (e.g. papilla preservation techniques [Checchi & Schonfeld, 1988; Dello Russo, 1981; Michaelides & Wilson, 1996; Takei, Han, Carranza, Kenney, & Lekovic, 1985]). The first two approaches achieve a reduction in PD by a combination of gingival recession and CAL gain along time. In the minimally resective flaps, a marginal soft tissue resection is also performed in order to eliminate the epithelium of the periodontal pocket and the inflamed tissue from the inner part of the flap. On the contrary, papilla preservation techniques are particularly indicated in the treatment of intra-bony defects with the objective of maximizing CAL gain by enhancing blood clot stability and, thus, reducing tissue contraction. As reported in a previous systematic review, the magnitude of clinical improvement may be related to the surgical flap design (Graziani et al., 2012). In this sense, greater CAL gain and less recession were observed when papilla preservation techniques were applied in the surgical treatment of intra-bony defects. Moreover, the adjunctive use of magnification and microsurgical instruments may also improve visibility of subgingival deposits and calculus removal. Currently, there is a paucity of literature on the use of magnification in non-surgical periodontal

Clinical Relevance

Scientific rationale for the study: Severe periodontitis (stages III and IV) is one of the most prevalent non-communicable chronic diseases affecting humans. Since in this clinical situation subgingival debridement may not be enough to control the disease, it is worthy to assess if a surgical access is needed.

Principal findings: The effectiveness of the mode of therapy is going to be dependent on the initial probing depth (PD). Surgery has shown benefits in terms of PD reduction when treating initially moderate and deep pockets. When combining the results for PD reduction and clinical attachment level (CAL) gain, subgingival debridement seems to be the choice at moderately deep pockets and access surgery at initially deep pockets.

Practical implications: In order to attain proper pocket reduction, access flaps might be more efficacious when aimed to treat initially deep pockets. However, subgingival debridement provides higher CAL gain in moderately deep pockets and less clinical attachment loss in shallow sites. Long-term results seem to be related with other factors different than the type of debridement provided (surgical or non-surgical), such as hygiene measures or maintenance protocols, as differences between therapies tend to disappear throughout time.

treatment (Nibali, Pometti, Chen, & Tu, 2015; Nibali, Yeh, Pometti, & Tu, 2018; Ribeiro et al., 2011).

Ideally, the least invasive treatment with the better biological cost-effectiveness ratio should be used to restore periodontal health. Furthermore, patients' perspective of therapy may differ from traditional clinical endpoints. Assessment of pain and discomfort experienced during and after therapy should complement conventional clinical measures (Hujoel, 2004).

Therefore, the aim of the present systematic review was to answer the following two PICO questions: "In patients with periodontitis (population), how effective are access flaps (intervention) as compared to subgingival debridement (comparison) in attaining PD reduction (primary outcome)?" (PICO 1) and "In patients with periodontitis (population), does the type of access flaps (intervention and control) impact PD reduction (primary outcome)?" (PICO 2).

2 | MATERIAL AND METHODS

2.1 | Protocol development and focused question

The protocol of this systematic review was made a priori, agreed upon all authors and registered in the PROSPERO International Prospective Register of Systematic Reviews hosted by the Centre for

Reviews and Dissemination, University of York, National Institute for Health Research (United Kingdom; CRD42019123077). This systematic review was reported according to the PRISMA (Preferred Reporting Items for Systematic Review and Meta-Analyses) statement (Moher, Liberati, Tetzlaff, & Altman, 2009).

2.1.1 | Eligibility criteria

Inclusion criteria (PICOS)

- Population: patients older than 18 with chronic or aggressive periodontitis based on the criteria described by Armitage (1999);
- Interventions: any type of access flap (PICO 1) or any access flap different to conventional MWF (PICO 2);
- Comparisons: subgingival debridement (PICO 1) or conventional MWF (PICO 2);
- Outcomes: changes in PD;
- Study design: randomized controlled clinical trials (RCTs) with at least 6 months of follow-up and a minimum of 10 patients per treatment group. Only RCTs were included to be able to make direct comparisons and, if possible, to be included in the meta-analyses.

Exclusion criteria

- Studies aiming at apically position the flaps;
- Studies using resective approaches;
- Studies aiming at using regenerative technologies in intra-osseous defects;
- Studies focusing only on furcated teeth.

2.1.2 | Type of intervention and comparisons

Studies were selected that included interventions for cause-related periodontal therapy. To answer question PICO 1, interventions in the test group included any access flap such as traditional open flap debridement (OFD; Kirkland, 1931), MWF or microsurgical access procedures, whereas interventions in the control group included subgingival debridement with or without adjunctive therapies, irrespective of the instrument used for the mechanical removal of biofilm. To answer question PICO 2, intervention in the test group included any AF different to conventional MWF, whereas interventions in the control group included conventional MWF.

2.1.3 | Type of outcomes

The primary outcome for assessing the efficacy of the treatment of periodontitis was PD reduction.

The following secondary outcomes were studied:

- CAL change,
- Changes in gingival or bleeding indices and plaque indices,

- Tooth loss,
- Mean number of residual pockets (PD > 3 mm and/or PD > 3 mm with BOP or PD > 4 mm and/or PD > 4 mm with BOP) at the end of therapy,
- Percentage of closed pockets (PD ≤ 3 mm or PD ≤ 4 mm) out of the total number of pockets at baseline (PD > 3 mm or PD > 4 mm),
- Percentage of cases with need for re-treatment,
- Harms and adverse effects such as swelling, root sensitivity, gingival recession, etc.,
- Patient-reported outcome measures (PROMs), such as pain, discomfort, satisfaction, etc.

2.2 | Information sources and search

2.2.1 | Electronic search

Three electronic databases were used as sources in the search for studies satisfying the inclusion criteria: (a) The National Library of Medicine (MEDLINE via PubMed); (b) Cochrane Central Register of Controlled Trials; and (c) Scopus. These databases were searched for studies published until 8 February 2019. The search was limited to human subjects and to studies reported in English. No more languages were considered due to the limited time for the preparation of this systematic review.

2.2.2 | Manual search

All reference lists of the selected studies and previously published systematic reviews were checked for cross-references. The following journals were hand-searched from year 2008 to 2018: Journal of Clinical Periodontology, Journal of Periodontology, The International Journal of Periodontics and Restorative Dentistry and Journal of Periodontal Research.

2.2.3 | Search strategy

The following search terms were used in PubMed.

Population

[MeSH terms]: (adult periodontitis[mh] OR chronic periodontitis[mh] OR aggressive periodontitis[mh] OR early onset periodontitis[mh] OR juvenile periodontitis[mh])
OR
[Text Words, Title]: periodontitis[tw].

Intervention

[MeSH terms]: surgical flap[mh] OR surgical flaps[mh]
OR
[Text Word, Title]: access flap[tw] OR access flaps[tw] OR open flap debridement[tw] OR open flap debridements[tw] OR modified

Widman flap[tw] OR modified Widman flaps[tw] OR Widman flap[tw] OR Widman flaps[tw] OR Kirkland flap[tw] OR Kirkland flaps[tw].

The search was combined as Population and Intervention

The search strategy applied for the Cochrane was '(adult periodontitis OR chronic periodontitis OR aggressive periodontitis OR early onset periodontitis OR juvenile periodontitis OR periodontitis) AND (surgical flap OR surgical flaps OR access flap OR open flap debridement OR modified Widman flap OR Widman flap OR Kirkland flap) AND (dental scaling OR root scaling OR root scalings OR scaling, dental OR scaling, root OR scaling, subgingival OR scaling, supragingival OR scalings, root OR subgingival scaling OR supragingival scaling OR debridement, nonsurgical periodontal OR debridement, periodontal OR debridement, periodontal pocket OR debridements, nonsurgical periodontal OR debridements, periodontal OR debridements, periodontal pocket OR scaling and root planing OR subgingival debridement) AND (clinical trials, randomized OR controlled clinical trials, randomized OR randomized controlled trial)'.

The search strategy used in SCOPUS was as follows (paradontitis OR parodontitis OR peridontitis OR periodontitis OR "periodontal disease" OR "dental loss" OR "furcation defects" OR "paradontal disease" OR paradontopathy OR paraodontopathy OR parodontopathy OR "parodontal disease" OR "parodontium disease" OR "parodontive tissue disease" OR "peridontal disease" OR "peridontal tissue disease" OR "peridontium disease" OR "periodontal atrophy" OR "periodontal Attachment loss" OR "periodontal diseases" OR "periodontal infection" OR "periodontium disease" OR periodontopathy) AND ("surgical flaps" OR "surgical flap" OR "access flap" OR "access flaps" OR "open flap debridement" OR "modified widman flap" OR "widman flap" OR "kirkland flap" OR "modified widman flaps" OR "widman flaps" OR "kirkland flaps") with the limit to dental and medical journals (LIMIT-TO (SUBJAREA "MEDI") OR LIMIT-TO (SUBJAREA,"DENT"))).

2.3 | Screening methods

Two reviewers (F.C. and A.M.) screened independently the titles and abstracts. The same reviewers selected full manuscripts of studies meeting the inclusion criteria, or those with insufficient data in the title and abstract to make a clear decision. Any disagreement was resolved by discussion with a third reviewer (I.S.). The inter-reviewer reliability (percentage of agreement and kappa correlation coefficient) of the full-text analysis was calculated.

2.4 | Data extraction

Two different reviewers (I.S. and E.M.) performed duplicate data extraction. When data were incomplete or missing, authors of studies were contacted for clarification. If agreement could not be reached, data were excluded until further clarification was available. When the results of a study were published more than once, the data were

used upon necessity. If a study was comparing more than two arms, the data from the groups of interest were extracted.

2.5 | Quality assessment (risk of bias in individual studies)

A quality assessment of the included RCTs was performed according to the Cochrane Collaboration risk of bias tool for randomized trials (Rob2 updated in October 2018) (Higgins et al., 2016) and the CONSORT statement (Moher et al., 2012). Five main quality domains were assessed: randomization process, deviation from the intended intervention including blinding of participants and personnel (performance bias) and blinding of outcome assessment (detection bias), incomplete outcome data (attrition bias), measurement of the outcome and selective outcome reporting (reporting bias). These parameters were rated to be in low risk of bias if all the criteria were met.

2.6 | Risk of bias across studies

The publication bias was evaluated using a Funnel plot and the Egger's linear regression method for the primary outcome. A sensitivity analysis of the meta-analysis results was also performed for this outcome (Tobias & Campbell, 1999).

2.7 | Data analyses

To summarize studies, they were combined in order to perform meta-analyses (MAs) reporting weighted mean differences (WMDs) and 95% confidence intervals (CI) to compare the effect of subgingival debridement and AFs (PICO 1), and of different types of AFs (PICO 2) for the primary and secondary outcomes. For a better interpretation of the clinical effect of each procedure, PD and CAL were reported together. Subgroup analyses were performed on the selected outcome variables using the study follow-up (short [≤ 12 months] or long [> 12 months]) and the initial PD as explanatory variables. Furthermore, the absolute effect of each intervention was reported through weighted mean effects (WMEs) with their respective 95% CI. The statistical heterogeneity among studies was assessed using the Q test based on chi-square statistics (Cochran, 1954) as well as the I^2 index (Higgins, Thompson, Deeks, & Altman, 2003) in order to know the percentage of variation in the global estimate that is attributable to heterogeneity. Study specific estimates were pooled with both the fixed and random-effect models (DerSimonian & Laird, 1986). If a significant heterogeneity was found, then the random-effect model results were presented. A Forest Plot was created to illustrate the effects of the different studies and the global estimation. STATA® (StataCorp LP), and OpenMeta [Analyst] intercooled software was used to perform all analyses. Statistical significance was defined as a p value $< .05$.

3 | RESULTS

3.1 | Search

Figure 1 depicts the flow chart summarizing the results of the search. The search rendered 2,117 titles, which, after evaluating their titles and abstracts, resulted in 56 articles for full-text analysis. After this analysis, 36 articles were included for data extraction, representing 18 investigations, since eight groups of papers reported the results of the same material at different time points, or different outcomes from the same population. Agreement for title and abstract screening was 98.68% ($\kappa = 0.72$; 95% CI [0.62–0.82]) and 91.07% ($\kappa = 0.80$; 95% CI [0.64–0.96]) for full text. The reasons for excluding the remaining studies are detailed in Table S1.

3.2 | Description of selected studies

Table 1 depicts the methodological characteristics of the selected studies. Out of the 18 investigations, 15 RCTs had a split-mouth design and 3 a parallel design (Ribeiro et al., 2011; Serino et al., 2001; Trombelli, Simonelli, Schincaglia, Cucchi, & Farina, 2012). Ten studies compared two different arms and eight investigations three or more treatment approaches. The first PICO question (AFs vs. subgingival debridement) was studied in 13 investigations and the second PICO question (comparison between different AFs) in six RCTs. One study provided data and results for

both PICO questions (Lindhe & Nyman, 1985). There were four investigations evaluating the treatment of localized intra-bony defects (Renvert, Nilvéus, Dahlén, Slots, & Egelberg, 1990; Renvert, Nilveus, & Egelberg, 1985; Ribeiro et al., 2011; Trombelli et al., 2012; Wennström, Wennström, & Lindhe, 1986) and one additional publication reporting the results for intra-bony defects from an investigation treating the whole mouth (Isidor, Attström, & Karring, 1985).

Most of the studies performed subgingival debridement prior to surgery (12 out of 18), as part of the initial periodontal therapy, and received a second round of subgingival debridement in the control group during the surgical phase, except in the study performed in Minnesota, in which quadrants assigned to the control group received only one round of subgingival debridement (Pihlstrom, McHuon, Oliphant, & Ortiz-Campos, 1983; Pihlstrom, Oliphant, & McHugh, 1984; Pihlstrom, Ortiz-Campos, & McHugh, 1981). In the remaining six investigations, subgingival debridement was performed as part of the surgical phase, but not before (Lindhe & Nyman, 1985; Lindhe et al., 1982; Polansky, Haas, Lorenzoni, Wimmer, & Pertl, 2003; Serino et al., 2001; Wennström et al., 1986; Westfelt et al., 1985).

Some studies presented the results in relation to the initial PD. While most of the studies that categorized initial pockets used the ranges 1–3, 4–6 and >6 mm (Becker et al., 1988; Hill et al., 1981; Lindhe & Nyman, 1985; Lindhe et al., 1982; Pihlstrom et al., 1981; Westfelt et al., 1985), the study from Kaldahl, Kalkwarf, Patil, Molvar, and Dyer (1996) subgrouped for 1–4, 5–6 and >6 mm initial pockets, and the study from Serino et al. (2001) presented data for 1–3, 4–5 and >5 mm initial pockets.




FIGURE 1 Flow chart depicting the article selection process [Colour figure can be viewed at wileyonlinelibrary.com]

TABLE 1 Methodological characteristics of the included studies

Authors/Year	Study Design and PICO question	Number of patients baseline (final) Age range (mean)	Longer follow-up reported (in parenthesis, previous follow-ups reported)	Intervention	Other interventions not included in this SR	Outcomes	Setting, location and funding	Comments (Teeth included in the study, data subgrouping by PD or CAL, other study characteristics)
Lindhe et al. (1982a,b; 1984)	RCT Split-mouth PICO 1	15 (11) 32–57 (47.9) years	60 months (24 months)	Initial therapy: OHI Test: MWF Control: SRP Maintenance protocol: OHI with supra and subgingival debridement every 2 weeks the first 6 months, every 3 months from 6 to 24 months. 4–6 months supragingival cleaning from 24 to 60 months	None	PI GI Δ PD Δ CAL	University, Sweden, supported by governmental research funding	All teeth included PD subgrouping: Initial PD 1–3 mm Initial PD 4–6 mm Initial PD > 6 mm non-molars 5-years data: frequency distributions of Δ CAL in patients with plaque score ≤10% and ≥50%
Pihlstrom et al. (1981, 1983, 1984)	RCT Split-mouth PICO 1	17 (10) 22–59 (43) years	78 months (48 months)	Initial therapy: OHI + SRP Test: MWF Control: no further therapy Maintenance protocol: OHI with supra- and subgingival instrumentation every 3–4 months	None	PI CI GI Δ PD Δ CAL	University, USA, funding not reported	All teeth included PD subgrouping: Initial PD 1–3 mm Initial PD 4–6 mm Initial PD > 6 mm
Svoboda et al. (1984)	RCT Split-mouth PICO 2	12 (12) 34–60 (44) years	6 months	Initial Therapy: OHI + SRP + OA (when needed) Test: OFD Control: MWF Maintenance protocol: Not specified	None	PI GI Δ PD Δ CAL Δ REC Mobility Rx BL	University, USA, funding not reported	Only anterior teeth included (canine-canine) All type of defects treated without distinction between horizontal or vertical defects
Lindhe and Nyman (1985)	RCT Split-mouth PICOs 1 & 2	15 (15) 42–59 (50) years	12 months	Initial therapy: OHI Test 1: MWF Test 2: OFD Control: SRP Maintenance protocol: prophylaxis every 2 weeks the first 3 months, and every 3 months from 3 to 12 months follow-up	None	PI GI BoP Δ PD Δ CAL Microbial analysis	University, Sweden, supported by governmental research funding	All teeth included PD subgrouping: Initial PD 1–3 mm Initial PD 4–6 mm Initial PD > 6 mm All type of defects treated without distinction between horizontal or vertical defects
Westfelt et al. (1985)	RCT Split-mouth PICO 1	16 (16) 35–65 years	6 months	Initial therapy: OHI Test: MWF Control: SRP Maintenance protocol: prophylaxis every 2 weeks	Test 2: Gingivectomy Test 3: AF Test 4: AF with bone recontouring Test 5: MWF with bone recontouring	PI BoP Δ PD Δ CAL	University, Sweden, supported by governmental research funding	All teeth included PD subgrouping: Initial PD 1–3 mm Initial PD 4–6 mm Initial PD > 6 mm

(Continues)

TABLE 1 (Continued)

Authors/Year	Study Design and PICO question	Number of patients baseline (final) Age range (mean)	Longer follow-up reported (in parenthesis, previous follow-ups reported)	Intervention	Other interventions not included in this SR	Outcomes	Setting, location and funding	Comments (Teeth included in the study, data subgrouping by PD or CAL, other study characteristics)
Isidor et al. (1984, 1985) and Isidor and Karring (1986)	RCT Split-mouth PICO 1	17 (16) 28–52 (40) years	60 months (6 and 12 months)	Initial therapy: OHI + SRP (without LA) Test: MWF Control: SRP (with LA) Maintenance protocol: prophylaxis every 2 weeks	Test 2: Reverse bevel flap	PI GI BoP Δ PD Δ CAL REC RxBL	University, Denmark, funding not reported	Only non-molar teeth included Angular bony defects separately reported (Isidor et al., 1985)
Wennström et al. (1986)	RCT Split-mouth PICO 1	16 (16) 14–29 (19.7) years	60 months	Initial Therapy: OHI Test: MWF Control: SRP (twice in 3 weeks) Maintenance protocol: Professional tooth cleaning every 4 weeks for 6 months. Later, professional tooth cleaning every 3 months for 2 years. During the last 3 years the patients were seen only by their regular dentists	None	PI GI BoP Δ PD Δ CAL Δ REC RxBL Microbial analysis	University, Sweden, supported by governmental research funding	All teeth included Juvenile and post-juvenile periodontitis. Focus on intra-bony defects (PD ≥ 5 mm + presence of intra-bony defects on radiographs)
Renvert, Garrett, et al. (1985), Renvert, et al. (1990)	RCT Split-mouth PICO 1	14 (12) 32–67 (NR) years	60 months (6 months)	Initial Therapy: OHI + SRP Test: OFD Control: SRP Maintenance protocol: After 6 weeks, polishing was performed. From 6 weeks up to 6 months, patients were seen every 6 weeks for tooth cleaning. From 6 to 60 months, tooth cleaning was performed every 6 months	None	PI BoP Δ PD Δ CAL Probing bone level	University, not specified, funding not reported	All teeth included Focus on intra-bony defects (2 separate proximal sites with radiographic intra-bony lesions + PD ≥ 6 mm)
Burgett et al. (1992), Hill et al. (1981), Ramfjord et al. (1987)	RCT Split-mouth PICO 1	90 (72) 24–68 (45) years	60 months (24 months)	Initial therapy: OHI + SRP + OA (OA only when needed) Test: MWF Control: SRP Maintenance protocol: polishing every week for 3–4 weeks, and prophylaxis with re-SRP (when needed) every 3 months	Test 2: Surgical pocket elimination Test 3: Subgingival curettage	PI CI GI Mobility Δ PD Δ CAL Tooth loss	University, USA, supported by governmental research funding	All teeth included PD subgrouping: Initial PD 1–3 mm Initial PD 4–6 mm Initial PD > 6 mm

(Continues)

TABLE 1 (Continued)

Authors/Year	Study Design and PICO question	Number of patients baseline (final) Age range (mean)	Longer follow-up reported (in parenthesis, previous follow-ups reported)	Intervention	Other interventions not included in this SR	Outcomes	Setting, location and funding	Comments (Teeth included in the study, data subgrouping by PD or CAL, other study characteristics)
Kaldahl, Kalkwarf, Patil, & Molvar, (1988, a, b, c), Kalkwarf et al. (1989), Kalkwarf, Kaldahl, & Patil, (1992), Kaldahl, Kalkwarf, Patil, Molvar, & Dyer, (a, b)	RCT Split-mouth PICO 1	82 (51) NR (43.5) years	84 months (24 and 36 months)	Initial therapy: OHI + CS (1 quadrant) + SRP (3 quadrants) Test: MWF Control: SRP (no further treatment after initial therapy) Maintenance protocol: OHI + supra- and subgingival debridement (when needed)	Test 2: OS Control 2: CS	PI BoP Δ PD Δ CAL REC PROMS	University, USA, supported by governmental research funding	All teeth included CS group follow-up discontinued at 2 years: teeth in CS group that showed ≥3 mm CAL loss from baseline during maintenance, were scaled and root planned and incorporated in SRP group PD subgrouping: Initial PD 1–4 mm Initial PD 5–6 mm Initial PD > 6 mm
Becker et al. (1988, 2001)	RCT Split-mouth PICO 1	16 (16) 30–57 (42) years	60 months (12 months)	Initial therapy: OHI + SRP (without LA) + OA Test: MWF Control: SRP (with LA) Maintenance protocol: prophylaxis weekly for 6 weeks, and SRP and polishing every 3 months	Test 2: OS	PI GI Δ PD Δ CAL REC Mobility Furcation lesions Tooth loss	Private practice, USA, funding not reported	Only posterior teeth included (premolars and molars) PD subgrouping: Initial PD 1–3 mm Initial PD 4–6 mm Initial PD > 6 mm
Serino et al. (2001)	RCT Parallel PICO 1	64 (45) NR	156 months	Initial Therapy: OHI Test: MWF Control: SRP Maintenance protocol: OHI and supra- and subgingival debridement every 3–4 months	None	PI BoP Δ PD Δ PAL Rx BL Tooth loss	University, Sweden, supported by governmental research funding and by company	Only non-molar teeth included PD subgrouping: Initial PD 1–3 mm Initial PD 4–5 mm Initial PD > 5 mm
Polansky et al. (2003)	RCT Split-mouth PICO 1	29 (29) 38–75 (54.6) years	12 months	Initial Therapy: OHI + supragingival scaling Test: MWF Control: SRP Maintenance protocol: supra- and subgingival (when needed) debridement every 3 months	Control 2: no further treatment after initial therapy (CS)	PI PBI Δ PD Δ CAL Mobility	University, Austria, funding not reported	Canines and premolars receiving telescopic crowns

(Continues)

TABLE 1 (Continued)

Authors/Year	Study Design and PICO question	Number of patients baseline (final) Age range (mean)	Longer follow-up reported (in parenthesis, previous follow-ups reported)	Intervention	Other interventions not included in this SR	Outcomes	Setting, location and funding	Comments (Teeth included in the study, data subgrouping by PD or CAL, other study characteristics)
Ribeiro et al. (2011)	RCT Parallel PICO 1	29 (27) 35–57 (45.43) years	6 months	<i>Initial Therapy:</i> OHI + SRP <i>Test:</i> MIST <i>Control:</i> SRP with microscope and micro instruments <i>Maintenance protocol:</i> supragingival prophylaxis every 15 days for the first month and monthly for 6 months	None	PI BoP Δ PD Δ CAL Δ REC PROMS Chair-time	University, Brazil, supported by governmental research funding and by foundation	Only single-rooted teeth included Focus on intra-bony defects (≥1 single-rooted tooth with PD ≥ 5 mm + BoP, CAL ≥ 5 mm and radiographic evidence of an intra-bony defect (depth ≥ 4 mm and width ≥ 2 mm))
Trombelli et al. (2012)	RCT Parallel PICO 2	28 (28) 36–62 (49) years	6 months	<i>Initial Therapy:</i> OHI + SRP <i>Test:</i> single-flap approach <i>Control:</i> double-flap approach <i>Maintenance protocol:</i> OHI + Supragingival plaque removal every month for 3 months, and at intervals according to personal needs thereafter	None	BoP Δ PD Δ CAL Δ REC	University, Italy, supported by governmental research funding	All teeth included Focus on intra-bony defects (presence of ≥1 site with PD ≥ 6 mm and radiographic depth ≥ 3 mm)
Reddy et al. (2014)	RCT Split-mouth PICO 2	13 (13) 22–60 (NR) years	6 months	<i>Initial Therapy:</i> OHI + SRP <i>Test:</i> OFD <i>Control:</i> MWF <i>Maintenance protocol:</i> Not specified	None	GI PI PD CAL REC GCI Mobility Furcation lesions DH	University, India, self-supported	Only posterior teeth included (premolars and molars) All type of defects treated with distinction between horizontal and vertical defects
Perumal et al. (2015)	RCT Split-mouth PICO 2	13 (13) 30–50 (NR) years	9 months	<i>Initial Therapy:</i> OHI + SRP <i>Test:</i> MWF (microsurgical approach) <i>Control:</i> MWF (conventional approach) <i>Maintenance protocol:</i> Not specified	None	GI PD RAL REC EHI Pain	University, India, self-supported	All teeth included All type of defects treated without distinction between horizontal or vertical defects
Chandra et al. (2016)	RCT Split-mouth PICO 2	22 (19) 25–45 (34.8) years	6 months	<i>Initial Therapy:</i> OHI + SRP + SPT <i>Test 1:</i> MWF with Microdissection needle <i>Test 2:</i> MWF with electrocautery tip <i>Control:</i> MWF with conventional scalpel <i>Maintenance protocol:</i> not specified	None	PI GI PD CAL PH EHI Blood loss Pain	University, India, self-supported	All teeth included All type of defects treated without distinction between horizontal or vertical defects

Abbreviations: AF, access flap; BoP, bleeding on probing; CAL Loss, clinical attachment loss; CAL, clinical attachment level; CI, calculus index; CS, coronal scaling; DH, dentinal hypersensitivity; EHI, early healing index; GCI, gingival contour index; GI, gingival index; LA, local anaesthesia; MIST, minimally invasive surgical technique; MWF, modified Widman flap; NR, not reported; OA, occlusal adjustment; OFD, open flap debridement; OHI, oral hygiene instructions; OS, osseous surgery; PD, probing depth; PH, papilla height; PI, plaque index; PROMS, patient-reported outcome measures; RAL, relative attachment level (measured from a stent to the bottom of the pocket); RCT, randomized controlled trial; REC, recession; Rx BL, radiographic bone level; SPT, supportive periodontal therapy; SR, systematic review; SRP, scaling and root planing; Sup, suppuration.

The resulting systematic review pooled data from 503 patients at baseline, from which 428 were analysed at the end of the investigation. Patients were followed for a mean period of 41.50 ± 39.90 months (range: 6–156 months).

3.3 | Risk of bias in individual studies

Rob2 tool was used to score the randomized clinical trials (Table 2). Serious methodological inadequacies associated with bias were found in almost all the studies for at least one domain. Only the study by Ribeiro et al. (2011) was at low risk of bias. The most frequent domains with risk of bias were incomplete reporting of the randomization process (sequence generation and allocation concealment) and examiner blinding (Table S2).

3.4 | Risk of bias across studies

No significant publication bias was observed for the primary outcome measure (PD reduction) when considering studies reporting the mean PD change (Lindhe et al., 1982; Polansky et al., 2003; Serino et al., 2001), or for studies reporting the PD change according to the initial PD distribution ($p > .10$). The sensitivity analysis showed that the exclusion of a single study did not substantially alter any estimate.

3.5 | Effects of interventions

3.5.1 | PD reduction and CAL change (PICO 1: access flap vs. subgingival debridement)

Table 3 depicts the meta-analyses comparing PD and CAL change between access flaps and subgingival debridement. The WME within each group for PD reduction and CAL change is depicted in Table 4.

All pockets

Access flap showed significant greater PD reduction when all categories of initial PD were combined irrespectively of the follow-up ($n = 3$; WMD = 0.71 mm; 95% CI 0.44, 0.98; $p < .001$). On the contrary, when combining all initial PD categories, subgingival debridement showed significant CAL gain, whereas access flap showed no change. The difference between both groups in terms of CAL change was significant ($n = 3$; WMD = -0.26 mm; 95% CI -0.39 , -0.14 ; $p < .001$). For studies focusing only on intra-bony defects, the PD reduction was significantly greater in the surgery group ($n = 4$; WMD = 0.49 mm; 95% CI 0.11, 0.86; $p = .01$), although no differences were observed in terms of CAL gain ($n = 4$; WMD = 0.07 mm; 95% CI -0.29 , 0.44; $p = .691$).

Shallow pockets (1–3 mm or 1–4 mm)

Non-significant changes were observed in the test or control groups in terms of PD reduction in shallow pockets (Figures 2 and 3). Although

PD reduction was significantly greater in the group receiving surgery at the short term ($n = 4$; WMD = 0.13 mm; 95% CI 0.07, 0.18; $p < .001$), this difference became non-significant at the long term ($n = 4$; WMD = -0.02 mm; 95% CI -0.09 , 0.05; $p = .648$). For this pocket category, both groups showed significant clinical attachment loss at the long term (not at short term for subgingival debridement), which was greater in the surgery group at all time points ($n = 7$; WMD = -0.43 mm; 95% CI -0.56 , -0.28 ; $p < .001$; prediction interval -0.81 , -0.04 ; at the short term and $n = 6$; WMD = -0.27 mm; 95% CI -0.34 , -0.20 ; $p < .001$, at the long term, respectively). For studies reporting the change in the frequency distribution of shallow pockets in the long term, the use of access flap provided an additional increase of 11.60%, which was statistically significant (95% CI 6.76, 16.45; $p < .001$).

Moderately deep pockets (4–6, 5–6 or 4–5 mm)

Both groups showed significant PD reduction in moderately deep pockets (Figures 2 and 3). As it happened with shallow pockets, PD reduction was significantly greater in the access flap group at the short term ($n = 4$; WMD = 0.34 mm; 95% CI 0.21, 0.46; $p < .001$). This additional reduction in PD for AFs over subgingival debridement amounted for 29.6%. Whereas the subgingival debridement group showed significant CAL gain only at the short term, the changes were not significant in the surgery group. When comparing both treatment modalities, the subgingival debridement group showed greater CAL gain in the short and the long terms, although this difference tended to mitigate with time ($n = 7$; WMD = -0.34 mm; 95% CI -0.46 , -0.22 ; $p < .001$, and $n = 6$; WMD = -0.12 mm; 95% CI -0.22 , -0.01 ; $p = .032$, respectively). For studies reporting the change in the frequency distribution of moderately deep pockets in the long term, the use of access flap provided an additional increase of 9.49%, which was statistically significant (95% CI 4.88, 14.10; $p < .001$).

Deep pockets (>6 mm or ≥ 6 mm)

Although both groups showed significant PD reduction in this pocket category, this reduction was significantly greater in areas receiving surgery in the short and long terms ($n = 4$; WMD = 0.67 mm; 95% CI 0.37, 0.97; $p < .001$, and $n = 4$; WMD = 0.39 mm; 95% CI 0.09, 0.70; $p = .012$, respectively) (Figures 2 and 3). The additional PD reduction for AFs over subgingival debridement amounted for 27.5% in the short term and 25.3% in the long term. As it happened with moderately deep pockets, the differences tended to be smaller with time. On the contrary, no significant differences among groups were observed for CAL gains at the short and long terms ($n = 7$; WMD = 0.19 mm; 95% CI -0.04 , 0.43; $p = .111$, and $n = 6$; WMD = 0.07 mm; 95% CI -0.15 , 0.29; $p = .524$, respectively). No differences among groups could be found for the change in the frequency distribution of this PD category.

Molar versus non-molar teeth

Only three studies compared the difference in treatment response between molar and non-molar teeth. In the study from Lindhe et al. (1982), it was observed that in surgically treated sites, PD reduction was significantly greater (for initial pockets of ≥ 4 mm) in single-rooted than in multi-rooted teeth. However, in the subgingival debridement group, the PD reduction was similar in both types of teeth. CAL loss was seen in sites

with initial PD < 4 mm irrespectively of the type of tooth. Similarly, CAL gain was observed for sites with initial PD > 6 mm, without differences between groups or between the type of tooth. In the study from Philstrom et al. (1984), it was observed that for initial pockets of 4–6 mm, molars showed greater PD and CAL irrespectively of the assigned treatment; and that for initial pockets >7 mm, AF resulted in less PD on non-molars than on molars, without significant differences for CALs, neither for tooth type nor for the method of therapy. Additionally, one study reported the results differentiating four tooth and site groupings: interproximal sites of single-rooted teeth, facial and lingual sites of single-rooted teeth, non-furcation sites of molar teeth and furcation sites of molar teeth (Kaldahl, Kalkwarf, Patil, & Molvar, 1990). While no differences were observed for PD reductions or CAL gains in shallow sites, PD reduction and CAL gain were greater in single-rooted teeth at moderately deep and initial deep pockets.

3.5.2 | PD reduction and CAL change (PICO 2: different access flaps)

Three studies compared the MWF to the Open Flap Debridement. No meta-analyses were performed for PD or CAL changes due to the heterogeneity found for these outcomes (mean changes, changes in proportions, data expressed as figures, subgrouping by initial PD, etc.). While two of these studies did not find any significant difference among both flap procedures (Lindhe & Nyman, 1985; Svoboda, Reeve, & Sheridan, 1984), one study reported better results for conventional MWF in terms of PD reduction and CAL gain (Reddy et al., 2014). Additionally, one study compared the MWF using conventional blades to the MWF using a micro-dissection needle or an electrocautery tip (Chandra, Savitharani, & Reddy, 2016) and another investigation compared the MWF using conventional instruments to the use of microsurgical instruments (Perumal, Ramegowda, Lingaraju, & Raja, 2015). In both studies, there was significant PD reduction and CAL gain without significant differences among interventions. Finally, one study compared the use of the single-flap approach (SFA) to the double-flap approach (DFA) in the surgical treatment of intra-bony defects with significant better results in terms of PD reduction and CAL gain for the SFA (Trombelli et al., 2012).

3.6 | Secondary outcomes

Changes in gingival or bleeding indices and in plaque indices (Table S3).

Bleeding on probing was assessed in 10 investigations (Isidor & Karring, 1986; Kaldahl et al., 1996; Kalkwarf, Kaldahl, Patil, & Molvar, 1989; Lindhe & Nyman, 1985; Perumal et al., 2015; Renvert, Garrett, Nilvéus, Chamberlain, & Egelberg, 1985; Renvert et al., 1990; Ribeiro et al., 2011; Serino et al., 2001; Trombelli et al., 2012; Wennström et al., 1986; Westfelt et al., 1985). Due to the high heterogeneity of studies analysing this outcome (PICO 1, PICO 2, studies focused exclusively on the treatment of intra-bony defects, etc.), no meta-analysis was performed. In general, both treatment groups showed similar and significant BOP reductions that slightly worsened with time in

long-term studies. In one study, the reduction of BOP was only significant in test group (Trombelli et al., 2012).

The gingival index was used to assess inflammation in eight investigations either as the Löe and Silness index (Becker et al., 2001, 1988; Isidor & Karring, 1986; Lindhe & Nyman, 1985; Lindhe et al., 1982; Reddy et al., 2014; Svoboda et al., 1984), the papillary bleeding index from Saxer et al. (Polansky et al., 2003) or as the gingival index from Lobene (Chandra et al., 2016). In summary, both treatment groups showed a significant reduction in mean inflammation scores and an increase in the percentage of sites without inflammation (score = 0) with no differences among groups.

Plaque indices were assessed in 13 out of the 18 included investigations, either as the plaque index from Silness & Löe (Becker et al., 2001; Chandra et al., 2016; Isidor & Karring, 1986; Lindhe & Nyman, 1985; Lindhe et al., 1982; Reddy et al., 2014; Svoboda et al., 1984) or as the percentage of surfaces harbouring plaque (Kaldahl et al., 1996; Renvert et al., 1990; Ribeiro et al., 2011; Serino et al., 2001; Wennström et al., 1986; Westfelt et al., 1985). In general, both treatment groups showed a significant reduction in mean plaque scores and an increase in the percentage of sites without plaque (score = 0) with no differences among groups.

3.6.1 | Tooth loss

Tooth loss was reported in seven investigations (Isidor & Karring, 1986; Kaldahl et al., 1990b; Pihlstrom et al., 1984; Ramfjord et al., 1987; Renvert et al., 1990; Serino et al., 2001; Wennström et al., 1986), with a range between 0% and 2.43% in periods from 60 to 156 months. Tooth loss was due to periodontal- and non-periodontal-related causes, and the most frequent teeth extracted were molars.

3.6.2 | Percentage of residual pockets (PD > 3 mm or PD > 4 mm with or without the assessment of BOP) or closed pockets (PD < 3 mm or PD < 4 mm with or without the assessment of BOP)

At the end of the therapy, none of the included studies in this systematic review evaluated the percentage of closed or open pockets using a composite outcome that included BOP in the equation together with PD. The percentage of residual sites with PD > 3 mm after treatment varied from 17% to 49% in the access flap group, and 20%–62% in the subgingival debridement group (Becker et al., 2001; Lindhe & Nyman, 1985; Lindhe et al., 1982a; Serino et al., 2001; Wennström et al., 1986).

3.6.3 | Harms, adverse effects and patient-reported outcomes (PROMs)

Harms and adverse effects related to the technique were not reported in any of the studies. The most common complications

TABLE 2 Cochrane tool for the qualitative assessment of randomized clinical trials

Authors/Year	Randomization process	Deviations from the intended interventions	Missing outcome data	Measurement of the outcome	Selection of the reported result	Overall risk of bias
Ramfjord, Knowles, Nissle, Burgett, & Shick, (1975)	?	?	-	-	?	-
Lindhe et al. (a)	?	?	+	-	?	-
Lindhe et al. (b)	?	?	+	-	?	-
Lindhe et al. (1984)	?	?	+	-	?	-
Pihlstrom et al. (1981)	?	+	+	-	?	-
Pihlstrom et al. (1983)	?	+	+	-	?	-
Pihlstrom et al. (1984)	?	+	+	+	?	?
Svoboda et al. (1984)	?	?	-	-	?	-
Lindhe and Nyman (1985)	?	?	-	-	?	-
Westfelt et al. (1985)	?	?	-	-	?	-
Isidor et al. (1984)	?	?	-	-	?	-
Isidor et al. (1985)	?	?	+	-	?	-
Isidor and Karring (1986)	?	?	+	-	?	-
Wennström et al. (1986)	?	?	+	-	?	-
Renvert, Garrett, et al. (1985)	?	?	+	-	?	-
Renvert et al. (1990)	?	?	+	-	?	-
Hill et al. (1981)	?	?	-	-	?	-
Ramfjord et al. (1987)	?	?	-	-	?	-
Burgett et al. (1992)	?	?	+	-	?	-
Kaldahl et al. (1988)	?	+	+	+	?	?
Kalkwarf et al. (1989)	?	+	+	+	?	?
Kaldahl et al. (a)	?	+	+	+	?	?
Kaldahl et al. (b)	?	+	+	+	?	?
Kaldahl et al. (c)	?	+	+	+	?	?
Kalkwarf et al. (1992)	?	+	+	-	?	-
Kaldahl et al. (a)	?	+	+	+	?	?
Kaldahl et al. (b)	?	+	+	+	?	?
Becker et al. (1988)	?	?	+	-	?	-
Becker et al. (2001)	?	?	+	-	?	-
Serino et al. (2001)	?	?	+	-	?	-
Polansky et al. (2003)	?	?	-	-	?	-
Ribeiro et al. (2011)	+	+	+	+	+	+
Trombelli et al. (2012)	?	+	+	+	?	?
Reddy et al. (2014)	?	?	-	-	?	-
Perumal et al. (2015)	?	?	-	-	?	-
Chandra et al. (2016)	+	?	+	+	+	?

Abbreviations: +, low risk of bias; -, high risk of bias; ?, some concerns.

reported during follow-up were the incidence of further attachment loss and the need for re-treatment. The weighted mean incidence of sites exhibiting attachment loss ≥ 2 mm or ≥ 3 mm by intervention group and by initial PD is depicted in Table 5. Due to the high heterogeneity found when reporting this outcome, no meta-analyses could be done to compare the risk for future attachment loss between both interventions. Just four studies reported the percentage

of patients or teeth in need for re-treatment during the study follow-up, with values between 0% and 14% in the access flap group, and from 8% to 29% in the subgingival debridement group (Kaldahl, Kalkwarf, Patil, Dyer, & Bates, 1988; Pihlstrom et al., 1984; Ramfjord et al., 1987; Serino et al., 2001).

Finally, PROMs were reported in four investigations. One study used a questionnaire to rate seven domains (Kaldahl et al., 1996a),

two studies used the visual analogue scale to rate pain (Chandra et al., 2016; Perumal et al., 2015) and one study evaluated the percentage of cases with hypersensitivity (Reddy et al., 2014). In general, no differences could be observed between groups.

4 | DISCUSSION

The results of this systematic review, based on 36 publications reporting data from 18 investigations, indicate a high variability in terms of treatment protocols and on how the outcomes were reported, so the results should be interpreted with caution. This systematic review focused on the clinical performance of subgingival debridement and AF and on the comparison of different surgical approaches for flap debridement in the treatment of periodontitis. The primary outcome was PD reduction. The results of the meta-analysis showed that for all pockets greater PD reduction is expected when doing an AF. In particular, AF has shown to offer consistent benefits over subgingival debridement in deep pockets on the short and long term. Due to lack of studies, it was impossible to draw definitive conclusions for the comparative efficacy of different surgical approaches.

Both treatments resulted in PD reduction in shallow sites with limited clinical relevance. Despite the fact that PD reduction was greater in sites treated with AF, both therapies resulted in CAL loss at the short and long term (0.2–0.8 mm). Nevertheless, subgingival debridement might have an advantage, since treatment can be limited to moderately deep and initial deep pockets, whereas surgery needs to include in many situations shallow sites adjacent to pockets (PD \geq 4 mm). These findings confirmed those of a previous systematic review on the same topic (Heitz-Mayfield et al., 2002).

In moderately deep pockets both treatment modalities showed significant PD reduction that was greater for AF in the short term. However, these differences disappeared and both procedures were comparable in terms of PD reduction after 1 year. CAL gain was only significant in the subgingival debridement at the short term and the differences between groups, although limited, were statistically significant at short and long term. The subgingival debridement group showed a decrease in CAL gain without major changes in PD in the long term, indicating that with time gingival recession increased. Based on these results, non-surgical periodontal therapy may be considered the ideal treatment option to restore periodontal health in patients affected by moderately deep pockets only (Badersten, Nilvéus, & Egelberg, 1981).

In deep pockets greater PD reduction was achieved by means of access therapy on the short and long term. However, this difference was progressively reduced over time and the added effect of access therapy was <0.5 mm in the long term. Moreover, no differences were found in terms of CAL change between the procedures neither at short- nor long term. We might consider that these results may be attributable to a greater gingival recession in the sites treated by means of AF. Recession in fact contributes to PD reduction and allows direct access to root surfaces during oral hygiene procedures. This could be beneficial to maintain periodontal stability over time

due to improved plaque control. However, in order to draw robust conclusions, data regarding residual PD rather than PD reduction would have been desirable. Furthermore, a study reporting for plaque accumulation after different treatments did not observe statistically significant differences between AF and SRP (Kaldahl, Kalkwarf, Patil, & Molvar, 1990).

The main objective of periodontal treatment is to control inflammation and thereby arrest attachment and tooth loss. This is achieved in part by reducing the severity and the number of pockets, creating an accessible environment for oral hygiene. These goals should be maintained over time to warrant periodontal stability (Pini Prato, Di Gianfilippo, & Wang, 2019). Interestingly, AF was more successful in increasing the number of shallow sites (11.6%). However, it is unclear whether this might translate into a better stability in the long term, since both treatments resulted in a consistent percentage of sites exhibiting further CAL loss. Even though AF was better at increasing the percentage of shallow sites compared to subgingival debridement, this might not have been enough to avoid disease progression at patient level. Unfortunately, we could not compare both treatments approaches since standard deviations were not provided in most of the studies reporting on attachment loss. Furthermore, both treatments were associated to a comparatively low frequency of tooth loss (0%–2.5%) in respect to people who discontinued periodontal treatment (Becker, Berg, & Becker, 1979; Harrel & Nunn, 2001).

Ideally a better outcome for periodontal treatment would be “pocket closure,” a composite variable including PD and BOP, since residual PD and BOP have been consistently associated to an increased odds for tooth loss and disease progression at site- and patient-level (Claffey & Egelberg, 1995; Matuliene et al., 2008). However, none of the included studies reported the changes in terms of “open/closed pockets,” which prevented us to select it as primary outcome. Additionally, we were not able to compare the frequency of residual pockets due to inconsistencies found on how this outcome was reported.

Only two studies (Lindhe et al., 1982; Pihlstrom et al., 1984) compared the efficacy of subgingival debridement and AFs between molars and non-molars. In general, a better performance on single-rooted teeth was observed for both treatments. This difference may be explained by accessibility and anatomical factors, such as the increased buccolingual dimension, the presence of anatomical abnormalities and the position in the arch, which could impair the efficacy of subgingival debridement, especially when deep pockets are present (Caffesse et al., 1986; Rateitschak-Pluss et al., 1992; Waerhaug, 1978). In this context, it is also important to remark that the most frequent lost teeth were molars.

We found a limited number of studies comparing different AFs for the treatment of periodontitis, and thus, no definitive conclusions can be drawn regarding a higher efficacy of any particular technique. It seems that traditional techniques such as MWF and OFD are equivalent. We also tried to evaluate whether there was an impact of minimally invasive approaches in comparison with traditional approaches in the surgical treatment of periodontitis. However, this has only been demonstrated in intra-bony defects,

TABLE 3 Summary of meta-analysis performed for probing depth (PD) and clinical attachment level (CAL) changes

Initial PD category (mm)	Number of studies	WMD (mm or %)	95% CI		p Value for WMD	Heterogeneity		
						I ² (%)	p Value	Method
<i>Studies including all type of defects</i>								
PD reduction (mean)								
All	3	0.71	0.44	0.98	<.001	63.6	.064	IV
Short term (≤12 months)								
Shallow pockets	4	0.13	0.07	0.18	<.001	34.2	.207	IV
Moderate pockets	4	0.34	0.21	0.46	<.001	40.5	.169	IV
Deep pockets	4	0.67	0.37	0.97	<.001	0	.582	IV
Long term (>12 months)								
Shallow pockets	4	-0.02	-0.09	0.05	.648	0	.739	IV
Moderate pockets	4	0.09	-0.01	0.18	.063	0	.897	IV
Deep pockets	4	0.39	0.09	0.70	.012	41.4	.163	IV
CAL gain (mean)								
All	3	-0.26	-0.39	-0.14	<.001	36.4	.208	IV
Short term (≤12 months)								
Shallow pockets	7 ^a	-0.43	-0.56	-0.29	<.001	55.6	.036	DL
Moderate pockets	7 ^a	-0.34	-0.46	-0.22	<.001	6.6	.378	IV
Deep pockets	7 ^a	0.19	-0.04	0.43	.111	0	.664	IV
Long term (>12 months)								
Shallow pockets	6	-0.27	-0.34	-0.20	<.001	50.8	.071	IV
Moderate pockets	6	-0.12	-0.22	-0.01	.032	0	.511	IV
Deep pockets	6	0.07	-0.15	0.29	.524	0	.983	IV
Proportions (long term)								
Change in % of shallow pockets	3 ^a	11.60	6.76	16.5	<.001	35.0	.215	IV
Change in % of moderate pockets	3 ^a	9.49	4.88	14.10	<.001	0	.472	IV
Change in % of deep pockets	3 ^a	0.92	-1.27	3.11	.413	33.6	.222	IV
<i>Studies including exclusively intra-bony defects</i>								
PD reduction (mean)								
All pockets (short and long tem)	4	0.49	0.11	0.86	.010	54.7	.085	IV
CAL gain (mean)								
All pockets (short and long tem)	4	0.07	-0.29	0.44	.691	0	.405	IV

Note: Positive values indicate a mean difference in favour of surgery.

Abbreviations: CAL, clinical attachment level; CI, confidence interval; DL, DerSimonian and Laird (random-effect) model; IV, inverse-variance weighted (fixed effect) model; WMD, weighted mean difference.

^aLindhe et al. (1985) compared two different access flaps (Modified Widman Flap or Modified Kirkland Flap) versus subgingival debridement.

where minimally invasive techniques performed better than conventional procedures, suggesting that surgical procedures aiming at primary flap closure may protect the surgical area and the blood clot in the early healing phases (Graziani et al., 2012). Moreover, when the analysis was limited to intra-bony defects, we found greater PD reduction in sites receiving access flap, although without differences in terms of CAL gain in the long term. The fact that these differences were maintained on the long term suggests that AF may be superior to subgingival debridement in the treatment of intra-bony defects. When different surgical approaches were compared, the SFA performed better than the DFA in terms of PD reduction and CAL gain (Trombelli et al., 2012).

It is important to remark that in most of the studies, subgingival debridement was performed as part of the initial therapy and repeated in the assigned quadrant after randomization. Re-treatment of residual pockets after initial phase by means of subgingival debridement may result in further PD reduction, although with a smaller effect than after the first instrumentation (Tomasi, Koutouzis, & Wennstrom, 2008). Performing subgingival debridement as part of the initial non-surgical cause-related therapy could have affected the results by reducing the additional benefit of the surgical procedures. Anyway, it should be considered that this approach simulates better the situation of the patients treated in the daily practice, in which re-instrumentation of residual pockets is a critical component of supportive periodontal therapy.

TABLE 4 Weighted mean effect (WME) of scaling and root planing (SRP) and Access flaps according to the initial probing depth

Initial PD category (mm)	Number of studies	WME (mm)			p Value	Heterogeneity			
		IV	DL	95% CI		I ²	p Value		
<i>Studies including all type of defects</i>									
PD Reduction									
Access Flap									
All (short and long tem)	3	-	1.70	0.89	2.52	<.001	95.1	<.001	
SRP									
All (short and long tem)	3	-	1.04	0.55	1.5	<.001	84.5	.002	
Short term (≤12 months)									
Access Flap									
Shallow pockets	4	-	0.26	0.10	0.43	<.001	96.3	<.001	
Moderate pockets	4	1.53	-	1.44	1.62	<.001	21.4	.282	
Deep pockets	4	2.94	-	2.76	3.13	<.001	26.1	.255	
SRP									
Shallow pockets	4	-	0.13	-0.01	0.27	.158	90.8	<.001	
Moderate pockets	4	-	1.08	0.81	1.35	<.001	88.7	<.001	
Deep pockets	4	-	2.13	1.57	2.70	<.001	79.2	<.001	
Long term (>12 months)									
Access Flap									
Shallow pockets	4	-	-0.03	-0.38	0.32	.163	97.5	<.001	
Moderate pockets	4	-	1.23	0.85	1.62	<.001	94.8	<.001	
Deep pockets	4	-	2.77	2.34	3.20	<.001	73.1	<.001	
SRP									
Shallow pockets	4	-	0.01	-0.30	0.32	.258	97-2	<.001	
Moderate pockets	4	-	1.15	0.82	1.47	<.001	94.2	<.001	
Deep pockets	4	-	2.07	1.18	2.96	<.001	92.7	<.001	
CAL gain									
Access Flap									
All (short and long term)	3	-	-0.1	-0.32	0.12	.460	71.4	.030	
SRP									
All (short and long term)	3	0.14	-	0.04	0.24	<.001	37.1	.204	
Short term (≤12 months)									
Access Flap									
Shallow pockets	7	-	-0.64	-0.90	-0.38	<.001	93.9	<.001	
Moderate pockets	7	-	0.22	-0.12	0.57	.206	92.7	<.001	
Deep pockets	7	-	1.31	0.90	1.71	<.001	85.7	<.001	
SRP									
Shallow pockets	6	-	-0.16	-0.38	0.06	.396	91.6	<.001	
Moderate pockets	6	-	0.53	0.22	0.83	<.001	91.9	<.001	
Deep pockets	6	-	1.16	0.67	1.65	<.001	83.4	<.001	
Long Term (>12 months)									
Access Flap									
Shallow pockets	6	-	-0.76	-1.07	-0.45	<.001	96.6	<.001	
Moderate pockets	6	-	0.16	-0.23	0.55	.663	94.8	<.001	
Deep pockets	6	-	1.25	0.81	1.70	<.001	82.4	<.001	

(Continues)

TABLE 4 (Continued)

Initial PD category (mm)	Number of studies	WME (mm)			p Value	Heterogeneity		
		IV	DL	95% CI		I ²	p Value	
SRP								
Shallow pockets	6	-	-0.43	-0.78	-0.08	<.001	97.4	<.001
Moderate pockets	6	-	0.36	-0.04	0.75	.195	95.8	<.001
Deep pockets	6	-	1.18	0.69	1.66	<.001	84.2	<.001
<i>Studies including exclusively intra-bony defects</i>								
PD Reduction								
Access Flap								
All	4	3.20	-	2.75	3.65	<.001	65.5	.034
SRP								
All	4	2.75	-	1.88	3.61	<.001	88.8	<.001
CAL gain								
Access Flap								
All	4	-	1.36	0.43	2.29	<.001	92.3	<.001
SRP								
All	4	-	1.43	0.57	2.29	<.001	73.8	<.001

Abbreviations: CAL, clinical attachment level; CI, confidence interval; DL, DerSimonian and Laird (random-effect) model; IV, inverse-variance weighted (fixed effect) model; PD, probing depth; SRP, scaling and root planing.

Short term changes (≤12 months) by initial PD




FIGURE 2 Short-term (≤12 months) changes in probing depth (PD) and clinical attachment level (CAL) by initial PD [Colour figure can be viewed at wileyonlinelibrary.com]


Long term changes (≥ 12 months) by initial PD

FIGURE 3 Long-term (>12 months) changes in probing depth (PD) and clinical attachment level (CAL) by initial PD [Colour figure can be viewed at wileyonlinelibrary.com]

Similarly, it must be underlined that in the present review only a limited number of studies performed subgingival debridement with magnification. Nowadays loupes and operative microscope are commonly used in high standard dental practice and may enhance the effect of subgingival debridement, overcoming its limitations in terms of accessibility and precision. In fact, it has been demonstrated that subgingival debridement under magnification has excellent clinical performance (Nibali et al., 2015; Ribeiro et al., 2011).

The available evidence suggests that the effects of non-surgical therapy are expressed over an extended period of time and that traditional re-evaluation at 3 months may mask the real potential of non-surgical treatment. In this sense, the healing potential of subgingival debridement has shown to last more than 9 months, depending on the maintenance protocol and patient compliance (Badersten, Nilveus, & Egelberg, 1984). All the studies included in this systematic review followed a strict maintenance protocol, which varied from intervals between every 2 weeks to every 6 months, including or not subgingival instrumentation. This may explain the positive results found in the studies included and the tendency to present similar results irrespectively of treatment modality (Axelsson & Lindhe, 1981; Lindhe, Westfelt, Nyman, Socransky, & Haffajee, 1984). Nevertheless, we must emphasize that the maintenance protocols used in many of the included studies might be difficult to extrapolate to the daily clinical practice.

In the present review, caution is required before interpreting the principal findings due to the high heterogeneity between studies,

mainly concerning the strategy of initial therapy, patient selection, teeth included in the analysis, the modalities and frequency of supportive periodontal treatment and the length of follow-up. Furthermore, most studies are out-dated and had a high risk of bias. One of the limitations of this review derives from the different treatment strategies, with some studies performing non-surgical subgingival debridement as part of the initial phase and others moving directly to surgery in the patients or quadrants assigned. No superiority of any strategy could be concluded, as long as subgroup meta-analysis was not possible due to lack of data. This may hamper the recommendations made on the basis of this review regarding the most appropriate timing for access flaps. Another important limitation is that in order to be more inclusive in the subgroup analysis, we combined studies using different initial PD categories. The shallow pockets from the "Nebraska studies" (1–4 mm) (Kaldahl et al., 1988, 1996) were pooled together with the 1–3 mm pockets of other included studies. Similarly, the deep pockets from Serino et al. (2001) (≥ 6 mm) were pooled together with the results for the ≥ 7 mm pockets of other investigations. This could have influenced the real effect of treatment, even though no significant heterogeneity was observed in most of the meta-analysis. Moreover, the inclusion of studies with at least 10 patients per treatment group in order to report more robust results forced us to exclude three studies, which although very unlikely, could have influenced the results. However, the effort of pooling together studies with similar characteristics

TABLE 5 Meta-analysis performed for the percentage of sites exhibiting attachment loss (ALoss) ≥ 2 mm or ≥ 3 mm by intervention group and by initial probing depth (PD)

Initial PD category (mm)	Number of studies	WMI (%)				p Value	Heterogeneity	
		IV	DL	95% CI	I^2		p Value	
AL ≥ 2 mm								
Access flap								
Shallow pockets	2 ^{a,b}	-	25.4	5.6	45.2	.012	98.2	<.001
Moderate pockets	2 ^{a,b}	-	21.3	8.2	34.4	.001	94.8	<.001
Deep pockets	2 ^{a,b}	10.3	-	4.0	16.6	.001	64.4	.094
SRP								
Shallow pockets	2 ^{a,b}	-	19.6	0.0	40.6	.066	99.0	<.001
Moderate pockets	2 ^{a,b}	-	16.1	6.0	26.2	.002	93.1	<.001
Deep pockets	2 ^{a,b}	15.7	-	7.5	24.0	<.001	0.0	.574
AL ≥ 3 mm								
Access flap								
Shallow pockets	2 ^{a,c}	-	6.5	0.0	17.9	.265	99.5	<.001
Moderate pockets	2 ^{a,c}	-	6.0	0.0	14.6	.169	98.1	<.001
Deep pockets	2 ^{a,c}	2.1	-	1.6	2.6	<.001	0.8	.315
SRP								
Shallow pockets	2 ^{a,c}	-	4.9	0.0	13.3	.255	99.3	<.001
Moderate pockets	2 ^{a,c}	-	4.9	0.0	11.0	.109	97.1	<.001
Deep pockets	2 ^{a,c}	3.2	-	2.6	3.8	<.001	0.0	.535

Note: Abbreviations: CI, confidence interval; DL, DerSimonian and Laird (random effect) model; IV, inverse-variance weighted (fixed effect) model; SRP, scaling and root planing; WMI, Weighted mean incidence.

^aRamfjord et al. (1987).

^bBecker et al. (2001).

^cKaldahl et al. (1996a).

allows us to describe the clinical performance of different modalities of periodontal treatment in different clinical conditions. This enlightens the fact that periodontal therapy should be tailored and personalized to accomplish patient-centred treatment needs focused on specific clinical conditions.

Within the limitations of the present systematic review (including the high risk of bias of the included studies), it can be concluded that:

- AF achieved larger PD reduction when compared to subgingival debridement in initially deep pockets (>6 mm or ≥ 6 mm). In this pocket category, no differences were detected between treatment groups for CAL gain.
- In moderately deep pockets (4–6, 5–6 or 4–5 mm), AF achieved greater PD reduction in the short term. However, subgingival debridement resulted in higher CAL gain.
- In shallow pockets (1–3 or 1–4 mm), AF resulted in significantly greater clinical attachment loss.
- None of the included studies reported outcomes in terms of a composite variable including PD and BOP. Just a few studies reported the percentage of residual pockets after therapy, not allowing comparing subgingival debridement and AFs for this outcome variable. Future studies should overcome this deficiency.

- Regarding the comparison of different AFs, no clear conclusions can be made due to the limited studies available.

ACKNOWLEDGEMENTS

The author would like to acknowledge Professor Lisa Heitz-Mayfield for her help with data extraction.

CONFLICT OF INTEREST

The authors report no conflicts of interest related to this study.

ORCID

Ignacio Sanz-Sánchez  <https://orcid.org/0000-0002-3698-4772>

Eduardo Montero  <https://orcid.org/0000-0003-2525-8529>

Ana Molina  <https://orcid.org/0000-0003-1443-1189>

Mario Aimetti  <https://orcid.org/0000-0003-0657-0787>

REFERENCES

- Aimetti, M. (2014). Nonsurgical periodontal treatment. *International Journal of Esthetic Dentistry*, 9(2), 251–267.
- Armitage, G. C. (1999). Development of a classification system for periodontal diseases and conditions. *Annals of Periodontology*, 4(1), 1–6. <https://doi.org/10.1902/annals.1999.4.1.1>
- Axelsson, P., & Lindhe, J. (1981). The significance of maintenance care in the treatment of periodontal disease. *Journal of Clinical*

- Periodontology*, 8(4), 281–294. <https://doi.org/10.1111/j.1600-051X.1981.tb02039.x>
- Badersten, A., Nilvéus, R., & Egelberg, J. (1981). Effect of nonsurgical periodontal therapy. I. Moderately advanced periodontitis. *Journal of Clinical Periodontology*, 8(1), 57–72. <https://doi.org/10.1111/j.1600-051x.1981.tb02024.x>
- Badersten, A., Nilveus, R., & Egelberg, J. (1984). Effect of nonsurgical periodontal therapy. II. Severely advanced periodontitis. *Journal of Clinical Periodontology*, 11(1), 63–76. <https://doi.org/10.1111/j.1600-051X.1984.tb01309.x>
- Becker, W., Becker, B. E., Caffesse, R., Kerry, G., Ochsenein, C., Morrison, E., & Prichard, J. (2001). A longitudinal study comparing scaling, osseous surgery, and modified widman procedures: Results after 5 years. *Journal of Periodontology*, 72(12), 1675–1684. <https://doi.org/10.1902/jop.2001.72.12.1675>
- Becker, W., Becker, B. E., Ochsenein, C., Kerry, G., Caffesse, R., Morrison, E. C., & Prichard, J. (1988). A longitudinal study comparing scaling, osseous surgery and modified Widman procedures. Results after one year. *Journal of Periodontology*, 59(6), 351–365. <https://doi.org/10.1902/jop.1988.59.6.351>
- Becker, W., Berg, L., & Becker, B. E. (1979). Untreated periodontal disease: A longitudinal study. *Journal of Periodontology*, 50(5), 234–244. <https://doi.org/10.1902/jop.1979.50.5.234>
- Buchanan, S. A., & Robertson, P. B. (1987). Calculus removal by scaling/ root planing with and without surgical access. *Journal of Periodontology*, 58(3), 159–163. <https://doi.org/10.1902/jop.1987.58.3.159>
- Burgett, F. G., Ramfjord, S. P., Nissle, R. R., Morrison, E. C., Charbeneau, T. D., & Caffesse, R. G. (1992). A randomized trial of occlusal adjustment in the treatment of periodontitis patients. *Journal of Clinical Periodontology*, 19(6), 381–387. <https://doi.org/10.1111/j.1600-051x.1992.tb00666.x>
- Caffesse, R. G., Sweeney, P. L., & Smith, B. A. (1986). Scaling and root planing with and without periodontal flap surgery. *Journal of Clinical Periodontology*, 13(3), 205–210. <https://doi.org/10.1111/j.1600-051X.1986.tb01461.x>
- Chandra, R. V., Savitharani, B., & Reddy, A. A. (2016). Comparing the outcomes of incisions made by Colorado microdissection needle, electrosurgery tip, and surgical blade during periodontal surgery: A randomized controlled trial. *Journal of Indian Society of Periodontology*, 20(6), 616–622. https://doi.org/10.4103/jisp.jisp_27_17
- Checchi, L., & Schonfeld, S. E. (1988). A technique for esthetic treatment of maxillary anterior infrabony lesions. *Quintessence International*, 19(3), 209–213.
- Claffey, N., & Egelberg, J. (1995). Clinical indicators of probing attachment loss following initial periodontal treatment in advanced periodontitis patients. *Journal of Clinical Periodontology*, 22(9), 690–696. <https://doi.org/10.1111/j.1600-051X.1995.tb00828.x>
- Cochran, W. G. (1954). The combination of estimates from different experiments. *Biometrics*, 10, 101–129. <https://doi.org/10.2307/3001666>
- Dello Russo, N. M. (1981). Use of the fiber retention procedure in treating the maxillary anterior region. *Journal of Periodontology*, 52(4), 208–213. <https://doi.org/10.1902/jop.1981.52.4.208>
- DerSimonian, R., & Laird, N. (1986). Meta-analysis in clinical trials. *Controlled Clinical Trials*, 7(3), 177–188. [https://doi.org/10.1016/0197-2456\(86\)90046-2](https://doi.org/10.1016/0197-2456(86)90046-2)
- Geisinger, M. L., Mealey, B. L., Schoolfield, J., & Mellonig, J. T. (2007). The effectiveness of subgingival scaling and root planing: An evaluation of therapy with and without the use of the periodontal endoscope. *Journal of Periodontology*, 78(1), 22–28. <https://doi.org/10.1902/jop.2007.060186>
- Graziani, F., Gennai, S., Cei, S., Cairo, F., Baggiani, A., Miccoli, M., ... Tonetti, M. (2012). Clinical performance of access flap surgery in the treatment of the intrabony defect. A systematic review and meta-analysis of randomized clinical trials. *Journal of Clinical Periodontology*, 39(2), 145–156. <https://doi.org/10.1111/j.1600-051X.2011.01815.x>
- Graziani, F., Karapetsa, D., Alonso, B., & Herrera, D. (2017). Nonsurgical and surgical treatment of periodontitis: How many options for one disease? *Periodontology 2000*, 75(1), 152–188. <https://doi.org/10.1111/prd.12201>
- Graziani, F., Karapetsa, D., Mardas, N., Leow, N., & Donos, N. (2018). Surgical treatment of the residual periodontal pocket. *Periodontology 2000*, 76(1), 150–163. <https://doi.org/10.1111/prd.12156>
- Harrel, S. K., & Nunn, M. E. (2001). Longitudinal comparison of the periodontal status of patients with moderate to severe periodontal disease receiving no treatment, non-surgical treatment, and surgical treatment utilizing individual sites for analysis. *Journal of Periodontology*, 72(11), 1509–1519. <https://doi.org/10.1902/jop.2001.72.11.1509>
- Heitz-Mayfield, L. J., Trombelli, L., Heitz, F., Needleman, I., & Moles, D. (2002). A systematic review of the effect of surgical debridement vs non-surgical debridement for the treatment of chronic periodontitis. *Journal of Clinical Periodontology*, 29 (Suppl 3), 92–102; discussion 160–102. <https://doi.org/10.1034/j.1600-051X.29.s3.5.x>
- Higgins, J. P., & Green, S. (2011). *Cochrane handbook for systematic reviews of interventions version 5.1.0*. Oxford, UK: Cochrane Collaboration. <https://doi.org/10.1097/01.prs.0000173862.20563.12>
- Higgins, J. P., Thompson, S. G., Deeks, J. J., & Altman, D. G. (2003). Measuring inconsistency in meta-analyses. *BMJ*, 327(7414), 557–560. <https://doi.org/10.1136/bmj.327.7414.557>
- Hill, R. W., Ramfjord, S. P., Morrison, E. C., Appleberry, E. A., Caffesse, R. G., Kerry, G. J., & Nissle, R. R. (1981). Four types of periodontal treatment compared over two years. *Journal of Periodontology*, 52(11), 655–662. <https://doi.org/10.1902/jop.1981.52.11.655>
- Hujoel, P. P. (2004). Endpoints in periodontal trials: the need for an evidence-based research approach. *Periodontology*, 2000(36), 196–204. <https://doi.org/10.1111/j.1600-0757.2004.03681.x>
- Isidor, F., Karring, T., & Attström, R. (1984). The effect of root planing as compared to that of surgical treatment. *Journal of Clinical Periodontology*, 11(10), 669–681. <https://doi.org/10.1111/j.1600-051x.1984.tb01315.x>
- Isidor, F., Attström, R., & Karring, T. (1985). Regeneration of alveolar bone following surgical and non-surgical periodontal treatment. *Journal of Clinical Periodontology*, 12(8), 687–696. <https://doi.org/10.1111/j.1600-051X.1985.tb00940.x>
- Isidor, F., & Karring, T. (1986). Long-term effect of surgical and non-surgical periodontal treatment. A 5-year clinical study. *Journal of Periodontal Research*, 21(5), 462–472. <https://doi.org/10.1111/j.1600-0765.1986.tb01482.x>
- Jepsen, S., Deschner, J., Braun, A., Schwarz, F., & Eberhard, J. (2011). Calculus removal and the prevention of its formation. *Periodontology 2000*, 55(1), 167–188. <https://doi.org/10.1111/j.1600-0757.2010.00382.x>
- Kaldahl, W. B., Kalkwarf, K. L., Patil, K. D., Dyer, J. K., & Bates, R. E. Jr (1988). Evaluation of four modalities of periodontal therapy. Mean probing depth, probing attachment level and recession changes. *Journal of Periodontology*, 59(12), 783–793. <https://doi.org/10.1902/jop.1988.59.12.783>
- Kaldahl, W. B., Kalkwarf, K. L., Patil, K. D., & Molvar, M. P. (1990a). Evaluation of gingival suppuration and supragingival plaque following 4 modalities of periodontal therapy. *Journal of Clinical Periodontology*, 17(9), 642–649.
- Kaldahl, W. B., Kalkwarf, K. L., Patil, K. D., & Molvar, M. P. (1990b). Responses of four tooth and site groupings to periodontal therapy. *Journal of Periodontology*, 61(3), 173–179. <https://doi.org/10.1902/jop.1990.61.3.173>
- Kaldahl, W. B., Kalkwarf, K. L., Patil, K. D., & Molvar, M. P. (1990c). Relationship of gingival bleeding, gingival suppuration, and supragingival plaque to attachment loss. *Journal of Periodontology*, 61(6), 347–351. <https://doi.org/10.1902/jop.1990.61.6.347>
- Kaldahl, W. B., Kalkwarf, K. L., Patil, K. D., Molvar, M. P., & Dyer, J. K. (1996a). Long-term evaluation of periodontal therapy: I. Response

- to 4 therapeutic modalities. *Journal of Periodontology*, 67(2), 93–102. <https://doi.org/10.1902/jop.1996.67.2.93>
- Kaldahl, W. B., Kalkwarf, K. L., Patil, K. D., Molvar, M. P., & Dyer, J. K. (1996b). Long-term evaluation of periodontal therapy: II. Incidence of sites breaking down. *Journal of Periodontology*, 67(2), 103–108. <https://doi.org/10.1902/jop.1996.67.2.103>
- Kalkwarf, K. L., Kaldahl, W. B., Patil, K. D., & Molvar, M. P. (1989). Evaluation of gingival bleeding following 4 types of periodontal therapy. *Journal of Clinical Periodontology*, 16(9), 601–608. <https://doi.org/10.1111/j.1600-051X.1989.tb02145.x>
- Kalkwarf, K. L., Kaldahl, W. B., & Patil, K. D. (1992). Patient preference regarding 4 types of periodontal therapy following 3 years of maintenance follow-up. *Journal of Clinical Periodontology*, 19(10), 788–793. <https://doi.org/10.1111/j.1600-051x.1992.tb02172.x>
- Kirkland, O. (1931). The suppurative periodontal pus pocket; its treatment by the modified flap operation. *Journal of the American Dental Association*, 18, 1462–1470.
- Laleman, I., Cortellini, S., De Winter, S., Rodriguez Herrero, E., Dekeyser, C., Quirynen, M., & Teughels, W. (2017). Subgingival debridement: End point, methods and how often? *Periodontology 2000*, 75(1), 189–204. <https://doi.org/10.1111/prd.12204>
- Lindhe, J., & Nyman, S. (1985). Scaling and granulation tissue removal in periodontal therapy. *Journal of Clinical Periodontology*, 12(5), 374–388. <https://doi.org/10.1111/j.1600-051X.1985.tb00928.x>
- Lindhe, J., Westfelt, E., Nyman, S., Socransky, S. S., & Haffajee, A. D. (1984). Long-term effect of surgical/non-surgical treatment of periodontal disease. *Journal of Clinical Periodontology*, 11(7), 448–458. <https://doi.org/10.1111/j.1600-051X.1984.tb01344.x>
- Lindhe, J., Westfelt, E., Nyman, S., Socransky, S. S., Heijl, L., & Bratthall, G. (1982a). Healing following surgical non-surgical treatment of periodontal disease: A clinical study. *Journal of Clinical Periodontology*, 9(2), 115–128. <https://doi.org/10.1111/j.1600-051X.1982.tb01227.x>
- Lindhe, J., Socransky, S. S., Nyman, S., Haffajee, A., & Westfelt, E. (1982b). "Critical probing depths" in periodontal therapy. *Journal of Clinical Periodontology*, 9(4), 323–336. <https://doi.org/10.1111/j.1600-051x.1982.tb02099.x>
- Matuliene, G., Pjetursson, B. E., Salvi, G. E., Schmidlin, K., Bragger, U., Zwahlen, M., & Lang, N. P. (2008). Influence of residual pockets on progression of periodontitis and tooth loss: Results after 11 years of maintenance. *Journal of Clinical Periodontology*, 35(8), 685–695. <https://doi.org/10.1111/j.1600-1051X.2008.01245.x>
- Michaelides, P. L., & Wilson, S. G. (1996). A comparison of papillary retention versus full-thickness flaps with internal mattress sutures in anterior periodontal surgery. *International Journal of Periodontics and Restorative Dentistry*, 16(4), 388–397.
- Moher, D., Hopewell, S., Schulz, K. F., Montori, V., Gøtzsche, P. C., Devereaux, P. J., ... Altman, D. G. (2012). CONSORT 2010 explanation and elaboration: Updated guidelines for reporting parallel group randomised trials. *International Journal of Surgery*, 10(1), 28–55. <https://doi.org/10.1016/j.ijsu.2011.10.001>
- Moher, D., Liberati, A., Tetzlaff, J., & Altman, D. G. (2009). Preferred reporting items for systematic reviews and meta-analyses: The PRISMA statement. *Journal of Clinical Epidemiology*, 62(10), 1006–1012. <https://doi.org/10.1016/j.jclinepi.2009.06.005>
- Nibali, L., Pometti, D., Chen, T. T., & Tu, Y. K. (2015). Minimally invasive non-surgical approach for the treatment of periodontal intrabony defects: A retrospective analysis. *Journal of Clinical Periodontology*, 42(9), 853–859. <https://doi.org/10.1111/jcpe.12443>
- Nibali, L., Yeh, Y. C., Pometti, D., & Tu, Y. K. (2018). Long-term stability of intrabony defects treated with minimally invasive non-surgical therapy. *Journal of Clinical Periodontology*, 45(12), 1458–1464. <https://doi.org/10.1111/jcpe.13021>
- Perumal, M. P. B., Ramegowda, A. D., Lingaraju, A. J., & Raja, J. J. (2015). Comparison of microsurgical and conventional open flap debridement: A randomized controlled trial. *Journal of Indian Society of Periodontology*, 19(4), 406–410. <https://doi.org/10.4103/0972-124X.156884>
- Pihlstrom, B. L., McHuon, R. B., Oliphant, T. H., & Ortiz-Campos, C. (1983). Comparison of surgical and nonsurgical treatment of periodontal disease A review of current studies and additional results after 6 1/2 years. *Journal of Clinical Periodontology*, 10(5), 524–541. <https://doi.org/10.1111/j.1600-051X.1983.tb02182.x>
- Pihlstrom, B. L., Oliphant, T. H., & McHugh, R. B. (1984). Molar and non-molar teeth compared over 6 1/2 years following two methods of periodontal therapy. *Journal of Periodontology*, 55(9), 499–504. <https://doi.org/10.1902/jop.1984.55.9.499>
- Pihlstrom, B. L., Ortiz-Campos, C., & McHugh, R. B. (1981). A randomized four-years study of periodontal therapy. *Journal of Periodontology*, 52(5), 227–242. <https://doi.org/10.1902/jop.1981.52.5.227>
- Pini Prato, G. P., Di Gianfilippo, R., & Wang, H. L. (2019). Success in periodontology: An evolutive concept. *Journal of Clinical Periodontology*, 46(8), 840–845. <https://doi.org/10.1111/jcpe.13150>
- Polansky, R., Haas, M., Lorenzoni, M., Wimmer, G., & Pertl, C. (2003). The effect of three different periodontal pre-treatment procedures on the success of telescopic removable partial dentures. *Journal of Oral Rehabilitation*, 30(4), 353–363. <https://doi.org/10.1046/j.1365-2842.2003.01057.x>
- Rabbani, G. M., Ash, M. M. Jr, & Caffesse, R. G. (1981). The effectiveness of subgingival scaling and root planing in calculus removal. *Journal of Periodontology*, 52(3), 119–123. <https://doi.org/10.1902/jop.1981.52.3.119>
- Ramfjord, S. P., Caffesse, R. G., Morrison, E. C., Hill, R. W., Kerry, G. J., Appleberry, E. A., ... Stults, D. L. (1987). 4 modalities of periodontal treatment compared over 5 years. *Journal of Clinical Periodontology*, 14(8), 445–452. <https://doi.org/10.1111/j.1600-051X.1987.tb02249.x>
- Ramfjord, S. P., & Nissle, R. R. (1974). The modified widman flap. *Journal of Periodontology*, 45(8), 601–607. <https://doi.org/10.1902/jop.1974.45.8.2.601>
- Ramfjord, S. P., Knowles, J. W., Nissle, R. R., Burgett, F. G., & Shick, R. A. (1975). Results following three modalities of periodontal therapy. *Journal of Periodontology*, 46(9), 522–526. <https://doi.org/10.1902/jop.1975.46.9.522>
- Rateitschak-Pluss, E. M., Schwarz, J. P., Guggenheim, R., Duggelin, M., & Rateitschak, K. H. (1992). Non-surgical periodontal treatment: Where are the limits? An SEM study. *Journal of Clinical Periodontology*, 19(4), 240–244. <https://doi.org/10.1111/j.1600-051X.1992.tb00460.x>
- Reddy, S., Bhowmik, N., Prasad, M. G. S., Kaul, S., Rao, V., & Singh, S. (2014). Evaluation of postsurgical clinical outcomes with/without removal of pocket epithelium: A split mouth randomized trial. *Journal of Indian Society of Periodontology*, 18(6), 751–761. <https://doi.org/10.4103/0972-124X.147413>
- Renvert, S., Garrett, S., Nilvéus, R., Chamberlain, A. D. H., & Egelberg, J. (1985). Healing after treatment of periodontal intraosseous defects: VI. Factors influencing the healing response. *Journal of Clinical Periodontology*, 12(9), 707–715. <https://doi.org/10.1111/j.1600-051X.1985.tb01396.x>
- Renvert, S., Nilvéus, R., Dahlén, G., Slots, J., & Egelberg, J. (1990). 5-year follow up of periodontal intraosseous defects treated by root planing or flap surgery. *Journal of Clinical Periodontology*, 17(6), 356–363. <https://doi.org/10.1111/j.1600-051X.1990.tb00031.x>
- Renvert, S., Nilvéus, R., & Egelberg, J. (1985). Healing after treatment of periodontal intraosseous defects. V. Effect of root planing versus flap surgery. *Journal of Clinical Periodontology*, 12(8), 619–629. <https://doi.org/10.1111/j.1600-051X.1985.tb00933.x>
- Ribeiro, F. V., Casarin, R. C. V., Palma, M. A. G., Júnior, F. H. N., Sallum, E. A., & Casati, M. Z. (2011). Clinical and Patient-centered outcomes after minimally invasive Non-surgical or surgical approaches for the treatment of intrabony defects: A randomized clinical trial. *Journal*

- of *Periodontology*, 82(9), 1256–1266. <https://doi.org/10.1902/jop.2011.100680>
- Roberts, F. A., & Darveau, R. P. (2015). Microbial protection and virulence in periodontal tissue as a function of polymicrobial communities: Symbiosis and dysbiosis. *Periodontology 2000*, 69(1), 18–27. <https://doi.org/10.1111/prd.12087>
- Sanz, I., Alonso, B., Carasol, M., Herrera, D., & Sanz, M. (2012). Nonsurgical treatment of periodontitis. *Journal of Evidence Based Dental Practice*, 12(3 Suppl), 76–86. [https://doi.org/10.1016/S1532-3382\(12\)70019-2](https://doi.org/10.1016/S1532-3382(12)70019-2)
- Serino, G., Rosling, B., Ramberg, P., Socransky, S. S., & Lindhe, J. (2001). Initial outcome and long-term effect of surgical and non-surgical treatment of advanced periodontal disease. *Journal of Clinical Periodontology*, 28(10), 910–916. <https://doi.org/10.1034/j.1600-051x.2001.028010910.x>
- Smiley, C. J., Tracy, S. L., Abt, E., Michalowicz, B. S., John, M. T., Gunsolley, J., ... Hanson, N. (2015). Evidence-based clinical practice guideline on the nonsurgical treatment of chronic periodontitis by means of scaling and root planing with or without adjuncts. *Journal of the American Dental Association*, 146(7), 525–535. <https://doi.org/10.1016/j.adaj.2015.01.026>
- Svoboda, P. J., Reeve, C. M., & Sheridan, P. J. (1984). Effect of retention of gingival sulcular epithelium on attachment and pocket depth after periodontal surgery. *Journal of Periodontology*, 55(10), 563–566. <https://doi.org/10.1902/jop.1984.1955.1910.1563>
- Takei, H. H., Han, T. J., Carranza, F. A. Jr, Kenney, E. B., & Lekovic, V. (1985). Flap technique for periodontal bone implants. Papilla preservation technique. *Journal of Periodontology*, 56(4), 204–210. <https://doi.org/10.1902/jop.1985.56.4.204>
- Tobias, A., & Campbell, M. J. (1999). Modelling influenza epidemics in the relation between black smoke and total mortality. A sensitivity analysis. *Journal of Epidemiology and Community Health*, 53(9), 583–584.
- Tomasi, C., Koutouzis, T., & Wennstrom, J. L. (2008). Locally delivered doxycycline as an adjunct to mechanical debridement at retreatment of periodontal pockets. *Journal of Periodontology*, 79(3), 431–439. <https://doi.org/10.1902/jop.2008.070383>
- Trombelli, L., Simonelli, A., Schincaglia, G. P., Cucchi, A., & Farina, R. (2012). Single-flap approach for surgical debridement of deep intraosseous defects: A randomized controlled trial. *Journal of Periodontology*, 83(1), 27–35. <https://doi.org/10.1902/jop.2011.110045>
- Van der Weijden, G. A., & Timmerman, M. F. (2002). A systematic review on the clinical efficacy of subgingival debridement in the treatment of chronic periodontitis. *Journal of Clinical Periodontology*, 29 (Suppl 3), 55–71; discussion 90–51. <https://doi.org/10.1034/j.1600-051X.29.s3.3.x>
- Waerhaug, J. (1978). Healing of the dento-epithelial junction following subgingival plaque control. II: As observed on extracted teeth. *Journal of Periodontology*, 49(3), 119–134. <https://doi.org/10.1902/jop.1978.49.3.119>
- Wennström, A., Wennström, J., & Lindhe, J. (1986). Healing following surgical and non-surgical treatment of juvenile periodontitis: A 5-year longitudinal study. *Journal of Clinical Periodontology*, 13(9), 869–882. <https://doi.org/10.1111/j.1600-051X.1986.tb02245.x>
- Westfelt, E., Bragd, L., Socransky, S. S., Haffajee, A. D., Nyman, S., & Lindhe, J. (1985). Improved periodontal conditions following therapy. *Journal of Clinical Periodontology*, 12(4), 283–293. <https://doi.org/10.1111/j.1600-051X.1985.tb0>

SUPPORTING INFORMATION

Additional supporting information may be found online in the Supporting Information section.

How to cite this article: Sanz-Sánchez I, Montero E, Citterio F, Romano F, Molina A, Aimetti M. Efficacy of access flap procedures compared to subgingival debridement in the treatment of periodontitis. A systematic review and meta-analysis. *J Clin Periodontol*. 2020;(Suppl. 22):282–302. <https://doi.org/10.1111/jcpe.13259>