AperTO - Archivio Istituzionale Open Access dell'Università di Torino

Characterization and growth under different storage temperatures of ropy slime-producing *Leuconostoc mesenteroides* isolated from cooked meat products

This is the author's manuscript

Original Citation:

Availability:

This version is available <http://hdl.handle.net/2318/1761088> since 2020-11-04T11:16:58Z

Published version:

DOI:10.4315/JFP-19-521

Terms of use:

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)

Journal of Food Protection

Characterization and growth under different storage temperatures of ropy slime-producing *Leuconostoc mesenteroides* isolated from cooked meat products --Manuscript Draft--

Manuscript Number:	JFP-19-521R2
Article Type:	Research Note
Section/Category:	Food Microbiology
Keywords:	Ropy slime; <i>Leuconostoc mesenteroides</i> ; meat spoilage; growth curves
Corresponding Author:	Luca Grispoldi Dipartimento di medicina Veterinaria, Università degli Studi di Perugia Perugia, ITALY
First Author:	Beniamino T. Cenci Goga
Order of Authors:	Beniamino T. Cenci Goga
	Paola Sechi
	Maria F. Iulietto
	Shahin Amirjalali
	Salvatore Barbera
	Musafiri Karama
	Sharif A Aly, DVM
	Luca Grispoldi
Manuscript Region of Origin:	ITALY
Abstract:	<p>The presence of lactic acid bacteria (LAB) can be detrimental when the abundant growth of slime-producing strains (<i>Lactobacillus</i> spp. and <i>Leuconostoc</i> spp.) causes spoilage of meat products. Two strains of LAB were isolated from vacuum packed, cooked hams, which had been withdrawn from the market for the so-called «ropy slime» defect and identified as <i>Leuconostoc mesenteroides</i>. In an attempt to define the behaviour of ropy slime-producing bacteria, two strains of <i>L. mesenteroides</i> were incubated in MRS broth at different storage temperatures and conditions of thermal abuse (4, 12, 20, 30, 37, 44 °C). Both strains showed a lack of growth at 44°C, a good level of development at 30 and 37°C and evident growth ability at low temperatures with a long stationary phase. In particular, the bacterial concentration at 4°C was above 10⁵ cfu ml⁻¹ after over 120 days of incubation. This research demonstrates that the refrigeration temperature for cooked meat products does not constitute a hurdle for ropy slime-producers and their subsequent ability to spoil.</p>

Received: November 4, 2019; Accepted: January 8, 2019; Published Online Early: January 2020

Beniamino T. Cenci-Goga, Paola Sechi, Maria F. Iulietto, Shahin Amirjalali, Salvatore Barbera, Musafiri Karama, Sharif S. Aly, Luca Grispoldi. Characterization and growth under different storage temperatures of ropy slime-producing *Leuconostoc mesenteroides* isolated from cooked meat products. *Journal of Food Protection*. <https://doi.org/10.4315/JFP-19-521>

This Online Early paper will appear in its final typeset version in a future issue of the *Journal of Food Protection*. This article has been accepted for publication and undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the Version of Record.

Running head: Growth dynamics of ropy slime producing *Leuconostoc mesenteroides*

Research note

Characterization and growth under different storage temperatures of ropy slime-producing *Leuconostoc mesenteroides* isolated from cooked meat products

Beniamino T. Cenci-Goga^{1,3}, Paola Sechi¹, Maria F. Iulietto¹, Shahin Amirjalali¹, Salvatore Barbera², Musafiri Karama³, Sharif S. Aly⁴, Luca Grispoldi^{1*}

¹Dipartimento di Medicina Veterinaria, Laboratorio di Ispezione degli Alimenti di Origine Animale, Università degli Studi di Perugia, 06126 Perugia, Italy

²Dipartimento di Scienze Agrarie, Forestali e Alimentari, Università di Torino, Italy

³Department of Paraclinical Sciences, University of Pretoria, Onderstepoort, South Africa

⁴UC Davis, School of Veterinary Medicine, CA, USA

Keywords: Ropy slime, *Leuconostoc mesenteroides*, meat spoilage, growth curves

* Corresponding author: Luca Grispoldi, Dipartimento di Medicina veterinaria, Università di Perugia, via San Costanzo, 4 - 06126 Perugia, Italy, tel.: +39 075 585 7935, fax: 7976, email: grisluca@outlook.it

ABSTRACT

The presence of lactic acid bacteria (LAB) can be detrimental when the abundant growth of slime-producing strains (*Lactobacillus* spp. and *Leuconostoc* spp.) causes spoilage of meat products. Two strains of LAB were isolated from vacuum packed, cooked hams, which had been withdrawn from the market for the so-called «ropy slime» defect and identified as *Leuconostoc mesenteroides*. In an attempt to define the behaviour of ropy slime-producing bacteria, two strains of *L. mesenteroides* were incubated in MRS broth at different storage temperatures and conditions of thermal abuse (4, 12, 20, 30, 37, 44 °C). Both strains showed a lack of growth at 44°C, a good level of development at 30 and 37°C and evident growth ability at low temperatures with a long stationary phase. In particular, the bacterial concentration at 4°C was above 10⁵ cfu ml⁻¹ after over 120 days of incubation. This research demonstrates that the refrigeration temperature for cooked meat products does not constitute a hurdle for ropy slime-producers and their subsequent ability to spoil.

HIGHLIGHTS

- Lactic acid bacteria can be detrimental when slime-producing strains cause meat spoilage.
- *L. mesenteroides* from cooked ham investigated at different temperature.
- Refrigeration temperature is not a hurdle for slime-producer strains

44 Meat spoilage is one of the most important causes of food waste and market rejection of meat
45 products (12, 38). The shelf-life of meat depends on the type of bacteria initially present and their
46 ability to grow on food. Favourable growth conditions during storage, such as storage temperature,
47 should be determined for each bacterial genus involved (16, 22). Knowledge of bacterial growth
48 kinetics is, therefore, considered the first step to prevent the development of spoilage microbiota on
49 meat products (8, 37). Although a large variety of species can be isolated from meat and meat
50 products, the majority of attention has focused on lactic acid bacteria (LAB), widely found in nature
51 and in the processing plants (30, 34). They are «Generally Regarded As Safe» (GRAS) micro-
52 organisms (18, 20, 30, 32) and are not only traditionally applied to the manufacture of fermented
53 meat products (4, 5, 7, 9, 48), such as salami, but are also used as biopreservatives (9, 43, 44, 46).
54 However, some strains of LAB are defined as specific spoilage organisms (SSO) and represent the
55 major cause of spoilage of vacuum and modified-atmosphere packed, cooked meat products (25,
56 30, 31, 35). Typical, detectable, organoleptic alterations are: off-flavours, discoloration, gas
57 production, an excessive decrease in pH and slime formation, resulting in the reduction of the
58 product's shelf-life (2). Spoilage can occur within the shelf-life period and this requires producers to
59 make withdrawals (17). Even though these bacteria are usually below detection level ($<10 \text{ cfu g}^{-1}$)
60 after packaging, the products may spoil quickly at a later stage (13, 27, 39). Among the sensory
61 changes, the accumulation of ropy slime on the surface of cooked meat products is probably one of
62 the most important and least tackled defects (28, 34). Ropy slime-producing LAB are able to grow
63 and survive at refrigeration temperatures, competing with other bacteria in meat products and meat
64 processing plants (24, 41). Consequently, the use of low temperatures in the preparation and storage
65 of meat products may not prevent the formation of ropy-slime, even though refrigeration storage
66 temperatures determine a longer shelf-life of the product (19). The constant bacterial growth is
67 favoured by the slime formation, which creates a barrier between the surrounding environment and
68 competitors (45). The slime is due to long-chain, high-molecular-mass, viscosifying or gelling,
69 exocellular polysaccharides (10).

Despite the many studies on the control of *Leuconostoc* spp. spoilage that have been carried out by the academia and the industry in recent years (11, 20, 23, 27, 36), there has been no agreed methodology nor qualified studies for the evaluation of growth dynamics of *L. mesenteroides*. This study was therefore intended to evaluate the behaviour of two slime-producing *L. mesenteroides* strains at different storage temperatures and thermal abuse conditions, isolated from vacuum packed cooked hams, which had been withdrawn from the market due to the so-called ropy slime defect.

MATERIALS AND METHODS

Origin of *L. mesenteroides* strains. Strains were isolated from commercial cooked ham presenting slime on the surface, during a thorough study on *Leuconostoc* spp. contamination of cooked meat products. Samples were taken from a batch of commercial, cooked ham, withdrawn from the market due to the so-called ropy slime defect in order to isolate and identify the causative agent. Cooked ham had been produced by a large Italian company that exports products to the US market. The production is done according to a traditional recipe, after selecting fresh pork thighs which have previously been defatted and deboned.

Isolation and identification of *L. mesenteroides*. Approximately 25 cm² of ham were aseptically swabbed and transferred to 225 ml of sterile, buffered, peptone water (PW, Oxoid, Milan, Italy, CM1049), and homogenized at room temperature in a stomacher (PBI International) for 1 min at low speed and 1 min at high speed. Serial decimal dilutions in buffered PW were prepared and triplicate 1 ml or 0.1 ml samples of appropriate dilutions were poured or spread on de Man, Rogosa, Sharpe (MRS) Agar (Oxoid, CM0361), added with Vancomycin (20 µg/ml) to create a selective media for vancomycin-resistant bacteria (29, 40) and incubated for 48 hours at 30°C in microaerophilic conditions and further purified by four steps of sequential streaking on MRS-Va agar. Circular shaped colonies, with a convex elevation, 3-4 mm high, 4-6 mm in diameter, and a smooth surface, an even margin and opaque density, were presumed to be *Leuconostoc* spp. and

95 subcultured on MSE Agar (Mayeux, Sandine and Elliker, Biolife, Milan, Italy) and incubated at
 96 30°C for 48 hours. Colonies grown on MSE Agar with a gelatinous appearance were primarily
 97 characterized by means of Gram staining, cell morphology, catalase reaction of 3% hydrogen
 98 peroxide (H₂O₂) and oxidase reactions. Gas production from glucose was evaluated on MRS broth
 99 (Oxoid, CM0359) with a Durham tube, after incubation in air at 30°C for 48 hours. Bacterial
 100 motility was detected using the hanging drop method.

101 Gram positive, catalase negative and oxidase negative, gas-producing, non-motile bacteria were
 102 transferred to MRS broth and incubated at 30°C for 24 hours and then stored as stock cultures
 103 at -80°C for further examination.

104 **Isolation and purification of DNA, oligonucleotide primers and identification by**
 105 **sequencing.** Template DNA was obtained using the method described by Cenci Goga et al. (6).
 106 Primer sequences, lengths of PCR amplification products and amplification conditions are listed in
 107 Table 1. Briefly, genus-specific primers Lu1r and Lu2 were used (MWG Biotech, Ebersberg, DE)
 108 (47) followed by *L. mesenteroides* identification by primers that target the 16S rRNA gene: L.mesF
 109 and L.mesR (MWG Biotech) (3, 26). Universal primers Y1 and Y2 for bacterial 16S rRNA gene
 110 were used as a positive control to ensure that the template DNA was correctly amplified (47). PCR
 111 products were visualized after agarose gel electrophoresis under UV illumination (15). The
 112 universal primers for eubacteria P27f and P1495r were used to amplify a 16S rRNA gene segment
 113 (9, 33). The amplicons were sent to the Microgem Lab (University of Naples, Italy) for purifying
 114 and sequencing. Isolates were allocated to a given species on percentages of sequencing identity
 115 and on visual inspection of the concordance, using the DIALIGN software
 116 (<http://dialign.gobics.de/chaos-dialign-submission>).

117 **Growth dynamics of *L. mesenteroides*.** Each strain examined for its growth curve was
 118 transferred in MRS broth (Oxoid, CM0359) tubes and incubated at 30°C for 48 hours to get a
 119 concentration of approximately 10⁸ cfu ml⁻¹. A dilution series to get a final concentration of 10⁴ cfu

ml⁻¹ was performed in MRS broth before incubation at different temperatures for appropriate periods. A Sanyo MIR-153 incubator (Moriguchi-City Osaka, Japan) was used to maintain storage temperatures of 4, 12, 20, 30, 37 and 44°C. Samples were taken in triplicate at 6, 12, 24, 36, and 48 h and then, every 24 h until the bacterial concentration fell below the detection limit (10 cfu ml⁻¹). At each sampling point, 1 ml of sample was transferred aseptically to 9 ml of Maximum Recovery Diluent (Oxoid, CM0733) for serial decimal dilutions. Duplicate samples of appropriate dilutions were poured or spread plated on MRS Agar (Oxoid, CM1153) and incubated at 30°C for 48 hours in jars (Oxoid) under microaerophilic conditions. Growth rates based on the viable counts on MRS agar were determined with a polynomial curve fitting. The sensitivity of the spread plate was 10² cfu ml⁻¹ and of the pour plate was 10 cfu ml⁻¹. The 95% confidence limit, as given by the classic formula $2s=2\sqrt{x}$ (1), ranged between ±37% and ±12% (*i.e.* plates with a number of cfu ranging from 30 to 300). Consequently, plates with less than 30 cfu were not considered for data analysis and when this applied to the lowest dilution, the results were recorded as <30 for the pour plate and <300 for the spread plate (4).

Analysis of the results. Each triplicate tube was examined in duplicate for each sampling, and all values were converted to log for microbiological analyses and analysed using GraphPad InStat, version 3.0b for Mac OS X. A Canonical Discriminant Analysis (CDA) was performed using 4 parameters to show the temperature action on the 2 strains: the intercept of the fourth-degree polynomial equation (a_0); day ($D_{\max}$) and log ($Y_{\max}$) at peak; growth rate (b).

A fourth-degree polynomial equation was used as an empirical model to fit the experimental data.

$$[1] \quad y=a_0+a_1x+a_2x^2+a_3x^3+a_4x^4$$

where: y = log population; x = time from 0 to 180 days; a_0 , a_1 , a_2 , a_3 , a_4 = coefficients of polynomial determined by the function «PROC REG» regression procedure in SAS 9.4 (42). The first derivative of [1] gives the maximum value for log population and days ($Y_{\max}$ $D_{\max}$), which

corresponds to the peak point of the growth curve. The growth rate from day 0 to $D_{\max}$ was calculated as the slope of a linear equation as in [2].

$$[2] \quad b=(y-a)/x$$

where: y = log concentration; x = time from 0 to $D_{\max}$ days; a = intercept or initial concentration for $d=0$. Data analysis was performed by SAS/STAT in SAS 9.4 (42) using a regression model (REG). The canonical correlation is a multivariate analysis of correlation. Canonical is the statistical term for analysing latent variables (which are not directly observed) that represent multiple variables (which are directly observed). A Canonical Discriminant Analysis (CDA) is a dimension-reduction technique related to principal component analysis and canonical correlation. In this study CDA finds linear combinations (canonical variables) of the quantitative variables (different temperatures) that have the highest possible multiple correlations with the groups and provide maximal separation between groups in much the same way that principal components summarize total variation. The canonical variable can show substantial differences between the groups, even though none of the original variables do.

RESULTS

Isolation and identification of *L. mesenteroides*. According to the phenotypic and genotypic findings, two strains of ropy slime-producing were identified as *L. mesenteroides*. The isolates collected were Gram positive and catalase negative, vancomycin resistant coccoid bacteria, producing gas and acid by fermenting glucose. The identification via PCR and DNA sequencing defined the strains as *L. mesenteroides*. The two strains of ropy slime-producing *L. mesenteroides* were, therefore, identified and classified as 649 and 650 (Laboratory collection ID).

Growth dynamics of *L. mesenteroides* 649 and 650. Growth curves are graphically represented in Figure 1. Both strains of *L. mesenteroides* (649 and 650) started from an initial concentration of approximately 10^4 cfu ml⁻¹. The highest population level (3.58 log cfu ml⁻¹ and 3.94 log cfu ml⁻¹ for *L. mesenteroides* 649 and 650, respectively) was reached after 4 hours at 44°C. The bacterial concentration was below the

detection limit after 24 and 48 hours for *L. mesenteroides* 650 and 649, respectively. At 37°C, the maximum population (8 log cfu ml⁻¹) was observed after 2.5 and 2.6 days for *L. mesenteroides* 650 and 649, respectively. After 6 days of storage, cell concentration was still over 5 log cfu ml⁻¹ and then rapidly fell below the detection limit.

At 30°C, the highest population level (9.22 log cfu ml⁻¹ and 9.40 log cfu ml⁻¹ for *L. mesenteroides* 649 and 650) occurred after 48 hours. Bacterial concentration was below the detection limit after 8 days and 13 days for *L. mesenteroides* 649 and 650, respectively.

At 20°C, the maximum population (8.53 log cfu ml⁻¹ and 8.56 log cfu ml⁻¹ for *L. mesenteroides* 649 and 650 respectively) was reached after 8,8 days and 7,7 days for *L. mesenteroides* 649 and 650. At 12°C, the maximum population (8,1 log cfu ml⁻¹ and 8.16 log cfu ml⁻¹ for *L. mesenteroides* 649 and 650, respectively) was reached after 15.4 days and 13.7 days for *L. mesenteroides* 649 and 650; the stationary phase was maintained for two weeks. Then, bacterial concentration was below the detection limit after 120 days of incubation for both strains. At 4°C, the maximum population (8,9 log cfu ml⁻¹ and 8.97 log cfu ml⁻¹ for *L. mesenteroides* 649 and 650, respectively) was reached after 42.8 days and 44.7 days for *L. mesenteroides* 649 and 650. Bacterial concentration was maintained over 5 log cfu ml⁻¹ after 120 days of incubation for both strains.

Figures 1 and 2 show the effect of the different storage temperatures on the growth of two *L. mesenteroides* strains.

Canonical Discriminant Analysis. Table 2 indicates the estimated days of the growth peak (D_{max}), and log (Y_{max}) at peak and the growth rate (b) to reach the maximum viable cell concentration. With the exception of the abuse temperature of 44°C with an immediate decrease in the population, the maximum growth rate detected was at 30°C. The maximum length of complete inhibition was described during each temporal slot. However, viable cells were still detectable at 4°C. Table 3 shows the polynomial, descriptive parameters for both strains and the growth

temperature. Figure 2 (CDA) shows the effect of the temperature on the two strains. The first canonical variable explained 99% of the among-class separation and all 4 parameters contributed significantly. Four groups (12°C-20°C-30°C; 37°C; 4°C and 44°C) are clearly identified, mostly by $D_{\max}$ and $Y_{\max}$. The univariate statistics result in R^2 values ranging from 0.903 for $Y_{\max}$ to 0.999 for $D_{\max}$ and each variable is significant above the 0.005 level.

DISCUSSION

The microbiota of many valuable foods, which are susceptible to bacterial spoilage, is usually dominated by LAB (14). Refrigerated meat products can show several defects, such as discoloration, gas production, off-odours, off-flavours, a decrease in pH and slime production (19, 22). Kalschne, D. L. et al. (22) observed the formation of milky exudates at 45 days of storage. The initial LAB population was approximately $1.98 \log \text{cfu g}^{-1}$, and after 45 days of storage raised values of $7.59 \log \text{cfu g}^{-1}$ at 4 °C and $8.25 \log \text{cfu g}^{-1}$ at 8 °C. Therefore, *Lactobacillus curvatus*, *Lactobacillus sakei* were identified as the dominant spoilage bacteria of sliced, vacuum-packed, cooked ham after 45 days of storage and *Leuconostoc mesenteroides* as a minor component (22). Raimondi et al (2019) described the microbiota of sliced, cooked ham, packaged in a modified atmosphere: a few days after packaging, the LAB population of the samples was $2.9 \log \text{cfu g}^{-1}$ on average, whereas the amount increased to $7.7 \log \text{cfu g}^{-1}$ in the samples from the end of their shelf-life and in those rejected due to spoilage phenomena.

The results of our research, on the other hand, give a detailed description of the growth behaviour of two *L. mesenteroides* strains isolated from cooked meat products. The optimum growth temperature in MRS broth is 30°C, with a short lag phase followed by fast growth. Strain 649 reached the highest concentration ($9.22 \log \text{cfu ml}^{-1}$) on day 2.2 and 650 ($9.40 \log \text{cfu ml}^{-1}$) on day 1.9. On the contrary, no growth was recorded at 44°C, whereas the ability to grow at refrigeration temperatures was well documented at 12°C and 4°C. Low storage temperatures determined a slowing down of bacterial growth. However, the rate was steady and very high concentrations (up to $8.9 \log \text{cfu ml}^{-1}$)

were reached (strain 649 on day 42.8 and 650 on day 44.7 at 4°C). These findings show that starting from an initial concentration of 4 log cfu ml⁻¹, the maximum population concentration reached over 8 log cfu ml⁻¹ for all storage temperature considered (except for 44°C). However, the length of the lag phase stretched to a greater or lesser extent. In addition, it demonstrated that viable cells are still detectable after several days of incubation at refrigeration temperature (>5 log cfu ml⁻¹ on day 120) and this supports the hypothesis of the abundant bacterial growth and subsequent accumulation of slime on the surface of meat products, even though the cold chain is maintained throughout product shelf life. CDA (Figure 2) showed that four groups (12°C-20°C-30°C; 37°C; 4°C and 44°C) are clearly identified and that the growth dynamics of *L. mesenteroides* at 4°C and 12°C are more similar to the growth dynamics at 20°C and 30°C (fastest growth rate and highest bacterial concentration in cfu ml⁻¹) than the abuse at 40°C. From the bacterial growth perspective, the refrigeration at 4°C or at a minor temperature abuse of 12°C is worse than a mismanagement at 37°C or at 44°C.

This research highlighted the growth ability of two strains of *L. mesenteroides* at refrigeration temperatures. Their capability to grow and persist at refrigeration temperatures makes the exponential increase of their total population possible, by exploiting the nutrients and producing abundant exopolysaccharides to create a favourable surrounding environment. The latter is facilitated by the low concentration of the competitive microbiota present on the product following heat treatment (21). Maintenance of the cold chain, of paramount importance for food safety, appears not to be an obstacle for the growth of *L. mesenteroides*, given the ability of these microorganisms to grow well at refrigeration temperatures.

Author Contributions: Conceptualization, BCG and PS; methodology BCG, PS and MFI, software LG and SB; analysis, MFI, PS and EB; supervision, BCG and MK; writing, PS, MFI and BCG.

Acknowledgments: The authors wish to express sincere appreciation to the members of Polyglot for the careful reading and editing of the manuscript

Conflicts of Interest: The authors declare no conflict of interest.

REFERENCES

1. Adams, M. R., and M. O. Moss. 2000. Food Microbiology. Royal Society of Chemistry, Cambridge, UK.
2. Audenaert, K., K. D'Haene, K. Messens, T. Ruysen, P. Vandamme, and G. Huys. 2010. Diversity of lactic acid bacteria from modified atmosphere packaged sliced cooked meat products at sell-by date assessed by PCR-denaturing gradient gel electrophoresis. p. 12-8. *In*, Food Microbiol, vol. 27. England.
3. Bounaix, M.-S., V. Gabriel, H. Robert, S. Morel, M. Remaud-Siméon, B. Gabriel, and C. Fontagné-Faucher. 2010. Characterization of glucan-producing *Leuconostoc* strains isolated from sourdough. *Int J Food Microbiol.* 144:1-9.
4. Cenci Goga, B. T., M. Karama, P. Sechi, M. F. Iulietto, S. Novelli, R. Selvaggini, and S. Mattei. 2015 Growth inhibition of selected microorganisms by an association of dairy starter cultures and probiotics. *Ital J Anim Sci.* 14.
5. Cenci Goga, B. T., P. V. Rossitto, P. Sechi, S. Parmegiani, V. Cambiotti, and J. S. Cullor. 2012. Effect of selected dairy starter cultures on microbiological, chemical and sensory characteristics of swine and venison (Dama dama) nitrite-free dry-cured sausages. *Meat Sci.* 90:599-606.
6. Cenci-Goga, B. T., M. Karama, P. V. Rossitto, R. A. Morgante, and J. S. Cullor. 2003. Enterotoxin Production by *Staphylococcus aureus* Isolated from Mastitic Cows. *J Food Protect.* 66:1693-1696.
7. Cenci-Goga, B. T., M. Karama, P. Sechi, M. F. Iulietto, L. Grispoldi, R. Selvaggini, M. Ceccarelli, and S. Barbera. 2018. Fate of selected pathogens in spiked «SALAME NOSTRANO» produced without added nitrates following the application of NONIT™ technology. *Meat Sci.* 139:247-254.
8. Cenci-Goga, B. T., M. Karama, P. Sechi, M. F. Iulietto, S. Novelli, and S. Mattei. 2014. Evolution under different storage conditions of anomalous blue coloration of Mozzarella cheese

- 274 intentionally contaminated with a pigment-producing strain of *Pseudomonas fluorescens*. *Journal of*
275 *Dairy Science*. 97:6708-6718.
- 276 9. Cenci-Goga, B. T., M. Karama, P. Sechi, M. F. Iulietto, S. Novelli, R. Selvaggini, and S.
277 Barbera. 2016. Effect of a novel starter culture and specific ripening conditions on microbiological
278 characteristics of nitrate-free dry-cured pork sausages. *Ital J Anim Sci*. 15:358-374.
- 279 10. De Vuyst, L., and B. Degeest. 1999. Heteropolysaccharides from lactic acid bacteria. *FEMS*
280 *Microbiol Rev*. 23.
- 281 11. Dror, B., A. Savidor, B. B. Salam, N. Sela, Y. Lampert, P. Teper-Bamnolker, A. Daus, S.
282 Carmeli, S. Sela, and D. Eshela. 2019. High Levels of CO₂ Induce Spoilage by *Leuconostoc*
283 *mesenteroides* by Upregulating Dextran Synthesis Genes. *Appl Environ Microbiol*. 85.
- 284 12. FAO. 2011. Global food losses and food waste – Extent, causes and prevention. *In*, Rome.
- 285 13. Fontana, C., P. S. Cocconcelli, and G. Vignolo. 2006. Direct Molecular Approach to
286 Monitoring Bacterial Colonization on Vacuum-Packaged Beef. *Appl Environ Microbiol*. 72:5618-
287 5622.
- 288 14. Geeraerts, W., V. Pothakos, L. De Vuyst, and F. Leroy. 2017. Diversity of the dominant
289 bacterial species on sliced cooked pork products at expiration date in the Belgian retail. *Food*
290 *Microbiol*. 65:236-243.
- 291 15. Grispoldi, L., M. Karama, F. Ianni, A. La Mantia, L. Pucciarini, E. Camaioni, R. Sardella, P.
292 Sechi, B. Natalini, and T. B. Cenci-Goga. 2019. The Relationship between *S. aureus* and Branched-
293 Chain Amino Acids Content in Composite Cow Milk. *Animals*. 9.
- 294 16. Grispoldi, L., P. A. Popescu, M. Karama, V. Gullo, G. Poerio, E. Borgogni, P. Torlai, G.
295 Chianese, A. G. Fermani, P. Sechi, and B. Cenci-Goga. 2019. Study on the growth and enterotoxin
296 production by *Staphylococcus aureus* in canned meat before retorting. *Toxins*. 11.
- 297 17. Hamasaki, Y., M. Ayaki, H. Fuchu, M. Sugiyama, and H. Morita. 2003. Behavior of
298 psychrotrophic lactic acid bacteria isolated from spoiling cooked meat products. *Appl Environ*
299 *Microbiol*. 69:3668-71.
- 300 18. Iulietto, M. F., P. Sechi, E. Borgogni, and B. T. Cenci Goga. 2016. Antibiotic susceptibility
301 profiles of ropy slime-producing *Leuconostoc mesenteroides* isolated from cooked meat products.
302 *Microbiology Research*. 7:4-7.
- 303 19. Iulietto, M. F., P. Sechi, E. Borgogni, and B. T. Cenci-Goga. 2015. Meat Spoilage: A
304 Critical Review of a Neglected Alteration Due to Ropy Slime Producing Bacteria. *Ital J Anim Sci*.
305 14:316-326.
- 306 20. Iulietto, M. F., P. Sechi, E. Borgogni, and B. T. Cenci-Goga. 2016. Antibiotic susceptibility
307 profiles of ropy slime-producing *Leuconostoc mesenteroides* isolated from cooked meat products.
308 *Microb Res*. 7:4-7.

- 309 21. Iulietto, M. F., P. Sechi, S. Mattei, S. Novelli, and B. T. Cenci Goga. 2014. Ropy slime
310 formation on meat products: an old problem, a new concern. p. 155. *In*, LXVIII Annual meeting of
311 the Italian Society for Veterinary Sciences, Università di Pisa.
- 312 22. Kalschne, D. L., R. Womer, A. Mattana, C. M. P. Sarmiento, L. M. Colla, and E. Colla.
313 2015. Characterization of the spoilage lactic acid bacteria in “sliced vacuum-packed cooked ham”.
314 *Braz J Microbiol.* 46:173-181.
- 315 23. Khorsandi, A., E. Ziaee, E. Shad, M. Razmjooei, M. H. Eskandari, and M. Aminlari. 2018.
316 Antibacterial Effect of Essential Oils against Spoilage Bacteria from Vacuum-Packed Cooked
317 Cured Sausages. *J Food Protect.* 81:1386-1393.
- 318 24. Korkeala, H. J., P. M. Makela, and H. L. Suominen. 1990. Growth temperatures of ropy
319 slime-producing lactic acid bacteria. *J Food Protect.* 53:793-794.
- 320 25. Koutsoumanis, K. 2009. Modeling food spoilage in microbial risk assessment. *J Food Prot.*
321 72:425-7.
- 322 26. Lee, H. J., S.-Y. Park, and J. Kim. 2000. Multiplex PCR-based detection and identification
323 of *Leuconostoc* species. *FEMS Microbiol Lett.* 193:243-7.
- 324 27. Li, X., C. Li, H. Ye, Z. Wang, X. Wu, Y. Han, and B. Xu. 2019. Changes in the microbial
325 communities in vacuum-packaged smoked bacon during storage. *Food Microbiol.* 77:26-37.
- 326 28. Makela, P. M., H. J. Korkeala, and J. J. Laine. 1992. Ropy slime-producing lactic acid
327 bacteria contamination at meat processing plants. *Int J Food Microbiol.* 17:27-35.
- 328 29. Mathot, A., M. Kihal, H. Prevost, and C. Divies. 1994. Selective Enumeration of
329 *Leuconostoc* on Vancomycin Agar Media. *Int Dairy J.*
- 330 30. Nychas, G.-J. E., P. N. Skandamis, C. C. Tassou, and K. P. Koutsoumanis. 2008. Meat
331 spoilage during distribution. *Meat Sci.* 78:77-89.
- 332 31. Nychas, G. J. E., and P. Skandamis. 2005. Fresh meat spoilage and modified atmosphere
333 packaging (MAP). p. 461-502. *In* S. J.N. (ed.), Improving the safety of fresh meat Cambridge Press,
334 UK.
- 335 32. Ogier, J. C., E. Casalta, C. Farrokh, and A. Saihi. 2008. Safety assessment of dairy
336 microorganisms: the *Leuconostoc* genus. p. 286-90. *In*, *Int J Food Microbiol*, vol. 126. Netherlands.
- 337 33. Osimani, A., C. Garofalo, L. Aquilanti, V. Milanovic, and F. Clementi. 2015. Unpasteurised
338 commercial boza as a source of microbial diversity. *Int J Food Microbiol.* 194:62-70.
- 339 34. Pothakos, V., F. Devlieghere, F. Villani, J. Björkroth, and D. Ercolini. 2015. Lactic acid
340 bacteria and their controversial role in fresh meat spoilage. *Meat Sci.* 109:66-74.

- 341 35. Pothakos, V., B. Taminiau, G. Huys, C. Nezer, G. Daube, and F. Devlieghere. 2014.
 342 Psychrotrophic lactic acid bacteria associated with production batch recalls and sporadic cases of
 343 early spoilage in Belgium between 2010 and 2014. *Int J Food Microbiol.* 191:157-63.
- 344 36. Raimondi, S., R. Luciani, T. M. Sirangelo, A. Amaretti, A. Leonardi, A. Ulrici, G. Foca, G.
 345 D'Auria, A. Moya, V. Zuliani, T. M. Seibert, J. Soltoft-Jensen, and M. Rossi. 2019. Microbiota of
 346 sliced cooked ham packaged in modified atmosphere throughout the shelf life Microbiota of sliced
 347 cooked ham in MAP. *Int J Food Microbiol.* 289:200-208.
- 348 37. Ray, B., and A. Bhunia. 2013. Fundamental Food Microbiology, Fifth Edition. Taylor &
 349 Francis.
- 350 38. Remenant, B., E. Jaffres, X. Dousset, M. F. Pilet, and M. Zagorec. 2015. Bacterial spoilers
 351 of food: behavior, fitness and functional properties. *Food Microbiol.* 45:45-53.
- 352 39. Rossitto, P. V., J. S. Cullor, J. Crook, J. Parko, P. Sechi, and B. T. Cenci-Goga. 2012.
 353 Effects of uv irradiation in a continuous turbulent flow uv reactor on microbiological and sensory
 354 characteristics of cow's milk. *J Food Protect.* 75:2197-2207.
- 355 40. Saccaro, D., C. Hirota, A. Tamime, and N. d. O. M. 2012. Evaluation of different selective
 356 media for enumeration of probiotic micro-organisms in combination with yogurt starter cultures in
 357 fermented milk. *Afr J Microbiol Res.* 6:2239-2245.
- 358 41. Samelis, J., A. Kakouri, and J. Rementzis. 2000. Selective effect of the product type and the
 359 packaging conditions on the species of lactic acid bacteria dominating the spoilage microbial
 360 association of cooked meats at 4°C. *Food Microbiol.* 17:329-340.
- 361 42. SAS. 2016. The SAS System for Windows, Release 9.4. In SAS Institute Inc, Cary, NC,
 362 USA. .
- 363 43. Sechi, P., M. F. Iulietto, S. Mattei, G. Traina, M. Codini, and B. Cenci Goga. 2014. Effect of
 364 a formulation of selected dairy starter cultures and probiotics on microbiological, chemical and
 365 sensory characteristics of swine dry-cured sausages. *J Biotechnol.* 185.
- 366 44. Silva, C. C. G., S. P. M. Silva, and S. C. Ribeiro. 2018. Application of Bacteriocins and
 367 Protective Cultures in Dairy Food Preservation. *Front Microbiol.* 9:594-594.
- 368 45. Ullrich, M. 2009. Bacterial Polysaccharides: Current Innovations and Future Trends. Caister
 369 Academic Press.
- 370 46. Varsha, K. K., and K. M. Nampoothiri. 2016. Appraisal of lactic acid bacteria as protective
 371 cultures. *Food Control.* 69:61-64.
- 372 47. Yost, C. K., and F. M. Nattress. 2000. The use of multiplex PCR reactions to characterize
 373 populations of lactic acid bacteria associated with meat spoilage. *Lett Appl Microbiol.* 31(2):129-33.

374 48. Zhao, F., G. Zhou, K. Ye, S. Wang, X. Xu, and C. Li. 2014. Microbial changes in vacuum-
375 packed chilled pork during storage. *Meat Sci.* 100:145-149.
376

377

FIGURE LEGEND

378 FIGURE 1. Growth curves of strains 649 (____) and 650 (---) at 4-12-20-30-37-44°C (log cfu ml⁻¹,
379 average of three replications).

380 FIGURE 2. Axis 1-2 by canonical discriminant analysis of the effect of temperature on the growth
381 of *L. mesenteroides* strains 649 and 650.

382

TABLE 1. Primer sets used in this study.

DNA target	Primer	Sequence (5'-3')	Size (bp)	PCR conditions	References
Bacterial 16S rRNA	P27f P1495r	GAG AGT TTG ATC CTG GCT CAG CTA CGG CTA CCT TGT TAC GA	1100	1 cycle at 95 °C for 5', 35 cycles at 94 °C for 30", 50 °C for 45" and 72 °C for 2' and final extension at 72 °C for 10'	(9, 33)
Bacterial 16S rRNA	Y1 Y2	TGGCTCAGAACGAACGCTGGCCCG CCCACTGCTGCCTCCCGTAGGAGT	350	1 cycle at 94 °C for 3' followed by 30 cycles at 94 °C for 45", 55°C for 45" and 72°C for 1 min and final extension at 72 °C for 10'	(47)
<i>Leuconostoc</i> spp. 16S rRNA	Lu1r Lu2	CCACAGCGAAAGGTGCTTGAC GATCCATCTCTAGGTGACGCCG	175	1 cycle at 94 °C for 3' followed by 30 cycles at 94 °C for 45", 55°C for 45" and 72°C for 1 min and final extension at 72 °C for 10'	(47)
<i>L. mesenteroides</i> 16S rRNA	L.mes-f L.mes-r	AACTTAGTGTCGCATGAC AGTCGAGTTACAGACTACAA	110	1 cycle at 94 °C for 5', followed by 30 cycles at 94°C for 1', 60°C for 1', 72°C for 2" and final extension of 72 °C for 10'	(3, 26)

TABLE 2. Kinetic parameters.

Strain	Temperature °C	R ²	Kinetic parameters			
			D _{max} (days)	Y _{max} (log)	b (Δlog/day)	Dtot (days)
L. m. 649	4	0.9392	42.8	8.901	0.123	179
	12	0.8752	15.4	8.099	0.177	117
	20	0.7712	8.8	8.533	0.260	80
	30	0.9308	2.2	9.220	2.157	14
	37	0.9779	2.6	8.039	1.804	11
	44	1.0000	0.2	3.587	1.855	2
L. m. 650	4	0.9504	44.7	8.966	0.117	179
	12	0.8977	13.7	8.160	0.122	117
	20	0.8094	7.7	8.563	0.310	80
	30	0.9602	1.9	9.401	2.662	14
	37	0.9950	2.5	8.112	1.886	11
	44	1.0000	0.2	3.939	3.534	1

D_{max}: days; Y_{max}: log cfu ml⁻¹; b: growth rate; Dtot: length of the experiment in days

TABLE 3. Polynomial coefficients.

Strain	Temperature °C	R ²	a0	a1	a2	a3	a4
L. m. 649	4	0.9392	3.6375	0.2966	-0.0054	0.00003	-75 E-9
	12	0.8752	5.3856	0.4009	-0.0179	0.00023	-95 E-8
	20	0.7712	6.2350	0.5802	-0.0435	0.00087	-539 E-8
	30	0.9308	4.4547	5.0347	-1.6578	0.17387	-0.0059
	37	0.9779	3.3841	4.3766	-1.3325	0.14459	-0.0057
	44	1.0000	3.1761	3.9834	-10.8990	6.02439	-0.9839
L. m. 650	4	0.9504	3.7246	0.2924	-0.0054	0.00004	-88 E-9
	12	0.8977	6.4905	0.2724	-0.0132	0.00017	-697 E-9
	20	0.8094	6.1725	0.6857	-0.0577	0.00122	-778 E-8
	30	0.9602	4.2432	6.0930	-2.1975	0.23564	-0.0080
	37	0.9950	3.3326	4.6729	-1.5040	0.17864	-0.0075
	44	1.0000	3.1661	7.6398	-21.4817	12.9166	-2.2407

a0, a1, a2, a3, a4: coefficients of polynomial determined by the function «PROC REG» regression procedure in SAS 9.4

Figure 1

Figure 2

[Click here to access/download](#)

Tracked Changes
Blank 1.docx

Click here to access/download
Response to Reviewers
Blank 2.docx

[Click here to access/download](#)

Copyright Release
Blank 3.docx