



AperTO - Archivio Istituzionale Open Access dell'Università di Torino

Anaerobic digestates from sewage sludge used as fertilizer on a poor alkaline sandy soil and on a peat substrate: Effects on tomato plants growth and on soil properties

This is a pre print version of the following article:
Original Citation:

Availability:

This version is available http://hdl.handle.net/2318/1762460

since 2020-11-11T17:18:31Z

Published version:

DOI:10.1016/j.jenvman.2020.110767

Terms of use:

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)

1 Title

2	Anaerobic digestates from sewage sludge used as fertilizer on a poor alkaline sandy
3	soil and on a peat substrate: effects on tomato plants growth and on soil properties
4	Authors
5	Giulio Cristina ^{a†} , Enrico Camelin ^{a†} , Tonia Tommasi ^{a*} , Debora Fino ^a and Massimo Pugliese ^{b,c*} .
6	^a Department of Applied Science and Technology (DISAT), Politecnico di Torino, Corso Duca degli
7	Abruzzi 24, 10129 Torino (TO) – Italy
8	^b Agricultural, Forestry and Food Science Department (DISAFA), University of Torino, Largo Paolo
9	Braccini 2, 10095 Grugliasco (TO) – Italy
10	^c AGROINNOVA – Centre of Competence for the Innovation in the Agro-Environmental Sector,
11	University of Torino, Largo Paolo Braccini 2, 10095 Grugliasco (TO) – Italy
12	
13	[†] These authors contributed equally to the work
14	
15	* Corresponding authors:
16	Tonia Tommasi
17	Phone: +39 011 090 4774; e-mail: tonia.tommasi@polito.it
18	Department of Applied Science and Technology (DISAT), Politecnico di Torino
19	Corso Duca degli Abruzzi 24, 10129 Torino (TO) – Italy
20	
21	Massimo Pugliese
22	Phone: + 39 011 670 8545; email: massimo.pugliese@unito.it
23	Agricultural, Forestry and Food Science Department (DISAFA), University of Torino
24	Largo Paolo Braccini 2, 10095 Grugliasco (TO) – Italy
25	

26 Abstract

Anaerobic digestates from sewage sludge (SSADs) are a by-product of the wastewater treatment 27 28 process that still preserves a certain agronomic interest for its richness in plant nutrients and organic 29 matter. Fertilizing properties of two liquid and two dewatered SSADs were tested on tomato plants (Solanum lycopersicum L.). Pot experiments were performed on sandy soil and peat substrate under 30 greenhouse conditions with a SSADs application rate of 170 kg N/ha over a period of three months. 31 Beneficial effects of SSADs were reported on different growth parameters, revealing an increase in 32 33 biomass and height up to 37.5 and 6-folds over untreated control. No phytotoxic effect occurred on SSAD-exposed plants. Chemical analysis of soils treated with SSADs showed enrichment of macro-34 and micro-nutrients as well as organic matter. In some cases, the chemical characterization of leaves 35 revealed an enhancement of uptaken macronutrients. This study contributed in general to deepen the 36 37 knowledge on the short-term growing season fertilizing effects of SSAD. Despite the treatment dosage was calculated only on nitrogen requirements, the study highlighted the importance of the 38 other nutrients and organic matter on plant growth. 39

40

Keywords: waste management; nitrogen; soil organic matter; nutrient recycling; sewage sludge;
tomato plants.

44 1. Introduction

Globally, the demand of the three primary plant nutrients used for soil fertilization (N, P_2O_5 and K_2O) is 45 increasing (Vanotti et al., 2019). In 2015, the total fertilizer nutrient demand was around 184 Mt and, 46 by the end of 2020, it is expected to overcome 200 Mt (FAO, 2017). The production processes of 47 48 these fertilizers are very expensive in terms of energy (ammonia) and non-renewable resources (phosphorus and potassium), with heavy environmental costs (Li et al., 2009). Ammonia production is 49 mainly performed via the Haber-Bosch process which requires a large amount of fossil fuel (Basosi et 50 al., 2014). Phosphate rock is the principal raw material exploited in the production of nearly all 51 phosphate fertilizers (Fixen and Johnston, 2012; Reijnders, 2014). This non-renewable resource may 52 53 contain many toxic heavy metals such us As, Hg, Ni, V (Mortvedt, 1995), Cd, Cr, Cu, Pb, Zn (Sabiha-Javied et al., 2009), fluorine (Mirlean and Roisenberg, 2007) and uranium (Schnug and Lottermoser, 54 2013). The P_2O_5 extraction can cause environmental pollution by contaminants accumulating in air, 55 soil, and water bodies around the manufacturing place (Mirlean et al., 2008; Sabiha-Javied et al., 56 2009). It has been observed that these impurities can persist into phosphate fertilizers, provoking a 57 subsequent accumulation in agricultural soils (De López Camelo et al., 1997). Potassium derives from 58 non-renewable resources like minerals such as sylvite, sylvinite, hartsalz and langbeinite (Fixen and 59 60 Johnston, 2012). Furthermore, world distribution of phosphorous and potassium mines is not uniform: 45% of global phosphate rock is concentrated in Morocco and the Western Sahara (Fixen and 61 Johnston, 2012). 62

Within a circular economy perspective, the reuse of sewage sludge (SS) as fertilizer is an interesting scenario. SS can be defined as "the residue generated from the treatment of wastewater" (Smith et al., 2009). This matrix is a valuable source in terms of plant nutrients: a study conducted on 240 dried samples from Pennsylvania revealed an average N, P and K content of 4.74%, 2.27%, and 0.31%, respectively (Stehouwer et al., 2000). Furthermore, SS can contain many micronutrients (e.g. Ca, Mg,

Abbreviations: A_N: assimilation; ANRE: Apparent Nitrogen Recovery Efficiency; ANUE: Agronomic Nitrogen Use Efficiency; C: centrifuged SSAD; CEC: cation exchange capacity; CCI: Chlorophyll Content Index; C_i: CO₂ concentration in substomatal cavity; CRF: controlled release fertilizer; D: dried SSAD; D.M.: dry matter; EC: electrical conductivity; EmC: Emerging Contaminants; EU: European Union; g_s: stomatal conductance; IRGA: infra-red gas analyzer; M: mineral fertilizer; OM: organic matter; P: primary SSAD; QL: quantification limits; S: secondary SSAD; SS: sewage sludge; SSAD: anaerobic digestate from sewage sludge; T: non-treated, control thesis; WWTP: wastewater treatment plant.

S, Fe, Mn, Cu, Zn and B) which are important for plant growth, but usually not included in most commercial fertilizers (Warman and Termeer, 2005). The percentage of the nutrients appears low, but it is important to underline that every year a huge amount of wastewater is produced. An empirical study revealed that approximately 330 km³ of municipal wastewater are produced worldwide yearly (Mateo-Sagasta et al., 2015). Therefore, also the SS production has dramatically risen thanks to policies dealing with the improvement of wastewater treatment and of standard quality of effluents, such as the E.U. directive 91/271/EEC (Council of the European Communities, 1991a).

The considerable presence of organic carbon and organic matter in SS is another strength of its reuse (Alvarenga et al., 2015; Mateo-Sagasta et al., 2015). In fact, land application of organic matter (OM) improves soil physical properties such as cation exchange capacity (CEC), soil structure, soil moisture content and retention (Epstein, 2002). Furthermore, the addition of SS can enhance the amount of organic carbon in soils (Kladivko and Nelson, 1979; Perez-Espinosa et al., 1999) and thus reverse the current reduction of organic matter in soils (known as *SOM decline*) (Schulze and Freibauer, 2005).

Today, SS is classified as waste and its safe disposal represents a very important issue in waste 82 83 management (Epstein, 2002; Singh and Agrawal, 2008). The four main destinations of SS are incineration, landfilling, composting and agricultural use. In Italy, according to data of 2010 (Eurostat, 84 2019), the majority of SS is sent to landfill (50.8 %), while 34.7% is reused in agriculture, 4% is 85 86 incinerated and 10.4% is sent to other destinations. The Council Directive 86/278 (Council of the 87 European Communities, 1986) regulates the agricultural SS reuse in Europe to prevent soil 88 contamination. In fact, this practice has three principal problems that limit its unconditioned use: biological risk, heavy metal contamination and contamination by organic pollutants. The biological risk 89 is principally represented by pathogens such as Salmonella spp., Escherichia coli (enterotoxigenic 90 and enteropathogenic variants), Campylobacter spp., Clostridium spp., and Yersinia spp. (Arthurson, 91 2008); stabilization treatments can reduce significantly their presence in SS and are mandatory before 92 subsequent SS applications (Dumontet et al., 1999). For instance, one of the most diffused 93 stabilization techniques is anaerobic digestion (Liu et al., 2012), in which the reduction of pathogens, 94 putrescence and odor is coupled with biogas production, allowing energy recovery (Epstein, 2002). 95

Heavy metal content (normally represented by Cd, Cr, Cu, Hg, Ni, Pb, Zn) can be abated by means of 96 chemical (e.g. chelating addition), physical (e.g. electroremediation) or biological 97 (e.g. 98 vermicomposting) treatments (Camargo et al., 2016). Finally, some organic pollutants (e.g. pesticides, 99 antibiotics and hormones) can be volatilized or degraded through biotic or abiotic processes (Harrison et al., 2006). "Concerning organic pollutants, their abatement is trickier. Indeed, class of emerging 100 101 contaminants (EmC) in wastewater is increasingly gaining more interest within the organic 102 compounds. EmC include molecules such as endocrine disrupting compounds (e.g. hormones), 103 pharmaceutically active compounds (e.g. antibiotics), illicit drugs and pesticides (Fijalkowski, 2019). 104 EmC abatement is becoming even more required both on the effluent of WWTPs with advanced treatments (e.g. activated carbon absorption, advanced oxidation processes, reverse osmosis) and on 105 sewage sludge (Gadupudi et al., 2019). Some studies affirmed that anaerobic digestion is the 106 107 stabilization strategy ensuring the best EmC removal, especially when the sludge is pretreated (e.g. 108 via ozonation) (Neumann et al., 2016). However, further studies are still required to improve the 109 performances and reduce the costs of these techniques, which nowadays are rarely applied at WWTP 110 level since they are money and/or time consuming (Camargo et al., 2016). The abovementioned EU 111 directive regulates the SS soil application in the EU and establishes threshold values of some of these 112 pathogens and pollutants in SS.

113 On the basis of these opportunities and threats related to SS, this work aims to deepen the 114 knowledge about SS fertilizing effects over time in terms of nutrients and OM on a poor alkaline sandy 115 soil. This kind of soil was selected because: i) nutrient depletion constrains plant growth to depend on 116 treatment application; ii) a high pH both hinders the nutrient adsorption and reduce the metal 117 bioavailability (Alvarenga et al., 2016); iii) sandy-textured soil lacks nutrients and has low water-118 holding capacity. These results were compared to the one obtained with a richer peat substrate. Pot 119 experiments were performed on tomato plants (Solanum lycopersicum L.) in a greenhouse to 120 evaluate nutrient provision of anaerobic digestates from sewage sludge (SSADs). Tomato plant was 121 chosen because: i) it is one of the most exploited vegetables crop (Jones Jr, 2008); ii) there is an increasing interest on alternative nutrient sources for this crop (Zucco et al., 2015); iii) it has a high 122 fertilizer requirements (Zucco et al., 2015); iv) plenty of scientific literature is available for this crop 123

124 (Jones Jr, 2008). In this work, no analysis on pathogens was carried out since anaerobic digestion is considered one of the safest technologies for pathogen reduction in SS (Epstein, 2002). 125 126 Nevertheless, this aspect may be taken into consideration in future researches. Concerning the use of 127 SS in agronomic experiments, a lack in details about SS typology is provided. Indeed, in many work no detail on stabilization strategy is provided (Bakshi et al., 2019), or the kind of SS digestion is not 128 specified (Hossain et al., 2015). In the present work, the digestates used derived from the same 129 WWTP and were obtained with consequent treatments (Cristina et al., 2019). As far as we know, this 130 131 is the first example of use of four different and consequent SSADs to fertilize tomato plants. The paper examines agronomic parameters of tomato plants, the nutrient distribution in soil and nutrient 132 absorption by plants after the application of SSADs. Furthermore, numerous plants physiological 133 parameters were evaluated over a span of three months in order to better understand the effects in a 134 135 time course approach.

- 136 2. Materials and methods
- 137 2.1. Characterizations

138 2.1.1.Anaerobic digestates from sewage sludge

Four SSADs were used in the experiment: two liquid (primary (P) and secondary (S)) and two solid (centrifuged (C) and dried (D)). Physical and chemical characterization of the SSADs is described in a previous work (Cristina et al., 2019); characterization of the four SSADs is reported in Supplementary Material (Section I - Table S1).

143 2.1.2. Cultivation substrates

Two types of substrates were used: a sandy soil and a commercial peat substrate (**Table 1**). The sandy soil was sampled within 20 and 100 cm depth in Grugliasco (TO), Italy ($45^{\circ}03'58.4"N$, 7°35'32.9"E). Analytical methods used for characterization of the sandy soil and the peat substrate are specified in Supplementary materials (Section II). Based on the distribution of the particle size (sand: 94% ± 2; silt: 3% ± 1; clay: 3% ± 1), the selected soil was classified as sandy (Buol et al., 2011). Based on ARPAV soil analysis (Arpa Veneto, 2007), the soil was considered alkaline (8.2 ± 150 0.16), very poor in OM (0.38 \pm 0.12% < 0.8%) and very poor in macronutrients such as nitrogen (0.29 151 \pm 0.09 g/kg < 0.5 g/kg), phosphorous (1.8 \pm 1.3 mg/kg < 7 mg/kg), potassium (18 \pm 1 mg/kg < 40 152 mg/kg) and magnesium (15 \pm 5 mg/kg < 50 mg/kg). On the other hand, content of calcium (675 \pm 27 153 mg/kg < 1000 mg/kg) and some microelements such as iron (2.5 mg/kg < 6.7 \pm 1.1mg/kg < mg/kg 20) 154 and manganese (2 mg/kg < 6.5 \pm 3.0 mg/kg < mg/kg 10) resulted normal.

155 2.2. Experimental set-up

156 A greenhouse experiment was performed over three months during the summer season in a greenhouse of the Centre of Competence AGROINNOVA – University of Torino, located in Grugliasco 157 158 (TO), Italy. The experimental campaign was carried out with commercial plastic pots of 2.5 L (Ø 17 cm, height 20 cm, surface area 0.227 m²). Four types of SSADs (P, S, C, D) were applied as 159 treatments, and compared to a commercial fertilizer (M) (NPK 22-5-6 + 2MgO, "Osmocote Topdress", 160 ICL, Israel) and an untreated control (T). The experiment was designed in a completely randomized 161 162 block, with 15 replications per each thesis. The same experimental set-up was adopted on the two 163 cultivation substrates (sandy soil and peat substrate). Each treatment was applied at the dosage of 164 170 kg N/ha, in line with the European Nitrates Directive (Council of the European Communities, 165 1991b). Moreover, this application rate was chosen as it showed the best results in a preliminary 166 study (Cristina et al., 2019). Three untreated seeds of tomato (Solanum lycopersicum L. cv. Beefsteak, "Furia sementi", Parma, Italy) were sown in each pot. Automatic sprinkler irrigation was set three 167 times a day. Ten days after sowing a thinning was conducted and the best plant from each pot was 168 169 kept. At the end of each month, five pre-selected replicates of each treatment were removed to carry 170 out all the measurements.

171 2.3. Measurement of plant parameters

At the end of every month, the five removed replicates were examined. Firstly, height was measured, then, leaves, inflorescences and fruits were counted, if present. After that, the Chlorophyll Content Index (CCI) was evaluated with a SPAD 502 chlorophyll meter (CCM-200, Opti Sciences, Inc., Hudson, NH, USA) using the method described in the previous work (Cristina et al., 2019). One

176 month after sowing, it was not possible to measure CCI on the sand specimen because the minimum leaves size was not satisfied. At the end of the second month, assimilation (A_N), stomatal 177 conductance (g_s) and CO₂ concentration in substomatal cavity (C_i) were measured by the means of 178 179 an Infrared Gas Analyzer (IRGA, ADC, Hoddesdon, UK). These measurements were performed on 180 three fully formed leaves in each replicate. The selected leaves had to be non-senescing, at the same 181 physiological age (in the middle part of the plant, considering the third to fourth leaf from the shoot 182 apex) and directly exposed to sunlight. After all the measurements were taken, each plant was 183 subsequently cut and immediately weighed to record the fresh biomass value. In order to evaluate the mean dry biomass, each plant was dried at 105°C for at least 72 hours. Subsequently, agronomic 184 nitrogen use efficiency (ANUE) was calculated as: 185

$ANUE = \frac{(Dry \ biomass \ treated \ samples - Dry \ biomass \ control \ samples)}{Amount \ of \ nitrogen \ applied \ in \ treated \ samples}$

186 2.4. Chemical analysis

Substrates were chemically characterized at the end of the second month, once the aerial plant part had been cut. Chemical analyses were performed on samples from the treatment with SSADs (P, S, C, D) as well as on minerally fertilized ones (M) and untreated control (T). The samples were collected excluding the upper 3 cm of topsoil and the rhizosphere area. The analyses were performed with the same methods exploited for the chemical characterization of substrates prior to the experiment (see Supplementary material - Section II).

193 Chemical analyses of the leaves were conducted at the end of the second month, after the biomass 194 measurement, in order to assess the content of nitrogen, phosphorus and potassium in the leaves. In the case of the samples from the sandy soil, the measurements were performed on samples treated 195 196 with one liquid digestate (P), one solid digestate (D) and the mineral fertilizer (M). It was not possible 197 to analyze samples from the negative control (T) due to the low biomass production. On the peat 198 substrate, it was possible to evaluate N-P-K content not only in the P, D, and M samples, but also in the negative control ones (T). The plant samples were firstly processed with a humid digestion 199 200 protocol (Mills and Jones Jr, 1996). Then, nitrogen was measured through the Kjeldahl method,

201 phosphorus was evaluated through colorimetry (molybdovanadate method) and potassium was 202 quantified by Atomic Absorption Spectroscopy (AAS). Finally, the N, P and K percentages were used 203 to calculate the mean total element present in the epigean part of the plant using the following 204 formula:

$$\frac{dry \ sample \ biomass \ (g)}{N, \ P, \ K \ in \ sample \ (\%)} * \frac{100}{1000} = total \ N, P, K \ presence \ in \ single \ sample \ (mg)$$

205 2.5. Statistical analysis

The experimental data were subjected to statistical analyses. Two-way ANOVA was used to compare the average results of different treatments on plant measurements. Differently, one-way ANOVA was used to compare the mean results of different treatments on the chemical analyses of soils and leaf nutrient content. After the ANOVA, Tukey's post-hoc test (P < 0.05) was performed. The statistical software R (version 3.5.1 - Feather Spray - 2018) was used for all statistical analysis.

211 **3. Results**

3.1. Plant measurements

3.1.1.Dry biomass and ANUE

On the sandy soil at the end of the first month, the dry biomass of the tomato plants grown with 214 digestates did not show any difference between each other. Despite the absence of significant 215 216 differences, it must be pointed out that biomass of S was 28.7 and 12.7-folds higher than control (T) 217 and mineral fertilizer (M), respectively. At the end of second month, all digestates (P, S, C, D) showed a dry biomass production significantly higher (26.7, 33, 35.3 and 37.5-folds, respectively) than control. 218 219 At the same time, S, C and D showed a higher biomass than mineral fertilizer (2.9, 3.1 and 3.3-folds, 220 respectively).. At the end of third month, dewatered SSADs proved to be the most productive 221 treatments, with C and D displaying the highest yields (10.23 g and 10.97 g). Their biomasses 222 doubled mineral fertilizer one (5.13 g), which was only comparable to the biomass produced by plants treated with SSADs after two months. Furthermore, C and D yields were 16 and 17-folds higher than 223 T (0.64 g), respectively (**Figure 1.A**). 224

225 On the peat substrate, no significant differences between treatments were appreciable within the 226 same month. The only significant differences emerged between biomass values between three 227 different months (**Figure 1.B**).

Results of ANUE (**Table S2 and S3**) did not satisfy the requirements of Levene test during ANOVA, hence no significant differences could be reported reasonably. However, it must be pointed out that ANUE values of SSADs in sandy soil were up to 23, 3.5 and 2.3-folds higher than mineral ones after one, two and three months after sowing, respectively. In the case of substrate, the value did not follow any trend, also considering that in some cases control samples resulted more massive than treated ones.

234 3.1.2.Height

On the sandy soil, no differences in plant height were present at the end of the first month. 235 236 Nevertheless, S treatment revealed the tallest tomato plants, up to 2.6 and 2.5-folds higher than T 237 and M. In the second month, all SSADs-treated plants were significantly taller than control and 238 mineral fertilizer, with D treatment displaying a height 6 and 2.1-folds higher than T and M, 239 respectively. After three months, the mean height of T was still the lowest. The mean height of the 240 plants grown on P and D was comparable to plants grown on mineral fertilizer. Plants grown with S 241 and C treatments had a statistically higher height than mineral fertilizer (M). It is worth highlighting that 242 the mean heights of the plants grown on all digestates was at least 3.5-folds higher than the control 243 ones (Figure 2.A).

On the peat substrate, no significant differences were observed between the different treatments within the same month. The only significant differences emerged between the height of the samples between three different months (data not shown).

247 3.1.3.Leav

3.1.3. Leaves and inflorescences

After the first month, the plants grown on sandy soil in presence of D and C treatments showed a number of leaves comparable to control and minerally fertilized plants. On the other hand, samples from liquid SSADs (P and S) revealed a higher mean leaf number than control. After two months, the leaves number on plants grown with digestates was significantly higher only than negative control plants. At the end of the experiment, samples from S and D treatments showed the highest number of leaves, which were not statistically different from samples from C treatment. Plants grown with P had similar number of leaves than C and mineral fertilizer, while leaves number in negative control was still the lowest one (**Figure 2.B**).

With regards to the number of inflorescences, no plant on sandy soil showed flowers one month after sowing. At the end of the second month, plants in T and M were still not revealing any flower. Differently, P, S, C and D had some inflorescences, but no significant difference between treatments was present. At the end of the experiment, negative control plants still did not show any flower. Plants treated with P and S had a number of inflorescences statistically similar to mineral fertilizer. The highest number of inflorescences was found on C and D treatments (**Figure 2.C**).

As regards the number of leaves and inflorescences of plants grown on peat substrate, no differences between treatments at the same month were highlighted by statistical analysis (data not shown).

3.1.4. Chlorophyll Content Index (CCI)

On sandy soil, leaves dimension after one month was too small to measure CCI. At the end of second month leaves of plants treated with P, S and C showed a CCI higher than control and comparable to mineral fertilizer. The mean CCI value of plants grown with D digestate was statistically higher than mineral fertilizer (M) but comparable to the others SSADs. CCI measures performed at the end of third month revealed a substantial decrease in CCI values registered in all SSADs and in mineral fertilizer, whose values were not significantly different from the control. The only significant difference emerging at the endpoint was between P and mineral fertilizer measure. (**Figure 2.D**).

272 On peat substrate, the only differences were recorded between the CCI measure of D and P at the 273 end of second month, and S and control at the end of the third month (data not shown).

3.1.5. Infra-red gas analyzer (IRGA)

As regards IRGA measurements, on sandy soil the lowest A_N value was found in control, where significantly higher values were recorded on C and S. Detailed results are reported in **Table S4**.

277 3.2. Chemical analysis

3.2.1. Substrates analyses

Results of chemical analyses performed on the sandy soil after two months from treatments 279 application are summarized in Table 2A. SSADs showed all an intermediate mean pH included 280 281 between control (8.3) and mineral fertilizer samples (8.0). OM was significantly higher in P, C and D treatments than in S, mineral fertilizer and control. As expected, values of organic carbon showed a 282 283 trend similar to OM. Total nitrogen (Kieldahl) was lower in control, mineral fertilizer and S than P. C and D treatments. All results of nitrite analysis were below quantification limits (QL). Nitrates were 284 detectable only in S, C and D treatments, showing very low concentrations (between 1 and 4 mg/kg) 285 with respect to M sample (60 mg/kg). Organic nitrogen values were roughly similar to total Kjeldahl 286 287 nitrogen ones. Regarding C/N ratio, the lowest value was calculated in control and mineral fertilizer, 288 while all SSADs revealed higher values. Olsen phosphorus was below QL in T and M samples; differently, phosphorous content in samples treated with SSADs was higher. The lowest value of 289 290 exchangeable calcium was observed in S samples followed by negative control, D, P, C and mineral 291 fertilizer. A great difference in exchangeable sodium content was found between negative control 292 samples and all the treatments. Available zinc ranged between 0.21 mg/kg in control samples, and 293 1.00 mg/kg in D ones, with samples treated with liquid SSADs and mineral fertilizer showing an 294 intermediate behavior. Digestates showed intermediate values of CEC, included between control 295 (2.81 cmol/kg) and mineral fertilizer samples (3.58 cmol/kg). Values of electrical conductivity, ammonia nitrogen (NH⁴⁺), exchangeable K, exchangeable Mg, available Mn and available Cu did not 296 297 show any significant difference between treatments on sandy soil.

Results of chemical analyses performed on peat substrate two months after treatments application are summarized in **Table 2B**. pH values ranged from a minimum of 6.6 (M) to a maximum of 7.4 (D). Total Kjeldahl nitrogen was lower in control samples and gradually increased along with the dry matter of SSADs; the highest value was displayed by mineral fertilizer samples. Organic nitrogen values were analogous to total Kjeldahl nitrogen in terms of values, trend and differences between

treatments. Nitrites, nitrates, extractable Mn, Cu and Zn were below detection limits. All other
 parameters did not show any significant difference.

305 3.2.2.Leaf analysis

On sandy soil, chemical characterization of leaves showed a concentration of nitrogen and potassium in P and D significantly lower than mineral fertilizer samples. As regards phosphorous, no significant difference emerged. The total nitrogen accumulated in leaves in D plants was significantly higher than in P ones. The mean phosphorous uptake by plants was significantly different across D, P and M samples. Finally, the potassium uptaken in leaves did not show significant differences between thesis (**Table 3A**).

On plants grown on peat substrate, concentrations and total uptake of both nitrogen and potassium on control, P and D were statistically similar to each other, but they resulted lower in comparison with mineral fertilizer ones. Concentration and total uptake of phosphorous in leaves, control showed the lowest values while D samples the highest ones (**Table 3B**).

316 4. Discussion

4.1. Agronomic and physiological evaluations

318 For many years extensive studies and reviews have shown that soil and plant benefit from SS. 319 Indeed, SS is a good source of macro and micro nutrients as well as of OM; this enhances soil fertility 320 and, as a consequence, crop production even in a more effective way than commercial fertilizers 321 (Singh and Agrawal, 2008). The results of the present work were in agreement with literature and the 322 better performances of SS compared to inorganic fertilizers have been confirmed. Table 4 shows 323 technical details and results of other works dealing with SS treatment of tomato plant with pot 324 experiments. It is important to notice that not only SSAD application rate was considerably lower in 325 the present work, but also that the results obtained were remarkably higher. For instance, biomass 326 and height of treated tomato plants at two months after sowing was up to 37.5 and 6-folds higher, 327 respectively, than control plants (corresponding to an increase of 3652% and 500%), results never reached before in other works on tomato plants. Interestingly, fertilizing performances of SSAD also 328

329 overcame the ones of mineral fertilizer, especially one month after sowing, when S treatment revealed 330 biomass and height of tomato plants up to 12.7 and 2.5-folds higher than M. From here on out, 331 differences between SSAD treatments and M samples were less accentuated, probably because 332 nutrients release of the mineral fertilizer was faster after an initial "lag" phase. As a corollary, biomass values were reflected by ANUE ones, which were higher than the ones reported in literature for 333 334 tomato plants grown in pot under greenhouse conditions treated with a 10-folds higher nitrogen 335 application (Wang et al., 2013). Improvement in terms of leaves number and chlorophyll content were 336 less intense, but still higher than the examples reported in literature (Bakshi et al., 2019; Elloumi et al., 337 2016; He et al., 2016; Hossain et al., 2015).

338 To a broader extent, results of the present study in terms of biomass and plant height can be 339 compared to other works conducted with a similar experimental setup but exploiting different model 340 species. In order to biomass, the general trend was an increase in dry matter ranging usually between 341 4 (Capsicum annuum L.; Pascual et al., 2008) and 16-folds (Triticum aestivum L.; Eid et al., 2019) 342 more than untreated control. The findings of the present work confirmed and went beyond these 343 results, considering also that the most used SS application rates ranged between the dosage used in 344 this work and a 35-folds higher one (Eid et al., 2019). On the other hand, the improvements in plant 345 height were in line with the results obtained by Eid and colleagues on cucumber (Cucumis sativus L.) 346 (2017) and wheat (Triticum aestivum L.) (2019), reporting a stem length improvement up to 3 and 6-347 folds, respectively, over untreated control. The only case with a striking higher biomass production 348 was described for the sunflower (Heliantus annuus L), whose production increased up to 125-folds 349 more than the untreated control. However, the SS dosage was up to 35-folds higher than the present 350 study. Moreover improvement in terms of height was comparable to the present work (Bourioug et al., 351 2018). Taking into account the works using SS dosages comparable to 170 kg N/ha, the majority 352 were open field experiments. For instance, triticale (X Triticosecale Wittmack) (Kchaou et al., 2018) 353 revealed a biomass increase of 2-folds. Furthermore, results of the present work corroborate positive 354 effects on biomass of SS application on soils poor in nutrients (Walter et al., 2000) and strongly alkaline (Zuo et al., 2019). 355

356 SSAD application on tomato crops resulted also in an augmented number of leaves and 357 inflorescences with respect to control and mineral fertilizer. Moreover, inflorescences number of 358 SSAD-treated plants increased from 2 to 3-folds over the last month. These findings were in general 359 agreement with other results reported on tomato grown in presence of SS (Bakshi et al., 2019), 360 despite the higher treatment dosages.

361 Number of leaves and inflorescences are developmental parameters considered also with other plant 362 species when testing the fertilizing effects of SSAD. For instance, Eid and colleagues (2017) 363 registered on cucumber a boost in the number of leaves of more than 2-folds, which is in line with the 364 results of the present work. Similar outcomes have been reported in terms of number of flowers in common bean (Phaseouls vulgaris L.) (Fernández-Lugueño et al., 2010) and marigold (Tagetes erecta 365 366 L.)(Solanki et al., 2017) grown in SS dosages lower and higher, respectively, than the present work. 367 In contrast with these results, Tarig and co-workers (2012) described a decrease up to 60% in flowers 368 number in Dahlia x hortensis, whose growth had probably been compromised by an excessive SS dosage. 369

370 Results of the present work confirmed the positive effects of SS application on net photosynthesis 371 (Bourioug et al., 2018; Pascual et al., 2008) and chlorophyll content. Leaf chlorophyll content was 372 directly correlated with indirect chlorophyll measurements such as readings through SPAD and CCI-373 meters (Xiong et al., 2015), whose value can be compared to each other with the equations proposed 374 by Parry and colleagues (2014). Application of SSAD improved chlorophyll content values of tomato 375 plants grown on sandy soil at the end of second month, as well as dry biomass and net 376 photosynthesis (A_N) . This beneficial effect has been already observed also in sunflower (Bourioug et al., 2018), sorghum (Alvarenga et al., 2016) and triticale (Kchaou et al., 2018). On the other hand, 377 378 literature provides examples of reduction of leaf chlorophyll content in tomato after treatment with SS 379 (Elloumi et al., 2016), which is probably due to the excessive heavy metals presence in the used SS 380 (Singh and Agrawal, 2007). However, this aspect was likely not linked with the reduction of chlorophyll content over time observed in the present study. Indeed, this phenomenon has been already 381 382 observed in other SS-treated plant species, such as common bean (Fernández-Luqueño et al., 2010). 383 A possible explanation of this reduction in CCI at the end of the third month could be the deficiency of

nutrients in soil. A second hypothesis for CCI decrease has been proposed by de Oliveira and co-384 workers (2017): after the initial blooming of the plant, gradual degradation of chlorophyll occurs due to 385 386 the beginning of the fruit development phase, which induces a metabolic change in the plant, with a 387 more sustained nutrients accumulation in the fruit. Taking into account the relationship between leaf nitrogen and chlorophyll content (Xiong et al., 2015), a third justification for CCI decrease can be 388 provided by the so-called nitrogen dilution curve. In fact, biomass increase in tomato plant was 389 390 accompanied by a reduction in nitrogen concentration (and, consequently in chlorophyll content) 391 because the structural compartment (lower in N%) becomes proportionally more massive than 392 metabolic active one (higher in N%) (Tei et al., 2002).

393 4.2. Chemical analysis

394 4.2.1. Substrates analyses

The application of SS on soil can affect different physical and chemical soil characteristics (Epstein, 2002). Likewise, many changes were documented in this experiment (both on sandy soil and on peat substrate) two months after treatments application. Although peat substrate was low in nutrient content, it showed a consistently higher amount of microelement than sandy soil. Moreover, peat substrate has many other advantages such as lightweight, high water holding capacity and high air space (Gruda et al., 2016). All these peculiarities most probably contributed to the minor differences registered on peat substrate.

Soil analysis results revealed a change in soil pH after the treatments application. Many works 402 403 reported an increase (Bayoumi Hamuda et al., 2009; Ferreiro-Domínguez et al., 2011) or a decrease 404 (Mosquera-Losada et al., 2016; Singh and Agrawal, 2007) in soil pH. In the present work, acidification 405 occurred in treated sandy soil samples, probably due to both the lower pH of SSADs and the nitrogen mineralization (Rasouli-Sadaghiani and Moradi, 2014). In particular, the nitrification process (NH₄⁺ \rightarrow 406 NO_3) (Stamatiadis et al., 1999) induces the release of H⁺ in soil solution media and the leaching of 407 NO₃⁻ by water (Whitehead, 1995). Another conceivable theory for soil acidification in SSAD-treated 408 409 samples could be the generation of organic acids during sewage sludge mineralization (Angin et al.,

2012; Bourioug et al., 2018). Additionally, the low buffering capacity might be yet another conceivable
effect occurring in the sandy soil case.

412 Electrical conductivity values (both on sandy soil and on peat substrate) did not statistically change 413 after treatments application unlike many other works (Bourioug et al., 2018; Singh and Agrawal, 414 2007), likely due to the consistently lower SSAD application rates. Nevertheless, it must be pointed 415 out that, concerning sandy soil, EC values in M were approximatively doubled compared to SSAD 416 ones, which in turns were somewhat higher than control. High EC of M might be due to the 417 particularly higher concentration of nitrates, likely released as bioavailable form nitrogen by the 418 commercial fertilizer. However, these relatively elevate nitrate amounts were likely not necessary, as confirmed by the better growth parameters and ANUE values of tomato plants growing on SSAD 419 420 amended soil. On the contrary, excess of nitrates may result in undesired drawbacks such as 421 leaching and hyperaccumulation in plant tissues, feature in agreement with the foliar analyses. EC 422 values in SSAD treatments was probably influenced by sodium presence in the digestates, which 423 however did not affect the physiological parameters of tomato plants as confirmed by IRGA 424 measurements.

The thesis of a possible increasing of soil OM in soils treated with SSADs (Kladivko and Nelson, 1979; Perez-Espinosa et al., 1999) was confirmed by the present work. Despite the OM percentage was very low in all samples, the value in SSADs treated theses was higher than control and mineral fertilizer. This may partially justify the better performances of treated samples in term of biomass and height, according to the well-known soil OM benefits on plants growth (Bot and Benites, 2005).

430 CEC significantly increased in SSADs-treated soil, which was probably caused by the OM increment. 431 This effect is even more pronounced on alkaline soils (Bohn et al., 2001) and similar results were 432 found in other works (Angin et al., 2012; Ferreiro-Domínguez et al., 2011).

Total N, available P, exchangeable Ca and Na and available Fe and Zn concentrations increased in the sandy soil amended with SSADs due to their higher concentration in SS (Singh and Agrawal, 2007).

436 Two months after treatments application, N_{Tot} (Kjeldahl) was higher in C and D than liquid SSADs (P 437 and S), probably due to their solid form that plausibly induced a slower release, both on sandy soil

and peat substrate. Other studies revealed that total soil nitrogen can persist in higher concentrations
also for longer periods after SSAD treatment application (Bourioug et al., 2015). Anyway, all samples
showed a total N content lower than before digestates application. It means that a remarkable part of
nitrogen both already present in sandy soil and added with digestates was absorbed, transformed or
leached after two months.

443 The significant variation in N and OM content in treated sandy soils changed C/N ratio. The results 444 obtained with SSADs were still low (< 9; Arpa Veneto, 2007), but higher than in control and mineral 445 fertilizer. The small changes in C/N and the relatively low values across treatments likely indicated that nitrogen mineralization could have prevailed over microbial immobilization. Therefore, nitrogen in 446 SSAD treated samples was surely bioavailable and used efficiently plants, as also confirmed by 447 ANUE values. However, it should be also noticed that mineralisation was likely a slow nitrogen 448 449 release process, as evidenced by soil nitrate soil and leaf nitrogen analyses. Indeed, these evidenced 450 that nitrogen was much more bioavailable in M treatments, but not efficiently utilizable, according to 451 ANUE values.

452 In all SSADs treated soils, the available P was higher than control and mineral fertilizer. Considering 453 that the different dosages were normalized on N dosage per each thesis, the difference in P content 454 between the samples treated with SSADs can be explained by the different percentages of P in the four SSADs. This diversity could also explain the differences among different treatments on 455 456 physiological parameters of tomato. Moreover, the addition of OM probably enhanced the availability 457 of P in soil treated with SSADs (Fekri et al., 2011). In fact, this can increase the abundance and the 458 activity of microorganisms, favoring P capture (Nobile et al., 2019). Similar results in increase of soil P 459 were obtained by Singh and Agrawal (2007) and Walter and colleagues (2000).

For what it concerns K, no differences were registered in soil after digestates application, probably
due to their low concentration in this macronutrient. These results agree with other works (Bourioug et
al., 2015; Walter et al., 2000).

463 Many SS are rich of Ca due to the stabilization by means of liming (Epstein, 2002). Although the 464 SSADs exploited in this work did not undergo Ca addition at WWTP level, its content was pretty high 465 (> 4.64% D.M.). Considering the medium content in the initial soil, exchangeable Ca increased in

466 some cases in treated soils, confirming the results of Ferreiro Dominguez and Singh (Ferreiro-467 Domínguez et al., 2011; Singh and Agrawal, 2007).

A significant increase of exchangeable Na was measured in all treated soils due to the sodium percentage in SSAD and confirmed by two abovementioned works (Ferreiro-Domínguez et al., 2011; Singh and Agrawal, 2007). The excess of Na is a well-known limiting factor for plants growing (Jones Jr., 2012) but Na has been recently defined as a "new beneficial element" (Morgan, 2000) that, in small quantities, can increase tomato yields (Jones Jr., 2012).

The consistent presence of Fe and Zn in SSADs likely provoked the increase in their concentration in sandy soil, confirming the results of Angin and colleagues (2012).

475 4.2.2.Leaf analysis

In some cases, in literature the use of SS enhanced the percentage of macronutrients in leaves 476 (Angin et al., 2012; Zuo et al., 2019), in other ones no change took place (Kotecki et al., 2014; Pinna 477 478 et al., 2009) and still in other ones concentration increased only for some nutrients (Bakshi et al., 2019; De Andres et al., 2010). This work belongs to the third category, since only foliar P% and total 479 480 uptaken P of control plants grown on peat substrate were significantly lower than SSADs ones. On 481 sandy soil, content of uptaken P was significantly higher in D and P treatments, which was likely 482 influenced by the phosphorous amount in the initial application. Nevertheless, no significant 483 differences emerged in foliar P% despite the difference in uptaken P content between SSADs and mineral fertilizer: probably, the controlled nutrient release of the mineral fertilizer compensated the 484 485 higher quantity of P in the SSADs. Moreover, it could be inferred that differences in foliar 486 macronutrient content could have been appreciated between control and treated samples on sandy soil. However, the too low biomass of untreated samples made impossible this investigation. 487

The total amount of N and K uptaken in leaves had varied results. On sandy soil, D samples revealed a significantly higher N content than P ones due to the different biomass production. Concerning plants grown on peat substrate, P and D showed a nitrogen plant uptake similar to negative control, but lower than mineral fertilizer, likely due to the characteristics of the fertilizer, such as the controlled nutrient release and the presence of readily bioavailable nitrogen forms. As regards K, despite its

higher amount in mineral fertilizer, total K uptaken in leaves did not result significantly different
between the treatments applied on sandy soil, due to the different aboveground biomass production.
On the other hand, on peat substrate, the K_{Extractable} content of plant with mineral fertilizer was the
highest considering the similar biomass production.

497 5. Conclusions

In the present work, pot experiments under greenhouse conditions on two different substrates were 498 performed to evaluate fertilizing effects of four different SSADs over a time span of three months. The 499 500 application of these digestates clearly highlighted beneficial effects on different growth parameters of 501 tomato plants, especially when cultivated on a sandy, alkaline and poor (in nutrient and OM) soil. For 502 instance, it is important to point out that plant biomass and height reached values up to 37.5 and 6-503 folds, respectively, higher than untreated control; additionally, SSAD-treated plants showed values of 504 biomass and height up to 12.7 and 2.5-folds, respectively, higher than mineral treatment, indicating 505 that SSAD could be a valuable alternative to mineral fertilizers to boost fertility in poor and sandy soils 506 . Moreover, the present work confirmed the thesis of the enhancement of soil OM with the use of 507 SSAD. Furthermore, it is important to notice the increments of some macro- (nitrogen, phosphorous 508 and calcium) and micro-nutrients (iron and zinc) in sandy soil, showing significant differences with 509 respect to untreated control. Nevertheless, some of the registered values were low and it can be 510 reasonably assumed that most of nutrients had already been assimilated to let the plant grow. This 511 aspect was confirmed by leaves analysis, which showed a remarkable uptake in N, P and K by 512 tomato plants. With respect to these macronutrients, it is worth emphasizing that the experiment was 513 designed to administer plants, across the different treatments, the same nitrogen dosage as sludge 514 application rate is usually based on plants nitrogen requirements. However, the differences in SSADs 515 composition implied a remarkable imbalance in terms of other nutrients and OM. Hence, we can 516 assume that these differences likely influenced plant growth, providing consistent differences between 517 different theses.

518 Future work should include on one side a deeper analysis of the issues tackled in the present paper, 519 and on the other hand it should consider also related aspects. Concerning the formers, chemical characterization of the treated substrates and plants should be carried out in a time-course fashion, allowing to properly describe the mass balance of the elements and their release dynamics over time, and, consequently, a more detailed evaluation of fertilizing indexes such as ANRE and ANUE. As regards the latter ones, soil application of SSAD should be explored both analyzing the presence of organic pollutants (e.g. antibiotics, EDC) as well as considering microbiological aspects, such as the effects on microbial communities and the study of metagenomics and metatranscriptomics traits (e.g. antibiotics resistance genes).

527 Despite reserves and resources for N, P and K appear adequate for the near future, it is necessary to 528 find less impactful solutions to produce fertilizers in the short term. In this way, the reuse of SS can 529 reduce the negative effects connected by the extraction, manufacturing and the use of mineral 530 fertilizers derived from non-renewable resources. Furthermore, this experiment showed how the 531 positive effects of SSADs are emphasized if applied on a poor alkaline soil.

532 Acknowledgements

Authors wish to thank Mr. Umberto Rossi for supporting this research with the admirable and ethical aim to help people in either desertification areas and developing countries, in order to overcome issues related to poverty and climate change.

536 Credits for icons in Graphical Abstract: Icon made by Freepik from www.flaticon.com.

537 Funding

538 This research was partially supported by the EU Horizon2020 NUTRIMAN project (GA 818470) to 539 University of Torino – Agroinnova.

540 References

Alvarenga, P., Farto, M., Mourinha, C., Palma, P., 2016. Beneficial Use of Dewatered and Composted
Sewage Sludge as Soil Amendments: Behaviour of Metals in Soils and Their Uptake by Plants.
Waste and Biomass Valorization 7, 1189–1201. https://doi.org/10.1007/s12649-016-9519-z

- Alvarenga, P., Mourinha, C., Farto, M., Santos, T., Palma, P., Sengo, J., Morais, M.C., Cunha-Queda,
- 545 C., 2015. Sewage sludge, compost and other representative organic wastes as agricultural soil 546 amendments: Benefits versus limiting factors. Waste Management 40, 44–52.

547 https://doi.org/10.1016/j.wasman.2015.01.027

548 Angin, I., Aslantas, R., Kose, M., Karakurt, H., Ozkan, G., 2012. Changes in chemical properties of

549 soil and sour cherry as a result of sewage sludge application. Horticultural Science 39, 61–66.

550 https://doi.org/10.17221/4/2011-HORTSCI

- Arpa Veneto, 2007. Capitolo 13 Agrelan: un sistema di interpretazione dell'analisi del terreno, in:
 L'interpretazione Delle Analisi Del Terreno Strumento per La Sostenibilità Ambientale. Padova,
 pp. 62–67.
- Arthurson, V., 2008. Proper sanitization of sewage sludge: A critical issue for a sustainable society.

555 Applied and Environmental Microbiology 74, 5267–5275. https://doi.org/10.1128/AEM.00438-08

- 556 Bakshi, M., Liné, C., Bedolla, D.E., Stein, R.J., Kaegi, R., Sarret, G., Pradas del Real, A.E., Castillo-
- 557 Michel, H., Abhilash, P.C., Larue, C., 2019. Assessing the impacts of sewage sludge

amendment containing nano-TiO2 on tomato plants: A life cycle study. Journal of Hazardous
Materials 369, 191–198. https://doi.org/10.1016/j.jhazmat.2019.02.036

- Basosi, R., Spinelli, D., Fierro, A., Jez, S., 2014. Mineral nitrogen fertilizers: Environmental impact of
 production and use. Fertilizers: Components, Uses in Agriculture and Environmental Impacts 3–
 43.
- 563 Bayoumi Hamuda, H.E.A.F., Orosz, E., Horváth, M., Palágyi, A., Szederné Baranyai, B., Patkó, I.,
- 564 Kecskés, M., 2009. Effect of sewage sludge on soil properties, the growth of Lycopersicon
- s65 esculentum L. and the properties of the rhizosphere in a model experiment. Agrokemia es
- 566 Talajtan 58, 325–342. https://doi.org/10.1556/Agrokem.58.2009.2.12
- 567 Bohn, H.L., O'Connor, G.A., McNeal, B.L., 2001. Soil Chemistry, 3rd ed.
- Bot, A., Benites, J., 2005. The importance of soil organic matter, FAO Soils Bulletin.
- 569 https://doi.org/10.1080/03650340214162

- 570 Bourioug, M., Alaoui-Sehmer, L., Laffray, X., Benbrahim, M., Aleya, L., Alaoui-Sossé, B., 2015.
- 571 Sewage sludge fertilization in larch seedlings: Effects on trace metal accumulation and growth
- 572 performance. Ecological Engineering. https://doi.org/10.1016/j.ecoleng.2015.01.031
- Bourioug, M., Krouna, M., Abouabdillah, A., Harraq, A., Bouabid, R., Aleya, L., 2018. Sewage sludge
 used as organic manure in Moroccan sunflower culture: Effects on certain soil properties, growth
 and yield components. Science of The Total Environment 627, 681–688.
- 576 https://doi.org/10.1016/j.scitotenv.2018.01.258
- Buol, S.W., Southard, R.J., Graham, R.C., McDaniel, P.A., 2011. Soil genesis and classification, sixth
 edit. ed, John Wiley & Sons, Ltd., Publication. John Wiley & Sons, Inc.
- Camargo, F.P., Sérgio Tonello, P., Dos Santos, A.C.A., Duarte, I.C.S., 2016. Removal of Toxic Metals
 from Sewage Sludge Through Chemical, Physical, and Biological Treatments—a Review. Water,
 Air, and Soil Pollution 227, 433. https://doi.org/10.1007/s11270-016-3141-3
- Council of the European Communities, 1991a. Council Directive 91/271/EEC of 21 May 1991
 concerning urban waste-water treatment. Official Journal of the European Communities L 135,
 40–52.
- 585 Council of the European Communities, 1991b. Council Directive 91/676/EEC of 12 December 1991
- 586 concerning the protection of waters against pollution caused by nitrates from agricultural
- sources. Official Journal of the European Communities L 375, 1–8.
- 588 Council of the European Communities, 1986. Council Directive 86/278/EEC of 12 June 1986 on the 589 protection of the environment, and in particular of the soil, when sewage sludge is used in 590 agriculture. Official Journal of the European Communities L 181, 6–12.
- Cristina, G., Camelin, E., Pugliese, M., Tommasi, T., Fino, D., 2019. Evaluation of anaerobic
 digestates from sewage sludge as a potential solution for improvement of soil fertility. Waste
 Management 99, 122–134. https://doi.org/10.1016/j.wasman.2019.08.018
- 594 De Andres, E.F., Tenorio, J.L., Walter, I., 2010. Biomass production and nutrient concentration of
- kenaf grown on sewage sludge-amended soil. Spanish Journal of Agricultural Research 8, 472.

596 https://doi.org/10.5424/sjar/2010082-1202

597 De López Camelo, L.G., De Miguez, S.R., Marbán, L., 1997. Heavy metals input with phosphate 598 fertilizers used in Argentina. Science of the Total Environment 204, 245–250.

599 https://doi.org/10.1016/S0048-9697(97)00187-3

- de Oliveira, M.O.A., Alves, S.M. de F., Freitas, E. de F.M., de Faria, H.F.L., Lisboa, F.C., 2017.
- 601 Relative chlorophyll index on doses of nitrogen fertilization for cherry tomato culture. African
- Journal of Agricultural Research 12, 2946–2953. https://doi.org/10.5897/ajar2016.12051
- Dumontet, S., Dinel, H., Baloda, S.B., 1999. Pathogen Reduction in Sewage Sludge by Composting
- and Other Biological Treatments: A Review. Biological Agriculture and Horticulture 16, 409–430.
- 605 https://doi.org/10.1080/01448765.1999.9755243
- Eid, E.M., Alrumman, S.A., El-Bebany, A.F., Fawy, K.F., Taher, M.A., Hesham, A.E.-L., El-Shaboury,
- G.A., Ahmed, M.T., 2019. Evaluation of the potential of sewage sludge as a valuable fertilizer for
 wheat (Triticum aestivum L.) crops. Environmental Science and Pollution Research 26, 392–401.
 https://doi.org/10.1007/s11356-018-3617-3
- Eid, E.M., Alrumman, S.A., El-Bebany, A.F., Hesham, A.E.L., Taher, M.A., Fawy, K.F., 2017. The
- 611 effects of different sewage sludge amendment rates on the heavy metal bioaccumulation, growth
- and biomass of cucumbers (Cucumis sativus L.). Environmental Science and Pollution Research
- 613 24, 16371–16382. https://doi.org/10.1007/s11356-017-9289-6
- Elloumi, N., Belhaj, D., Jerbi, B., Zouari, M., Kallel, M., 2016. Effects of sewage sludge on bio-
- accumulation of heavy metals in tomato seedlings. Spanish Journal of Agricultural Research 14,
- 616 e0807. https://doi.org/10.5424/sjar/2016144-9210
- 617 Epstein, E., 2002. Land application of sewage sludge and biosolids. CRC Press.
- Eurostat, 2019. Sewage sludge production and disposal.
- 619 FAO, 2017. World fertilizer trends and outlook to 2020 Summary report. Food and Agriculture
- 620 Organization of United Nations.

- 621 Fekri, M., Gorgin, N., Sadegh, L., 2011. Phosphorus desorption kinetics in two calcareous soils
- amended with P fertilizer and organic matter. Environmental Earth Sciences 64, 721–729.

623 https://doi.org/10.1007/s12665-010-0892-9

- 624 Fernández-Luqueño, F., Reyes-Varela, V., Martínez-Suárez, C., Salomón-Hernández, G., Yáñez-
- 625 Meneses, J., Ceballos-Ramírez, J.M., Dendooven, L., 2010. Effect of different nitrogen sources
- on plant characteristics and yield of common bean (Phaseolus vulgaris L.). Bioresource
- 627 Technology 101, 396–403. https://doi.org/10.1016/j.biortech.2009.07.058
- 628 Ferreiro-Domínguez, N., Rigueiro-Rodríguez, A., Mosquera-Losada, M.R., 2011. Response to
- 629 sewage sludge fertilisation in a Quercus rubra L. silvopastoral system: Soil, plant biodiversity and
- 630 tree and pasture production. Agriculture, Ecosystems and Environment.
- 631 https://doi.org/10.1016/j.agee.2011.02.009
- Fixen, P.E., Johnston, A.M., 2012. World fertilizer nutrient reserves: A view to the future. Journal of
 the Science of Food and Agriculture 92, 1001–1005. https://doi.org/10.1002/jsfa.4532
- Gruda, N., Caron, J., Prasad, M., Maher, M., 2016. Growing Media, in: Lal, R. (Ed.), Encyclopedia of
- 635 Soil Science, Third Edition. CRC Press, pp. 1053–1058. https://doi.org/10.1081/E-ESS3-
- 636 120053784
- Harrison, E.Z., Oakes, S.R., Hysell, M., Hay, A., 2006. Organic chemicals in sewage sludges. Science
 of the Total Environment. https://doi.org/10.1016/j.scitotenv.2006.04.002
- He, S., Li, A., Lei, W., 2016. Effect of Sewage Sludge and its Biomass Composting Product on the
- 640 Soil Characteristics and N2O Emission from the Tomato Planting Soil. International Journal of
- 641 Agriculture and Biology 18, 501–508. https://doi.org/10.17957/IJAB/15.0113
- Hossain, M.K., Stretzov, V., Nelson, P.F., 2015. Comparative Assessment of the Effect of Wastewater
 Sludge Biochar on Growth, Yield and Metal Bioaccumulation of Cherry Tomato. Pedosphere 25,
- 644 680–685. https://doi.org/10.1016/S1002-0160(15)30048-5
- Jones Jr., J.B., 2012. Plant Nutrition and Soil Fertility Manual, Plant Nutrition and Soil Fertility Manual.
- 646 https://doi.org/10.1201/b11577

- Jones Jr, J.B., 2008. Tomato plant culture, Second edi. ed. CRC Press.
- 648 Kchaou, R., Baccar, R., Bouzid, J., Rejeb, S., 2018. Agricultural use of sewage sludge under sub-
- humid Mediterranean conditions: effect on growth, yield, and metal content of a forage plant.
- 650 Arabian Journal of Geosciences 11, 746. https://doi.org/10.1007/s12517-018-4103-4
- Kladivko, E.J., Nelson, D.W., 1979. Changes in soil properties from application of anaerobic sludge.
 Journal of the Water Pollution Control Federation 51, 325–332.
- 653 Kotecki, A., Kozak, M., Malarz, W., Kotecki, A., 2014. Wpływ nawożenia osadem ściekowym na
- 654 rozwój, plonowanie i skład chemiczny spartiny preriowej (spartina pectinata link.). Journal of
- 655 Elementology 19, 1021–1036. https://doi.org/10.5601/jelem.2014.19.3.725
- Li, L., XianJin, H., Hui, W., YuanHua, D., Sisi, X., 2009. Estimation of environmental costs of chemical
 fertilizer utilization in China. Acta Pedologica Sinica 46, 63–69.
- Liu, S., Zhu, N., Li, L.Y., 2012. The one-stage autothermal thermophilic aerobic digestion for sewage
 sludge treatment: Stabilization process and mechanism. Bioresource Technology 104, 266–273.
 https://doi.org/10.1016/j.biortech.2011.11.041
- Mateo-Sagasta, J., Raschid-Sally, L., Thebo, A., 2015. Global Wastewater and Sludge Production,
- Treatment and Use, in: Wastewater. Springer Netherlands, Dordrecht, pp. 15–38.
- 663 https://doi.org/10.1007/978-94-017-9545-6_2
- Mills, H.A., Jones Jr, J.B., 1996. Plant analysis handbook II: A practical sampling, preparation,
 analysis, and interpretation guide.
- 666 Mirlean, N., Baisch, P., Machado, I., Shumilin, E., 2008. Mercury contamination of soil as the result of
- long-term phosphate fertilizer production. Bulletin of Environmental Contamination and
 Toxicology 81, 305–308. https://doi.org/10.1007/s00128-008-9480-z
- 669 Mirlean, N., Roisenberg, A., 2007. Fluoride distribution in the environment along the gradient of a
- 670 phosphate-fertilizer production emission (southern Brazil). Environmental Geochemistry and
- 671 Health 29, 179–187. https://doi.org/10.1007/s10653-006-9061-1

- Morgan, L., 2000. Beneficial elements for hydroponics: A new look at plant nutrition. Growing Edge.
- 673 Mortvedt, J.J., 1995. Heavy metal contaminants in inorganic and organic fertilizers. Fertilizer research
- 674 43, 55–61. https://doi.org/10.1007/BF00747683
- Mosquera-Losada, M.R., Ferreiro-Domínguez, N., Daboussi, S., Rigueiro-Rodríguez, A., 2016.
- 676 Sewage sludge stabilisation and fertiliser value in a silvopastoral system developed with
- 677 Eucalyptus nitens Maiden in Lugo (Spain). Science of the Total Environment 566–567, 806–815.
- 678 https://doi.org/10.1016/j.scitotenv.2016.05.003
- Nobile, C., Houben, D., Michel, E., Firmin, S., Lambers, H., Kandeler, E., Faucon, M.P., 2019.
- 680 Phosphorus-acquisition strategies of canola, wheat and barley in soil amended with sewage
- 681 sludges. Scientific Reports 9, 1–11. https://doi.org/10.1038/s41598-019-51204-x
- Parry, C., Blonquist, J.M., Bugbee, B., 2014. In situ measurement of leaf chlorophyll concentration:
- 683 Analysis of the optical/absolute relationship. Plant, Cell and Environment 37, 2508–2520.
- 684 https://doi.org/10.1111/pce.12324
- Pascual, I., Avilés, M., Aguirreolea, J., Sánchez-Díaz, M., 2008. Effect of sanitized and non-sanitized
- sewage sludge on soil microbial community and the physiology of pepper plants. Plant and Soil
- 687 310, 41–53. https://doi.org/10.1007/s11104-008-9626-0
- Perez-Espinosa, A., Moreno-Caselles, J., Moral, R., Perez-Murcia, M.D., Gomez, I., 1999. Effect of
 sewage sludge and cobalt treatments on tomato fruit yield, weight, and quality. Journal of Plant
 Nutrition 22, 379–385. https://doi.org/10.1080/01904169909365635
- Pinna, M.V., Giunta, F., Nardi, S., Pusino, A., 2009. Effects of a municipal sewage sludge amendment
- on triasulfuron soil sorption and wheat growth. Journal of Agricultural and Food Chemistry 57,
- 693 11249–11253. https://doi.org/10.1021/jf902659f
- Rasouli-Sadaghiani, M.H., Moradi, N., 2014. Effect of poultry, cattle, sheep manures and sewage
- sludge on N mineralisation. Chemistry and Ecology 30, 666–675.
- 696 https://doi.org/10.1080/02757540.2014.889122

- Reijnders, L., 2014. Phosphorus resources, their depletion and conservation, a review. Resources,
- 698 Conservation and Recycling 93, 32–49. https://doi.org/10.1016/j.resconrec.2014.09.006
- 699 Sabiha-Javied, Mehmood, T., Chaudhry, M.M., Tufail, M., Irfan, N., 2009. Heavy metal pollution from
- phosphate rock used for the production of fertilizer in Pakistan. Microchemical Journal 91, 94–
- 701 99. https://doi.org/10.1016/j.microc.2008.08.009
- Schnug, E., Lottermoser, B.G., 2013. Fertilizer-derived uranium and its threat to human health.
- Environmental Science and Technology 47, 2433–2434. https://doi.org/10.1021/es4002357
- Schulze, E.D., Freibauer, A., 2005. Environmental science: Carbon unlocked from soils. Nature 437,
 205–206. https://doi.org/10.1038/437205a
- Singh, R.P., Agrawal, M., 2008. Potential benefits and risks of land application of sewage sludge.
- 707 Waste Management. https://doi.org/10.1016/j.wasman.2006.12.010
- Singh, R.P., Agrawal, M., 2007. Effects of sewage sludge amendment on heavy metal accumulation
 and consequent responses of Beta vulgaris plants. Chemosphere 67, 2229–2240.
- 710 https://doi.org/10.1016/j.chemosphere.2006.12.019
- 711 Smith, K.M., Fowler, G.D., Pullket, S., Graham, N.J.D., 2009. Sewage sludge-based adsorbents: A
- review of their production, properties and use in water treatment applications. Water Research
- 713 43, 2569–2594. https://doi.org/10.1016/j.watres.2009.02.038
- 714 Solanki, P., Kalavagadda, B., Akula, B., Sharma, S.H.K., Reddy, D.J., 2017. Effect of Sewage Sludge
- on Marigold (Tagetes erecta). International Journal of Current Microbiology and Applied
- 716 Sciences 6, 825–831. https://doi.org/10.20546/ijcmas.2017.606.097
- 717 Stamatiadis, S., Doran, J.W., Kettler, T., 1999. Field and laboratory evaluation of soil quality changes
- resulting from injection of liquid sewage sludge. Applied Soil Ecology 12, 263–272.
- 719 https://doi.org/10.1016/S0929-1393(99)00007-4
- 720 Stehouwer, R.C., Wolf, A.M., Doty, W.T., 2000. Chemical monitoring of sewage sludge in
- 721 Pennsylvania: Variability and application uncertainty. Journal of Environmental Quality 29, 1686–

- 722 1695. https://doi.org/10.2134/jeq2000.00472425002900050041x
- Tariq, U., ur Rehman, S., Khan, M.A., Younis, A., Yaseen, M., Ahsan, M., 2012. Agricultural and
 municipal waste as potting media components for the growth and flowering of Dahlia hortensis
- 725 "Figaro." Turkish Journal of Botany 36, 378–385. https://doi.org/10.3906/bot-1109-16
- Tei, F., Benincasa, P., Guiducci, M., 2002. Critical nitrogen concentration in processing tomato, in:
- 727 European Journal of Agronomy. https://doi.org/10.1016/S1161-0301(02)00096-5
- 728 Vanotti, M.B., García-González, M.C., Molinuevo-Salces, B., Riaño, B., 2019. Editorial: New
- Processes for Nutrient Recovery From Wastes. Frontiers in Sustainable Food Systems 3, 1–2.
 https://doi.org/10.3389/fsufs.2019.00081
- Walter, I., Cuevas, G., García, S., Martínez, F., 2000. Biosolid effects on soil and native plant
 production in a degraded semiarid ecosystem in central Spain. Waste Management and
 Research. https://doi.org/10.1034/j.1399-3070.2000.00111.x
- Wang, Y., Liu, F., Jensen, L.S., de Neergaard, A., Jensen, C.R., 2013. Alternate partial root-zone
 irrigation improves fertilizer-N use efficiency in tomatoes. Irrigation Science 31, 589–598.
- 736 https://doi.org/10.1007/s00271-012-0335-3

739

Warman, P.R., Termeer, W.C., 2005. Evaluation of sewage sludge, septic waste and sludge compost
applications to corn and forage: Ca, Mg, S, Fe, Mn, Cu, Zn and B content of crops and soils.

Bioresource Technology 96, 1029–1038. https://doi.org/10.1016/j.biortech.2004.09.014

- 740 Whitehead, D.C., 1995. Grassland nitrogen. CAB INTERNATIONAL, Wallingford.
- Xiong, D., Chen, J., Yu, T., Gao, W., Ling, X., Li, Y., Peng, S., Huang, J., 2015. SPAD-based leaf
- nitrogen estimation is impacted by environmental factors and crop leaf characteristics. Scientific
 Reports 5, 13389. https://doi.org/10.1038/srep13389
- Zucco, M.A., Walters, S.A., Chong, S.K., Klubek, B.P., Masabni, J.G., 2015. Effect of soil type and
 vermicompost applications on tomato growth. International Journal of Recycling of Organic
- 746 Waste in Agriculture 4, 135–141. https://doi.org/10.1007/s40093-015-0093-3

- Zuo, W., Gu, C., Zhang, W., Xu, K., Wang, Y., Bai, Y., Shan, Y., Dai, Q., 2019. Sewage sludge
- amendment improved soil properties and sweet sorghum yield and quality in a newly reclaimed
- mudflat land. Science of The Total Environment 654, 541–549.
- 750 https://doi.org/10.1016/j.scitotenv.2018.11.127
- 751
- 752

Table 1. Physiscal and chemical anlalysis of soil and peat used in the present work. CEC: Cation-Exchange Capacity; AAS: Atomic Absorption

754 Spectroscopy.

Sand	ly soil		Peat substrate								
Parameter	Unit	Value	Parameter	Unit	Value						
Stones	-	absent	Stones		-						
Sand (2.0 - 0.020 mm)	%	94 ± 2	Sand (2.0 - 0.020 mm)		-						
Silt (0.020 - 0.002 mm)	%	3 ± 1	Silt (0.020 - 0.002 mm)		-						
Clay (< 0.002 mm)	%	3 ± 1	Clay (< 0.002 mm)		-						
Texture	-	sandy	Texture		-						
рН	-	8.2 ± 0.16	рН	-	6.2 ± 0.1						
Electrical conductivity	dS/m	0.131 ± 0.018	Electrical conductivity	dS/m	0.722 ± 0.146						
Organic matter	%	0.38 ± 0.12	Organic matter		-						
Organic carbon	%	0.22 ± 0.07	Organic carbon		-						
N - Tot (Kjeldahl)	g/kg	0.29 ± 0.09	N - Tot (Kjeldahl)	%	0.42 ± 0.06						
N - NO ₂	mg/kg	< 0,2	N - NO ₂	mg/l	< QL						
N - NO ₃	mg/kg	6.33 ± 1.53	N - NO ₃ ⁻	mg/l	30.4 ± 7.2						
$N - NH_4^+$	mg/kg	3 ± 1	$N - NH_4^+$	mg/l	1.3 ± 0.3						
N - Org	g/kg	0.29 ± 0.09	N - Org	%	0.4 ± 0.40						
C/N		7.6 ± 0.2	C/N		-						
P _{Olsen}	mg/kg	1.8 ± 1.3	P _{extractable}	mg/l	8.1 ± 2.3						
K _{exchangeable}	mg/kg	18 ± 1	K _{extractable}	mg/l	41.1 ± 6.8						
Mg _{exchangeable}	mg/kg	15 ± 5	Mg extractable	mg/l	28 ± 7						
Ca exchangeable	mg/kg	675 ± 27	Ca extractable	mg/l	36 ± 8						
Na _{exchangeable}	mg/kg	6 ± 3	Na extractable	mg/l	16 ± 11						
Fe available	mg/kg	6.7 ± 1.1	Fe extractable	mg/l	0.79 ± 0.21						
Mn _{available}	mg/kg	6.5 ± 3.0	Mn _{extractable}	mg/l	0.15 ± 0.04						
Cu available	mg/kg	0.69 ± 0.29	Cu extractable	mg/l	< QL						
Zn _{available}	mg/kg	0.47 ± 0.29	Zn extractable	mg/l	0.02 ± 0.00						
CEC	cmol/kg	3.65 ± 0.35	CEC		-						

Table 2. Chemical characterization performed two months after treatments application on sandy soil (A) and on peat substrate (B). Data are expressed as mean \pm standard deviation. Asterisks mean significant differences according to ANOVA test (*, **, *** differences between means significant at *P* ≤ 0.05, 0.01 and 0.001, respectively). CEC, cation exchange capacity; QL, quantification limit.

758 **A**

Parameter	Unit of measure	С	ontr (T)	ol		Pr	ima (P)	у		Seco	onda (S)	iry		Cent (rifug C)	jed		D	ried (D)			Minera	l fer (M)	tiliser	
рН	-	8.3	±	0.1	***	8.2	±	0.1	***	8.2	±	0.1	***	8.1	±	0.1	***	8.1	±	0.1	***	8.0	±	0.1	***
Electrical conductivity	dS/m	0.155	±	0.020)	0.219	±	0.032		0.201	±	0.010	I	0.197	±	0.023		0.198	±	0.025		0.399	±	0.146	
Organic matter	%	0.16	±	0.01	***	0.24	±	0.01	***	0.18	±	0.02	***	0.25	±	0.02	***	0.26	±	0.01	***	0.16	±	0.02	***
Organic carbon	%	0.09	±	0.00	***	0.14	±	0.00	***	0.11	±	0.01	***	0.14	±	0.01	***	0.15	±	0.00	***	0.10	±	0.01	***
N - Tot (Kjeldahl)	g/kg	0.17	±	0.01	***	0.19	±	0.01	***	0.15	±	0.01	***	0.20	±	0.00	***	0.22	±	0.01	***	0.17	±	0.01	***
N - NO ₂	mg/kg	< QL				< QL				< QL				< QL				< QL				< QL			
N - NO ₃	mg/kg	< QL				< QL				1	±	1	*	4	±	4	*	2	±	1	*	60	±	40	*
N - NH4 ⁺	mg/kg	< QL				< QL				< QL				1	±	1		2	±	0		1	±	0	
N - Org	g/kg	0.17	±	0.01	***	0.19	±	0.01	***	0.15	±	0.01	***	0.20	±	0.00	***	0.22	±	0.01	***	0.17	±	0.01	***
C/N	-	5.5	±	0.2	**	7.3	±	0.6	**	6.9	±	0.5	**	7.3	±	0.5	**	7.1	±	0.3	**	5.8	±	1.0	**
P _{Olsen}	mg/kg	< QL				4.2	±	0.5	**	10.3	±	0.8	**	15.4	±	4.6	**	36.2	±	11.4	**	< QL			
K exchangeable	mg/kg	14	±	3		11	±	3		12	±	1		12	±	2		9	±	1		13	±	2	
Mg exchangeable	mg/kg	11	±	2		21	±	7		26	±	3		25	±	4		22	±	4		25	±	2	
Ca exchangeable	mg/kg	524	±	26	*	594	±	25	*	491	±	62	*	626	±	94	*	579	±	48	*	646	±	62	*
Na exchangeable	mg/kg	16	±	1	***	35	±	2	***	33	±	3	***	26	±	3	***	32	±	3	***	27	±	5	***
Fe available	mg/kg	5.7	±	0.1	***	7.6	±	0.5	***	9.3	±	0.2	***	11.2	±	0.7	***	12.5	±	0.4	***	6.1	±	0.21	***
Mn _{available}	mg/kg	5.1	±	0.4		14.8	±	16.6		33.7	±	2.5		35.8	±	27.1		20.5	±	28.1		5.2	±	0.21	
Cu available	mg/kg	0.40	±	0.08		0.47	±	0.13		0.60	±	0.06		0.85	±	0.12		0.96	±	0.29		0.40	±	0.01	
Zn available	mg/kg	0.21	±	0.03	***	0.36	±	0.04	***	0.37	±	0.02	***	0.73	±	0.08	***	1.00	±	0.07	***	0.43	±	0.05	***
CEC	cmol/kg	2.81	±	0.13	*	3.32	±	0.17	*	2.83	±	0.33	*	3.47	±	0.43	*	3.24	±	0.20	*	3.58	±	0.32	*

 $QL: N - NO_2^- = 0.2 \text{ mg/kg}; N - NO_3^- = 1 \text{ mg/kg}; N - NH_4^+ = 1 \text{ mg/kg}; P = 1 \text{ mg/kg}.$

B

Parameter	Unit of measure	С	ontr (T)	ol		Prim	Primary SSAD (P)		Secor	Secondary SSAD (S)				Centrifuged SSAD (C)				Dried SSAD (D)				Mineral (M)			
рН	-	7.0	±	0.4	*	6.7	±	0.3	*	7.2	±	0.2	*	6.9	±	0.2	*	7.4	±	0.3	*	6.6	±	0.2	*
Electrical conductivity	dS/m	0.235	±	0.040		0.436	±	0.220		0.183	±	0.038		0.495	±	0.134		0.225	±	0.074		0.523	±	0.202	
N - Tot (Kjeldahl)	% D.M.	0.23	±	0.03	*	0.24	±	0.03	*	0.25	±	0.03	*	0.28	±	0.04	*	0.31	±	0.04	*	0.32	±	0.04	*
N - NO ₂	mg/l	< QL				< QL				< QL				< QL				< QL				< QL			
N - NO ₃ ⁻	mg/l	1.0	±	0.9		1.5	±	0.5		1.0	±	0.2		2.6	±	0.4		1.2	±	0.3		2.6	±	1.7	
N - NH4 ⁺	mg/l	< QL				< QL				< QL				< QL				< QL				< QL			
N - Org	% D.M.	0.22	±	0.03	*	0.24	±	0.03	*	0.25	±	0.03	*	0.28	±	0.04	*	0.31	±	0.04	*	0.32	±	0.04	*
P extractable	mg/l	0.4	±	0.1		1.2	±	0.4		1.2	±	0.5		0.6	±	0.3		1.1	±	0.1		< QL			
K extractable	mg/l	2.9	±	0.3		2.0	±	0.3		2.4	±	0.8		2.4	±	0.4		5.0	±	2.2		2.6	±	0.2	
Mg extractable	mg/l	8	±	2		22	±	16		5	±	2		24	±	12		6	±	3		26	±	17	
Ca extractable	mg/l	13	±	4		26	±	16		12	±	3		33	±	14		14	±	3		32	±	17	
Na extractable	mg/l	24	±	2		29	±	7		17	±	3		32	±	4		23	±	7		31	±	4	
Fe extractable	mg/l	1.17	±	0.26		0.52	±	0.46		0.80	±	0.13		0.52	±	0.30		0.73	±	0.06		0.28	±	0.11	
Mn extractable	mg/l	< QL				< QL				< QL				< QL				< QL				0.03	±	0.01	
Cu extractable	mg/l	< QL				< QL				< QL				< QL				< QL				< QL			
Zn extractable	mg/l	< QL				< QL				< QL				< QL				< QL				0.02	±	0	

QL: $N - NO_2^- = 0.05 \text{ mg/l}; N - NH_4^+ = 0.06 \text{ mg/l}; P = 0.3 \text{ mg/l}; Mn = 0.03 \text{ mg/l}; Cu = 0.03 \text{ mg/l}; Zn = 0.02 \text{ mg/l}.$

Table 3. Results of leaves analyses performed after two months after treatments application on sandy soil (A) and on peat substrate (B). Different

⁷⁶⁷ letters indicate differences between treatments that are significant at P < 0.05 (Tukey HSD). Data are expressed as mean ± standard deviation.

A

Parameter	Unit of measure	Prima	ary (P)	SSAD		Drie		Mineral fertilizer (M)								
N	%	1.10	±	0.05	b	1.35	±	0.28	b	2.95	±	0.36	а			
IN	Total (mg)	46.64	±	7.45	b	81.08	±	15.63	а	60.83	±	9.26	ab			
'n	%	0.14	±	0.01	а	0.16	±	0.01	а	0.13	±	0.02	а			
μ	Total (mg)	5.97	±	1.25	b	9.61	±	1.26	а	2.56	±	0.33	С			
ĸ	%	1.46	±	0.38	b	1.40	±	0.18	b	3.63	±	0.57	а			
n	Total (mg)	61.61	±	16.41		84.54	±	10.95		74.24	±	7.52				

B

Parameter	Unit of measure	Co	ontro (T)	ol	Primary SSAD (P)					Dri	Dried SSAD (D)					Mineral fertilizer (M)					
N	%	1.26	±	0.08	b	1.32	±	0.04	b	1.29	±	0.19	b	2.05	±	0.36	а				
	Total (mg)	319.01	±	11.24	b	323.86	±	20.48	b	360.41	±	53.17	b	550.22	±	102.49	а				
'n	%	0.23	±	0.01	b	0.29	±	0.02	ab	0.31	±	0.04	а	0.27	±	0.02	ab				
P	Total (mg)	58.02	±	7.29	b	69.82	±	2.92	ab	85.44	±	9.77	а	72.95	±	8.78	ab				
K	%	1.83	±	0.04	b	1.92	±	0.04	b	1.74	±	0.21	b	2.53	±	0.27	а				
n	Total (mg)	465.11	±	37.08	b	470.69	±	24.49	b	486.97	±	57.85	b	677.03	±	53.15	а				

773 Table 4. Comparison of the results from other works in literature on the effects of treatment with sewage sludge on tomato plants. Application

dosages are shown as reported in the original works; values in brackets indicate how many folds more is the SS application rate with respect to the

present study. n.a., not available.

	<u> </u>		Kind of	Cultivation		Differences with	respect to untreated cont	rol	
Cultivar	ss typology	Dosage	experiment	substrate	Biomass increase	Plant height	Leaves and inflorescences	Chlorophyll content	Reference
Cherry	Digested ^a	10 t/ha (2X)	Pot experiment Greenhouse 16 weeks	Chromosol	Dry biomass: + 20%	10 weeks: + 50% 13 weeks: + 20% 15 weeks: + 7%	n.a.	n.a.	Hossain et al., 2015
Red Robin	n.a.	SS:soil 1:10 (65X)	Pot experiment Growth chamber 120 days	Loamy soil	Fresh biomass Stem: + 70% Leaves: + 142%	+ 43%	Leaves: + 33% Flowers: +130%	Chlorophyll ^b : a: + 18.3% b: + 34.8%	Bakshi et al., 2019
Rio Grande	Aerobically digested	2.5%; 5.0%; 7.5% (11X; 22X; 33X)	Pot experiment Greenhouse 30 days	Sandy soil	<i>Dry biomass:</i> + 180%; + 280%; +140%	n.a	n.a.	<i>Chlorophyll a+b^b:</i> + 17.5%; - 40%; - 68.5%	Elloumi et al., 2016
n.a.	Aerobically digested	400 - 800 kg N/ha (2.35X; 4.7X)	Pot experiment Greenhouse 90 days	Clay soil	Dry biomass: + 18.6% + 29.6%	+ 19.2%; + 24.5%	n.a.	n.a.	He et al. 2016
Beefsteak	Anaerobically digested (4 typologies: P, S, C, D)	170 kg N/ha	Pot experiment Greenhouse 120 days	Sandy soil	Dry biomass up to + 3652% (D treatment, II month)	up to + 500% (D treatment, II month)	Leaves: up to + 180% (S treatments, I month) Flowers: not observed in untreated control	<i>CCI:</i> up to + 172% (D treatment, II month)	This work
				Peat substrate	Dry biomass: up to + 70% (C treatment, I month)	up to + 24% (P treatment, I month)	n.a.	<i>CCI:</i> up to + 64% (D treatment, III month)	_

^a In this work, no details about the typology of digestion are provided.

^b In these works, leaf chlorophyll content was evaluated with methods based on extraction with organic solvents followed by spectrophotometrical quantification.