

AperTO - Archivio Istituzionale Open Access dell'Università di Torino

**What is Climate-Smart Forestry? A definition from a multinational collaborative process focused on mountain regions of Europe**

**This is the author's manuscript**

*Original Citation:*

*Availability:*

This version is available <http://hdl.handle.net/2318/1778117> since 2021-03-07T13:02:12Z

*Published version:*

DOI:10.1016/j.ecoser.2020.101113

*Terms of use:*

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)

# What is Climate-Smart Forestry?

## A definition from a multinational collaborative process focused on mountain regions of Europe

**Authors:** Euan Bowditch<sup>1</sup>; Giovanni Santopuoli<sup>2,3,\*</sup>; Franz Binder<sup>4</sup>; Miren del Rio<sup>5</sup>; Nicola La Porta<sup>6,7</sup>; Tatiana Klavankova<sup>8</sup>; Jerzy Lesinski<sup>9</sup>; Renzo Motta<sup>10</sup>; Maciej Pach<sup>9</sup>; Pietro Panzacchi<sup>3,11</sup>; Hans Pretzsch<sup>12</sup>; Christian Temperli<sup>13</sup>; Giustino Tonon<sup>11</sup>; Melanie Smith<sup>1</sup>; Violeta Velikova<sup>14</sup>; Andrew Weatherall<sup>15</sup>; Roberto Tognetti<sup>2,3,7</sup>

1: Inverness College UHI, University of the Highlands and Islands, 1 Inverness Campus, Inverness IV2 5NA, UK

2: Dipartimento di Agricoltura, Ambiente e Alimenti, Università degli Studi del Molise, Via Francesco De Sanctis, I-86100, Campobasso, Italy

3: Centro di Ricerca per le Aree Interne e gli Appennini (ARIA), Università degli Studi del Molise, via Francesco De Sanctis 1, 86100 Campobasso

4: Sachgebiet Schutzwald und Naturgefahren, Bayerische Landes-anstalt für Wald und Forstwirtschaft, Am Hochanger 11, 85354 Freising, Germany

5: INIA, Forest Research Centre, Department of Silviculture and Forest Management, Crta. La Coruña km 7.5, 28040 Madrid, Spain. IUFOR, Sustainable Forest Management Research Institute, University of Valladolid & INIA

6: IASMA Research and Innovation Centre, Fondazione Edmund Mach, Department of Sustainable Agroecosystems and Bioresources, San Michele all'Adige, Trento, Italy

7. The EFI Project Centre on Mountain Forests (MOUNTFOR), San Michele a/Adige, Trento, Italy

8: SlovaKGlobe, Slovak University of Technology, Institute of Forest Ecology, Slovak Academy of Sciences, Vazovova 5, Bratislava 812 43, Slovak Republic,

9: Department of Forest Biodiversity. University of Agriculture. Al. 29-Listopada 46. 31-425 Krakow, Poland

10: Università degli Studi di Torino, DISAFA, L.go Braccini 2, I-10095 Grugliasco, TO (Italy)

11: Facoltà di Scienze e Tecnologie, Libera Università di Bolzano, Piazza Università 1, 39100, Bolzano/Bozen, Italy

12: Technische Universität München. Hans-Carl-von-Carlowitz-Platz 2. D-85354 Freising, Germany

13: Swiss Federal Institute for Forest, Snow and Landscape Research WSL - Forest Resources and Management. Zürcherstrasse 111 CH-8903 Birmensdorf, Switzerland

14: Institute of Plant Physiology and Genetics - Bulgarian Academy of Sciences, Acad. G. Bonchev Str., Bl.21 1113, Sofia, Bulgaria

15: National School of Forestry, University of Cumbria, Lake District Campus, Ambleside, Cumbria, LA22 9BB, UK

## **Abstract**

1  
2 Climate-Smart Forestry (CSF) is an emerging branch of sustainable forest management that aims to manage  
3 forests in response to climate change. Specific CSF strategies are viewed as a way forward for developing  
4 suitable management responses and enhancing the provision of ecosystem services. However, there is  
5 currently a lack of comprehensive and cohesive assessment to implement CSF. This paper describes the  
6 step-by-step process that developed a comprehensive and shared definition of CSF, and the process for  
7 selecting indicators that assess the “climate-smartness” of forest management. Adaptation, mitigation and  
8 social dimensions are the core focus of the CSF definition, which recognises the need to integrate and avoid  
9 development of these aspects in isolation. An iterative participatory process was used with a range of  
10 experts in forest-related fields from the CLIMO project, this was subsequently supported by a network  
11 analysis to identify sustainable forest management indicators important to CSF. The definition developed  
12 here, is an important first step in to promote CSF that will aid practice in the forestry sector. It can be used  
13 as a template across Europe, tailored to local contexts. Further work communicating CSF to practitioners  
14 and policy-makers will create a CSF practice and culture that will help to safeguard future forest economies  
15 and communities.  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29

30 **Keywords: sustainable forest management, adaptation, mitigation, social dimension, bioeconomy,**  
31 **indicator**  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

# 1. Introduction

In recent years the term Climate-Smart Forestry (CSF) has become increasingly common in forestry circles permeating into academic and sector-specific culture (Hansen et al., 2010; Jantke et al., 2016; Nabuurs et al., 2018; Yousefpour et al., 2018). The CSF concept is emerging as an important next step in furthering the goals of Sustainable Forest Management (SFM) and the sector's response to the threat of climate change.

The Intergovernmental Panel on Climate Change (IPCC) places great emphasis on forests and associated activities, as crucial for mitigating the impacts of climate change (IPCC, 2014). In order to meet the Paris Agreement goals, considerable contributions will be required from forests, as nations are simultaneously aiming to reduce forest degradation and enhance carbon sinks (Rockström et al., 2017). As a consequence, forests can be viewed as both an issue and a solution, with healthy and widely beneficial forests the ultimate goal.

Forestry has been placed at the forefront of action to mitigate climate change through afforestation and carbon sequestration efforts (Bastin et al., 2019). Subsequent management strategies will be significant, as these will determine and regulate emissions, as well as producing ecosystem services that will support and enhance dependent communities. The visibility of action through afforestation provides clear metrics and mitigation goals that currently align with the aims of many governments that are pledging to massive planting commitments without future planning for these new forests. Planting alone does not guarantee mitigation; careful establishment and subsequent management is required to support the forests, communities and economies. CSF definition and indicators could be vital important for guiding climate-smart decision-making. Moreover, they could be components of a broader landscape-scale strategy, which continuously develops with complex socio-ecological systems to achieve sustainable development goals (Denton et al., 2014). Coordination and collaboration between stakeholders to create a shared vision is paramount to achieving such goals (Folke et al., 2010; Kok et al., 2014). Often institutional and technical capacities need to develop with working concepts such as CSF to help assess and promote effective application by stakeholders.

## 1.1. Climate-Smart Forestry definition

27 The climate-smart concept originated with agriculture (FAO, 2010a) and recognises the need for “an  
1  
28 *approach that helps to guide actions needed to transform and reorient agricultural systems to effectively*  
3  
4  
529 *support development and ensure food security in a changing climate.”* This approach, which is adopted by  
6  
730 Nabuurs et al. (2018) with regard to forestry, describes three main objectives using climate-smart  
8  
931 agriculture as a template: (i) sustainably increasing production and incomes; (ii) adapting and building  
10  
11  
1232 resilience; and (iii) reducing or removing GHG emissions (FAO, 2010a; Zilberman et al., 2018). Nabuurs and  
13  
1433 colleagues (2018) focus on the third objective of reducing or removing GHG emissions through forestry.  
15  
16  
1734 A Web of Science database query with the keywords ‘Climate-Smart Forest’ resulted in 106 publications  
18  
19  
2035 since 2000 (Figure S1) with only four articles explicitly addressing the subject of ‘Climate-Smart Forestry’ in  
21  
2236 Europe as the main focus of the articles (Jandl et al., 2018; Nabuurs et al., 2018, 2017; Yousefpour et al.,  
23  
2437 2018). CSF literature over the last eighteen years addressed a number of issues including GHG emissions  
25  
2638 (Yousefpour et al., 2018) carbon sequestration with a strong cluster of publications on REDD+ (Nabuurs et  
27  
28  
2939 al., 2018; Vass and Elofsson, 2016), land use and habitat change (Pussinen et al., 2009; Smiraglia et al.,  
30  
3140 2016) impact on tree species diversity/distribution and forest structure (Del Río et al., 2016), wildfire  
32  
33  
3441 regimes (Fernandes, 2013; Sousa-Silva et al., 2018) and the resulting effects on management and decision-  
35  
3642 making (Hansen et al., 2010). Beyond the CSF concept, there is an emerging area focusing on mobile  
37  
38  
3943 applications that collect forest data through citizen science and professional monitoring networks (Yang et  
40  
4144 al., 2015). A thorough analysis of the CSF related literature is provided in Appendix A of supplementary  
42  
4345 material.  
44  
45  
4646 No widely adopted definition currently exists for CSF, therefore, the term is being interpreted in a number  
47  
48  
4947 of ways, which mostly focus on the reduction of greenhouse gas (GHG) emissions and effective carbon  
50  
5148 sequestration, as the core mitigation actions (Nabuurs et al., 2018; Yousefpour et al., 2018). Socio-  
52  
53  
5449 ecological systems have mostly not been addressed so far in CSF with the majority of studies based on  
55  
5650 modelling techniques mainly addressing carbon sequestration, substitution and climate impacts (Nabuurs  
57  
5851 et al., 2018; Yousefpour et al., 2018). For this reason, practical guidance for assessing and implementing  
59  
60  
61  
62  
63  
64  
65

52 CSF is needed to support forest regions and to promote climate-smart management within the context of  
1  
23 diverse social-ecological systems that they support (Folke et al., 2016; Melnykovich et al., 2018).  
3  
4

54 **1.2. From SFM indicators to CSF indicators**  
6  
7

85 Forests provide valuable ecosystem services that contribute to social capital, support rural economies and  
9  
10 generate significant income for communities (Biber et al., 2015; Melnykovich et al., 2018). Folke et al.  
11  
12 (2016) highlighted that social and ecological interests are inextricably intertwined, shaping one another  
13  
14 along spatial and temporal trajectories, which feedback to influence opinion and behaviour. These changes  
15  
16 to external effects, such as climate change, require shedding legacy social-ecological systems to be replaced  
17  
18 by new social-ecological systems capable of responding to current and emerging challenges that threaten  
19  
20 human well-being. Social-ecological legacy and adaptability present as much an uncertainty and  
21  
22 unpredictability, as climate change impacts, therefore forest managers rely on structures, expertise and  
23  
24 guidance in which to operate to minimise unwanted outcomes. SFM Criteria and Indicators (C&I), payment  
25  
26 for ecosystem services (PES) and certification schemes represent such tools and expert guidance. In  
27  
28 particular, C&I are a widely-applied policy tool for monitoring, assessing and reporting on SFM and for  
29  
30 supporting the definition of forest management priorities and targets (Santopuoli et al., 2016;  
31  
32 Wijewardana, 2008; Wolfslehner and Baycheva-Merger, 2016).  
33  
34  
35  
36  
37  
38  
39  
40 C&I for SFM may be required in areas where the capacity to collect, measure, record and assess data is  
41  
42 challenging. Here guidance can increase the visibility and highlight the importance of the services that  
43  
44 forests provide to the wider landscape. In a similar way, a definition and indicators for CSF, can enable  
45  
46 forestry professionals to respond to current uncertainties in forest management. Processes such as SFM  
47  
48 recognise that our understanding of forest ecosystems, socio-economic conditions, technologies and  
49  
50 stakeholder priorities will continually change. Therefore, indicators must be flexible enough to adapt to  
51  
52 emerging challenges and demands (Linser et al., 2018), but also highlight when new guidance is needed and  
53  
54 management should be reassessed.  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

76 This paper presents, for the first time, a broad definition of CSF, which can be subjected to further testing  
1  
277 and verification. The definition was developed through an iterative deliberative process that involved  
3  
478 confrontation among a multidisciplinary group of experts from 28 countries (<http://climo.unimol.it/>)  
5  
6  
779 participating in the COST action CA 15226, Climate-Smart Forestry in Mountain Regions (CLIMO).  
8  
980 Developed for policy-makers and practitioners, the CSF definition initially aimed to guide European  
10  
11  
1281 forestry, but can also be used as a template in other areas of the world. Moreover, the paper describes the  
13  
1482 process used to select indicators, enabling assessment of the climate-smartness of forests in support of  
15  
1683 SFM and limiting the negative impacts induced by climate change. Section 2 describes the definition  
17  
18  
1984 building process as well as the methodology for the network analysis of the SFM indicators. The results of  
20  
2185 the network analysis and the CSF definition are described in the section 3.1 and 3.2 respectively. Section 4  
22  
2386 discusses the implications for European forestry and climate policy, while outlining further work and next  
24  
25  
2687 steps. Section 5 covers conclusions.

## 28 2. Material and methods

### 32 3.1. The definition process

35  
3690 The CSF definition process involved participants of the EU COST Action CLIMO during and between three  
37  
3891 separate meetings (Table S1). Deliberative processes are used in diverse environments from shaping of  
39  
4092 international policy to decision-making at a community level with the aim to build common consensus on a  
41  
42  
4393 wide range of perspectives, leading to collective agreements and actions (Dietz, 2013; Wolf and Klein,  
44  
4594 2007). Development of the CSF definition for CSF and relevant indicators for monitoring and assessment  
46  
4795 (Figure 1), aims to build a guiding framework for future research, practice and policy.

49  
50  
5196 The multidisciplinary background of CLIMO forest professionals fostered an iterative and discursive  
52  
5397 participatory approach that has proven effective for forest governance and wider decision-making  
54  
5598 processes. This enabled participants to share views on forest management and CSF, identifying important  
56  
57  
5899 linkages between SFM and forest ecosystem services that would prove beneficial for CSF.

100 Säynäjoki (2014) suggests that confrontation of perspectives is a key stage for groups that aim to elicit and  
1  
101 examine individual reasoning, allowing for broader reflection on complex issues. Therefore, the definition  
3  
4  
102 was tested and challenged over the series of meetings (Table S1) until a consensus was reached with the  
5  
6  
103 current version. These meetings were part of a longer process planned to take the definition from an initial  
8  
104 concept inspired by Climate-Smart Agriculture and the Forest Europe C&I for SFM, through this initial  
10  
11  
105 definition. Further amendments are envisioned after consultation with policymakers and practitioners with  
12  
13  
106 the ultimate aim of developing a CSF toolkit for forest managers.  
15  
16

107 [Insert Figure 1 here]  
18  
19

108 At the inaugural meeting (Trento, Italy, February 2017), Working Group (WG1) was formed to develop a  
21  
22  
109 comprehensive definition for CSF and to identify criteria for CSF assessment and monitoring to generate  
23  
24  
110 baselines for future research, practice and policy. Working Group 1 was the largest of four working groups  
26  
27  
111 with a diverse range of participants with backgrounds ranging from forest management to ecophysiology  
28  
29  
112 and ecology, from soil science to forest policy, as well as genetics and landscape management (Table S2 -  
30  
31  
113 Appendix B, in Supplementary material). In the first meeting, the general aim and methodology for  
33  
34  
114 reaching a definition and selecting criteria were discussed. A sub-group of WG1, with members from the  
35  
36  
115 COST Action CORE group and participants from other working groups, was established with the specific task  
37  
38  
116 of developing a CSF definition. This sub-group met in Sofia (Bulgaria) in September 2017 to produce a  
40  
41  
117 preliminary (incomplete) draft of a one-page definition. This was shared by email for consultation and  
42  
43  
118 amendment first with the sub-group participants who had not been able to attend, and second with all  
45  
46  
119 members of the COST Action between October 2017 and February 2018. In February 2018, the sub-group  
47  
48  
120 reconvened at the annual CLIMO meeting in Sofia to come up with a first full draft that could be presented  
49  
50  
121 during the plenary session to all participants.  
52  
53

122 At the end of the third meeting, the definition was challenged, in an open discussion, by representatives  
55  
56  
123 from each WG by providing comments, suggestions and amendments that had been compiled and  
57  
58  
124 synthesised from each separate WG. These were presented to WG1 and discussed until a  
60  
61  
62  
63  
64  
65

125 consensus/agreement was reached to either integrate, modify or reject the proposed changes. This  
1  
126 confrontational method allowed all suggestions to be put forward for debate, defence and negotiation,  
3  
127 which is an increasingly necessary stage for contentious issues and areas of uncertainty (Mavrommati et al.,  
4  
5  
6  
128 2017; Pellow, 1999; Runhaar et al., 2016). This important step ensured that expectations, inherent bias and  
8  
129 agendas were challenged, so that the definition can be judged to be developed through a comprehensive,  
10  
11  
130 transparent, legitimate and effective process.  
12  
13

131 A high number of CLIMO members actively participated in the development of the definition. Overall, 48  
14  
15  
16  
132 members contributed from 19 countries with a range of expertise and management experience (Table S2 -  
18  
19  
133 Appendix B, in Supplementary material).  
20  
21

## 134 **2.2. Determining criteria and indicators**

22  
23  
24

135 Since the 1990s, many international processes developed principles, C&I to support SFM worldwide  
25  
26  
27  
136 (Castañeda, 2000). In Europe, the pan-European set of C&I for SFM is considered one of the most important  
29  
30  
137 tools for assessing, monitoring and reporting on SFM, and is implemented widely among European  
31  
32  
138 countries (Santopuoli et al., 2016; Wolfslehner and Baycheva-Merger, 2016).  
34

139 Indicators for assessing CSF include adaptation and mitigation strategies, and also consider the multiple  
35  
36  
37  
140 benefits that forests provide to society. The CLIMO partners agreed to identify CSF indicators, from the  
38  
39  
141 pan-European set (Table 1), that address adaptation and mitigation, but at the same time observe a  
40  
41  
42  
142 number of forest ecosystem services, which are important for supporting smart decision-making in forestry.  
44  
45  
143 For this reason, the Common International Classification of Ecosystem Services (hereafter CICES) version 4  
46  
47  
144 (Haines-Young and Potschin, 2012) was used to select the most relevant forest-related ecosystem services  
48  
49  
145 (Table 2). Maintaining ecosystem services is crucial for supporting human well-being, however the quality  
50  
51  
146 of these services relies on healthy forest environments and sustainable forest management. For this  
52  
53  
147 reason, all ecosystem services were considered equally important in this study.  
54  
55  
56

148 [Insert Table 1 here]  
57  
58  
59  
60  
61  
62  
63  
64  
65

149 During the meeting in Sofia 2017 participants were asked to (i) define whether or not the SFM indicators  
1  
150 supported monitoring of adaptation and/or mitigation forest management strategies; and (ii) identify the  
2  
3  
4  
151 forests ecosystem services that can be monitored by indicators previously identified. Beyond the SFM  
5  
6  
152 indicators, four additional forestry indicators (i.e., management system, slenderness coefficient, vertical  
7  
8  
153 and horizontal distribution of tree crowns) were included in the process by CLIMO participants, due to their  
9  
10  
154 accessibility through National Forest Inventory data and field surveys. Participant choices were organized in  
11  
12  
13  
1455 a matrix, within which forest ecosystem services were columns, and SFM indicators were rows. Three  
15  
16  
156 matrices were created, one for adaptation, one for mitigation, and one for adaptation and mitigation  
17  
18  
157 combined, namely CSF (see Figure S2, S3, and S4 appendix C).  
19  
20  
21

2158 [Insert Table 2 here]  
22  
23  
24

259 The Analytic Network Analysis was implemented to identify the most relevant indicators for assessing CSF.  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

166 Additionally, key arguments from CLIMO participants in response to the group's discussion and debate of  
167 indicators, iteratively contributed to shape the CSF definition.  
168 Results were displayed as a network through a fuzzy cognitive map carried out with NetDraw software  
169 (Borgatti et al., 2002). Indicators and forest ecosystem services are the nodes of the network, while the ties  
170 reflect the suitability of indicators to monitor adaptation, mitigation or both, as well as the forest  
171 ecosystem services they address. In the two-mode network, the ties are unidirectional flowing, from the  
172 indicators to ecosystem services, therefore no connections between nodes of the same entity (i.e., among

173 indicators) exist. Further details about network analysis and structure can be found in the Annex C in  
1  
274 Supplementary material.

### 175 **3. Results**

#### 176 **3.1. Network analysis**

177 Twenty-three forest-related ecosystem services were selected and used in the CSF assessment (Table 2).

178 Twelve out of 23 belong to the regulating section, six to the provisioning and five to the cultural section.

179 However, regulating (i.e., sequestration, biological control, and refugia) and provisional services (i.e., raw  
180 materials and PBP) were highly connected with most of the indicators (Figure 2 and Figure S3). By contrast,  
181 cultural ecosystem services, such as spiritual, recreation, tourism and aesthetic, were slightly less  
182 connected.

183 Results show that a total of 29 indicators were selected by participants as suitable to assess adaptation and  
184 mitigation, then were subsequently used to perform the network analysis. Twenty-five of these indicators  
185 came from the original 34 for SFM, plus the four new indicators identified by the participants (Table 2).

186 Seven indicators of criterion 6 – socio-economic function (i.e., 6.1 forest holdings, 6.2 contribution of forest  
187 sector to GDP, 6.3 net revenue, 6.4 expenditure for services, 6.5 forest sector force, 6.6 occupational safety  
188 and health, and 6.11 cultural and spiritual values) and two indicators of criterion 3 -forest productivity (i.e.,  
189 3.3 non-wood goods, and 3.4 services) were not selected. This highlights that many socio-economic aspects  
190 were lost when considering adaptation and mitigation as central pillars of CSF (for the group of experts  
191 consulted).

192 The centrality variables (Table 3) confirms this trend showing that indicators 6.8 trade in wood and 6.1  
193 accessibility, are two of the most peripheral indicators. Nevertheless, nine out of 29 indicators are central  
194 to the network, showing a degree of 23, which means that they have connections with all the 23 forest  
195 ecosystem services considered in this study.

196 Moreover, contrary to the expectations, indicators 1.4 carbon stock, 1.2 growing stock and 4.5 deadwood,  
197 are positioned peripherally in the network due to their linkages with regulating services and poor

198 connection with other forest ecosystem services. Conversely, the new indicators provided by CLIMO  
199 participants obtained higher and medium centrality values.

200 [insert Table 3 here]

201 Four groups have been identified based on the centrality values (Table 3) and the position of nodes in the  
202 network (Figure 2).

203 1<sup>st</sup> core group indicators

204 This group includes 12 indicators with higher centrality values (Table 3), positioned centrally in the  
205 network. Strong linkages are demonstrated to indicators in assessing the preservation of forest resources,  
206 such as indicators 1.1 forest area and 4.7 forest fragmentation. Most of the indicators of criterion 4 –  
207 biodiversity conservation - are included in this group, such as 4.3 naturalness, 4.1 tree species composition  
208 and 4.9 protected forests. Moreover, the group includes indicators which address planning and promoting  
209 sustainable timber production, such as 3.5 management forest plan, 6.9 energy from wood and 6.7 wood  
210 consumption. Monitoring these core indicators will be necessary in order to promote CSF through the  
211 adaptation and mitigation by forestry practices.

212 [insert figure 2 here]

213 2<sup>nd</sup> core group indicators:

214 This group centres upon indicators of criterion 2 – forest health and vitality, including 2.2 soil condition and  
215 2.3 defoliation, as well as indicators of criterion 4 – forest biodiversity including 4.2 regeneration, 4.6  
216 genetic resources, and 4.8 threatened tree species. Indicator 5.1 protective forests can be considered an  
217 overarching indicator for this group. Results show that these indicators have higher centrality values and  
218 similar capacity to monitor both provisioning and regulating services. Interestingly, this group has also  
219 some connections to cultural ecosystem services suggesting that management under CSF would directly  
220 enhance cultural services of the forest ecosystem. For these reasons they are considered core group even  
221 though they occupy a peripheral position in the network.

222 1<sup>st</sup> peripheral indicators:

1  
2  
223 This group includes indicators that demonstrate the influence of silvicultural interventions at stand level  
4  
224 which effect tree competition, tree growth and forest production (3.1 increment and felling, 1.3 age  
6  
7  
225 structure, 1.2 growing stock, 1.4 carbon stock and 4.5 deadwood). These indicators are mostly linked with  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

231 2<sup>nd</sup> peripheral indicators:

232 This group is composed of a single indicator, 6.10 accessibility for recreation, mainly due to the marginal  
233 position rather than the relatively low centrality values. However, the marginal position on one hand and  
234 the power of connection with cultural ecosystem services on the other identifies accessibility as peripheral  
235 indicator. This does not diminish its important role for assessing CSF, particularly in mountain forests for  
236 which forest roads are crucial not only for recreation purposes but also to ensure harvesting activities (i.e.,  
237 sanitary interventions), counteract forest fires and reduce forest management costs.

238 Results highlighted perspectives of CLIMO participants with regard to monitoring and promoting CSF.

239 During the analysis the main themes which emerged from the participatory process and network analysis  
240 (Figure 3), are summarised below:

- 241 • *CSF is not limited to regulating ecosystem services.*
- 242 • *Strengthening of adaptation and mitigation measures to support forest management strategies.*
- 243 • *Forest health and vitality are crucial for protection and maintaining other forest functions.*
- 244 • *Maintaining forest biodiversity is key to counteracting climate change.*
- 245 • *CSF has to maintain and enhance the provision of wider forest ecosystem services.*

- *Integration of social dimensions is key for implementing climate-smart forest management.*

[insert Figure 3 here]

### **3.2. Climate-smart forestry definition**

The working group participants developed a CSF definition that included five sections: a brief overarching CSF definition, sections on adaptation, mitigation and social dimensions and a concise summary statement about CSF (Box 1).

[insert Box 1 here]

## **4. Discussion**

### **4.1. CSF definition and indicators**

CSF continues to develop, as a concept and in practice, but enhancing and facilitating clear implementation pathways could be the difference between forest communities either thriving or declining. The underpinnings of CSF are already integrated into literature, policy and practice through established and accepted frameworks (i.e., adaptive management, mitigation and ecosystem services) under the SFM umbrella, which provides a platform to develop targeted and relevant climate-smart expertise. Climate change is challenging current management systems with wide reaching impacts for forests and societies; these challenges are heightened when considering vulnerable areas such as mountainous regions.

Therefore, rapid action that fosters greater understanding of climate induced changes will require meaningful cooperation between practitioners and policy-makers. Communication of relevant advances in the field is needed to address these issues with targeted knowledge and management approaches that can be applied and adapted to local areas. In order to continually refine the assessment of CSF indicators and to promote communication among policy-makers, forest researchers and practitioners, a CSF approach that develops understanding and impact needs to be established.

In section 3.1, indicators considered central to adaptation and mitigation for forests (carbon stock and growing stock: see Figure 2) were identified as peripheral for CSF, whereas core group indicators were more explicitly related to human activities, management actions and interventions (fragmentation, wood

271 consumption, wood energy and management plans etc.). These focus on the combined processes and  
1  
272 interactions between forests and human intervention through forestry activities. However, indicators such  
2  
3  
4  
273 as trade in wood, roundwood, as well as increment and felling fall within the 1<sup>st</sup> peripheral group despite  
5  
6  
274 their association with management. Linser et al. (2018) stress that the discussion around SFM will  
7  
8  
275 continually evolve and respond to socio-economic aspects and emerging challenges. Therefore, subsets of  
9  
10  
11  
276 indicators might be necessary to link with other sectors or tailored to fit particular issues such as climate  
12  
13  
14  
277 change. CSF indicators may change with the input from more diverse range of stakeholders or more  
15  
16  
278 focussed geographical ranges. This paper differs from those of Nabuurs (2018; 2017) and Yousefpour  
17  
18  
279 (2018) as it includes adaptation and mitigation as broad frameworks, and emphasises the importance of  
19  
20  
280 the social dimensions of forestry. Additionally, this study includes a core set of CSF indicators that could  
21  
22  
23  
281 guide managers and policy-makers to more climate-smart practices, enhancing climate adaptation of  
24  
25  
282 forests and the provision of ecosystem services. Central indicators promote forest biodiversity conservation  
26  
27  
283 and sustainable management of forest resources that fit with the demand of both the forest sector and  
28  
29  
30  
284 wider public.  
31  
32  
33  
285 Jandl et al. (2018) conclude that the production of long living wood products (i.e., construction timber) is a  
34  
35  
36  
286 favourable strategy for CSF wherein carbon is stored in longer standing trees and products. This supports  
37  
38  
287 the centrality of the wood consumption indicator, which takes into account the use of wood and the types  
39  
40  
288 of products that would be favourable to CSF. However, some issues are related to the risk to long-term  
41  
42  
289 standing trees from episodic disturbances and the fact that long-standing timber slows carbon capture in  
43  
44  
45  
290 comparison to new forests (Harmon and Campbell, 2017; Jandl et al., 2018). The relationship between tree  
46  
47  
291 carbon and resulting soil carbon capture is viewed as positive for CSF even though tree carbon capture  
48  
49  
50  
292 slows with time, soil carbon is generally recognised to increase. In spite of this widely accepted view, Ji et  
51  
52  
293 al. (2017) present evidence for accumulation of soil carbon peaks in temperate broadleaf forests in stands  
53  
54  
294 around 50 years of age during the pre-mature stage, with general decreases thereafter. This further  
55  
56  
295 emphasises the importance of adapting CSF recommendations to regional or local context. Scenarios of CSF  
57  
58  
296 are based upon trade-offs and decision-making processes, suitable to the socio-economic systems and  
59  
60  
61  
62  
63  
64  
65

297 ecological lifecycles. As an example, early thinning can be viewed as an integral stage of forest production  
1  
298 to supply bioenergy (woodfuel), substituting the use of fossil fuels, while supporting growth of higher  
2  
3  
4  
299 quality and economically valuable final timber crop (roundwood) (Bowditch et al., 2019; Jandl et al., 2018;  
5  
6  
300 Röser et al., 2011). Managing woody biomass, for example, may reduce the risk of large catastrophic  
8  
301 wildfires and alter the flow rates of watercourses during extreme weather conditions (droughts, heat  
10  
302 waves, rainstorms and strong winds) to reduce flooding risk. The potential impact of a wood energy  
11  
12  
13  
303 indicator through management and use would explain the importance within the network analysis.  
15  
16  
304 Developing CSF resources to contextualise the indicators, especially at a local level, will require collection of  
18  
19  
305 data from long-term monitoring plots, creation of new plots to evaluate targeted attributes at both the  
20  
21  
306 stand and landscape level, which will enable the analyses of trends in CSF indicators to identify priority  
22  
23  
307 areas for adaptation and mitigation. Trends in CSF indicators could also be mapped out through the  
25  
26  
308 trajectory of scenarios that either implement CSF or alternative management actions (Yousefpour et al.,  
27  
28  
309 2010). Other tools, such as the European network of marteloscope sites, are being used to gather historic  
30  
31  
310 and baseline data, as well as being deployed for forestry training activities (Santopuoli et al., 2019). These  
32  
33  
311 sites have the potential to play a crucial role in testing SFM indicators through the simulation of various  
34  
35  
312 forestry scenarios, as well as maintaining and evaluating time series data from sites that quantify  
37  
38  
313 vulnerability to climate change.  
39  
40  
314 Mina et al. (2017) observed that climate change projections under different management scenarios have  
41  
42  
43  
44 highly variable impacts on the provision of ecosystem services in mountains areas, subject to regional and  
45  
46  
47 site conditions, as well as future climate responses. Therefore, establishment of connected and comparable  
48  
317 demonstration sites and permanent forest plots focused upon collection of CSF indicator data over  
49  
50  
518 geographically distributed areas will aid monitoring of vital climate change impacts on forest ecosystems  
52  
53  
519 (e.g., Instruction on Site Characteristic in measurements in Forests (ICOS)). Nevertheless, long-term plots  
54  
55  
520 can only approximate certain trade-offs without greater accountability of disturbances and pressures that  
56  
57  
521 change responses (Locatelli et al., 2017). In some areas, such as mountain regions, vulnerability to climate  
58  
59  
60  
622 change and everyday pressures will place forestry at the centre of an integrated social-ecological support  
61  
62  
63  
64  
65

323 system that enhances economic opportunity and boosts resilience to the uncertainty of climate induced  
1  
324 changes (Schultz et al., 2015). Therefore, planning and management strategies would benefit from co-  
3  
4  
325 development with local populations to ensure investment and recognition of human-forest key  
5  
6  
326 relationship.  
8  
9

## 10 **4.2. Implementation**

11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52

Each country and region will interpret definitions, principles and indicators differently, as diverse forest types and tree species, ecosystems, socio-economic conditions, and people will vary. These will be especially prevalent in critical or more sensitive areas, such as European mountains (Lexer and Bugmann, 2017). As the need for action against climate change heightens, greater guidance will be needed at all scales from international policy to individual forest managers. A keystone development to bridging the gap between policy and practice will be effective engagement with forest managers, forest communities and different forest owner types to assess the accessibility of CSF indicators. Implementing locally adapted CSF through diverse interactive methods aims to help sustainable communities develop alongside a productive forest resource. Andersson and Keskitalo (2018) state that actions must be achieved through social rather than environmental logics, as internal institutional systems, motivations and incentives will often steer and limit adaptation conceived by external drivers and influences. Mitigation strategies that dominate high-level policy should be balanced and aligned with development of lower level actions, so regional nuance of both the culture and managed environment are captured, and reflected in *'applied or working'* policy. Additionally, over-dependency on carbon sequestration and storage as a panacea to climate-induced changes could seriously overshadow adaptive capacity on regional and local scales inhibiting small innovations and planning strategies that could contribute cumulatively to the global issue (Bull et al., 2018; Frame et al., 2018; Thornton and Comberti, 2017).

53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

CSF adaptation measures can include the aim to improve management of specific species mixtures to maintain production under a changing climate, as well as maintaining or increasing associated biodiversity (Del Río et al., 2016, 2014; Nabuurs et al., 2018; Pretzsch et al., 2017), which could simultaneously support the social dimensions of these forests for resilient communities (Armatas et al., 2016; Brunner and Grêt-

349 Regamey, 2016; Seidl et al., 2016). A mixed species approach can be used to buffer but not entirely protect  
1  
350 against conditions resulting from climate change, however, the complementarity of mixed species  
3  
4  
351 outperforms block mixtures under most climate change scenarios (Paul et al., 2019). Many forests will be in  
5  
6  
352 vulnerable climatic zones with concentrated climate induced changes impacting species mixes, habitat  
8  
9  
353 suitability and ultimately shifting distribution ranges. Nevertheless these changes may favour and provide  
10  
11  
354 new production opportunities for some regions (Lindner et al., 2014). This may represent an important  
12  
13  
355 revenue source for local economies, especially for communities suffering from depopulation, as well as a  
15  
16  
356 lack of services and infrastructure (structural and business) that supports community development.  
17  
18  
357 However, most predictions cite the reduction of species range and consequently lower production as more  
20  
21  
358 probable (Liang et al., 2016), which makes conservation and enhancement of current native forest mixtures  
22  
23  
359 important.  
24  
25  
360 One way of furthering resilience and adaptability of native forest diversity is to improve connectivity and  
27  
28  
361 migration corridors of key species and forest structures to sustain the availability of seed sources, as well as  
30  
31  
362 genetic variation (Yang et al., 2015). Such adaptation and response to climate induced change embraces the  
32  
33  
363 multifunctionality of forests, which must be reflected in management approaches and cultural integration  
34  
35  
364 (Halofsky et al., 2018). In Finland, forest owners show a willingness to participate in PES schemes aimed at  
37  
38  
365 reducing wider-landscape risks of pests and diseases, as long as it provides some management flexibility  
39  
40  
366 and does not unnecessarily prohibit forest operations that produce revenue and improved end crop yields  
42  
43  
367 (Sheremet et al., 2018). This approach demonstrates a step towards customising PES to fit individuals and  
44  
45  
368 regional differences through a common goal that binds forest owners over large landscapes (Curtin, 2014).  
46  
47  
369 A similar tailored PES model could be used for CSF guided by indicators and local priorities that taps into  
49  
50  
370 important issues for managers.  
51  
52  
371 CSF mitigation focusses upon carbon sequestration and storage, timber product use, bioenergy growth and  
54  
55  
372 use, and the interactions between lifecycles to optimise carbon neutral activity. Therefore, reconciling and  
56  
57  
373 aligning these sometimes-competing activities will be a crucial step. However, the focus on tangible metrics  
59  
60  
374 currently dominating the CSF (Nabuurs et al., 2018, 2017) could be a limiting factor in developing future CSF

375 strategies. Carbon sequestration is a now widely accepted substantive mitigation mechanism, as trade-offs  
1  
376 in decision-making and resultant management activities are measurable, such as yield increment, revenue  
3  
4  
377 variability and the use of various silvicultural approaches. Greater recognition of forest planning and  
5  
6  
378 diverse forestry approaches in mitigation strategies should be integrated into CSF. In fact, these actions  
8  
379 underpin carbon sequestration and storage, as well as adaptation issues, which will cumulatively impact the  
10  
11  
380 rest of the supply chain and forest-dependent communities who are supported economically and socially  
12  
13  
381 by a range of ecosystem services (Brang et al., 2014; Colloff et al., 2016; Nagel et al., 2017).  
15  
16

382 Ecosystem services assessment has been identified as a tool that could link stakeholders with management  
18  
19  
383 practices and CSF governance, a visible action and beneficial outcome that strongly relates to social  
20  
21  
384 dimensions of CSF. Therefore, PES, conservation partnerships and bridging organisations with local  
22  
23  
385 stakeholder expertise could play an important capacity building role (Cockburn et al., 2016; Lange et al.,  
25  
26  
386 2016; Rouillard et al., 2015). Creating and maintaining social capital, such as knowledge exchange,  
27  
28  
387 upskilling, technological development and use, as well as better monitoring and reporting to aid adaptive  
30  
31  
388 learning (Curtin, 2014; Lawrence, 2017) could be key components of assessing the social dimensions of CSF.  
32  
33  
389 The CSF definition and subsequent development of applicable C&I guidance aim to support livelihoods of  
34  
35  
390 sustainable communities based on production, conservation and well-being.  
37  
38

391 Key features to creating an implementation pathway for CSF should include management and connectivity  
40  
41  
392 of large datasets for long-term monitoring to track and understand the biophysical processes and changing  
42  
43  
393 trends. These features will be supported by openly accessible data, guidance on use, and landscape scale  
45  
46  
394 data that can distil into customisable tools for managers, as well as into coherent points of collaboration  
47  
48  
395 and transparency between policy and practice (Nabuurs et al., 2019).  
49  
50

### 396 **4.3. Perspectives** 52

397 European policy should aim to be comprehensive and flexible enough to include broad actions and  
53  
54  
55  
398 strategies with locally tailored solutions for regions and individual countries. A survey by Sousa-Silva et al.  
56  
57  
399 (2018) explored forest managers views on climate change over seven European countries. These managers  
58  
59  
400 recognised the potential impact of climate change but had little awareness of how to respond to threats or  
60  
61

401 implement adaptive management measures. This is supported by Coll et al. (2018) who's work identified  
1  
402 forest managers lack of knowledge regarding adaptability and trade-offs to environmental change in mixed-  
3  
403 species forests. A CSF toolkit that uses the definition and indicators identified through the analysis  
4  
5  
6  
404 presented here could be used as a base to address the knowledge/information gap between science and  
8  
405 practice.  
10

11  
12  
406 Developing CSF from the bottom-up using experience of forest managers will be central to shaping best  
13  
14  
407 practice from current adaptive management implementation and applied learning. On the international  
16  
408 level, forestry is increasingly viewed as one of the most effective ways to mitigate climate change (Bastin et  
18  
19  
409 al., 2019; Chazdon and Brancalion, 2019; FAO, 2010b), however, the discussion between forest accounting  
20  
21  
410 being based upon past management activities and prospective management is ongoing (Grassi et al., 2018).

23  
24  
411 Establishing different management approaches that deviate from the status quo has been challenging in  
25  
26  
412 regard to PES, as integrating clear and trusted verification processes and documenting an agreed baseline  
27  
28  
413 from where alternative management diverges is difficult to implement on a wide-scale (Kang et al., 2019).

30  
31  
414 In this context an appropriate mix of traditional and novel indicators of climate change impacts on  
32  
33  
415 European forests have been proposed to improve the prediction of stand dynamics and forest productivity  
34  
35  
416 (Bussotti and Pollastrini, 2017). Under the Paris Agreement a flexible approach toward baseline accounting  
37  
38  
417 of past management that supports changes that increase production, forest health and forest community  
39  
40  
418 resilience, would reflect a middle ground or best of both world's scenarios.  
42

43  
44  
419 Mismatches in communication and failed initiatives often originate from narrow scope and a lack of vital  
45  
46  
420 information. Therefore, combining social sciences with more traditional areas of forest science could help  
47  
48  
421 avoid issues that perpetuate problems (Duckett et al., 2016). Recognising and integrating novel factors such  
49  
50  
422 as social territories and scales, based upon core social-ecological systems, could identify key intervention  
52  
53  
423 points that help enhance our ability to manage forests and avoid future spatial, temporal and political  
54  
55  
424 mismatches (Fischer, 2018). In Melnykovich et al. (2018) community members showed interest in  
56  
57  
425 designing and implementing the sustainable forest policy measures that managed the provision of  
59  
60  
426 ecosystem services and enhanced well-being. However, these aspirations were tempered by the need to  
61

427 prioritise income generation and daily management actions. This underlines the need to develop resilience  
1  
428 with local community sustainable development goals, as well as balance long-term policy goals with short-  
3  
429 term needs of the locality (Bull et al., 2018). An approach that operates through co-design and production  
5  
430 may have a greater chance of creating CSF framework with bespoke set-ups for different regions,  
8  
431 ownership and forest types.

432 CSF definitions, guidance and tools generated by such projects as CLIMO could help refine the approach to  
13  
433 forestry as a more sophisticated mitigation measure encouraging socio-economic growth and innovation.

434 CSF should endeavour to take a systemic approach rather than treating individual symptoms, looking to the  
18  
435 long-term health and resilience of forest ecosystems, which includes people as a central component.

436 Ollikainen (2014) states that the EU Bioeconomy Action Plan fails to take account of the forestry sector's  
23  
437 link to climate change and superficially addresses the land use management dynamics that are vital for  
26  
438 developing climate-smart policy. Such criticism highlights crucial gaps in the connectivity of European  
28  
439 strategies (Forestry, Bioeconomy, and Climate etc.) that could potentially be addressed and informed by  
31  
440 further developed CSF definitions, guidelines and toolkits, which aim to enhance the science/policy  
33  
441 interface. However, Hodge et al. (2017) emphasises that bioeconomy already acts as a bridging concept,  
35  
442 bringing forestry and climate change closer together rather than dividing them into separate branches that  
38  
443 operate in isolation. Using an umbrella concept, such as bioeconomy, which can be inclusive and  
40  
444 comprehensive, can also embed broad-based approaches that overlook key implementation issues in  
42  
445 specific sectors.

446 Additionally, such concepts as bioeconomy are dominated by economic, resource use and commodity  
48  
447 concerns, which have the tendency to neglect social and ecological considerations (Karvonen et al., 2017),  
50  
448 such as high-level governance and local participation (Pülzl et al., 2014). These have been recognised as key  
52  
449 elements of the CSF approach and, therefore, should be highlighted as important considerations for  
55  
450 translating bioeconomy impacts into relatable regional versions. Bioeconomy could be an important  
57  
451 concept in halting the reduction of biodiversity and ecosystem services flows, which rely on balancing the

452 effective integration of social-ecological, cultural and economic dimensions alongside the management of  
1  
453 natural resources that are subject to uncertain changes (Marchetti et al., 2014). Therefore, CSF as defined  
3  
454 here, offers a legitimate and inclusive discourse linking energy, carbon, production, biodiversity  
5  
6  
455 conservation and resilient communities; supported by scientific evidence and metrics that can provide  
8  
456 implementation options and guide management goals. Market inclusion and transparency of climate  
10  
457 impact and risk to key environments from increasing climate-induced changes will be crucial to avoiding  
12  
458 economic recession, therefore markets need to take more responsibility and support PES and CSF or risk  
15  
459 further loss of vital ecosystem services (Griffin, 2020). To achieve such goals specific evidence and  
17  
460 management roadmaps are needed to give policy the necessary teeth to affect wide-ranging change that  
20  
461 will reach and work with individual forest managers.

#### 462 **4.4. Limitations and future work**

463 When approaching any type of definition or guidance for a wide range of stakeholders over large  
28  
464 geographical areas there will be limitations, as well as scope for improvement and further development.  
31  
465 Strengthening our definition and set of indicators for CSF will be the next steps, as the participatory process  
33  
466 was represented by a group of mostly forest research professionals, although these were geographically  
35  
467 dispersed with a range of expertise, which substantiates the broad European approach. Including forestry  
38  
468 industry professionals and practitioners in the refinement of the definition and indicators is the next step  
40  
469 and currently forest managers and professionals in 20 countries of the CLIMO network are responding to  
42  
470 the CSF definition and indicators to inform the next iteration and potential steps for tailoring at regional  
45  
471 levels.

472 Aligning the CSF indicators with available data from European forest monitoring networks and National  
50  
473 Forest Inventories data will allow for creation of baselines and historical analysis of trends toward climate  
53  
474 smartness and the corresponding management actions. Lorente et al. (2018) created a web-based platform  
55  
475 to display and communicate a set forest indicators for climate change. However, they acknowledged  
57  
476 further refinement at the regional level and greater consideration of socio-economic factors, as well as

477 partnership and engagement with practitioners would enhance the impact and utility of the indicators.  
1  
478 Indicators for the transition from a federal to a hybrid social forestry governance structure in the USA were  
3  
479 developed focussing upon partnership, collaboration and institutional innovations (Abrams et al., 2019).  
5  
6  
480 This work provides the opportunity to map changes in governance patterns of forest units over spatial and  
8  
481 temporal scales but also allows for analysis of capacity building and institutional infrastructure required to  
10  
482 support these transitions. However, the authors also recognised the importance of understanding how  
12  
483 changes in governance influenced and impacted ecological change to enhance the stewardship function  
15  
484 increasing resilience and ecological integrity (Cannon et al., 2018). Such steps will be key for CSF to  
17  
485 establish a sound platform of communication and dissemination between practice and policy, shaping clear  
20  
486 linkages that can be understood and interpreted by both managers and policy-makers.  
22  
23  
487 Finer detail of data collection to recognise the importance of local management systems that work within  
25  
488 SFM indicators has been highlighted in northern Italy by demonstrating the complex socio-economic  
27  
489 dynamics of coppice management systems that support a historic and wider economy (Riccioli et al., 2019).  
30  
490 Despite the peripheral location of cultural indicators in this paper the definition reconciles the importance  
32  
491 of social dimensions including socio-economics in supporting and promoting CSF. Further work around  
35  
492 cultural and socio-economic indicators will be required perhaps as a linking sub-set of indicators that  
37  
493 facilitates the key channel of communication between managers and policy-makers.  
39  
40

## 494 **5. Conclusion**

495 Promoting CSF will require more time to build effective working relationships between stakeholders and  
45  
496 policymakers cultivating the trust needed to realise these strategies, recommendations and best practice  
47  
497 guidance (Lange et al., 2016). Improved communication and use of expansive participatory methods to  
50  
498 engage and interact with forest managers will play a key role in this process and naturalising what may be  
52  
499 seen by many, as "*another recycled concept*" into practice and common vocabulary of the practitioner.  
54  
500 Advancing CSF processes, beginning with baseline definitions, testing of indicators through key sites,  
57  
501 monitoring and experimental data will be influential in developing and progressing climate change policy,  
59  
502 such as the Paris Agreement. This paper offers a potential roadmap and process to implementing CSF  
61

503 starting from the definition and indicators with the aim of expanding our understanding of management  
1  
504 and decision-making challenges that can be ultimately refined into tool that delivers on SFM goals. This  
3  
4  
505 paper is an important first step in offering a CSF definition and indicators that can be used as a template on  
5  
6  
506 a European level and adapted to localities to provide much needed guidance for managing more resilient  
8  
507 forests and practicing CSF. However, engaging forest managers and wider stakeholders of forest  
10  
11  
508 communities will be central to developing, testing and refining further steps that aim to produce a valuable  
12  
13  
509 resource for forest practitioners to enhance the management of their forests in response to future  
15  
16  
510 uncertainty and growing demands on these forests.  
17  
18

## 511 **Acknowledgements**

21  
22

23 This study generated from the COST (European Cooperation in Science and Technology) Action CLIMO  
24  
25 (Climate-Smart Forestry in Mountain Regions - CA15226) financially supported by the EU Framework  
26  
27 Programme for Research and Innovation HORIZON 2020. We are grateful to all people who took part at  
28  
29  
30 meetings providing helpful contribution in the CSF definition process and indicator selection.  
31  
32

33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

517 **References**

1  
2  
3  
518 Abrams, J., Huber-Stearns, H., Gosnell, H., Santo, A., Duffey, S., Moseley, C., 2019. Tracking a Governance  
4  
5  
519 Transition: Identifying and Measuring Indicators of Social Forestry on the Willamette National Forest.  
6  
7  
520 Society and Natural Resources 1–20. doi:10.1080/08941920.2019.1605434  
8  
9  
10  
11  
521 Andersson, E., Keskitalo, E.C.H., 2018. Adaptation to climate change? Why business-as-usual remains the  
12  
13  
522 logical choice in Swedish forestry. *Global Environmental Change* 48, 76–85.  
14  
15  
523 doi:10.1016/J.GLOENVCHA.2017.11.004  
16  
17  
18  
1924 Armatas, C.A., Venn, T.J., McBride, B.B., Watson, A.E., Carver, S.J., 2016. Opportunities to utilize traditional  
20  
21  
525 phenological knowledge to support adaptive management of social-ecological systems vulnerable to  
22  
23  
526 changes in climate and fire regimes. *Ecology and Society* 21, art16. doi:10.5751/ES-07905-210116  
24  
25  
26  
527 Bastin, J.-F., Finegold, Y., Garcia, C., Mollicone, D., Rezende, M., Routh, D., Zohner, C.M., Crowther, T.W.,  
28  
29  
528 2019. The global Tree Restoration potential. *Science* 365, 76–79. doi:10.1126/science.aax0848  
30  
31  
32  
529 Biber, P., Borges, J., Moshhammer, R., Barreiro, S., Botequim, B., Brodrechtová, Y., Brukas, V., Chirici, G.,  
33  
34  
530 Cordero-Debets, R., Corrigan, E., Eriksson, L., Favero, M., Galev, E., Garcia-Gonzalo, J., Hengeveld, G.,  
35  
36  
531 Kavaliauskas, M., Marchetti, M., Marques, S., Mozgeris, G., Navrátil, R., Nieuwenhuis, M., Orazio, C.,  
37  
38  
532 Paligorov, I., Pettenella, D., Sedmák, R., Smreček, R., Stanislovaitis, A., Tomé, M., Trubins, R., Tuček, J.,  
39  
40  
533 Vizzarri, M., Wallin, I., Pretzsch, H., Sallnäs, O., 2015. How Sensitive Are Ecosystem Services in  
41  
42  
534 European Forest Landscapes to Silvicultural Treatment? *Forests* 6, 1666–1695. doi:10.3390/f6051666  
43  
44  
45  
46  
47  
535 Borgatti, S.P., Everett, M.G., Freeman, L.C., 2002. *Ucinet 6 for windows: Software for social network*  
48  
49  
536 *analysis*. Harvard, MA: Analytic Technologies.  
50  
51  
52  
537 Bowditch, E.A.D., McMorran, R., Bryce, R., Smith, M., 2019. Perception and partnership: Developing forest  
53  
54  
538 resilience on private estates. *Forest Policy and Economics* 99. doi:10.1016/j.forpol.2017.12.004  
55  
56  
57  
539 Brang, P., Spathelf, P., Larsen, J.B., Bauhus, J., Boncina, A., Chauvin, C., Drossler, L., Garcia-Guemes, C.,  
58  
59  
60  
540 Heiri, C., Kerr, G., Lexer, M.J., Mason, B., Mohren, F., Muhlethaler, U., Nocentini, S., Svoboda, M.,  
61  
62  
63  
64  
65

541 2014. Suitability of close-to-nature silviculture for adapting temperate European forests to climate  
1 change. *Forestry* 87, 492–503. doi:10.1093/forestry/cpu018  
542  
3  
4  
543 Brunner, S.H., Grêt-Regamey, A., 2016. Policy strategies to foster the resilience of mountain social-  
6 ecological systems under uncertain global change. *Environmental Science & Policy* 66, 129–139.  
7  
544 doi:10.1016/J.ENVSCI.2016.09.003  
8  
9  
545  
11  
12  
546 Bull, G.Q., Boedhihartono, A.K., Bueno, G., Cashore, B., Elliott, C., Langston, J D, Riggs, R.A., Sayer, J.,  
14  
15  
547 Langston, James D, 2018. Global forest discourses must connect with local forest realities,  
16  
17  
548 *International Forestry Review*.  
19  
20  
549 Bussotti, F., Pollastrini, M., 2017. Traditional and Novel Indicators of Climate Change Impacts on European  
22  
23  
550 *Forest Trees*. *Forests* 8. doi:10.3390/f8040137  
24  
25  
551 Cannon, J.B., Barrett, K.J., Gannon, B.M., Addington, R.N., Battaglia, M.A., Fornwalt, P.J., Aplet, G.H., Cheng,  
27  
28  
552 A.S., Underhill, J.L., Briggs, J.S., Brown, P.M., 2018. Collaborative restoration effects on forest  
29  
30  
553 structure in ponderosa pine-dominated forests of Colorado. *Forest Ecology and Management* 424,  
32  
33  
554 191–204. doi:10.1016/j.foreco.2018.04.026  
34  
35  
555 Castañeda, F., 2000. Criteria and indicators for sustainable forest management: International processes,  
37  
38  
556 current status and the way ahead. *Unasylva* 51, 34–40.  
40  
41  
557 Chazdon, R., Brancalion, P., 2019. Restoring forests as a means to many ends. *Science*.  
43  
44  
558 doi:10.1126/science.aax9539  
45  
46  
559 Cockburn, J., Rouget, M., Slotow, R., Roberts, D., Boon, R., Douwes, E., O'Connell, S., Downs, C.T.,  
48  
49  
560 Mukherjee, S., Musakwa, W., Mutanga, O., Mwabvu, T., Odindi, J., Odindo, A., Proche, J., Urban,  
51  
52  
561 Ramdhani, S., Ray-Mukherjee, J., Sershen, Schoeman, M.C., Smit, A.J., Wale, E., Willows-Munro, S.,  
53  
54  
562 2016. How to build science-action partnerships for local land-use planning and management: Lessons  
55  
56  
563 from Durban, South Africa. *Ecology and Society* 21. doi:10.5751/ES-08109-210128  
58  
59  
564 Coll, L., Ameztegui, A., Collet, C., Löf, M., Mason, B., Pach, M., Verheyen, K., Abrudan, I., Barbati, A.,  
61  
62  
63  
64  
65

565 Barreiro, S., Bielak, K., Bravo-Oviedo, A., Ferrari, B., Govedar, Z., Kulhavy, J., Lazdina, D., Metslaid, M.,  
1  
566 Mohren, F., Pereira, M., Peric, S., Rasztoivits, E., Short, I., Spathelf, P., Sterba, H., Stojanovic, D., Valsta,  
3  
4  
567 L., Zlatanov, T., Ponette, Q., 2018. Knowledge gaps about mixed forests: What do European forest  
5  
6  
568 managers want to know and what answers can science provide? *Forest Ecology and Management*  
8  
9  
569 407, 106–115. doi:10.1016/J.FORECO.2017.10.055  
10  
11  
12  
570 Colloff, M.J., Doherty, M.D., Lavorel, S., Dunlop, M., Wise, R.M., Prober, S.M., 2016. Adaptation services and  
13  
14  
571 pathways for the management of temperate montane forests under transformational climate change.  
15  
16  
572 *Climatic Change* 138, 267–282. doi:10.1007/s10584-016-1724-z  
18  
19  
20  
573 Curtin, C.G., 2014. Resilience design: toward a synthesis of cognition, learning, and collaboration for  
21  
22  
574 adaptive problem solving in conservation and natural resource stewardship. *Ecology and Society* 19,  
23  
24  
575 15.  
26  
27  
28  
576 Del Río, M., Pretzsch, H., Alberdi, I., Bielak, K., Bravo, F., Brunner, A., Condés, S., Ducey, M.J., Fonseca, T.,  
29  
30  
577 von Lüpke, N., Pach, M., Peric, S., Perot, T., Souidi, Z., Spathelf, P., Sterba, H., Tijardovic, M., Tomé, M.,  
31  
32  
578 Vallet, P., Bravo-Oviedo, A., 2016. Characterization of the structure, dynamics, and productivity of  
33  
34  
579 mixed-species stands: review and perspectives. *European Journal of Forest Research* 135, 23–49.  
35  
36  
580 doi:10.1007/s10342-015-0927-6  
37  
38  
39  
40  
581 Del Río, M., Schütze, G., Pretzsch, H., 2014. Temporal variation of competition and facilitation in mixed  
41  
42  
582 species forests in Central Europe. *Plant Biology* 16, 166–176. doi:10.1111/plb.12029  
43  
44  
45  
46  
583 Denton, F., Wilbanks, T.J., Abeyasinghe, A.C., Burton, I., Gao, Q., Lemos, M.C., Masui, T., O’Brien, K.L.,  
47  
48  
584 Warner, K., 2014. Climate-resilient pathways: Adaptation, mitigation, and sustainable development.  
49  
50  
585 *Climate Change* 1101–1131.  
51  
52  
53  
586 Dietz, T., 2013. Bringing values and deliberation to science communication. *Proceedings of the National*  
54  
55  
587 *Academy of Sciences of the United States of America* 110, 14081–7. doi:10.1073/pnas.1212740110  
56  
57  
58  
59  
588 Duckett, D., Feliciano, D., Martin-Ortega, J., Munoz-Rojas, J., 2016. Tackling wicked environmental  
59  
60  
61  
62  
63  
64  
65

589 problems: The discourse and its influence on praxis in Scotland. *Landscape and Urban Planning* 154,  
1 44–56. doi:10.1016/j.landurbplan.2016.03.015  
2  
3  
4  
591 FAO, 2010a. “Climate-Smart” Agriculture Policies, Practices and Financing for Food Security, Adaptation and  
6 Mitigation. Rome.  
7  
8  
9  
10  
11 593 FAO, 2010b. Working with countries to tackle climate change through sustainable forest management  
12 Managing forests for climate change. Rome.  
13  
14  
15  
16  
17 595 Fernandes, P.M., 2013. Fire-smart management of forest landscapes in the Mediterranean basin under  
18 global change. *Landscape and Urban Planning* 110, 175–182. doi:10.1016/j.landurbplan.2012.10.014  
19  
20  
21  
22 597 Fischer, A.P., 2018. Forest landscapes as social-ecological systems and implications for management.  
23 *Landscape and Urban Planning* 177, 138–147. doi:10.1016/j.landurbplan.2018.05.001  
24  
25  
26  
27 599 Folke, C., Biggs, R., Norström, A. V., Reyers, B., Rockström, J., 2016. Social-ecological resilience and  
28 biosphere-based sustainability science. *Ecology and Society* 21. doi:10.5751/ES-08748-210341  
29  
30  
31  
32 601 Folke, C., Carpenter, S.R., Walker, B., Scheffer, M., Chapin, T., Rockström, J., 2010. Resilience Thinking:  
33 Integrating Resilience, Adaptability and Transformability. *Ecology and Society* 15, 20–29.  
34  
35  
36  
37  
38 603 Frame, B., Lawrence, J., Ausseil, A.-G., Reisinger, A., Daigneault, A., 2018. Adapting global shared socio-  
39 economic pathways for national and local scenarios. *Climate Risk Management* 21, 39–51.  
40  
41  
42  
43 605 doi:10.1016/j.crm.2018.05.001  
44  
45  
46 606 Grassi, G., Pilli, R., House, J., Federici, S., Kurz, W.A., 2018. Science-based approach for credible accounting  
47 of mitigation in managed forests. *Carbon Balance and Management* 13. doi:10.1186/s13021-018-  
48  
49  
50  
51 608 0096-2  
52  
53  
54 609 Griffin, P., 2020. Energy finance must account for extreme weather risk. *Nature Energy* 1–3.  
55  
56  
57  
58  
59 610 Haines-Young, R., Potschin, M., 2012. Common International Classification of Ecosystem Services (CICES):  
60 Consultation on Version 4. Copenhagen, Denmark.  
61  
62  
63  
64  
65

- 612 Halofsky, J.E., Peterson, D.L., Prendeville, H.R., 2018. Assessing vulnerabilities and adapting to climate  
1  
613 change in northwestern U.S. forests. *Climatic Change* 146, 89–102. doi:10.1007/s10584-017-1972-6  
2  
3  
4  
5  
614 Hansen, L., Hoffman, J., Drews, C., Mielbrecht, E., 2010. Designing climate-smart conservation: Guidance  
6  
7  
8  
615 and case studies: Special section. *Conservation Biology* 24, 63–69. doi:10.1111/j.1523-  
9  
10  
11  
12  
13  
14  
617 Harmon, M.E., Campbell, J.L., 2017. Managing carbon in the forest sector, in: *People, Forests, and Change:  
15  
16  
17  
618 Lessons from the Pacific Northwest*. Island Press-Center for Resource Economics, pp. 161–173.  
18  
19  
20  
21  
619 doi:10.5822/978-1-61091-768-1\_12  
22  
23  
24  
25  
26  
27  
28  
29  
620 Hodge, D., Brukas, V., Giurca, A., 2017. Forests in a bioeconomy: bridge, boundary or divide? *Scandinavian  
30  
31  
32  
621 Journal of Forest Research* 32, 582–587. doi:10.1080/02827581.2017.1315833  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
622 IPCC, 2014. Summary for Policymakers, *Climate Change 2014: Synthesis Report*. Contribution of Working  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65  
623 Groups I, II and III to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change.  
624 doi:10.1017/CBO9781107415324  
625  
626  
627  
628  
629  
630  
631  
632  
633  
634  
635
- Jandl, R., Ledermann, T., Kindermann, G., Freudenschuss, A., Gschwantner, T., Weiss, P., 2018. Strategies  
for climate-smart forest management in Austria. *Forests* 9, 1–15. doi:10.3390/f9100592
- Jantke, K., Müller, J., Trapp, N., Blanz, B., 2016. Is climate-smart conservation feasible in Europe? Spatial  
relations of protected areas, soil carbon, and land values. *Environmental Science & Policy* 57, 40–49.  
doi:10.1016/j.envsci.2015.11.013
- Ji, Y., Guo, K., Fang, S., Xu, X., Wang, Z., Wang, S., 2017. Long-term growth of temperate broadleaved  
forests no longer benefits soil C accumulation. *Scientific Reports* 7, 42328. doi:10.1038/srep42328
- Kang, M.J., Siry, J.P., Colson, G., Ferreira, S., 2019. Do forest property characteristics reveal landowners'  
willingness to accept payments for ecosystem services contracts in southeast Georgia, U.S.? *Ecological  
Economics* 161, 144–152. doi:10.1016/j.ecolecon.2019.02.016
- Karvonen, J., Halder, P., Kangas, J., Leskinen, P., 2017. Indicators and tools for assessing sustainability

636 impacts of the forest bioeconomy. *Forest Ecosystems* 4, 2. doi:10.1186/s40663-017-0089-8  
1  
2  
637 Kok, K., Bärlund, I., Flörke, M., Holman, I., Gramberger, M., Sendzimir, J., Stuch, B., Zellmer, K., 2014.  
4  
638 European participatory scenario development: strengthening the link between stories and models.  
6  
7  
639 *Climatic Change* 128. doi:10.1007/s10584-014-1143-y  
8  
9  
10  
640 Lange, A., Siebert, R., Barkmann, T., 2016. Incrementality and Regional Bridging: Instruments for Promoting  
11  
12  
641 Stakeholder Participation in Land Use Management in Northern Germany. *Society and Natural*  
14  
15  
642 *Resources* 29, 868–879. doi:10.1080/08941920.2015.1122135  
16  
17  
18  
643 Lawrence, A., 2017. Adapting through practice: Silviculture, innovation and forest governance for the age of  
19  
20  
644 extreme uncertainty. *Forest Policy and Economics* 79, 50–60. doi:10.1016/j.forpol.2016.07.011  
22  
23  
645 Lexer, M.J., Bugmann, H., 2017. Mountain forest management in a changing world. *European Journal of*  
25  
26  
646 *Forest Research* 136, 981–982. doi:10.1007/s10342-017-1082-z  
27  
28  
29  
647 Liang, J., Crowther, T.W., Picard, N., Wisser, S., Zhou, M., Alberti, G., Schulze, E.-D., McGuire, A.D., Bozzato,  
30  
31  
648 F., Pretzsch, H., De-Miguel, S., Paquette, A., Hérault, B., Scherer-Lorenzen, M., Barrett, C.B., Glick, H.B.,  
33  
34  
649 Hengeveld, G.M., Nabuurs, G.-J., Pfautsch, S., Viana, H., Vibrans, A.C., Ammer, C., Schall, P., Verbyla,  
35  
36  
650 D., Tchebakova, N., Fischer, M., Watson, J. V, Chen, H.Y.H., Lei, X., Schelhaas, M.-J., Lu, H., Gianelle, D.,  
37  
38  
651 Parfenova, E.I., Salas, C., Lee, E., Lee, B., Kim, H.S., Bruelheide, H., Coomes, D.A., Piotta, D.,  
40  
41  
652 Sunderland, T., Schmid, B., Gourlet-Fleury, S., Sonké, B., Tavani, R., Zhu, J., Brandl, S., Vayreda, J.,  
42  
43  
653 Kitahara, F., Searle, E.B., Neldner, V.J., Ngugi, M.R., Baraloto, C., Frizzera, L., Bałazy, R., Oleksyn, J.,  
44  
45  
654 Zawila-Niedźwiecki, T., Bouriaud, O., Bussotti, F., Finér, L., Jaroszewicz, B., Jucker, T., Valladares, F.,  
47  
48  
655 Jagodzinski, A.M., Peri, P.L., Gonmadje, C., Marthy, W., O’Brien, T., Martin, E.H., Marshall, A.R.,  
49  
50  
656 Rovero, F., Bitariho, R., Niklaus, P.A., Alvarez-Loayza, P., Chamuya, N., Valencia, R., Mortier, F., Wortel,  
51  
52  
657 V., Engone-Obiang, N.L., Ferreira, L. V, Odeke, D.E., Vasquez, R.M., Lewis, S.L., Reich, P.B., 2016.  
54  
55  
658 Positive biodiversity-productivity relationship predominant in global forests. *Science* 354.  
56  
57  
659 doi:10.1126/science.aaf8957  
58  
59  
60  
660 Lindner, M., Fitzgerald, J.B., Zimmermann, N.E., Reyer, C., Delzon, S., van der Maaten, E., Schelhaas, M.-J.,  
62  
63  
64  
65

- 661 Lasch, P., Eggers, J., van der Maaten-Theunissen, M., Suckow, F., Psomas, A., Poulter, B., Hanewinkel,  
1  
662 M., 2014. Climate change and European forests: What do we know, what are the uncertainties, and  
3  
4  
663 what are the implications for forest management? *Journal of Environmental Management* 146, 69–83.  
5  
6  
664 doi:10.1016/J.JENVMAN.2014.07.030  
8  
9
- 665 Linser, S., Wolfslehner, B., Asmar, F., Bridge, S.R.J., Gritten, D., Guadalupe, V., Jafari, M., Johnson, S., Laclau,  
11  
12  
666 P., Robertson, G., 2018. 25 years of criteria and indicators for sustainable forest management: Why  
13  
14  
667 some intergovernmental C & I processes flourished while others faded. *Forests* 9, 1–23.  
16  
668 doi:10.3390/f9090515  
18  
19
- 669 Locatelli, B., Lavorel, S., Sloan, S., Tappeiner, U., Geneletti, D., 2017. Characteristic trajectories of ecosystem  
21  
22  
670 services in mountains. *Frontiers in Ecology and the Environment* 15, 150–159. doi:10.1002/fee.1470  
23  
24
- 671 Lorente, M., Gauthier, S., Bernier, P., Ste-Marie, C., 2018. Tracking forest changes: Canadian Forest Service  
26  
27  
672 indicators of climate change. *Climatic Change* 1–15. doi:10.1007/s10584-018-2154-x  
29  
30
- 673 Marchetti, M., Vizzarri, M., Lasserre, B., Sallustio, L., Tavone, A., 2014. Natural capital and bioeconomy:  
32  
33  
674 challenges and opportunities for forestry. *Annals of Silvicultural Research* 38, 62–73.  
34  
35
- 675 Mavrommati, G., Borsuk, M.E., Howarth, R.B., 2017. A novel deliberative multicriteria evaluation approach  
37  
38  
676 to ecosystem service valuation. *Ecology and Society* 22. doi:10.2307/26270123  
40  
41
- 677 Melnykovich, M., Nijnik, M., Soloviy, I., Nijnik, A., Sarkki, S., Bihun, Y., 2018. Social-ecological innovation in  
43  
44  
678 remote mountain areas: Adaptive responses of forest-dependent communities to the challenges of a  
45  
46  
679 changing world. *Science of The Total Environment* 613–614, 894–906.  
48  
49  
680 doi:10.1016/J.SCITOTENV.2017.07.065  
50  
51
- 681 Mendoza, G.A., Prabhu, R., 2000. Development of a methodology for selecting criteria and indicators of  
53  
54  
682 sustainable forest management: A case study on participatory assessment. *Environmental*  
55  
56  
683 *Management* 26, 659–673. doi:10.1007/s002670010123  
58  
59
- 684 Mina, M., Bugmann, H., Cordonnier, T., Irauschek, F., Klopčič, M., Pardos, M., Cailleret, M., 2017. Future  
61  
62  
63  
64  
65

- 685 ecosystem services from European mountain forests under climate change. *Journal of Applied Ecology*  
686 54, 389–401. doi:10.1111/1365-2664.12772  
687  
688  
689 Nabuurs, G.-J., Delacote, P., Ellison, D., Hanewinkel, M., Hetemäki, L., Lindner, M., Ollikainen, M., 2017. By  
690 2050 the mitigation effects of EU forests could nearly double through climate smart forestry. *Forests*  
691 8, 1–14. doi:10.3390/f8120484  
692  
693 Nabuurs, G.-J., Verkerk, H., Schelhaas, M., Ramon, J., Trasobares, A., Cienfiala, E., 2018. Climate-Smart  
694 Forestry : quantification of mitigation impacts in three case regions in Europe Outline - Concept of  
695 Climate-Smart Forestry - Three cases regions in Europe. Brussels.  
696  
697 Nabuurs, G.-J., Verweij, P., Eupen, M. Van, Pérez-Soba, M., Helga, P., Hendriks, K., 2019. Brief  
698 Communication: Next-generation information to support a sustainable course for European forests.  
699 *Nature Sustainability* 2, 815–818.  
700  
701 Nagel, L.M., Palik, B.J., Battaglia, M.A., D’amato, A.W., Guldin, J.M., Swanston, C.W., Janowiak, M.K.,  
702 Powers, M.P., Joyce, L.A., Millar, C.I., Peterson, D.L., Ganio, L.M., Kirschbaum, C., Roske, M.R., 2017.  
703 Adaptive Silviculture for Climate Change: A National Experiment in Manager-Scientist Partnerships to  
704 Apply an Adaptation Framework. *Journal of Forestry* 115, 167–178. doi:10.5849/jof.16-039  
705  
706 Ollikainen, M., 2014. Forestry in bioeconomy – smart green growth for the humankind. *Scandinavian  
707 Journal of Forest Research* 29, 360–366. doi:10.1080/02827581.2014.926392  
708  
709 Paul, C., Brandl, S., Friedrich, S., Falk, W., Härtl, F., Knoke, T., De, C.P., 2019. Climate change and mixed  
710 forests: how do altered survival probabilities impact economically desirable species proportions of  
711 Norway spruce and European beech? *Annals of Forest Science* 76, 1–15. doi:10.1007/s13595-018-  
712 0793-8  
713  
714 Pellow, D.N., 1999. Negotiation and Confrontation: Environmental Policymaking Through Consensus.  
715 *Society & Natural Resources* 12, 189–203. doi:10.1080/089419299279696  
716  
717 Pretzsch, H., Rötzer, T., Forrester, D.I., 2017. Modelling Mixed-Species Forest Stands, in: *Mixed-Species*

709 Forests. Springer Berlin Heidelberg, Berlin, Heidelberg, pp. 383–431. doi:10.1007/978-3-662-54553-  
1  
~~7~~10 9\_8  
3  
4  
~~5~~11 Pülzl, H., Kleinschmit, D., Arts, B., 2014. Bioeconomy-an emerging meta-discourse affecting forest  
6  
7  
~~7~~12 discourses? Bioeconomy-an emerging meta-discourse affecting forest discourses? Scandinavian  
8  
9  
~~10~~13 Journal of Forest Research 29, 386–393. doi:10.1080/02827581.2014.920044  
11  
12  
~~13~~14 Pussinen, A., Nabuurs, G.J., Wieggers, H.J.J., Reinds, G.J., Wamelink, G.W.W., Kros, J., Mol-Dijkstra, J.P., de  
14  
15  
~~15~~15 Vries, W., 2009. Modelling long-term impacts of environmental change on mid- and high-latitude  
16  
17  
~~16~~16 European forests and options for adaptive forest management. Forest Ecology and Management 258,  
18  
19  
~~17~~17 1806–1813. doi:10.1016/j.foreco.2009.04.007  
20  
21  
22  
~~18~~18 Riccioli, F., Fratini, R., Marone, E., Fagarazzi, C., Calderisi, M., Brunialti, G., 2019. Indicators of sustainable  
24  
25  
~~19~~19 forest management to evaluate the socio-economic functions of coppice in Tuscany, Italy. Socio-  
26  
27  
~~20~~20 Economic Planning Sciences 100732. doi:10.1016/j.seps.2019.100732  
29  
30  
~~21~~21 Rockström, J., Gaffney, O., Rogelj, J., Meinshausen, M., Nakicenovic, N., Schellnhuber, H.J., 2017. A  
32  
33  
~~22~~22 roadmap for rapid decarbonization. Science 355, 1269–1271. doi:10.1126/science.aah3443  
34  
35  
~~23~~23 Röser, D., Sikanen, L., Asikainen, A., Parikka, H., Väätäinen, K., 2011. Productivity and cost of mechanized  
37  
38  
~~24~~24 energy wood harvesting in Northern Scotland. Biomass and Bioenergy 35, 4570–4580.  
40  
41  
~~25~~25 doi:10.1016/j.biombioe.2011.06.028  
42  
43  
~~26~~26 Rouillard, J.J., Ball, T., Heal, K. V., Reeves, A.D., 2015. Policy implementation of catchment-scale flood risk  
45  
46  
~~27~~27 management: Learning from Scotland and England. Environmental Science and Policy 50, 155–165.  
47  
48  
~~28~~28 doi:10.1016/j.envsci.2015.02.009  
50  
51  
~~29~~29 Runhaar, H.A.C., Uittenbroek, C.J., van Rijswijk, H.F.M.W., Mees, H.L.P., Driessen, P.P.J., Gilissen, H.K.,  
53  
54  
~~30~~30 2016. Prepared for climate change? A method for the ex-ante assessment of formal responsibilities  
55  
56  
~~31~~31 for climate adaptation in specific sectors. Regional Environmental Change 16, 1389–1400.  
58  
59  
~~32~~32 doi:10.1007/s10113-015-0866-2  
60  
61  
62  
63  
64  
65

- 733 Santopuoli, G., di Cristofaro, M., Kraus, D., Schuck, A., Lasserre, B., Marchetti, M., 2019. Biodiversity  
 1  
~~734~~ conservation and wood production in a Natura 2000 Mediterranean forest. A trade-off evaluation  
 3  
~~735~~ focused on the occurrence of microhabitats. *iForest - Biogeosciences and Forestry* 76–84.  
 5  
~~736~~ doi:10.3832/ifor2617-011  
 8  
 9
- ~~737~~ Santopuoli, G., Ferranti, F., Marchetti, M., 2016. Implementing Criteria and Indicators for Sustainable Forest  
 11  
~~738~~ Management in a Decentralized Setting: Italy as a Case Study. *Journal of Environmental Policy and*  
 13  
~~739~~ *Planning* 18. doi:10.1080/1523908X.2015.1065718  
 14  
 15  
 16  
 17
- ~~740~~ Santopuoli, G., Requardt, A., Marchetti, M., 2012. Application of indicators network analysis to support  
 18  
~~741~~ local forest management plan development: a case study in Molise, Italy. *iForest - Biogeosciences and*  
 21  
~~742~~ *Forestry* 5, 31–37. doi:10.3832/ifor0603-009  
 22  
 23  
 24  
 25
- ~~743~~ Schultz, L., Folke, C., Österblom, H., Olsson, P., 2015. Adaptive governance, ecosystem management, and  
 26  
~~744~~ natural capital. *Proceedings of the National Academy of Sciences* 112, 7369–7374.  
 27  
~~745~~ doi:10.1073/pnas.1406493112  
 29  
 30  
 31  
 32
- ~~746~~ Seidl, R., Spies, T.A., Peterson, D.L., Stephens, S.L., Hicke, J.A., 2016. Searching for resilience: Addressing the  
 33  
~~747~~ impacts of changing disturbance regimes on forest ecosystem services. *Journal of Applied Ecology* 53,  
 36  
~~748~~ 120–129. doi:10.1111/1365-2664.12511  
 37  
 38  
 39  
 40
- ~~749~~ Sheremet, O., Ruokamo, E., Juutinen, A., Svento, R., Hanley, N., 2018. Incentivising Participation and Spatial  
 42  
~~750~~ Coordination in Payment for Ecosystem Service Schemes: Forest Disease Control Programs in Finland.  
 43  
~~751~~ *Ecological Economics* 152, 260–272. doi:10.1016/j.ecolecon.2018.06.004  
 44  
 45  
 46  
 47  
 48
- ~~752~~ Smiraglia, D., Ceccarelli, T., Bajocco, S., Salvati, L., Perini, L., 2016. Linking trajectories of land change, land  
 50  
~~753~~ degradation processes and ecosystem services. *Environmental Research* 147, 590–600.  
 51  
~~754~~ doi:10.1016/j.envres.2015.11.030  
 52  
 53  
 54  
 55  
 56
- ~~755~~ Sousa-Silva, R., Verbist, B., Lomba, Â., Valent, P., Suškevičs, M., Picard, O., Hoogstra-Klein, M.A., Cosofret,  
 58  
~~756~~ V.C., Bouriaud, L., Ponette, Q., Verheyen, K., Muys, B., 2018. Adapting forest management to climate  
 59  
 60  
 61  
 62  
 63  
 64  
 65

- 757 change in Europe: Linking perceptions to adaptive responses. *Forest Policy and Economics* 90, 22–30.  
1  
258 doi:10.1016/j.forpol.2018.01.004  
3  
4  
759 Thornton, T.F., Comberti, C., 2017. Synergies and trade-offs between adaptation, mitigation and  
6  
760 development. *Climate Change* 140, 5–18. doi:10.1007/s10584-013-0884-3/Published  
8  
9  
10  
761 Vass, M.M., Elofsson, K., 2016. Is forest carbon sequestration at the expense of bioenergy and forest  
11  
12 products cost-efficient in EU climate policy to 2050? *Journal of Forest Economics* 24, 82–105.  
13  
14  
762 doi:10.1016/J.JFE.2016.04.002  
15  
16  
17  
763  
18  
764 Wijewardana, D., 2008. Criteria and indicators for sustainable forest management: The road travelled and  
19  
20 the way ahead. *Ecological Indicators* 8, 115–122. doi:10.1016/j.ecolind.2006.11.003  
21  
22  
23  
765  
24  
766 Wolf, S.A., Klein, J.A., 2007. Enter the working forest: Discourse analysis in the Northern Forest. *Geoforum*  
25  
26  
767 38, 985–998. doi:10.1016/j.geoforum.2007.03.009  
27  
28  
29  
768 Wolfslehner, B., Baycheva-Merger, T., 2016. Evaluating the implementation of the Pan-European Criteria  
30  
31 and indicators for sustainable forest management - A SWOT analysis. *Ecological Indicators* 60, 1192–  
32  
33 1199. doi:10.1016/j.ecolind.2015.09.009  
34  
35  
36  
37  
771 Wolfslehner, B., Vacik, H., 2011. Mapping indicator models: From intuitive problem structuring to  
38  
39 quantified decision-making in sustainable forest management. *Ecological Indicators* 11, 274–283.  
40  
41  
772 doi:10.1016/j.ecolind.2010.05.004  
42  
43  
44  
773  
45  
774 Wolfslehner, B., Vacik, H., Lexer, M.J., 2005. Application of the analytic network process in multi-criteria  
46  
47 analysis of sustainable forest management. *Forest Ecology and Management* 207, 157–170.  
48  
49  
775 doi:10.1016/j.foreco.2004.10.025  
50  
51  
52  
776  
53  
777 Yang, J., Pedlar, J.H., McKenney, D.W., Weersink, A., 2015. The development of universal response  
54  
55 functions to facilitate climate-smart regeneration of black spruce and white pine in Ontario, Canada.  
56  
57  
778  
58  
779 *Forest Ecology and Management* 339, 34–43. doi:10.1016/J.FORECO.2014.12.001  
59  
60  
780  
61  
62  
63  
64  
65

781 Realizing Mitigation Efficiency of European Commercial Forests by Climate Smart Forestry. Scientific  
1  
2  
3 Reports 8, 1–11. doi:10.1038/s41598-017-18778-w

4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

783 Yousefpour, R., Hanewinkel, M., Le Moguédec, G., 2010. Evaluating the suitability of management  
784 strategies of pure Norway spruce forests in the Black Forest area of Southwest Germany for  
785 adaptation to or mitigation of climate change. Environmental Management 45, 387–402.  
786 doi:10.1007/s00267-009-9409-2

787 Zilberman, D., Goetz, R., Garrido, A., Lipper, L., Mccarthy, N., Asfaw, S., Giacomo Branca, 2018. Climate  
788 Smart Agriculture. Building resilience to climate change. doi:10.1007/978-3-319-61194-5

790 **List of captions of figures and tables**

1  
2  
791  
4  
792  
6  
7  
793  
8  
9  
10  
794  
11  
12  
795  
14  
15  
796  
16  
17  
797  
18  
19  
798  
21  
22  
799  
24  
25  
800  
26  
27  
801  
29  
30  
802  
32  
33  
803  
34  
35  
804  
36  
37  
38  
805  
40  
806  
42  
43  
807  
44  
45  
46  
808  
47  
48  
809  
50  
51  
810  
52  
53  
811  
54  
55  
812  
57  
58  
813  
59  
60  
61  
62  
63  
64  
65

Figure 1 – Process, and interaction within the CLIMO project, for building consensus on definitions of CSF in mountain regions. The iterative deliberative process conducted in CLIMO with a confrontation stage that engaged all working groups to account for any issues, difference of opinions and new information.

Figure 2 – Network analysis of CSF indicators relevance to adaptation and mitigation. The map shows the suitable indicators for assessing climate-smart forestry in mountain ecosystems. This map represents the merged map considering both, adaptive and mitigation issues, according the point of view of CLIMO participants. Indicators are clustered in core and peripheral groups in order to display their relevant importance to provide information about the climate-smartness of forests.


Figure 3 – Schematic representation of primary and secondary, core and peripheral, groups of indicators useful to assess and to support the development of climate-smart forestry management guidelines (Climate eye diagram). Black circle are the main themes/threads that support CSF and their indicators

Table 1 – Set of Indicators for SFM selected from the Forest Europe C&I set (<https://foresteurope.org/sfm-criteria-indicators/>), considered suitable for assessing the provision of ecosystem services, according the view of CLIMO participants. New 4 indicators have been included during the CLIMO project meetings.


Table 2: Ecosystem Services selected from CICES database (CICES classification, version4, 2012; <https://cices.eu>) and considered useful to monitor and assess Climate-Smart Forestry, according the view of CLIMO participants.

Table 3 - Centrality values. Degree is the number of ties that link each indicator with forest ecosystem services; Betweenness represents the number of times one indicator in the network is “between” other indicators on the causal paths. Closeness calculates the farness and normalized closeness centrality and variants of each vertex and gives the overall network closeness centralization. K-core is a maximal group of actors, all of which are connected to some number (k) of other members of the entity. Group show the cluster identified in this study, based on the centrality values and position of nodes in the network.

814 Figure 1




816  
817  
818  
819 Figure 2



830 Figure 3

831

2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29



30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

**Table 1: Set of Indicators for SFM selected from the Forest Europe C&I set (<https://foresteurope.org/sfm-criteria-indicators/>), considered suitable for assessing the provision of ecosystem services, according the view of CLIMO participants. New 4 indicators have been included during the CLIMO project meetings. Nine indicators were not selected (3.3 No-wood goods; 3.4 Services; 6.1 Forest holding; 6.2 Contribution of forest sector to GDP; 6.3 Net revenue; 6.4 Expenditures for services; 6.5 Forest sector force; 6.6 Occupational safety and health; 6.11 Cultural and spiritual values).**

<b>Criteria</b>	<b>Indicator name</b>	<b>Label</b>	<b>Description (based on the MCPFE 2002 &amp; Madrid 2015)</b>
C1: Forest Resources and Global Carbon Cycles	1.1 Forest area	<b>Forest area</b>	Area of forests and other wooded land, classified by forest type and by availability for wood supply, and share of forest and other wooded land in total land area
	1.2 Growing stock	<b>Growing stock</b>	Growing stock on forest and other wooded land, classified by forest type and by availability for wood supply
	1.3 Age structure and/or diameter distribution	<b>Forest structure</b>	Age structure and/or diameter distribution of forest and other wooded land, classified by availability for wood supply
	1.4 Carbon stock	<b>Carbon stock</b>	Carbon stock and carbon stock changes in forest biomass, forest soils and in harvested wood products
C2: Forest Health and Vitality	2.1 Deposition of air pollutants	<b>Air pollutants</b>	Deposition and concentration of air pollutants on forest and other wooded land
	2.2 Soil condition	<b>Soil condition</b>	Chemical soil properties (pH, CEC, C/N, organic C, base saturation) on forest and other wooded land related to soil acidity and eutrophication, classified by main soil types
	2.3 Defoliation	<b>Defoliation</b>	Defoliation of one or more main tree species on forest and other wooded land in each of the defoliation classes
	2.4 Forest damage	<b>Forest damage</b>	Forest and other wooded land with damage, classified by primary damaging agent (abiotic, biotic and human induced)
C3: Forest Biological Diversity	3.1 Increment and felling	<b>Increment/felling</b>	Balance between net annual increment and annual felling of wood on forest available for wood supply
	3.2 Roundwood	<b>Roundwood</b>	Quantity and market value of roundwood
	3.5 Forests under management plans	<b>Management plan</b>	Proportion of forest and other wooded land under a management plan or equivalent
C4: Forest Biological Diversity	4.1 Tree species composition	<b>Diversity</b>	Area of forest and other wooded land, classified by number of tree species occurring
	4.2 Regeneration	<b>Regeneration</b>	Total forest area by stand origin and area of annual forest regeneration and expansion
	4.3 Naturalness	<b>Naturalness</b>	Area of forest and other wooded land by class of naturalness (“undisturbed by man”, by “semi-natural” or by “plantations”)
	4.4 Introduced tree species	<b>New species</b>	Area of forest and other wooded land dominated by introduced tree species

16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57 5  
58 6  
59 7  
60  
61  
62  
63  
64  
65

C4: Forest Biological Diversity	4.5 Deadwood	<b>Deadwood</b>	Volume of standing deadwood and of lying deadwood on forest and other wooded land
	4.6 Genetic resources	<b>Genetic resources</b>	Area managed for conservation and utilisation of forest tree genetic resources (in situ and ex situ genetic conservation) and area managed for seed production
	4.7 Landscape pattern	<b>Fragmentation</b>	Area of continuous forest and of patches of forest separated by non-forest lands
	4.8 Threatened forest species	<b>Threatened species</b>	Number of threatened forest species, classified according to IUCN Red List categories in relation to total number of forest species
	4.9 Protected forests	<b>Protected area</b>	Area of forest and other wooded land protected to conserve biodiversity, landscapes and specific natural elements, according to MCPFE categories
C5: Protective Function (Soil and water 5	5.1 Protective forests - soil, water and other ecosystem functions, and infrastructures	<b>Protective forests</b>	Area of forest and other wooded land designated to prevent soil erosion, preserve water resources, maintain other protective functions, protect infrastructure and managed natural resources against natural hazards
C6: Socioeconomic Functions	6.7 Wood consumption	<b>Wood consumption</b>	Consumption per person of wood and products derived from wood
	6.8 Trade in wood	<b>Trade in wood</b>	Imports and exports of wood and products derived from wood
	6.9 Energy from wood resources	<b>Wood energy</b>	Share of wood energy in total primary energy supply, classified by origin of wood
	6.10 Accessibility for recreation	<b>Accessibility</b>	The use of forests and other wooded land for recreation in terms of right of access, provision of facilities and intensity of use
New indicators added by CLIMO participants	Management system	<b>Forestry</b>	Forest area classified according the silvicultural system adopted: coppice system; even-aged system (clear cut or shelterwood), uneven-aged system (selection system)
	Slenderness coefficient	<b>Slenderness</b>	The ratio of tree total height to diameter outside bark at 1.3 m above ground level
	Vertical distribution of tree crowns	<b>Vertical crowns</b>	Distribution of tree crown in the vertical space. It can be measure in terms of layers (one, two, multiple), or in terms of ratio between tree height and crown length. Canopy space filling and can be expressed in measure of density of tree crowns, such as crown area, tree crown diameter. It can be also expressed in measure of density of trees, such as trees per hectare, basal area per hectare (in this case the horizontal distribution refers to the tree).
	Horizontal distribution of tree crowns	<b>Horizontal crowns</b>	

**Table 2: Ecosystem Services selected from CICES database (CICES classification, version4, 2012; <https://cices.eu>) and considered useful to monitor and assess Climate-Smart Forestry, according the view of CLIMO participants.**

Ecosystem service	Label	CICES Section	CICES Division
Recreation and mental and physical health	<b>Recreation</b>	<b>Cultural</b>	Intellectual and experientially
Tourism	<b>Tourism</b>		Symbolic
Aesthetic appreciation and inspiration for culture, art and design	<b>Aesthetic</b>		Symbolic
Spiritual experience and sense of place	<b>Spiritual</b>		Symbolic
Protection of cultural heritage	<b>Cultural heritage</b>		Symbolic
Primary biomass production	<b>PBP</b>	<b>Provisioning</b>	Materials
Food	<b>Food</b>		Nutrition
Timber, fuel, fibre	<b>Raw materials</b>		Materials
Fresh water	<b>Fresh water</b>		Nutrition
Pharmaceuticals and bio-chemicals	<b>Medicinal</b>		Materials
Genetic resources	<b>Genetic resources</b>		Materials
Production of atmospheric oxygen	<b>Photosynthesis</b>	<b>Regulating</b>	Regulations biotic environment
Soil formation and retention	<b>Soil formation</b>		Regulations physical environment
Nutrient cycling	<b>Nutrient cycling</b>		Regulations biotic environment
Water cycling	<b>Water cycling</b>		Flow regulation
Maintenance of genetic diversity	<b>Gene pool</b>		Regulations biotic environment
Habitats for species	<b>Refugia</b>		Regulations biotic environment
Purification of water and air	<b>Purification</b>		Wastes regulation
Carbon sequestration and storage	<b>Sequestration</b>		Regulations physical environment
Moderation of natural disturbances, e.g., flood alleviation	<b>Disturbances</b>		Flow regulation
Erosion prevention and maintenance of soil health	<b>Erosion</b>		Flow regulation
Pollination	<b>Pollination</b>		Regulations biotic environment
Biological control	<b>Biological control</b>		Regulations biotic environment

5 Table 3: Centrality values. Degree is the number of ties that link each indicator with forest ecosystem  
 1 6 services; Betweenness represents the number of times one indicator in the network is “between” other  
 2 7 indicators on the causal paths. Closeness calculates the farness and normalized closeness centrality and  
 3 8 variants of each vertex and gives the overall network closeness centralization. K-core is a maximal group of  
 4 9 actors, all of which are connected to some number (k) of other members of the entity. Group show the  
 5 10 cluster identified in this study, based on the centrality values and position of nodes in the network.  
 6 7

8	9	10	11	12	13	14
	<b>Selected indicator</b>	<b>Degree</b>	<b>Betweenness</b>	<b>Closeness</b>	<b>K-core</b>	<b>Group</b>
10	1.1 - Forest area	23	30.1	79	14	1 <sup>st</sup> -core group
11	2.1 - Deposition of air pollutants	23	30.1	79	14	1 <sup>st</sup> -core group
12	2.4 - Forest damage	23	30.1	79	14	1 <sup>st</sup> -core group
13	3.5 - Management plan	23	30.1	79	14	1 <sup>st</sup> -core group
14	4.1 - Diversity	23	30.1	79	14	1 <sup>st</sup> -core group
15	4.3 - Naturalness	23	30.1	79	14	1 <sup>st</sup> -core group
16	4.4 - New species	23	30.1	79	14	1 <sup>st</sup> -core group
17	4.9 - Protected forests	23	30.1	79	14	1 <sup>st</sup> -core group
18	(Climo) - Forestry	23	30.1	79	14	1 <sup>st</sup> -core group
19	4.7 - Fragmentation	21	25.2	83	14	1 <sup>st</sup> -core group
20	6.7 - Wood consumption	19	20.9	87	14	1 <sup>st</sup> -core group
21	6.9 - Wood energy	19	17.3	87	14	2 <sup>nd</sup> -core group
22	2.2 - Soil condition	18	15	89	14	2 <sup>nd</sup> -core group
23	2.3 - Defoliation	18	15	89	14	2 <sup>nd</sup> -core group
24	5.1 - Protective forests	18	15	89	14	2 <sup>nd</sup> -core group
25	(Climo) - Vertical crowns	18	18.3	89	14	1 <sup>st</sup> -peripheral group
26	(Climo) - Horizontal crowns	17	16.5	91	13	1 <sup>st</sup> -peripheral group
27	1.3 - Forest structure	15	10.1	95	14	1 <sup>st</sup> -peripheral group
28	4.2 - Regeneration	13	7.8	99	13	2 <sup>nd</sup> -core group
29	4.5 - Deadwood	11	5.2	103	11	1 <sup>st</sup> -peripheral group
30	4.6 - Genetic resources	11	5.5	103	11	2 <sup>nd</sup> -core group
31	1.4 - Carbon stock	10	3.9	105	10	1 <sup>st</sup> -peripheral group
32	3.1 - Increment/fellings	10	3.8	105	10	1 <sup>st</sup> -peripheral group
33	1.2 - Growing stock	9	3	107	9	1 <sup>st</sup> -peripheral group
34	4.8 - Threatened species	9	3.2	109	9	2 <sup>nd</sup> -core group
35	(Climo) - Slenderness	9	2.9	109	9	1 <sup>st</sup> -peripheral group
36	6.1 - Accessibility	6	1.4	121	6	2 <sup>nd</sup> -peripheral group
37	3.2 - Roundwood	4	0.5	119	4	1 <sup>st</sup> -peripheral group
38	6.8 - Trade in wood	1	0	125	1	1 <sup>st</sup> -peripheral group

47 11  
 48 12  
 49  
 50 13  
 51  
 52  
 53  
 54  
 55  
 56  
 57  
 58  
 59  
 60  
 61  
 62  
 63  
 64  
 65

14 Box 1 – *Climate-Smart Forestry in mountain regions definition*

1  
2  
3  
4 **Climate-smart forestry** is sustainable adaptive forest management and governance to protect  
5 and enhance the potential of forests to adapt to, and mitigate climate change. The aim is to  
6 sustain ecosystem integrity and functions and to ensure the continuous delivery of ecosystem  
7 goods and services, while minimising the impact of climate-induced changes on mountain forests  
8 on well-being and nature’s contribution to people.  
9

10  
11  
12  
13 **Adaptation** measures of forests that maintain or improve their ability to grow under current and  
14 projected climatic conditions and increase their resistance and resilience. The adaptive capacity  
15 to changes in climate and to the timing and size of climate-induced disturbances (e.g., fire,  
16 extreme storm events, pests and diseases) can be enhanced by promoting genetic, compositional,  
17 structural, and functional diversity at both stand and landscape scales. This includes facilitating  
18 natural regeneration and planting of native as well as non-native tree species, genetic variants  
19 and individuals that are considered to be adapted to future conditions. Increased connectivity  
20 assists the migration of forest species.  
21  
22  
23

24  
25  
26 **Mitigation** of climate change by forests is a combination of carbon sequestration by trees, carbon  
27 storage by forest ecosystems, especially soils, and forest derived products, such as structural  
28 timber, and by carbon substitution - directly by replacing fossil fuels with bioenergy and indirectly  
29 through use of wood to substitute for higher carbon footprint materials.  
30  
31

32  
33  
34 The **social dimension** of forestry holds many aspects, from the involvement of stakeholders from  
35 local communities, and their conflicts over land use or for the access to skills and technology, to  
36 global forest governance challenges. Climate change may jeopardize forest ecosystem functioning  
37 and brings social and economic consequences for people, which may modify priorities of  
38 ecosystem services at various scales. Assessment for ecosystem services could be a tool making  
39 this process more efficient with respect to indicators relevant for governance regime and actors  
40 involved.  
41  
42  
43

44  
45 In summary, **climate-smart forestry** should enable both forests and society to transform, adapt to  
46 and mitigate climate-induced changes.  
47  
48

19 **Supplementary material**

1  
2

30 **Appendix A: Synthesis of the “climate-smart forestry” literature review**

4  
5

6 **The road to CSF concept**

7  
8

9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

Publications involving the search parameters of ‘Climate-Smart Forestry’ in the Web of Science database, accounts for 106 publications since 2000 (Figure S1).




Figure S1: Number of publications from 2001-2019 that include ‘Climate-Smart Forestry’ as search terms in Web of Science. Various searches (Scopus, google scholar, Web of science) has yielded different results, ranging from 82 – 106 publications using the same search parameters. Date search performed: 03/03/2020.

Nitschke and Innes (2008) model the response of forest ecosystem structure to potential fire regimes and species under climate-induced changes, highlighting the need to integrate climate-smart strategies into forest management planning and application. Jantke et al. (2016) compare Natura 2000 sites with levels of soil carbon and agricultural land values, which show a correlation between low land values and high soil carbon content and argue that such criteria could be a template for climate-smart conservation strategies. However, this study restricts the interpretation solely to agricultural values and thus omits wider economic, environmental and social potential. Yang et al. (2015) focus on the potential impact of species ‘*adaptational lag*’ to changes in and maintaining the productive stock and genetic variety of black spruce

36 (*Picea mariana* (Mill.) BSP.); they emphasise that important mixed forest structures can be achieved  
1  
237 through targeted climate-smart seed movements that mitigate against slow or unresponsive adaptation  
3  
438 processes. Halofsky et al. (2018) strongly advocate climate-smart strategies that facilitate rapid practical  
5  
6  
739 implementation, in response to the increasing frequency of extreme events.  
8  
9  
1040 Acceptance and adoption of climate-smart strategies and responses are dependent on people that drive  
11  
1241 policy, implement management and communities that rely on forests services. The potential for both  
13  
14  
1542 disruption and enhancement of the socio-economic landscape is wide-ranging. However, transnational  
16  
1743 forest policy continually fails to connect to the local working realities, which are critical for developing  
18  
1944 coherent policies that can adapt to the diversity and complexity of local contexts (Bull et al., 2018). Winkel  
20  
21  
2245 and Sotirov (2016) highlight that forest policy suffers from symbolic rhetoric to serve sectoral interests  
23  
2446 (economic and institutional competition) but rarely achieves any substantive progress. Common pathways  
25  
26  
2747 to realising change focusses on socio-ecological systems by connection to local ecosystems, promoting  
28  
2948 dialogue between stakeholders, fostering social innovation and providing an institutional framework to  
30  
3149 facilitate new structures (Biggs et al., 2010).  
32  
33  
3450 Socio-ecological systems can be viewed as a way to release economic and social potential (Fischer, 2018),  
35  
36  
3751 however, effective indicators for social values and cultural ecosystem services are still undeveloped and  
38  
3952 difficult to assess (Maes et al., 2016). Social values can be used indicatively within specific spatial areas and  
40  
41  
4253 help establish main priorities but usually require further clarification and verification (Chan et al., 2012;  
43  
4454 Tenerelli et al., 2016). However, such approaches could provide a frame for effective social indicators on a  
45  
4655 landscape level. Much like ecosystem management, CSF is underpinned by the interactions between  
47  
48  
4956 various dynamic components aiming to interlink at multiple levels and facilitating transformation through  
50  
5157 more adaptive, integrated and collaborative approaches (Folke et al. 2010). As forest managers, policy-  
52  
5358 makers and the wider population are central to promoting and developing the CSF process through socio-  
54  
55  
5659 ecological systems, participatory decision-making and social-innovation are considered crucial pathways to  
57  
5860 achieving these aims (Biggs et al. 2010). Greater expansion of socio-ecological systems and CSF will  
59  
60  
6161 continue with the emergence of new knowledge and problems, therefore the scope of CSF should be  
62  
63  
64  
65

62 responsive, as well as planned to best adapt to specific needs when time arrives. Blatter et al. (2017)  
1  
263 acknowledge that forest managers require appropriate management strategies to produce diverse services  
3  
4  
54 that the public require. They weighted their indicators to represent management functions considered  
5  
6  
765 more important in a local or regional context, such as mountain forests, and recognised the protective  
8  
966 function as the highest priority. This approach provides a basic flexibility to take into account more area-  
10  
11  
1267 specific requirements and changing management objectives over diverse landscapes. However, this  
13  
1468 approach should be investigated further to refine and expand the flexibility.  
15  
16  
1769 In summary, these papers on CSF highlight the need for collaboration across boundaries and over large  
18  
1970 spatial scales to ensure the productivity and functionality of forests into the future (Anton et al., 2010;  
20  
21  
2271 Curtin, 2014; Johansson, 2016). Aligning climate-smart characteristics of forestry with management  
23  
2472 practices and communicating clear methods to managers remains an elusive step for integrating climate  
25  
2673 change management responses with current management and guidance. Addressing these gaps is integral  
27  
28  
2974 to this project, as well as recognising the current shortfall of work around social-ecological systems of CSF,  
30  
3175 which is a key feature in harmonising new management approaches and knowledge into current systems.  
32  
33

## 3476 REFERENCES

35  
36

- 3777 Biggs, R., Westley, F.R. & Carpenter, S., 2010. Navigating the back loop:Fostering social innovation and  
38  
39  
4078 transformation in ecosystem management. *Ecology and Society*, 15(2), p.part 9.  
41  
42  
4379 Blatter, C., Lemm, R., Thees, O., Lexer, M.J. & Hanewinkel, M., 2017. Management of ecosystem services in  
44  
4580 mountain forests: Review of indicators and value functions for model based multi-criteria decision analysis.  
46  
47  
4881 *Ecological Indicators*, 79.  
49  
50  
5182 Bull, G.Q., Boedihartono, A.K., Bueno, G., Cashore, B., Elliott, C., Langston, J D, Riggs, R.A., Sayer, J. &  
52  
5383 Langston, James D, 2018. Global forest discourses must connect with local forest realities. *International*  
54  
55  
5684 *Forestry Review*, 20(2):160-166.  
57  
58  
5985 Chan, K.M.A., Satterfield, T. & Goldstein, J., 2012. Rethinking ecosystem services to better address and  
60  
6186 navigate cultural values. *Ecological Economics*, 74, pp.8–18.  
62  
63  
64  
65

87 Folke, C., Carpenter, S.R., Walker, B., Scheffer, M., Chapin, T. & Rockström, J., 2010. Resilience Thinking:  
1  
288 Integrating Resilience, Adaptability and Transformability. *Ecology and Society*, 15(4), pp.20–29.  
3  
4  
589 Fischer, A.P., 2018. Forest landscapes as social-ecological systems and implications for management.  
6  
7  
90 *Landscape and Urban Planning*, 177, pp.138–147.  
8  
9  
10  
1191 Halofsky, J.E., Peterson, D.L. & Prendeville, H.R., 2018. Assessing vulnerabilities and adapting to climate  
12  
1392 change in northwestern U.S. forests. *Climatic Change*, 146(1–2), pp.89–102.  
14  
15  
1693 Jantke, K., Müller, J., Trapp, N. & Blanz, B., 2016. Is climate-smart conservation feasible in Europe? Spatial  
17  
1894 relations of protected areas, soil carbon, and land values. *Environmental Science & Policy*, 57, pp.40–49.  
19  
20  
2195 Maes, J., Liqueste, C., Teller, A., Erhard, M., Paracchini, M.L., Barredo, J.I., Grizzetti, B., Cardoso, A., Somma,  
22  
23  
2496 F., et al., 2016. An indicator framework for assessing ecosystem services in support of the EU Biodiversity  
25  
2697 Strategy to 2020. *Ecosystem Services*, 17, pp.14–23. Available at:  
27  
2898 <http://www.sciencedirect.com/science/article/pii/S2212041615300504>.  
29  
30  
31  
3299 Nitschke, C.R. & Innes, J.L., 2008. Integrating climate change into forest management in South-Central  
33  
34100 British Columbia: An assessment of landscape vulnerability and development of a climate-smart  
35  
36101 framework. *Forest Ecology and Management*, 256(3), pp.313–327  
37  
38  
39  
40102 Tenerelli, P., Demšar, U. & Luque, S., 2016. Crowdsourcing indicators for cultural ecosystem services: A  
41  
42103 geographically weighted approach for mountain landscapes. *Ecological Indicators*, 64, pp.237–248.  
43  
44  
45104 Winkel, G. & Sotirov, M., 2016. Whose integration is this? European forest policy between the gospel of  
46  
47105 coordination, institutional competition, and a new spirit of integration. *Environment and Planning C:*  
48  
49106 *Government and Policy*, 34, pp.496–514.  
50  
51  
52  
53107 Yang, J., Pedlar, J.H., McKenney, D.W. & Weersink, A., 2015. The development of universal response  
54  
55108 functions to facilitate climate-smart regeneration of black spruce and white pine in Ontario, Canada. *Forest*  
56  
57109 *Ecology and Management*, 339, pp.34–43.  
58  
59  
60  
61110

111 **Appendix B**

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

Table S1: List of CLIMO meetings related to development of definition and indicators

Meeting	Date	Action	Comments
<b>Trento, Italy</b>	February, 2017	Identification of a sub-group to work on definition of Climate-Smart Forestry. First discussion of three components, comprising mitigation, adaptation and social dimensions.	Membership of definitions sub-group commenced with those expressing a strong interest in participation, but continued with filling of gaps in the composition. For example, ensuring that experts representing other Working Groups (not just Working Group 1) were included.
<b>Sofia, Bulgaria</b>	September, 2017	Three-day workshop to establish an initial definition of Climate-Smart Forestry. Sub-group included Chair, Vice Chair and Project Manager, Working Group 1 Leader and members of the other Working Groups.	A one-page initial draft definition was prepared. Contribution at this stage led to co-authorship of definition.
<b>Virtual (email)</b>	October, 2017 - February, 2018	One-page initial draft definition shared for comment and amendment with entire COST Action.	Contributions at this stage resulted in amendments to the draft and inclusion in co-authorship.
<b>Sofia, Bulgaria</b>	February, 2018	Working Group 1 sub-group refinement of draft and presentation to Open Meeting for further/final criticism and amendment.	A contribution at this stage that resulted in amendment to the draft brought inclusion in co-authorship.

115 Table S2: List of participants which attended to the meetings. In bold the co-authors of the manuscript.

Surname	Name	Country	Background	Trento Feb-17	Sofia Sept-17	Virtual Oct-17-Feb-18	Sofia Feb-18
Antonucci	Serena	Italy	Forest ecology	x			
Azevedo	Joao	Portugal	Forest economics				x
Bielak	Kamil	Poland	Forest management	x			
<b>Binder</b>	<b>Franz</b>	<b>Germany</b>	<b>Silviculture</b>	x	x	x	x
<b>Bowditch</b>	<b>Euan</b>	<b>United Kingdom</b>	<b>Forest policy</b>			x	x
Cherubini	Paolo	Switzerland	Forest ecology				x
Chianucci	Francesco	Italy	Forest management				x
Coll	Lluis	Spain	Forest management	x			x
Čurović	Željka	Montenegro	Landscape architecture				x
Čurović	Milić	Montenegro	Forest ecology				x
Dalponte	Michele	Italy	Remote sensing	x			
<b>delRio</b>	<b>Miren</b>	<b>Spain</b>	<b>Forest modeling</b>				x
Di Lella	Stefania	Italy	Forest ecology	x			
Dimopoulos	Panayotis	Greece	Forest ecology	x			
Dinca	Lucian	Romania	Forest soils	x			
Ditmarová	Ľubica	Slovakia	Forest ecophysiology	x			x
Fayvush	Georgi	Armenia	Forest botany				x
Frizzera	Lorenzo	Italy	Forest ecology	x			
Gianelle	Damiano	Italy	Remote sensing	x			
Heinze	Berthold	Austria	Forest genetics	x			x
Ilieva	iliana	Bulgaria	Forest communication		x		
Kašanin-Grubin	Milica	Serbia	Forest soils				x
<b>Kluvankova</b>	<b>Tatiana</b>	<b>Slovakia</b>	<b>Forest economics</b>			x	x
Kurylyak	Viktor	Ukraine	Silviculture	x			
<b>La Porta</b>	<b>Nicola</b>	<b>Italy</b>	<b>Forest pathology</b>	x	x	x	x
Lavadinovic	Vera	Serbia	Forest economics	x			
<b>Lesinski</b>	<b>Jerzy</b>	<b>Poland</b>	<b>Forest biodiversity</b>		x	x	x
Marshall	John	Sweden	Forest ecophysiology				x
Meszaros	Ilona	Hungary	Forest ecophysiology	x			
<b>Motta</b>	<b>Renzo</b>	<b>Italy</b>	<b>Silviculture</b>	x			x
Neroj	Bozydar	Poland	Forest inventory	x			
<b>Pach</b>	<b>Maciej</b>	<b>Poland</b>	<b>Forest management</b>	x	x	x	x
<b>Panzacchi</b>	<b>Pietro</b>	<b>Italy</b>	<b>Forest ecology</b>	x	x	x	x
<b>Pretzsch</b>	<b>Hans</b>	<b>Germany</b>	<b>Forest growth</b>			x	x
Pšidová	Eva	Slovakia	Forest ecophysiology				x

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

Puletti	Nicola	Italy	Forest inventory	x			
Radoglou	Kalliopi	Greece	Forest ecophysiology				x
<b>Santopuoli</b>	<b>Giovanni</b>	<b>Italy</b>	<b>Forest management</b>	x	x	x	x
<b>Smith</b>	<b>Melanie</b>	<b>United Kingdom</b>	<b>Forest policy</b>	x	x	x	x
Snorrason	Arnor	Iceland	Forest management	x			
Spathelf	Peter	Germany	Forest growth				x
Stojnic	Srdjan	Serbia	Forest ecophysiology				x
<b>Temperli</b>	<b>Christian</b>	<b>Switzerland</b>	<b>Forest inventory</b>	x	x	x	x
<b>Tognetti</b>	<b>Roberto</b>	<b>Italy</b>	<b>Forest ecophysiology</b>		x	x	x
<b>Tonon</b>	<b>Giustino</b>	<b>Italy</b>	<b>Forest ecology</b>	x	x		x
Torresan	Chiara	Italy	Forest management				
Tsonev	Tsonko	Bulgaria	Forest ecophysiology		x		x
<b>Velikova</b>	<b>Violeta</b>	<b>Bulgaria</b>	<b>Forest ecophysiology</b>	x	x	x	x
<b>Weatherall</b>	<b>Andrew</b>	<b>United Kingdom</b>	<b>Forest ecology</b>	x	x	x	x

121 **Appendix C**

1  
2

122 **Description of network analysis implementation**

4  
5

123 The analytic network analysis allows to assess pairwise relationships between objects and entities of a  
7  
124 network to identify trends and patterns. In this study, the entities are two important tools that deal with  
9  
125 forest management, namely the indicators set of sustainable forest management (SFM) developed by  
11  
126 Forest Europe in 1998 and recently updated in Madrid 2015 ([https://foresteurope.org/sfm-criteria-](https://foresteurope.org/sfm-criteria-indicators/)  
14  
127 [indicators/](https://foresteurope.org/sfm-criteria-indicators/)), and the forest-related ecosystem services according to the CICES classification version 4, 2012  
16  
128 (<https://cices.eu>). The objects of the network are, on one side the SFM indicators, and on the other side  
19  
129 the ecosystem services. The analysis aims to assess and displays the connection between SFM indicators  
21  
130 and forest-related ecosystem services according to the preferences of people involved in the project  
23  
131 meeting. All the objects were equally considered in this study, and connections were delineated  
26  
132 considering the suitability of indicators to provide useful information for assessing adaptation and  
28  
133 mitigation forest management.

31  
32

134 Analyzing the structure of network is possible to observe some centrality measures which allow to find the  
34  
135 most important nodes in a network.

36  
37

136 Some of the most commonly centrality parameters are:

39  
40

- 137 • Network size (Ns), which corresponds to the number of nodes forming the network (eqn. 1). In this  
42  
138 context, higher values of Ns display high variability in the number of indicators (29) and ecosystem  
44  
139 services (23) identified by the participants as useful to assess the climate-smart forestry.

47  
48

140  $N_s = \sum \text{nodes} \quad (1)$

50  
51

- 141 • Network density (Nd), which corresponds to the proportion of existing lines when compared to all  
53  
142 the possible lines (eqn. 2). Higher values of Nd reflect the complexity of the network and, in this  
54  
143 context, of social values. It reflects the non-trivial topological features as those the relationships  
56  
144 between SFM indicators and forest-related ecosystem services.

59  
60

61  
62

63  
64  
65

145  $Nd = \frac{\Sigma(ties)}{Rows*Columns} \% \quad (2)$

146 • The centrality parameters, such as the “Degree”, which reflects the number of relationships for  
 147 each node. Higher values of Degree indicate the central position of the node within the network,  
 148 reflecting the importance given by people to this node. The Degree is the sum of Indegree and  
 149 Outdegree values, for ecosystem services and Indicators respectively. The Indegree is the number  
 150 of ties that each square node receives while the Outdegree is the number of ties that each circle  
 151 node sends to other nodes. ‘Centrality’ identifies the node’s core importance within the network.  
 152 This is particularly relevant in this context, because it allows to highlight the key elements that  
 153 support CLIMO participants in the development of CSF definition.

154 • Betweenness centrality is based on the number of times a node in the network is “between” other  
 155 nodes on the causal paths. The Betweenness reflects on how many links depend on this particular  
 156 node (eqn. 4):

157 
$$CB(v) = \sum_{s \neq v \neq t \in V} \frac{\partial st(v)}{\partial st}$$

158 where  $\partial st$  is the number of shortest paths from s to t, and  $\partial st(v)$  is the number of shortest paths from s  
 159 to t that pass through a vertex v.


160 • K-core consists in the identification of particular subsets of the network. A k-core is a maximal  
 161 group of actors, all of which are connected to some number (k) of other members of the group. If  
 162 an actor has ties to a sufficient number of members of a group, they may feel tied to that group,  
 163 even if they do not know many, or even most members. It may be that identity depends on  
 164 connection, rather than on immersion in a sub-group.

165 **The main output of the analytic network analysis is displayed in the figure S2, S3, and S4.**




16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65


**Figure S3: Suitable Indicators for monitoring the MITIGATION of Climate-Smart Forestry in mountain ecosystems. Square blue nodes represent the Ecosystem Services, while circle red nodes represent the Indicators for Sustainable Forest Management.**




**The indicators listed on the top, left corner are not considered by CLIMO participants for the mitigation to climate change.**









**Declaration of interests**

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

The authors declare the following financial interests/personal relationships which may be considered as potential competing interests: