

# Lead poisoning from Ayurvedic treatment: a further case

CATALINA CIOCAN<sup>1,2</sup>, IHAB MANSOUR<sup>1</sup>, ALESSANDRO BENEDEUCE<sup>1</sup>, RICCARDO CORGIAT LOIA<sup>1</sup>, NICOLÒ MILANESIO<sup>1</sup>, MICHAEL DECLEMENTI<sup>1</sup>, ALESSANDRO GODONO<sup>1</sup>, GIACOMO GARZARO<sup>1,2</sup>, ENRICO PIRA<sup>1,2</sup>

<sup>1</sup>University of Torino, Department of Public Health and Pediatrics, Turin, Italy

<sup>2</sup>Città della Salute e della Scienza di Torino University Hospital, Occupational Health Service, Turin, Italy

**KEY WORDS:** Ayurvedic; lead poisoning; anaemia; chelation; drug safety

## SUMMARY

**Purpose:** We report a case of a 30 years old Indian sailor with microcytic anemia (Hb 9.9), persistent abdominal pain, emesis, dark stool, hyperchromic urine, latent jaundice and asthenia. Lead intoxication was confirmed (blood lead value of 102 µg/dL). The patient assumed Ayurvedic medicines in the previous months. Ayurveda is an ancient form of Indian traditional popular medicine aiming to re-establish health and body function through herbal preparations, heavy metals are often added. Our purpose was to treat the patient and to establish the source of poisoning. **Methods:** After testing blood and urine lead concentration of other 3 crew members and analyzing over than 150 products used on the ship professional exposure was excluded. We analyzed the two Ayurvedic drugs assumed by the patient with an Inductively Coupled Plasma Mass Spectrometry. The patient underwent three chelation cycles with Calcium Disodium Ethylenediaminetetraacetic acid (EDTA) while monitoring blood and urinary lead levels. **Results:** The final blood lead level at discharge, after three chelation cycles, was 36.27 µg/dL. One of the two drugs contained extremely high concentrations of lead and mercury. The three different mixtures of this preparation showed lead concentrations of 12,638.54 mg/kg (Sample 2A), 23,043.02 mg/kg (Sample 2B), 21,352.97 mg/kg (Sample 2C); these levels are much higher than the highest values reported in literature for the indian soil (32 mg/kg). **Conclusions:** This case and similar cases raise the safety alert on complementary and alternative medicines; Ayurvedic medicine users should be carefully informed about potential risks and signs of poisoning.

Ayurveda is a form of traditional medicine native to India which has been practiced extensively in the subcontinent for more than 2000 years (1) and is used approximately by 80% of Indians (2). Recently, Ayurveda is gaining popularity as a complementary treatment. As a result, approximately 40% of adults in the United States report using complementary and alternative medicines (CAMs) (3). Increasing numbers could probably be observed in countries

where Ayurveda is not yet so practiced, due to work migration (4). Ayurvedic formulations are based on herbal products but often include heavy metals, intentionally added to restore health and normal body function, as part of Rasa Shastra (5).

Contrary to allopathic medicine, manufacturing and access to CAMs are poorly regulated. Ayurvedic medicines in Western countries are not considered as drugs. Rather, as food supplements, they

Received 2.9.2020 - Accepted 14.10.2020

Corresponding author: Dr. Michael Declementi, Via Zuretti 29, 10126 Turin, Italy - E-mail: michael.declementi@unito.it

 open access [www.lamedicinadellavoro.it](http://www.lamedicinadellavoro.it)

are available without medical prescription through multiple sources including the Internet. Although even some physicians overlook possible risks associated with ayurvedic treatments, it is estimated that over 20% of ayurvedic medications distributed by U.S. and Indian companies contain toxic metals (5); some preparations have been found to contain lead and/or mercury exceeding by 100 to 10,000 times the limits set for food (6).

From 2000 to 2003, the Centers for Disease Control reported 12 cases of lead poisoning in adults associated with Ayurvedic medication intake occurring in five different states (7). Less than 20 scientific reports on lead poisoning could be found through scientific literature analysis of the past 10 years (Mesh key words: “lead poisoning” AND “Ayurvedic medication”). The contamination of herbal formulations with metals presents potential health risks. The most common diagnosis is lead poisoning, which is frequently revealed by the investigation of anaemia.

Lead is considered as an inhibitor of haem synthesis and a haemolysis promotor. Lead related anaemia can be normocytic or microcytic mainly due to the inhibition of delta aminolevulinic acid dehydratase and ferrochelatase. As a result, protoporphyrin accumulates in erythrocytes forming zinc protoporphyrin. Lead levels exceeding 30 µg/dL determine elevated blood levels of free erythrocyte protoporphyrin and zinc protoporphyrin. Anaemia develops when blood lead levels exceed 50 to 60 µg/dL. Lead has a prevalent urinary excretion, the non-excreted portion being deposited in bones. Other characteristic manifestations of lead poisoning include gastrointestinal and neurological symptoms (8).

In Western countries, industrial lead exposure control played a key role in minimizing job exposure (9) and in the last decades cases of saturnism requiring hospitalization were mostly non-occupational, due to contaminated wine (10). We recently had the chance to admit to our Department a young Indian sailor with high levels of blood lead. During a residential period in his country from 14/10/2019 to 31/01/2020 the patient presented lower back pain due to the recurrence of a pilonidal cyst, previously experienced in 2008 and 2013. In both cases, he was treated with Ayurvedic topic medications while, for

this last recrudescence, his Ayurvedic practitioner (referred to as “shaman”) prescribed an oral formulation. In February, the patient returned to Italy and rejoined his crew in Genoa (Italy) with whom he carried out different maintenance activities on board. At the beginning of March, due to persistent abdominal pain, he underwent gastroenterological examination and a few days later he presented emesis, dark stool, hyperchromic urine, latent jaundice and asthenia and was referred to Hospital Galliera in Genoa. The first blood test showed hemoglobin 9.9 mg/dL, total bilirubin 3.5 mg/dL (direct bilirubin 1.1 mg/dL), SGOT 90 mg/dL, SGPT 160 mg/dL (Table 1). Esophagogastroduodenoscopy was negative. The only finding in the chest and abdominal CT was hepatic steatosis. At the end of April, the patient underwent haematological counselling for microcytic anaemia associated with hyperferritinemia and reticulocytosis. A screening for lead poisoning showed a blood lead value of 102 µg/dL. On 8/6/2020, after passing the COVID-19 hospital prevention protocol (1), the patient came to our observation: during the previous three months the ayurvedic treatment had been withdrawn, and symptoms gradually disappeared thereafter. In April and May 2020, the patient did not undergo any blood tests.

During the first hospitalization in our department, we excluded occupational exposure. Blood and urine lead levels of other crew members were in a normal range and over 150 products used on board resulted lead free. At the admission in our department, the patient was asymptomatic, and the main haemato-chemical values were within the respective reference intervals (Table 1-2), whereas the basal lead values in blood (B-Pb) and urine samples were still high (74.61 µg/dL and 94.7 µg/L, respectively).

In consideration of the B-Pb (12, 13), a chelation protocol was initiated with the administration of Calcium Disodium Ethylenediaminetetraacetic acid (EDTA) 2 g in 500 ml of physiological solution (PS) in 12 hours for the first day, followed by four days of EDTA 1 g at the same dilution and infusion speed. Hematic and urinary lead variation during three chelation cycles is reported in Figure 1. B-Pb at discharge was 36.27 µg/dL.

The sailor referred that he used to assume an oral intake of 1 dose (about one spoon) of “product 1” mixed with 1 dose of each of the three parts of “product 2”, twice a day (products 1, 2A, 2B, 2C).

Our Industrial Toxicology Laboratory analysed 0.5 g of each product sample: we added 2.5 ml of concentrated Nitric Acid to each sample and after 1h of treatment in a furnace at 80°C we made up to 25 ml, we diluted the solutions (1:1000) and used an inductively coupled plasma mass spectrometry (ICP-MS) searching for arsenic, mercury, copper, chromium, manganese, iron, nickel and lead.

Finally, we compared our values with data in the literature regarding the highest values in different soil samples of Uttar Pradesh, our patient’s home region; results are summarized in Table 3 (14-16).

## CONCLUSION

Laboratory findings confirmed the source of poisoning. The absence of mercury-related symptoms despite a high content of mercury in the samples 2A, 2B and 2C could be explained by the species of the metal, as inorganic mercury is known to be poorly absorbed by the gastrointestinal tract. A peculiar aspect of this case is that haemoglobin (Hb) usually takes several months to return to normal values in the absence of chelation while our patient had a fast normalization of Hb values in less than three months, before arrival in our hospital (Table 2). This case and similar cases of lead poisoning caused by Ayurveda practice raise the safety alert of complementary and alternative medicines. The increasing number of users of Ayurvedic drugs

**Table 1** - Blood values of biomarkers of lead intoxication

	At admission (08/06/2020)	At discharge (10/07/2020)	Reference values
ALA Dehydratase, U.I./ml	11.5	20.8	20.0 – 40-0
Blood Lead Level (ICP-MS), µg/dL	74.61	42.2	0.50 – 10.0
Baseline Urinary Lead (ICP-MS), µg/dL	94.70	93.6	5.00 – 30.00
Total Hemoglobin, g/dL	12.80	13.9	14.00 – 18.00
Zinc protophorphyrin – ZP, µg/dL Hb	0.66	0.52	< 2.00
ALAU - Delta-aminolevulinic acid, mg/g creat	14.89	1.96	< 4.50
CPU – Coproporphyrin, µg/dL creat	423	3	< 100.00

**Table 2** - Selected haemato-chemical values

	Initial values (11/04/2020)	At admission (08/06/2020)	At discharge (10/07/2020)	Reference values
Creatinine, mg/dL	0.9	0.93	1.04	0.60 – 1.30
Sodium, mEq/L	139	143	142	135 – 145
Potassium, mEq/L	4.6	4.4	3.8	3.5 – 5.0
Leukocytes, No.x10 <sup>9</sup> /L	9.73	6.51	5.99	4.00 – 10.00
Erythrocytes, No.x10 <sup>12</sup> /L	3.23	4.58	5.12	4.50 – 5.80
Haemoglobin, g/dL	8.40	13.0	14.1	13.5 – 18.00
Hematocrit, %	25.5	41.0	42.4	40.0 – 52.0
Mean corpuscular volume, fL	78.9	90	83	79 – 96
Platelet Count, No.x10 <sup>9</sup> /L	300	270	223	140 – 450




Figure 1 - Blood and urine lead levels during chelation therapy

Table 3 - Concentration of analysed elements in the environment and in the ayurvedic drug taken by the patient

	Highest soil concentration in literature (mg/kg)*	Concentration in sample product 1 (mg/kg)	Concentration in sample product 2A (mg/kg)	Concentration in sample product 2B (mg/kg)	Concentration in sample product 2C (mg/kg)
Lead	520	2.31	12,638.54	23,043.02	21,352.97
Mercury	0.49	2.54	89,076.70	126,640.27	145,710.64
Arsenic	12	0.35	134.94	103.98	122.30
Copper	323	3.96	1,511.04	2,168.79	2,356.80
Chromium	0.9	1.14	8.86	7.68	12.04
Manganese	45.1	26.34	146.74	172.53	149.75
Iron	389	288.04	14,990.34	16,498.80	15,767.59
Nickel	100	0.66	7.24	8.02	9.21

\* Values referred to different sites of Uttar Pradesh region, except for mercury that is referred to whole India

worldwide calls for establishing standardized labeling and regulations of manufacturing and distribution of these treatments. Moreover, consumers should be properly informed about potential risks and clinical signs of poisoning. We believe it is important to stress these concepts and to be extremely vigilant, as the number of case reports is dwindling in the last ten years, while this work proves that the problem persists.

## REFERENCES

1. Datta-Mitra A, Ahmed O Jr.: Ayurvedic medicine use and lead poisoning in a child: a continued concern in the United States. *Clin Pediatr (Phila)* 2015; 54: 690-692. doi:10.1177/0009922814553397
2. Saper RB, Kales SN, Paquin J, et al. Heavy metal content of ayurvedic herbal medicine products. *JAMA*. 2004; 292: 2868-2873. doi:10.1001/jama.292.23.2868
3. Eisenberg DM, Davis RB, Ettner SL, et al. Trends in alternative medicine use in the United States, 1990-1997: results of a follow-up national survey. *JAMA* 1998; 280: 1569-1575. doi:10.1001/jama.280.18.1569
4. Mucci N, Traversini V, Giorgi G, et al. Migrant Workers and Physical Health: An Umbrella Review. *Sustainability* 2019; 11: 232
5. Mikulski MA, Wichman MD, Simmons DL, et al. Toxic metals in ayurvedic preparations from a public health lead poisoning cluster investigation. *Int J Occup Environ Health* 2017; 23: 187-192. doi:10.1080/10773525.2018.1447880
6. Gunturu KS, Nagarajan P, McPhedran P, Goodman TR, Hodsdon ME, Strout MP. Ayurvedic herbal medicine and lead poisoning. *J Hematol Oncol* 2011; 51. doi: 10.1186/1756-8722-4-51. PMID: 22185092; PMCID: PMC3259062
7. Centers for Disease Control and Prevention (CDC). Lead poisoning associated with ayurvedic medications—five states, 2000-2003. *MMWR Morb Mortal Wkly Rep* 2004; 53: 582-584
8. Kales SN, Christophi CA, Saper RB. Hematopoietic toxicity from lead-containing Ayurvedic medications. *Med Sci Monit* 2007; 13(7):CR295-CR298
9. Pira E, Garzaro G, De Cillis E, et al. Evolution of the concept OS&H from the second Post-war to today: From prescriptive system to assessment and management of risks in system quality -The extended model in collaboration with large Facilities. Evolution of multidisciplinary culture of safety and OS&H. *GEAM* 2018; 55: 16-20
10. Perrelli G, Capellaro E, Pira E, et al. Further cases of lead poisoning from wine. *Am J Ind Med* 1984; 5: 377-381. doi:10.1002/ajim.4700050506
11. Garzaro G, Clari M, Ciocan C, et al. COVID-19 infection and diffusion among the healthcare workforce in a large university-hospital in northwest Italy. *Med Lav* 2020; 111: 184-194. doi:10.23749/mdl.v111i3.9767
12. Flora SJ, Pachauri V. Chelation in metal intoxication. *Int J Environ Res Public Health* 2010; 7: 2745-2788. doi:10.3390/ijerph7072745
13. Schroeder AP, Tilleman JA, DeSimone II EM. Lead Toxicity and Chelation Therapy, *US Pharm* 2015; 40: 40-44
14. Singh S, Raju NJ, Nazneen S. Environmental risk of heavy metal pollution and contamination sources using multivariate analysis in the soils of Varanasi environs, India. *Environ Monit Assess* 2015; 187: 345. doi:10.1007/s10661-015-4577-4
15. Raj D, Maiti SK. Sources, toxicity, and remediation of mercury: an essence review. *Environ Monit Assess*. 2019; 191:566. doi:10.1007/s10661-019-7743-2
16. Adimalla N. Heavy metals pollution assessment and its associated human health risk evaluation of urban soils from Indian cities: a review. *Environ Geochem Health* 2020; 42:173-190. doi:10.1007/s10653-019-00324-4

NO POTENTIAL CONFLICT OF INTEREST RELEVANT TO THIS ARTICLE WAS REPORTED BY THE AUTHORS



## LA RACCOLTA COMPLETA DELLA NOSTRA RIVISTA DIVENTA DIGITALE

Procede la digitalizzazione della raccolta completa della nostra rivista (“Il Lavoro” dal 1901 al 1924 e poi “La Medicina del Lavoro” dal 1925 in poi). Oltre agli ultimi 20 anni, già accessibili nel sito della casa editrice ([www.lamedicinadellavoro.it](http://www.lamedicinadellavoro.it)), i volumi pubblicati nei primi 50 anni (1901-1950) sono ora disponibili sia in formato fotografico (.tiff o .jpg) che come documento ricercabile (.pdf). Tutti i volumi disponibili saranno online e ad accesso libero subito dopo la firma di una convenzione tra la struttura (SIML - Società Italiana di Salute sul Lavoro) e l'Università degli Studi di Milano (proprietaria dell'archivio storico della rivista). Nel frattempo, chi intendesse realizzare progetti sulla storia della medicina del lavoro utilizzando i volumi pubblicati nella prima metà del XX secolo è pregato di contattarci ([redazione\[at\]lamedicinadellavoro.it](mailto:redazione[at]lamedicinadellavoro.it)).

## THE COMPLETE COLLECTION OF OUR JOURNAL GOES DIGITAL

The digitalization of the complete collection of our journal (“Il Lavoro” and then “La Medicina del Lavoro” from the 16th volume onwards) is advancing. In addition to the last 20 years already accessible from the journal's site ([www.lamedicinadellavoro.it](http://www.lamedicinadellavoro.it)), all the volumes published during the first 50 years (1901-1950) are now available both in photographic format (.tiff and .jpg) and as a searchable document (.pdf). Open access will be provided soon once an agreement is signed between the property (SIML – Italian Society of Occupational Health) and the University of Milan (owner of the journal's historical archive). Meanwhile, scholars willing to implement projects on the history of Occupational Medicine relying on the volumes published during the first half of the 20th century are kindly requested to contact us ([redazione\[at\]lamedicinadellavoro.it](mailto:redazione[at]lamedicinadellavoro.it)).

