

AperTO - Archivio Istituzionale Open Access dell'Università di Torino

Effect of withering process on the evolution of phenolic acids in winegrapes: A systematic review

This is the author's manuscript

Original Citation:

Availability:

This version is available <http://hdl.handle.net/2318/1798296> since 2023-02-22T16:28:47Z

Published version:

DOI:10.1016/j.tifs.2021.08.004

Terms of use:

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)

Journal Pre-proof

Effect of withering process on the evolution of phenolic acids in winegrapes: A systematic review

Giulia Scalzini, Simone Giacosa, Susana Río Segade, Maria Alessandra Paissoni, Luca Rolle



PII: S0924-2244(21)00490-8

DOI: <https://doi.org/10.1016/j.tifs.2021.08.004>

Reference: TIFS 3510

To appear in: *Trends in Food Science & Technology*

Received Date: 24 March 2021

Revised Date: 15 July 2021

Accepted Date: 11 August 2021

Please cite this article as: Scalzini, G., Giacosa, S., Río Segade, S., Paissoni, M.A., Rolle, L., Effect of withering process on the evolution of phenolic acids in winegrapes: A systematic review, *Trends in Food Science & Technology* (2021), doi: <https://doi.org/10.1016/j.tifs.2021.08.004>.

This is a PDF file of an article that has undergone enhancements after acceptance, such as the addition of a cover page and metadata, and formatting for readability, but it is not yet the definitive version of record. This version will undergo additional copyediting, typesetting and review before it is published in its final form, but we are providing this version to give early visibility of the article. Please note that, during the production process, errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

© 2021 Published by Elsevier Ltd.

Effect of withering process on the evolution of phenolic acids in winegrapes: A systematic review

Giulia SCALZINI, Simone GIACOSA, Susana RÍO SEGADE*, Maria Alessandra PAISSONI,
Luca ROLLE

*Università degli Studi di Torino, Dipartimento di Scienze Agrarie, Forestali e Alimentari. Largo
Paolo Braccini 2, 10095 Grugliasco (TO), Italy.*

*Corresponding author: susana.riosegade@unito.it; +39-011-670-3989

Abstract

Background: Withering processes have a strong impact on the phenolic composition of winegrapes and related wines. Even if attention has often been focused on other phenolics, a great number of studies have included phenolic acids because of their quantitative and qualitative modifications during dehydration-withering processes. A systematic review that provides a concise overview of the extensive literature available on this important topic is lacking.

Scope and approach: This review identified 39 articles to answer the research question: *What changes occur in phenolic acids after the withering process of winegrapes?* The expected contribution of this systematic review is to have a full view of the worthwhile recent findings on these often-overlooked compounds to manage the technological process with new awareness and to find eventual weaknesses in this field to highlight new questions and research directions.

Key findings and conclusions: The research yielded useful results for withering and winemaking management. Phenolic acids are well represented in certain special productions to be proposed as markers of wine authenticity. Hydroxycinnamic and hydroxybenzoic acids and their derivatives are often affected differently by the withering process. Their evolution during grape withering is a

27 complex phenomenon that is affected by many varietal and technological variables. In particular, the
28 withering conditions applied and grape genotype play an important role in the changing amounts of
29 phenolic acids, but there is still much to be understood, especially related to the combined effect of
30 both factors through genetic responses to environmental stresses and their respective chemical
31 implications.

32

33 *Keywords:* Phenolic composition; Phenolic acids; Withering process; Grape dehydration; Special
34 wines.

Journal Pre-proof

Effect of withering process on the evolution of phenolic acids in winegrapes: A systematic review

Giulia SCALZINI, Simone GIACOSA, Susana RÍO SEGADE*, Maria Alessandra PAISSONI,
Luca ROLLE

*Università degli Studi di Torino, Dipartimento di Scienze Agrarie, Forestali e Alimentari. Largo
Paolo Braccini 2, 10095 Grugliasco (TO), Italy.*

*Corresponding author: susana.riosegade@unito.it; +39-011-670-3989

Abstract

Background: Withering processes have a strong impact on the phenolic composition of winegrapes and related wines. Even if attention has often been focused on other phenolics, a great number of studies have included phenolic acids because of their quantitative and qualitative modifications during dehydration-withering processes. A systematic review that provides a concise overview of the extensive literature available on this important topic is lacking.

Scope and approach: This review identified 39 articles to answer the research question: *What changes occur in phenolic acids after the withering process of winegrapes?* The expected contribution of this systematic review is to have a full view of the worthwhile recent findings on these often-overlooked compounds to manage the technological process with new awareness and to find eventual weaknesses in this field to highlight new questions and research directions.

Key findings and conclusions: The research yielded useful results for withering and winemaking management. Phenolic acids are well represented in certain special productions to be proposed as markers of wine authenticity. Hydroxycinnamic and hydroxybenzoic acids and their derivatives are often affected differently by the withering process. Their evolution during grape withering is a

27 complex phenomenon that is affected by many varietal and technological variables. In particular, the
28 withering conditions applied and grape genotype play an important role in the changing amounts of
29 phenolic acids, but there is still much to be understood, especially related to the combined effect of
30 both factors through genetic responses to environmental stresses and their respective chemical
31 implications.

32

33 *Keywords:* Phenolic composition; Phenolic acids; Withering process; Grape dehydration; Special
34 wines.

35

Journal Pre-proof

36 1. Introduction

37 The techniques currently utilized in winemaking and grape treatment for wine production are
38 the result of decades of history, experience, and tradition. Scientific research has answered many
39 practical questions, introduced important innovations, and allowed winemakers to manage traditional
40 processes with new awareness.

41 Among the oenological techniques, the withering of winegrapes is one of the most strongly
42 related to the cultural history and the climate peculiarities of the territories (Mencarelli & Tonutti,
43 2013). As a consequence, this process may produce wines with very different sensory features: (i)
44 sweet wines, such as the botrytized *Sauternes* and *Tokaj*, the Mediterranean Italian *Passito*, the
45 special icewines from Canada, the traditional wines made from *Pedro Ximénez* sun-dried grapes, or
46 the straw wines called "*Vin de Paille*" and produced in the French Jura region; (ii) reinforced wines,
47 sweet but more alcoholic than conventional wines, produced in territories, such as *Marsala*, where
48 the traditional wines were historically added with alcohol during overseas shipping for the English
49 trade; (iii) fortified wines, dry and full-bodied, such as the *Sforzato* or '*Sfursat*' from the heroic
50 viticulture of Valtellina slopes and the *Amarone* from the Italian Valpolicella region (Kallitsounakis
51 & Catarino, 2020; Scienza, 2013).

52 The withering process can be conducted on-vine, leaving the grapes on the plant subject to
53 atmospheric phenomena for a long time (e.g., botrytized wine, late harvest wines, or icewines), or
54 off-vine by subjecting the grapes to postharvest dehydration, which may occur outdoors or indoors,
55 under controlled conditions or not (Figueiredo-González et al., 2013a). In recent decades, many
56 authors have extensively studied wines produced from withered grapes because of their cultural and
57 economic relevance, highlighting important changes in berry metabolism throughout the withering
58 process (D'Onofrio et al., 2019; Esmaili et al., 2007; Zoccatelli et al., 2013).

59 Polyphenols, which are crucial compounds for wine quality, change during the grape
60 dehydration process (Harrison, 2018; Mencarelli et al., 2010). Several studies have been carried out
61 to better understand the changes in the content and composition of phenolic compounds during the

62 on- or off-vine withering of winegrapes (Corradini & Nicoletti, 2013; Figueiredo-González et al.,
63 2013a; Torchio et al., 2016). These authors highlighted the complexity of the evolution of phenolics
64 over the process, which seems to be the result of a composite balance between synthesis,
65 oxidation/loss, and concentration (Bonghi et al., 2012; De Rosso et al., 2016). This balance varies
66 greatly depending on the genetic features of each cultivar (Zenoni et al., 2016), the ripeness level and
67 mechanical properties of grapes (i.e., skin hardness; Rolle et al., 2009), the different dehydration
68 techniques applied (Constantinou et al., 2018), the management of three important environmental
69 factors (i.e., temperature, humidity, and airflow), and berry weight loss rate (Rolle et al., 2013).
70 Additionally, specific classes of phenolic compounds, although subjected to the same conditions, are
71 not affected in the same manner by the withering process (Toffali et al., 2011).

72 Phenolic acids are an important fraction of non-flavonoid compounds (Baderschneider &
73 Winterhalter, 2001) present in grapes and wines, which have attracted increasing interest over the
74 past ten years because of their potential health benefits (e.g., antioxidant, antibacterial, antiviral,
75 anticancerogenic, anti-inflammatory, and vasodilatory actions) (Babbar et al., 2015; Morales-Prieto
76 et al., 2020). Phenolic acids consist predominantly of two subgroups: hydroxycinnamic acids (HCAs)
77 and hydroxybenzoic acids (HBAs), of types C6-C3 and C6-C1, respectively (Monagas et al., 2005).
78 The chemical structures of the most represented phenolic acids and their average contents in the wines
79 are presented in **Table S1**.

80 HCAs represent the main class of phenolic compounds in white wines and the most represented
81 class of non-flavonoid phenolics in red wines, averaging approximately 130 and 60 mg/L,
82 respectively (Vanzo et al., 2007). They are synthesized through the shikimate pathway from
83 phenylalanine and tyrosine through the action of phenylalanine ammonia lyase (PAL), which is a key
84 enzyme that exhibits a relevant interconnection with the synthesis of other phenolic compounds
85 (Laura et al., 2019). Some of the most represented HCAs are *p*-coumaric, caffeic, ferulic, and sinapic
86 acids. They are usually present in grapes and wines as tartaric acid esters and diesters (HCTAs), such
87 as caftaric acid (caffeoyltartaric acid), coutaric acid (*p*-coumaroyltartaric acid), and fertaric acid
88 (feruloyltartaric acid) and rarely as sugar esters (Buiarelli et al., 2010; Ferrandino et al., 2012; Ong &

89 Nagel, 1978). In grapes, they are mainly located in the pulp and slightly in the skin (Garrido & Borges,
90 2013). The contents of HCA esters detected in grapes vary greatly for different cultivars; however,
91 caftaric acid is generally the major compound, followed by coumaric and ferulic acids (Waterhouse,
92 2002). During the fermentation process, the partial hydrolysis of esters occurs, with a reaction rate
93 depending on the pH and presence of specific enzymes, and free HCAs can be released and partially
94 esterified with ethanol, forming HCA ethyl esters (Somers et al., 1987). The oxidation of caftaric acid
95 by grape polyphenol oxidase (PPO) to the corresponding *o*-quinone and subsequent reaction with
96 glutathione through the –SH groups form a stable and colorless compound named GRP (Grape
97 Reaction Product), which plays a key role in the oxidation of other phenolic compounds. This
98 oxidation reaction starts to take place during the crushing and pressing of the grapes, causing must
99 browning (Cheynier et al., 1989a). HCAs and their derivatives are also known to have antimicrobial
100 properties, particularly against the wine spoilage lactic acid bacteria and yeasts (e.g., *Brettanomyces*
101 *bruxellensis*) (Harris et al., 2010; Sabel et al., 2017; Stivala et al., 2017), increasing cell membrane
102 permeability from wine (Campos et al., 2009) and interfering with the intracellular pH and
103 metabolism (Carmona et al., 2016). The presence of HCAs in wines markedly affects color quality
104 features in different ways: on one hand, they can contribute significantly to the browning of musts
105 due to their oxidation (Cheynier et al., 1989b); on the other hand, they can contribute to the color
106 stabilization of red wines due to the copigmentation effect or participate in the formation of acylated
107 anthocyanins and anthocyanin-derived pigments (i.e., pinotins and portisins) (Bloomfield et al., 2003;
108 He et al., 2012). Therefore, the decrease in the content of coumaroylated anthocyanins during
109 winemaking and aging processes due to hydrolytic and enzymatic reactions can release HCAs in
110 musts and wines (Monagas et al., 2005). In terms of wine mouthfeel qualities, these compounds have
111 been associated with astringency and bitterness perceptions (Garrido & Borges, 2013; Okamura &
112 Watanabe, 1981), but their levels seem not to be perceptible in wine (Vérette et al., 1988). However,
113 some more recent studies have pointed out that "puckering" astringency may be elicited by these
114 compounds and evidenced a synergism on bitterness perception (Gonzalo-Diago et al., 2014;
115 Hufnagel & Hoffman, 2008). Moreover, HCAs were identified as potential volatile precursors of

116 ethyl phenols produced by *Brettanomyces* metabolism, conferring unpleasant flavors to wine
117 (Vanbeneden et al., 2008). In particular, *p*-coumaric, ferulic, and caffeic acids could be enzymatically
118 decarboxylated to vinyl derivatives by a cinnamate decarboxylase and subsequently reduced by a
119 vinylphenol reductase, leading to the typical off-flavors described as “horse sweat,” “medicinal,”
120 “rancid,” and “barnyard” (Malfeito-Ferreira, 2018). While several microorganisms, including
121 *Saccharomyces cerevisiae*, can perform the enzymatic decarboxylation of HCAs, the populations of
122 *Brettanomyces* spp. have a particularly effective activity in the reduction step (Kheir et al., 2013).

123 HBAs represent a minor component in grapes and wines, averaging approximately 10–20 mg/L
124 in white wines and approximately 70 mg/L in red wines (Waterhouse & Teissedre, 1997; Waterhouse,
125 2002). Unlike HCAs, HBAs are not phenylpropanoids; they can be synthesized directly from the
126 shikimic acid pathway, even if PAL is not active (Laura et al., 2019). Para-hydroxybenzoic,
127 protocatechuic, vanillic, gallic, and syringic acids are the most abundant HBAs detected (Kallithraka
128 et al., 2009). Among them, gallic acid is considered the most important, as it is the precursor of all
129 hydrolyzable tannins (Neculescu et al., 2018) and for its relevant antioxidant activity (Kallithraka et
130 al., 2009). HBAs are mainly present in grape skins and pulp as glycosides, but gallic acid can also be
131 extracted from grape seeds in free form through the hydrolysis of tannin galloyl esters or encompassed
132 in condensed tannins (Revilla & Gonzalez-SanJose, 2003; Zou et al., 2002). In wines, HBAs are
133 mostly found in free form (Monagas et al., 2005), even though some authors have also reported ethyl
134 and methyl esters and glucose esters (Baderschneider & Winterhalter, 2001). With regard to sensory
135 contributions, some recent studies have pointed out a synergic astringency effect with other phenols
136 as reported for HCAs, but HBAs showed a higher affinity for salivary proteins than HCAs (Ferrer-
137 Gallego et al., 2014; Ferrer-Gallego et al., 2017). Moreover, Wang et al. (2020) highlighted a relevant
138 matrix effect of phenolic acids on wine aroma modulation, particularly focusing on the interaction
139 between gallic acid and free terpene compounds that inhibits volatile release.

140 When discussing the outcome of grape withering processes on phenolic compounds, attention
141 is often focused on major molecules, such as anthocyanins and tannins. Nevertheless, many
142 significant changes occur in the phenolic acids. In fact, their evolution throughout the withering

143 process is complex and affected by many variables. Many studies have addressed phenolic acid
144 evolution during grape withering, considering their importance in terms of abundance and wine
145 quality implications; however, a systematic review to summarize these findings is lacking.

146 Wines made from withered grapes are often identified as unique products of specific, limited
147 geographical areas. Considering their growing interest in competitive and global markets, several
148 studies have tried, through chemical analysis, to define characteristic compounds as quality markers
149 to differentiate these wines from other products. In many cases, the most abundant classes of phenolic
150 compounds that can be used as markers are HBAs or HCAs (Figueiredo-González et al., 2014a;
151 Loizzo et al., 2013; Panceri et al., 2015). The purpose of this paper is to provide an overview of the
152 scientific literature on this challenging topic. The approach used was aimed to focus on the evolution
153 of phenolic acids from both qualitative and quantitative points of view in withered grapes and wines
154 as objectively as possible. The expected contribution of this systematic review is twofold: (i) to have
155 a systematic full view of the worthwhile recent findings to manage the technological process with
156 new awareness, concerning the importance of these compounds, which are often overlooked; (ii) to
157 find eventual research gaps in this field to highlight new questions and research directions.

158

159 **2. Method**

160 To produce an accurate and concise overview of the extensive literature available on this theme,
161 a systematic review was performed on February 24, 2021, using the procedure based on the Preferred
162 Reporting Items for Systematic Reviews and Meta-Analyses (Moher et al., 2009).

163 A protocol was developed to provide explicit, rigorous, and transparent planning proceedings
164 and eligibility criteria for the identification and selection of the studies of interest.

165

166 **Research question**

167 The review addressed the following close-framed question:

168 *RQ - What changes occur in phenolic acids after the withering process of winegrapes?*

169 The question has been deemed suitable for carrying out a systematic review, with all the PICO
170 key elements specified, as illustrated in **Table 1**.

171

172 [Table 1]

173 **Search strategy**

174 The research was performed simultaneously on three digital sources of information: "Scopus"
175 (<https://www.scopus.com/>), "ScienceDirect" (<https://www.sciencedirect.com/>), and "Web of
176 Science" (<https://webofscience.com/>).

177 An initial list of search terms was developed, including keywords, synonyms, plural/singular
178 forms, and alternative words (**Table 2**).

179

180 [Table 2]

181

182 The set of terms was used to create the search strings, matching them with Boolean operators. The
183 search strings used for all the selected sources of information are listed in **Table 3**.

184

185 [Table 3]

186

187 The search fields selected were "all fields," "find articles with these terms" (i.e., all parts of the
188 documents excluding references), and "all fields" for "Scopus," "ScienceDirect," and "Web of
189 Science," respectively. The date range selected focused on articles published "from 2007 to the
190 present".

191

192 **Selection of relevant studies**

193 This query yielded 16,883 results, which were preliminarily selected and deduplicated. The
194 potentially relevant citations identified were 94, and a two-step screening was carried out.

195 **Step 1: Initial screening** The first stage consisted of an initial screening of the retrieved articles
196 based on the Title and Abstract, removing those not related to the evolution of phenolic acids
197 during the withering process of winegrapes.

198 **Step 2: Full-text review and data extraction** The full texts of the studies selected in the first
199 stage were then fully reviewed to determine if they were still eligible to undergo data
200 extraction.

201 The inclusion criteria applied for the article selection were:

- 202 . study on the phenolic fraction of withered winegrapes (*Vitis vinifera* L.) and wines;
- 203 . focus on phenolic acids;
- 204 . evolution of phenolic acids during on-vine and off-vine withering processes and
205 winemaking implications.

206 The flow diagram of the relevant study selection process, shown in **Figure 1**, allowed to obtain
207 39 suitable documents.

208 [Figure 1]

210 **Data extraction, quality assessment, and reporting phase**

211 In the extraction stage, all selected articles were assessed for methodological quality. The main
212 results were collected, described, and critically examined to extract the common trends, comparing
213 the findings regarding winegrapes used for on-vine withering and off-vine natural and controlled
214 withering, genetic-dependent effects toward thermo-hygrometric environmental conditions, and
215 winemaking implications. To produce a synthetic overview of the main findings pointed out by the
216 recent literature (from 2007 to the present) about withering conditions applied, three extraction tables
217 have been elaborated, ordering the studies based on the withering technique. **Table 4** includes the 14
218 most relevant findings on the modification of phenolic acids over the on-vine winegrape withering
219 process, with most of them regarding the production of ice or botrytized wines. **Table 5** briefly
220 describes the main outcomes of the 18 studies on the off-vine withering process under controlled

221 conditions, while **Table 6** shows a synthesis of the results of the seven studies regarding off-vine
222 natural withering conditions.

223

224 [Table 4]

225 [Table 5]

226 [Table 6]

227

228 **3. Results and Discussion**

229 As grape berry composition is affected by several metabolic and physicochemical mechanisms
230 during withering and the findings can vary depending on the withering process, the results were
231 divided into on- and off-vine and the latter further separated under controlled conditions or natural
232 dehydration (**Tables 4, 5, and 6**, respectively). This section highlights the different conclusions
233 reported in the selected studies regarding the changes in phenolic acids during the winegrape
234 withering process. The results are discussed in an attempt to understand the causes and establish
235 trends.

236

237 **3.1. Evolution of phenolic acid content over withering: the result of a complex balance**

238 In all the studies considered, phenolic acids were somehow affected by the withering process,
239 but the analysis of the literature revealed different positions among authors about the balance between
240 concentration, hydrolysis, synthesis, and oxidation/catabolism processes in winegrapes.

241 Some authors considered the increase in the content of phenolic acids as a consequence of the
242 prevalence of the concentration effect on the degradation reactions. Among these authors, Marquez
243 et al. (2012), assessing grape juice from off-vine dehydration at a constant temperature of 40 °C in
244 Tempranillo and Merlot grapes (**Table 5**), reported a substantial increase in the contents of HBAs
245 and selected HCTAs due to water evaporation and, possibly, the release from solid portions of the
246 grapes during withering, even if some of these compounds were degraded through enzymatic

247 browning reactions. Additionally, Torchio et al. (2016) studied the evolution of phenolic compounds
248 in grape pulp over the off-vine dehydration process at two different controlled temperature conditions
249 (18 and 28 °C) on cultivars Avanà, Chatus, and Nebbiolo (**Table 5**) and concluded that contents of
250 HCTAs increased in withered grapes compared to those of fresh samples significantly for Chatus and
251 Nebbiolo winegrapes. These results were probably due to a balance in favor of the concentration
252 effect, as suggested by the findings of Frangipane et al. (2007). Later, to avoid the weight loss variable
253 during the withering process, Frangipane et al. (2012) expressed the content of HCA in Roschetto juice
254 as mg/number of berries (**Table 5**), demonstrating in their experiment how the degradation effect
255 prevailed over the concentration effect. Regarding HBAs, contents of syringic and gallic acids
256 decreased during the withering process when expressed as mg/L juice, in agreement with previous
257 observations on Aleatico grapes (Frangipane et al., 2007; **Table 5**).

258 Lukić et al. (2016) (**Table 4**), focusing on the change in the physicochemical composition of
259 Gewürztraminer wines as a result of late and icewine harvest, hypothesized that the higher amounts
260 of ethyl cinnamates found in these wines could have been due to the concentration effect of HCAs as
261 a consequence of water loss. Accordingly, Panceri et al. (2013) observed an increase in the contents
262 of HCAs (mainly caffeic acid) in Merlot and Cabernet sauvignon grapes (**Table 5**). These authors
263 suggested that off-vine withering under controlled conditions at 7 °C, and low relative humidity could
264 promote the activity of cinnamoyl esterase, similar to that in the maturation of wines in bottles, where
265 HCA esters are hydrolyzed to the free form. However, in the same study, the authors observed a
266 divergence in the HBA amounts for the two grape varieties: Cabernet sauvignon showed an increase
267 in the content of these compounds during withering, whereas Merlot showed a decrease, suggesting
268 a genotype effect rather than a change in the metabolic pathways. More recently, Nievierowski et al.
269 (2021), studying the impact of off-vine withering in a naturally ventilated room on the quality of
270 Merlot grapes, found a significant increase in phenolic acid amounts, both for HBAs and HCAs,
271 particularly for gallic, *p*-hydroxybenzoic, caffeic, and caftaric acids (**Table 6**). Additionally,
272 Constantinou et al. (2017) estimated a sixfold increase in the HCA content and total amounts of HBAs
273 in both traditional Cypriot cultivars Xynisteri and Mavro undergoing an off-vine sun-drying short

274 process (**Table 6**). Interestingly, these authors pointed out some differences among the contents of
275 single HCAs within the subgroup. Many significant changes were observed in individual phenolic
276 compounds in a comprehensive comparative study on Xynisteri grapes (Constantinou et al., 2018),
277 and the increasing trend was common for all phenolic acids, but with different intensities and
278 prevalence of singular compounds within the HBA and HCA classes concerning the various
279 postharvest withering methods tested (**Table 6**). Additionally, the authors interpreted the significant
280 increase in phenolic acid content observed during withering to be higher than that caused by the
281 concentration effect, using four alternative dehydration methods, namely, multiple horizontal wires,
282 multiple vertical pallets, low greenhouse, and hot-air dryer. Serratos et al. (2008a) also suggested
283 the possibility of an additional increase in the amounts of these compounds in ways other than
284 concentration. Studying Pedro Ximenez grapes withered at temperatures of 40–50 °C (**Table 6**), as
285 in the typical sweet wine production in southern Spanish regions, the authors observed a 188%
286 increase in *p*-hydroxybenzoic acid content, suggesting an additional increase in the amount of these
287 compounds during off-vine sun-drying. However, other phenolic acids (i.e., gallic acid and HCTAs)
288 showed simultaneous degradation, probably due to oxidation or polymerization processes, leading to
289 a final increase lower than expected. In these peculiar conditions of high temperature, several authors
290 have highlighted the occurrence of browning enzymatic reactions involving phenolic acids
291 (Figueiredo-González et al., 2014a; Marquez et al., 2012; Serratos et al., 2008b). It is assumed that
292 the first step of the PPO enzymatic browning reaction consists of the oxidation of caftaric acid
293 (Serratos et al., 2011), which is very sensitive to atmospheric variation, particularly related to water
294 stress and UV light (Bellincontro et al., 2009).

295 Although numerous studies have shown an increase in the contents of phenolic acids,
296 supporting the assumption of the prevalence of the concentration effect or suggesting the possibility
297 of new formation over the withering process, other studies pointed out a decrease in their contents in
298 both on-vine and off-vine, controlled or not, withering conditions. Among the off-vine withering
299 studies (**Table 5**), Negri et al. (2017) found a decrease in amounts of many grape metabolites,
300 including HCTAs, in noble-rot botrytized Garganega grapes, suggesting that they were degraded by

301 fungal metabolism. The loss of phenolic acids in winegrapes infected with noble rot has been reported
302 in other white varieties. A reduction in the content of caftaric acid (−69%) was observed in Chenin
303 blanc grapes (Carbajal-Ida et al., 2016), and of HCAs in skins of botrytized Chardonnay grapes (Hong
304 et al., 2012), probably because of laccase and tyrosinase activities that could be responsible for the
305 oxidation of HCTAs (Dubernet et al., 1977). Nevertheless, in the early stages of the phenylpropanoid
306 pathway during *Botrytis cinerea* infection, increased production of HCAs has been highlighted
307 (Blanco-Ulate et al., 2015; Kallitsounakis et al., 2020). These results confirmed the complexity of the
308 balance that determines the final content of phenolic acids in these special wines.

309 Piano et al. (2013) observed a very slight increase in the HCTA content in the Piedmontese
310 variety Uvalino undergoing off-vine withering in a room at a constant temperature of 24 °C and
311 relative humidity of 30% (**Table 5**); however, fewer differences were observed in the contents of
312 these compounds in overripened grapes, all during a 21-day withering period. However, Nicoletti et
313 al. (2013) observed a decrease in the contents of phenolic acids during off-vine withering under all
314 the controlled conditions tested on Nebbiolo grapes harvested from vines defoliated at different times
315 (**Table 5**), except for the non-defoliated sample withered at 10 °C, confirming how the agronomic
316 techniques also affect the features of the withered grapes and wines. As a probable explanation for
317 the decrease in the contents of esters of HCAs observed during the off-vine natural withering process
318 of Garnacha tintorera (**Table 6**), Figueiredo-González et al. (2013b) hypothesized the participation
319 of these compounds in degradation reactions owing to their high suitability as substrates in some
320 other types of reactions (e.g., copigmentation or PPO browning), as well as anthocyanin acylation for
321 the free forms, as previously presumed in studies performed on-vine by Ivanova et al. (2011) and off-
322 vine by Toffali et al. (2011) (**Tables 4 and 5**). Nevertheless, lower contents in grapes did not lead to
323 lower contents in the final wines. Rusjan et al. (2017), who studied for the first time the impact of the
324 on-vine cutting of 1-year old canes with fruitful shoots during ripening, –the so-called *double*
325 *maturation raisonnée* (DMR)– on Merlot grapes (**Table 4**), found the lowest concentration of HCAs
326 in DMR-subjected grape skins when compared to that of the control, despite the presence of high

327 amounts of these compounds in the wine made from those grapes. By contrast, Budić-Leto et al.
328 (2017), in a study on the differentiation between Croatian dessert wine Prošek and dry wines based
329 on phenolic composition (**Table 5**), found significantly lower concentrations of phenolic acids in
330 Prošek dessert wines than in the dry wines.

331 These heterogeneous results can be explained by several factors: first, it has been shown that
332 the positive or negative final balance of HCAs and HBAs depends on the genotype (Mayén et al.,
333 1997; Zenoni et al., 2016), and several authors have indicated that different varieties subjected to the
334 same withering conditions can be affected differently by the process (Kilmartin et al., 2007; Marquez
335 et al., 2012; Torchio et al., 2016). Additionally, a very important part of the phenomenon explanation
336 is certainly attributable to the different withering conditions applied, both on-vine (Avizcuri-Inac et
337 al. 2018; Mikulic-Petrovsek et al., 2017) and off-vine (Bellincontro et al., 2009; Constantinou et al.,
338 2018; Frangipane et al., 2007; Nicoletti et al., 2013). Furthermore, the rate of weight loss can
339 significantly affect the complex balance dynamics (Bonghi et al., 2012; Mencarelli et al., 2010;
340 Panceri et al., 2015).

341

342 **3.2. Grape genetic regulation**

343 During withering, berry tissues are metabolically active, and several specific-activated
344 processes may affect the grape compositional features (Rizzini et al., 2009). Several studies have
345 examined the specific changes in gene expression and the complex integration of transcriptomic,
346 metabolomic, and proteomic levels during grape withering (Di Carli et al., 2011; Zamboni et al.,
347 2008; Zamboni et al., 2010). However, little information is available on the weight of genetic
348 regulation in determining the final concentration of phenolic acids in withered grapes. During the
349 study of chemical and biochemical changes in the off-vine process of Aleatico grapes at different
350 temperatures and weight loss rates, Mencarelli et al. (2010) found an upregulation of the expression
351 of PAL gene, which is active at the beginning of the phenolic acid pathway, in berries exposed to
352 temperatures of 10 ° and 20 °C, over dehydration at 10% weight loss. However, transcript abundance
353 of the PAL gene did not correlate well with metabolite abundance. Additionally, Bonghi et al. (2012)

354 evaluated phenolic compound metabolism and gene expression in the skins of Raboso Piave
355 winegrapes under off-vine controlled withering conditions related to slow and rapid rates of up to
356 10% and 30% weight loss. These authors highlighted the importance of the complex expression
357 regulation of numerous members of the *Vitis vinifera* L. PAL multigene family as a key step in the
358 multiple physiological responses to environmental stresses. In a comprehensive genetic/metabolomic
359 study, Zenoni et al. (2016) highlighted the complexity of the metabolomic and transcriptomic changes
360 during postharvest dehydration. Even though they found an upregulation of the expression of genes
361 encoding important enzymes that catalyze the earliest steps of phenylpropanoid biosynthesis, the
362 negative or positive final balance of HCAs/HBAs in several winegrape cultivars (Corvina,
363 Sangiovese, Merlot, Syrah, Oseleta, and Cabernet sauvignon) ongoing withering process was
364 genotype-dependent. This balance was negative in Corvina. Furthermore, the six varieties studied
365 were characterized by distinct gene expression profiles and modulation intensities: Corvina berries
366 showed the strongest and fastest response to postharvest withering, whereas Cabernet sauvignon
367 showed the weakest and slowest gene induction. In accordance with these findings, more recently,
368 the same authors assessed two different thermo-hygrometric conditions of off-vine withering with
369 natural or forced airflow on cultivar Corvina and found no significant differences in the accumulation
370 of HCAs/HBAs and their derivatives in response to different environmental stresses (Zenoni et al.,
371 2020).

372 Regarding the decrease in the concentration of phenolic acids observed by some authors during
373 the grape withering process, a possible explanation could be found from a genetic point of view. In
374 the transcriptomic studies performed by Zamboni et al. (2008) on Corvina winegrapes, the authors
375 described the upregulation of the expression of a chalcone isomerase gene, suggesting the activation
376 of the flavonoid pathway during the withering process, and two tags of polyphenol oxidase, indicating
377 a probable oxidation/polymerization of phenolic compounds. In accordance with the results of
378 Zamboni et al., Di Carli et al. (2011) described an upregulation of the activity of reactive oxygen
379 species (ROS)-scavenging enzymes during the slow withering of Corvina winegrapes. In particular,
380 different trends in ROS accumulation were observed between the veraison and ripening phases, and

381 a decrease in the abundance of PPO involved in browning reactions was observed during withering
382 with respect to the veraison stage. Moreover, as mentioned above, some authors hypothesized that
383 the decrease in the contents of HCAs in the late harvest phase is a result of their functions in some
384 types of reactions such as the synthesis of acylated anthocyanins (Figueiredo-González et al., 2013b;
385 Ivanova et al., 2011; Toffali et al., 2011). Although some genes involved in the first part of
386 anthocyanin acylation mechanisms have been found in transcriptomic studies (Bonpart et al., 2018;
387 Zamboni et al., 2008), further studies are needed to gain a better understanding of the role of genetic
388 regulation in these reactions and their important chemical implications.

389

390 **3.3. Winemaking implications: extraction and evolution**

391 Qualitative and quantitative evolution of phenolic acids has also been observed during the
392 winemaking of withered grapes. In particular, Marquez et al. (2014) evaluated the antioxidant activity
393 in relation to the phenolic profile during the winemaking of Cabernet sauvignon grapes off-vine
394 withered at a constant temperature of 40 °C and initial relative humidity of 20%. They observed an
395 overall increase in the concentration of HBAs, probably due to the early diffusion from the solid parts
396 of the grapes to their pulp during the withering process and later during subsequent maceration of the
397 skins. The peaks of the amounts of these compounds were observed after 48 h of maceration, as was
398 observed for HCA esters. These results are consistent with those previously reported for Tempranillo
399 and Merlot grapes by Marquez et al. (2012), in which small amounts of HCA esters were present in
400 the final wines, possibly owing to degradation reactions toward free forms and their suitability as
401 substrates for major reactions during maceration.

402 Few studies have been conducted to evaluate the influence of the grape withering process on
403 the phenolic composition of wines during aging and bottle storage. Figueiredo-González et al.
404 (2014b) studied the phenolic composition of naturally sweet and sweet fortified Garnacha tintorera-
405 based wines during 6 and 12 months of oak aging and observed an increase in the amounts of HBAs
406 and HCAs in both wines. This increase was probably due to release from oak wood and transfer to
407 the wine during the aging period or a consequence of the decrease in the contents of coumaroyl

408 anthocyanin forms to release HCAs. The authors also highlighted a simultaneous decrease in HCTA
409 amounts in the wines, suggesting the occurrence of slow hydrolysis reactions during the aging period.
410 Additionally, the esters could have participated in other reactions such as polymerization or
411 copigmentation with anthocyanins. These findings are in agreement with the previous results reported
412 by Karagiannis et al. (2000), who conducted one of the first studies on this topic on cultivar Muscat
413 lefko grapes from the island of Samos.

414 Panceri & Bordignon-Luiz (2017) observed an increase in the concentration of phenolic acids
415 (gallic, protocatechuic, *p*-coumaric acids) during 22 months of bottle ageing for Cabernet sauvignon
416 and Merlot wines made from grapes withered under controlled conditions, in accordance with the
417 findings of Issa-Issa et al. (2020) in Fondillón wines made from Monastrell on-vine overripe grapes.
418 The increased availability of free forms of HCAs may result in potential risk for the production of
419 unpleasant flavors because of their suitability as substrates for spoilage microorganisms (Kheir et al.,
420 2013). This possible risk, enhanced by the presence of residual sugars, necessitates rigorous
421 microbiological control by applying appropriate inactivation measures. Nevertheless, wines made
422 from withered grapes are often characterized by higher alcohol content than that of other styles of
423 wine, making the proliferation of spoilage microorganisms more difficult (Malfeito-Ferreira, 2018).

424

425 **3.4. The case of icewines: a discussed topic**

426 The evolution of phenolic acid content during the production of icewines is a peculiar and often
427 discussed topic. The first study on this issue was conducted by Kilmartin et al. (2007) on the
428 polyphenol composition of Canadian icewines to identify the quality markers of authentic icewines.
429 For example, "Faux" Riesling and Vidal icewines (harvested above -8°C) had up to 11- and 21-times
430 higher concentrations of total HCAs than those of "Real" icewines (harvested at temperatures of -8°C
431 $^{\circ}\text{C}$ or below). These findings agree with those reported by Tian et al. (2009a) who introduced a further
432 kind of "Faux" icewine for cultivar Vidal: a refrigerator-frozen juice, artificially refrigerated at -8°C ,
433 and a concentrated juice made using rotary evaporation at 66°C . The artificially concentrated juice

434 presented a concentration of phenolic acids 2.1-times higher than that of the refrigerator-frozen juice,
435 and notably, approximately 3.6-times higher than that of naturally frozen juice. These differences
436 were quite preserved in the icewine types produced from them, with little influence from the yeast
437 strain used. Furthermore, the influence of the harvest date on the phenolic acid content of icewines
438 was discussed by the same authors. Kilmartin (2007) pointed out that the earliest harvest dates
439 produced icewines with higher concentrations of HCAs in Riesling and Vidal winegrapes harvested
440 in Canada in 1999, 2000, and 2002, while Tian et al. (2009b) argued that the contents of HBAs and
441 HCAs increased as grape harvest time was delayed to produce icewines from the Vidal variety
442 harvested in China in 2005 and 2006. With regard to the considerations previously made on the
443 influence of withering conditions, it should be recalled that the traditional icewine production
444 technique is an on-vine withering process and is thus subjected to changes due to weather conditions
445 (e.g., freeze-thaw cycles) that could markedly affect the balance of phenolic acids in the berries.

446 Avizcuri-Inac et al. (2018) focused their efforts on the chemical and sensorial characterization
447 of sweet wines to understand the influence of the dehydration process on the features of several wines
448 made from on-vine withered grapes. The authors analyzed sweet wines obtained by using different
449 varieties and techniques and classified them into three clusters based on the phenolic compound
450 characteristics. The first cluster, characterized by the highest contents of HBAs, HCAs, and other
451 phenolic compounds, such as flavonols, included a late harvest wine and a natural icewine (both
452 involving Tempranillo red grapes). The second cluster, with intermediate amounts of phenolic
453 compounds, included two natural icewines (one from white and one from red grapes), one artificial
454 icewine (from Tempranillo red grapes), and a *supurao* wine (from Tempranillo and Grenache off-
455 vine dehydrated red grapes). The third cluster, which was described by the lowest values of phenolic
456 compounds, particularly for HBAs, included four icewines obtained by grape freezing in the chamber,
457 three produced from white grapes and one from red grapes. Although these findings could have been
458 influenced by variety, environmental and technological factors (such as grape pressing), they show
459 that the extraction and concentration of these compounds produced by freezing winegrapes were
460 lower than those in wines belonging to the other two clusters, produced with more traditional

461 techniques. The concentration of phenolic acids has been associated with the quality and authenticity
462 of icewines (Kilmartin et al., 2007; Tang et al., 2013; Tian et al., 2009a, 2009b). However, new
463 studies are needed to better understand the link between weather conditions and phenolic acid features
464 in the production of these special wines. This alternative approach can be challenging and technically
465 difficult, but will have very useful practical applications, such as a more focused choice of the harvest
466 period.

467

468 **4. Conclusive remarks and future perspectives**

469 The aim of the present systematic review was to produce a concise overview of the extensive
470 literature available on the evolution of phenolic acids in withering grapes and the wines produced
471 from them. Although phenolic acids are structurally simple molecules, their evolution over the
472 withering process is complex. In most cases, the amount of phenolic acids increased after withering,
473 both at high and low temperatures. However, in some studies, their concentrations decreased or
474 remained almost constant in grapes. In these previous studies, the authors considered a possible cause
475 of the decrease as the high suitability of phenolic acids as substrates for other types of reactions (e.g.,
476 copigmentation or PPO browning). Indeed, it is of fundamental importance to remember that the final
477 contents of phenolic compounds in withered grapes and the resulting wines are strongly dependent
478 on the balance between concentration, synthesis, and loss or degradation of these compounds. This
479 balance is particularly evident for phenolic acids in botrytized grapes (reported in section 3.1). Even
480 if at the end of the process the overall balance resulted in the loss of phenolic acids, probably owing
481 to tyrosinase and laccase activities, in the early stages of the phenylpropanoid pathway during *Botrytis*
482 *cinerea* infection an increased production of HCAs was highlighted. Moreover, HCAs, HBAs, and
483 their derivatives are often affected differently by the withering process, and in some cases, even single
484 compounds within the same class can present different trends. The contents of these compounds are
485 affected by many factors, including genotype and withering conditions. There is still much to
486 understand about the weight of the genetic control of the withering process, and new studies are

487 needed to investigate the berry genetic response to environmental stresses and the related chemical
488 repercussions. In this sense, the modulation of the water loss rate and intensity may be used to increase
489 the concentration of specific compounds. This knowledge can help manage the withering process
490 according to the desired oenological objective.

491 The winemaking implications are very interesting and well-studied: HBA and HCA ester
492 release seems to start during withering, from the solid parts of the grapes to the pulp, and to continue
493 thereafter during maceration, with a peak after 48 h of skin maceration. The free forms of HCAs tend
494 to increase throughout winemaking and aging. This information may be useful for practical
495 applications, allowing the management of technological processes with improved awareness. On one
496 hand, the increase in the contents of these compounds could involve copigmentation effects,
497 improved antioxidant activity, and health benefits; on the other hand, it may lead to increased risks,
498 such as must browning or off-flavor production, because of their role as substrates for spoilage
499 microorganisms.

500 Finally, several authors associated the concentration of phenolic acids with a quality and
501 authenticity marker in icewines, but further research should be carried out to better understand the
502 link between the weather/environmental conditions and the phenolic acid features of winegrapes in
503 on-vine withering processes.

504

505

506 **Funding**

507 This research did not receive any specific grant from funding agencies in the public, commercial, or
508 not-for-profit sectors.

509

510 **References**

511 Avizcuri-Inac, J. M., González Hernández, M., Rosáenz-Oroz, D., Martínez Ruiz, R., & Vaquero-
512 Fernández, L. (2018). Chemical and sensory characterisation of sweet wines obtained by different

- 513 techniques. *Ciência e Técnica Vitivinícola*, 33(1), 15-30.
514 <https://doi.org/10.1051/ctv/20183301015>
- 515 Babbar, N., Oberoi, H. S., & Sandhu, S. K. (2015). Therapeutic and nutraceutical potential of
516 bioactive compounds extracted from fruit residues. *Critical Reviews in Food Science and*
517 *Nutrition*, 55(3), 319-337. <https://doi.org/10.1080/10408398.2011.653734>
- 518 Baderschneider, B., & Winterhalter, P. (2001). Isolation and characterization of novel benzoates,
519 cinnamates, flavonoids, and lignans from Riesling wine and screening for antioxidant
520 activity. *Journal of Agricultural and Food Chemistry*, 49(6), 2788-2798.
521 <https://doi.org/10.1021/jf010396d>
- 522 Bellincontro, A., Nicoletti, I., Valentini, M., Tomas, A., De Santis, D., Corradini, D., & Mencarelli,
523 F. (2009). Integration of nondestructive techniques with destructive analyses to study postharvest
524 water stress of winegrapes. *American Journal of Enology and Viticulture*, 60(1), 57-65.
525 <https://www.ajevonline.org/content/60/1/57.short>
- 526 Blanco-Ulate, B., Amrine, K. C., Collins, T. S., Rivero, R. M., Vicente, A. R., Morales-Cruz, A.,
527 Doyle, C. L., Ye, Z., Allen, G., Heymann, H., Ebeler, S. E., & Ebeler, S. E. (2015). Developmental
528 and metabolic plasticity of white-skinned grape berries in response to *Botrytis cinerea* during
529 noble rot. *Plant Physiology*, 169(4), 2422-2443. <https://doi.org/10.1104/pp.15.00852>
- 530 Bloomfield, D. G., Heatherbell, D. A., & Nikfardjam, M. P. (2003). Effect of p-coumaric acid on the
531 color in red wine. *Mitt. Klosterneuburg*, 53(5-6), 195-198.
- 532 Bonghi, C., Rizzini, F. M., Gambuti, A., Moio, L., Chkaiban, L., & Tonutti, P. (2012). Phenol
533 compound metabolism and gene expression in the skin of wine grape (*Vitis vinifera* L.) berries
534 subjected to partial postharvest dehydration. *Postharvest Biology and Technology*, 67, 102-109.
535 <https://doi.org/10.1016/j.postharvbio.2012.01.002>
- 536 Bontpart, T., Ferrero, M., Khater, F., Marlin, T., Vialet, S., Vallverdù-Queralt, A., Pinasseau, L.,
537 Ageorges, A., Cheynier, V. & Terrier, N. (2018). Focus on putative serine carboxypeptidase-like
538 acyltransferases in grapevine. *Plant Physiology and Biochemistry*, 130, 356-366.
539 <https://doi.org/10.1016/j.plaphy.2018.07.023>
- 540 Budić-Leto, I., Zdunić, G., Gajdoš-Kljusurić, J., Mucalo, A., & Vrhovšek, U. (2017). Differentiation
541 between Croatian dessert wine Prošek and dry wines based on phenolic composition. *Journal of*
542 *Food Composition and Analysis*, 62, 211-216. <https://doi.org/10.1016/j.jfca.2017.05.015>

- 543 Buiarelli, F., Coccioli, F., Merolle, M., Jasionowska, R., & Terracciano, A. (2010). Identification of
544 hydroxycinnamic acid–tartaric acid esters in wine by HPLC–tandem mass spectrometry. *Food*
545 *Chemistry*, *123*(3), 827-833. <https://doi.org/10.1016/j.foodchem.2010.05.017>
- 546 Carbajal-Ida, D., Maury, C., Salas, E., Siret, R., & Mehinagic, E. (2016). Physico-chemical properties
547 of botrytised Chenin blanc grapes to assess the extent of noble rot. *European Food Research and*
548 *Technology*, *242*(1), 117-126. <https://doi.org/10.1007/s00217-015-2523-x>
- 549 Campos, F. M., Couto, J. A., Figueiredo, A. R., Tóth, I. V., Rangel, A. O., & Hogg, T. A. (2009).
550 Cell membrane damage induced by phenolic acids on wine lactic acid bacteria. *International*
551 *Journal of Food Microbiology*, *135*(2), 144-151.
552 <https://doi.org/10.1016/j.ijfoodmicro.2009.07.031>
- 553 Carmona, L., Varela, J., Godoy, L., & Ganga, M. A. (2016). Comparative proteome analysis of
554 *Brettanomyces bruxellensis* under hydroxycinnamic acid growth. *Electronic Journal of*
555 *Biotechnology*, *23*, 37-43. <https://doi.org/10.1016/j.ejbt.2016.07.005>
- 556 Cheynier, V., Rigaud, J., Souquet, J. M., Barillere, J. M., & Moutounet, M. (1989a). Effect of pomace
557 contact and hyperoxidation on the phenolic composition and quality of Grenache and Chardonnay
558 wines. *American Journal of Enology and Viticulture*, *40*(1), 36-42.
- 559 Cheynier, V., Basire, N., & Rigaud, J. (1989b). Mechanism of trans-caffeoyltartaric acid and catechin
560 oxidation in model solutions containing grape polyphenoloxidase. *Journal of Agricultural and*
561 *Food Chemistry*, *37*(4), 1069-1071. <https://hal.inrae.fr/hal-02722125>
- 562 Corradini, D., & Nicoletti, I. (2013). Changes in phenolic compounds. In Mencarelli, F., Tonutti, P.
563 (Eds.) *Sweet, Reinforced and Fortified Wines: Grape Biochemistry, Technology and Vinification*
564 (pp. 105-119). Chichester: John Wiley & Sons Ltd. <https://doi.org/10.1002/9781118569184.ch6>
- 565 Constantinou, S., Gómez-Caravaca, A. M., Goulas, V., Segura-Carretero, A., & Manganaris, G. A.
566 (2017). Metabolic fingerprinting of must obtained from sun-dried grapes of two indigenous
567 Cypriot cultivars destined for the production of ‘Commandaria’: A protected designation of
568 origin product. *Food Research International*, *100*, 469-476.
569 <https://doi.org/10.1016/j.foodres.2016.11.015>
- 570 Constantinou, S., Gómez-Caravaca, A. M., Goulas, V., Segura-Carretero, A., Koundouras, S., &
571 Manganaris, G. A. (2018). The impact of postharvest dehydration methods on qualitative attributes
572 and chemical composition of ‘Xynisteri’ grape (*Vitis vinifera*) must. *Postharvest Biology and*
573 *Technology*, *135*, 114-122. <https://doi.org/10.1016/j.postharvbio.2017.09.005>

- 574 D'Onofrio, C., Bellincontro, A., Accordini, D., & Mencarelli, F. (2019). Malic Acid as a potential
575 marker for the aroma compounds of Amarone winegrape varieties in withering. *American Journal*
576 *of Enology and Viticulture*, 70(3), 259-266. <https://doi.org/10.5344/ajev.2019.18071>
- 577 De Rosso, M., Soligo, S., Panighel, A., Carraro, R., Vedova, A. D., Maoz, I., Tomasi, D. & Flamini,
578 R. (2016). Changes in grape polyphenols (*V. vinifera* L.) as a consequence of post- harvest
579 withering by high- resolution mass spectrometry: Raboso Piave versus Corvina. *Journal of Mass*
580 *Spectrometry*, 51(9), 750-760. <https://doi.org/10.1002/jms.3835>
- 581 Di Carli, M., Zamboni, A., Pe, M. E., Pezzotti, M., Lilley, K. S., Benvenuto, E., & Desiderio, A.
582 (2011). Two-dimensional differential in gel electrophoresis (2D-DIGE) analysis of grape berry
583 proteome during postharvest withering. *Journal of Proteome Research*, 10(2), 429-446.
584 <https://doi.org/10.1021/pr1005313>
- 585 Dubernet, M., Ribereau-Gayon, P., Lerner, H. R., Harel, E., & Mayer, A. M. (1977). Purification and
586 properties of laccase from *Botrytis cinerea*. *Phytochemistry*, 16(2), 191-193.
587 [https://doi.org/10.1016/S0031-9422\(00\)86783-7](https://doi.org/10.1016/S0031-9422(00)86783-7)
- 588 Esmaili, M., Sotudeh-Gharebagh, R., Cronin, K., Mousavi, M. A. E., & Rezazadeh, G. (2007). Grape
589 drying: a review. *Food Reviews International*, 23(3), 257-280.
590 <https://doi.org/10.1080/87559120701418335>
- 591 Ferrandino A., Carra A., Rolle L., Schneider A., Schubert A. (2012). Profiling of hydroxycinnamoyl
592 tartrates and of acylated anthocyanins in the skin of 34 *Vitis vinifera* genotypes. *Journal of*
593 *Agricultural and Food Chemistry*, 60, 4931-4945. <https://doi.org/10.1021/jf2045608>
- 594 Ferrer-Gallego, R., Hernández-Hierro, J. M., Rivas-Gonzalo, J. C., & Escribano-Bailón, M. T.
595 (2014). Sensory evaluation of bitterness and astringency sub-qualities of wine phenolic
596 compounds: synergistic effect and modulation by aromas. *Food Research International*, 62, 1100-
597 1107. <https://doi.org/10.1016/j.foodres.2014.05.049>
- 598 Ferrer-Gallego, R., Hernández-Hierro, J. M., Brás, N. F., Vale, N., Gomes, P., Mateus, N., ... &
599 Escribano-Bailón, M. T. (2017). Interaction between wine phenolic acids and salivary proteins by
600 saturation-transfer difference nuclear magnetic resonance spectroscopy (STD-NMR) and
601 molecular dynamics simulations. *Journal of Agricultural and Food Chemistry*, 65(31), 6434-
602 6441. <https://doi.org/10.1021/acs.jafc.6b05414>
- 603 Figueiredo-González, M., Cancho-Grande, B., & Simal-Gándara, J. (2013a). Effects on colour and
604 phenolic composition of sugar concentration processes in dried-on or dried-off-vine grapes and

- 605 their aged or not natural sweet wines. *Trends in Food Science & Technology*, 31(1), 36-54.
606 <https://doi.org/10.1016/j.tifs.2013.02.004>
- 607 Figueiredo-González, M., Cancho-Grande, B., & Simal-Gándara, J. (2013b). Evolution of colour and
608 phenolic compounds during Garnacha Tintorera grape raisining. *Food Chemistry*, 141(3), 3230-
609 3240. <https://doi.org/10.1016/j.foodchem.2013.05.142>
- 610 Figueiredo-González, M., Regueiro, J., Cancho-Grande, B., & Simal-Gándara, J. (2014a). Garnacha
611 Tintorera-based sweet wines: Detailed phenolic composition by HPLC/DAD–ESI/MS
612 analysis. *Food Chemistry*, 143, 282-292. <https://doi.org/10.1016/j.foodchem.2013.07.120>
- 613 Figueiredo-González, M., Cancho-Grande, B., Simal-Gándara, J., Teixeira, N., Mateus, N., & De
614 Freitas, V. (2014b). The phenolic chemistry and spectrochemistry of red sweet wine-making and
615 oak-aging. *Food Chemistry*, 152, 522-530. <https://doi.org/10.1016/j.foodchem.2013.12.018>
- 616 Frangipane, M. T., Ceccarelli, A., Mencarelli, F., & Anelli, G. (2007). Study of phenolic compounds
617 in Aleatico grapes dried in a forced air tunnel. *Italian Journal of Food Science*, 19(2), 205-210.
- 618 Frangipane, M. T., Torresi, S., De Santis, D., & Massantini, R. (2012). Effect of drying process in
619 chamber at controlled temperature on the grape phenolic compounds. *Italian Journal of Food
620 Science*, 24(1), 26-31.
- 621 Garrido, J., & Borges, F. (2013). Wine and grape polyphenols - A chemical perspective. *Food
622 Research International*, 54(2), 1844-1858. <https://doi.org/10.1016/j.foodres.2013.08.002>
- 623 Gonzalo-Diago, A., Dizy, M., & Fernández-Zurbano, P. (2014). Contribution of low molecular
624 weight phenols to bitter taste and mouthfeel properties in red wines. *Food Chemistry*, 154, 187-
625 198. <https://doi.org/10.1016/j.foodchem.2013.12.096>
- 626 Güntert, M., Rapp, A., Takeoka, G. R., & Jennings, W. (1986). HRGC and HRGC-MS applied to
627 wine constituents of lower volatility. *Zeitschrift für Lebensmittel-Untersuchung und
628 Forschung*, 182(3), 200-204. <https://doi.org/10.1007/BF01042128>
- 629 Harris, V., Jiranek, V., Ford, C. M., & Grbin, P. R. (2010). Inhibitory effect of hydroxycinnamic
630 acids on *Dekkera spp.* *Applied Microbiology and Biotechnology*, 86(2), 721-729.
631 <https://doi.org/10.1007/s00253-009-2352-6>
- 632 Harrison, R. (2018). Practical interventions that influence the sensory attributes of red wines related
633 to the phenolic composition of grapes: a review. *International Journal of Food Science &
634 Technology*, 53(1), 3-18. <https://doi.org/10.1111/ijfs.13480>

- 635 He, F., Liang, N. N., Mu, L., Pan, Q. H., Wang, J., Reeves, M. J., & Duan, C. Q. (2012). Anthocyanins
636 and their variation in red wines II. Anthocyanin derived pigments and their color
637 evolution. *Molecules*, *17*(2), 1483-1519. <https://doi.org/10.3390/molecules17021483>
- 638 Hong, Y. S., Martinez, A., Liger-Belair, G., Jeandet, P., Nuzillard, J. M., & Cilindre, C. (2012).
639 Metabolomics reveals simultaneous influences of plant defence system and fungal growth in
640 *Botrytis cinerea*-infected *Vitis vinifera* cv. Chardonnay berries. *Journal of Experimental*
641 *Botany*, *63*(16), 5773-5785. <https://doi.org/10.1093/jxb/ers228>
- 642 Hufnagel, J. C., & Hofmann, T. (2008). Quantitative reconstruction of the nonvolatile
643 sensometabolome of a red wine. *Journal of Agricultural and Food Chemistry*, *56*(19), 9190-9199.
644 <https://doi.org/10.1021/jf801742w>
- 645 Issa-Issa, H., Guclu, G., Noguera-Artiaga, L., López-Lluch, D., Poveda, R., Kelebek, H., Selli, S., &
646 Carbonell-Barrachina, Á.A. (2020). Aroma-active compounds, sensory profile, and phenolic
647 composition of Fondillón. *Food Chemistry*, *316*, 126-353.
648 <https://doi.org/10.1016/j.foodchem.2020.126353>
- 649 Ivanova, V., Stefova, M., Vojnoski, B., Dörnyei, Á., Márk, L., Dimovska, V., Stafilov, T. & Kilar, F.
650 (2011). Identification of polyphenolic compounds in red and white grape varieties grown in R.
651 Macedonia and changes of their content during ripening. *Food Research International*, *44*(9),
652 2851-2860. <https://doi.org/10.1016/j.foodres.2011.06.046>
- 653 Kallithraka, S., Salacha, M. I., & Tzourou, I. (2009). Changes in phenolic composition and
654 antioxidant activity of white wine during bottle storage: Accelerated browning test versus bottle
655 storage. *Food Chemistry*, *113*(2), 500-505. <https://doi.org/10.1016/j.foodchem.2008.07.083>
- 656 Kallitsounakis, G., & Catarino, S. (2020). An overview on botrytized wines. *Ciência e Técnica*
657 *Vitivinicola*, *35*(2), 76-106. <http://hdl.handle.net/10400.5/20500>
- 658 Karagiannis, S., Economou, A., & Lanaridis, P. (2000). Phenolic and volatile composition of wines
659 made from *Vitis vinifera* cv. Muscat lefko grapes from the island of Samos. *Journal of Agricultural*
660 *and Food Chemistry*, *48*(11), 5369-5375. <https://doi.org/10.1021/jf000459c>
- 661 Kheir, J., Salameh, D., Strehaiano, P., Brandam, C., & Lteif, R. (2013). Impact of volatile phenols
662 and their precursors on wine quality and control measures of *Brettanomyces/Dekkera*
663 yeasts. *European Food Research and Technology*, *237*(5), 655-671.
664 <https://doi.org/10.1007/s00217-013-2036-4>

- 665 Kilmartin, P. A., Reynolds, A. G., Pagay, V., Nurgel, C., & Johnson, R. (2007). Polyphenol content
666 and browning of Canadian icewines. *Journal of Food Agriculture and Environment*, 5(3/4), 52.
667 <https://pubag.nal.usda.gov/catalog/723469>
- 668 Kunkee, R. E., & Amerine, M. A. (1968). Sugar and alcohol stabilization of yeast in sweet
669 wine. *Applied Microbiology*, 16(7), 1067-1075.
- 670 Laura, A., Moreno-Escamilla, J. O., Rodrigo-García, J., & Alvarez-Parrilla, E. (2019). Phenolic
671 compounds. In *Postharvest physiology and biochemistry of fruits and vegetables* (pp. 253-271).
672 Woodhead Publishing. <https://doi.org/10.1016/B978-0-12-813278-4.00012-9>
- 673 Loizzo, M. R., Bonesi, M., Di Lecce, G., Boselli, E., Tundis, R., Pugliese, A., Menichini, F. & Frega,
674 N. G. (2013). Phenolics, aroma profile, and in vitro antioxidant activity of Italian dessert passito
675 wine from Saracena (Italy). *Journal of Food Science*, 78(5), C703-C708.
676 <https://doi.org/10.1111/1750-3841.12110>
- 677 Lukić, I., Radeka, S., Grozaj, N., Staver, M., & Peršurić, Đ. (2016). Changes in physico-chemical
678 and volatile aroma compound composition of Gewürztraminer wine as a result of late and ice
679 harvest. *Food chemistry*, 196, 1048-1057. <https://doi.org/10.1016/j.foodchem.2015.10.061>
- 680 Malfeito-Ferreira, M. (2018). Two decades of “horse sweat” taint and *Brettanomyces* yeasts in wine:
681 Where do we stand now?. *Beverages*, 4(2), 32. <https://doi.org/10.3390/beverages4020032>
- 682 Marquez, A., Serratosa, M. P., Lopez-Toledano, A., & Merida, J. (2012). Colour and phenolic
683 compounds in sweet red wines from Merlot and Tempranillo grapes chamber-dried under
684 controlled conditions. *Food Chemistry*, 130(1), 111-120.
685 <https://doi.org/10.1016/j.foodchem.2011.07.010>
- 686 Marquez, A., Serratosa, M. P., & Merida, J. (2014). Antioxidant activity in relation to the phenolic
687 profile during the winemaking of sweet wines *Vitis vinifera* cv. Cabernet Sauvignon. *International*
688 *Journal of Food Science & Technology*, 49(9), 2128-2135. <https://doi.org/10.1111/ijfs.12521>
- 689 Mayén, M., Barón, R., Mérida, J., & Medina, M. (1997). Changes in phenolic compounds during
690 accelerated browning in white wines from cv. Pedro Ximenez and cv. Baladi grapes. *Food*
691 *Chemistry*, 58(1-2), 89-95. [https://doi.org/10.1016/S0308-8146\(96\)00218-X](https://doi.org/10.1016/S0308-8146(96)00218-X)
- 692 Mencarelli, F., Bellincontro, A., Nicoletti, I., Cirilli, M., Muleo, R., & Corradini, D. (2010). Chemical
693 and biochemical change of healthy phenolic fractions in winegrape by means of postharvest

- 694 dehydration. *Journal of Agricultural and Food Chemistry*, 58(13), 7557-7564.
695 <https://doi.org/10.1021/jf100331z>
- 696 Mencarelli, F., & Tonutti, P. (2013). *Sweet, reinforced and fortified wines: Grape biochemistry,*
697 *technology and vinification* (1st ed.). Chichester: John Wiley & Sons Ltd.
- 698 Mikulic-Petrovsek, M. M., Jug, T., Rescic, J., & Rusjan, D. (2017). Effects of partial dehydration
699 techniques on the metabolite composition in 'Refosk' grape berries and wine. *Turkish Journal of*
700 *Agriculture and Forestry*, 41(1), 10-22. [https://journals.tubitak.gov.tr/agriculture/issues/tar-17-](https://journals.tubitak.gov.tr/agriculture/issues/tar-17-41-1/tar-41-1-2-1609-65.pdf)
701 [41-1/tar-41-1-2-1609-65.pdf](https://journals.tubitak.gov.tr/agriculture/issues/tar-17-41-1/tar-41-1-2-1609-65.pdf)
- 702 Moher, D., Liberati, A., Tetzlaff, J., Altman, D.G., The PRISMA Group (2009). Preferred Reporting
703 Items for Systematic Reviews and Meta-Analyses: The PRISMA Statement. *PLoS Medicine*, 6(7):
704 e1000097. <https://doi.org/10.1371/journal.pmed1000097>
- 705 Monagas, M., Bartolome, B., & Gomez-Cordoves, C. (2005). Updated knowledge about the presence
706 of phenolic compounds in wine. *Critical Reviews in Food Science and Nutrition*, 45(2), 85-118.
707 <https://doi.org/10.1080/10408690490911710>
- 708 Morales-Prieto, N., Huertas-Abril, P. V., de Lerma, N. L., Pacheco, I. L., Pérez, J., Peinado, R., &
709 Abril, N. (2020). Pedro Ximenez sun-dried grape must: a dietary supplement for a healthy
710 longevity. *Food & Function*, 11, 4387-4402. <https://doi.org/10.1039/D0FO00204F>
- 711 Negri, S., Lovato, A., Boscaini, F., Salvetti, E., Torriani, S., Commisso, M., Danzi, R., Ugliano, M.,
712 Polverari, A., Tornielli, G.B. & Guzzo, F. (2017). The induction of noble rot (*Botrytis cinerea*)
713 infection during postharvest withering changes the metabolome of grapevine berries (*Vitis vinifera*
714 L., cv. Garganega). *Frontiers in Plant Science*, 8, 1002. <https://doi.org/10.3389/fpls.2017.01002>
- 715 Nicoletti, I., Bellincontro, A., De Rossi, A., De Sanctis, F., Tiberi, D., Pietromarchi, P., Botondi, R.,
716 Corradini, D. & Mencarelli, F. (2013). Postharvest dehydration of Nebbiolo grapes grown at
717 altitude is affected by time of defoliation. *Australian Journal of Grape and Wine Research*, 19(3),
718 358-368. <https://doi.org/10.1111/ajgw.12042>
- 719 Niculescu, V. C., Paun, N., & Ionete, R. E. (2018). *The Evolution of Polyphenols from Grapes to*
720 *Wines*. Justino, G.C. (Eds.); IntechOpen: London, UK.
- 721 Niewierowski, T. H., Veras, F. F., Silveira, R. D., Dachery, B., Hernandes, K. C., Lopes, F. C., ... &
722 Welke, J. E. (2021). Role of partial dehydration in a naturally ventilated room on the mycobiota,

- 723 ochratoxins, volatile profile and phenolic composition of Merlot grapes intended for wine
724 production. *Food Research International*, 110145. <https://doi.org/10.1016/j.foodres.2021.110145>
- 725 Okamura, S., & Watanabe, M. (1981). Determination of phenolic cinnamates in white wine and their
726 effect on wine quality. *Agricultural and Biological Chemistry*, 45(9), 2063-2070.
727 <https://doi.org/10.1271/bbb1961.45.2063>
- 728 Ong, B. Y., & Nagel, C. W. (1978). High-pressure liquid chromatographic analysis of
729 hydroxycinnamic acid—tartaric acid esters and their glucose esters in *Vitis vinifera*. *Journal of*
730 *Chromatography A*, 157, 345-355. [https://doi.org/10.1016/S0021-9673\(00\)92352-5](https://doi.org/10.1016/S0021-9673(00)92352-5)
- 731 Panceri, C. P., Gomes, T. M., De Gois, J. S., Borges, D. L., & Bordignon-Luiz, M. T. (2013). Effect
732 of dehydration process on mineral content, phenolic compounds and antioxidant activity of
733 Cabernet Sauvignon and Merlot grapes. *Food Research International*, 54(2), 1343-1350.
734 <https://doi.org/10.1016/j.foodres.2013.10.016>
- 735 Panceri, C. P., De Gois, J. S., Borges, D. L., & Bordignon-Luiz, M. T. (2015). Effect of grape
736 dehydration under controlled conditions on chemical composition and sensory characteristics of
737 Cabernet sauvignon and Merlot wines. *LWT-Food Science and Technology*, 63(1), 228-235.
738 <https://doi.org/10.1016/j.lwt.2015.02.014>
- 739 Panceri, C. P., & Bordignon- Luiz, M. T. (2017). Impact of grape dehydration process on the
740 phenolic composition of wines during bottle ageing. *Journal of Food Biochemistry*, 41(6), e12417.
741 <https://doi.org/10.1111/jfbc.12417>
- 742 Piano, F., Bertolone, E., Pes, D., Asproudi, A., & Borsa, D. (2013). Focusing on bioactive compounds
743 in grapes: stilbenes in Uvalino cv. *European Food Research and Technology*, 237(6), 897-904.
744 <https://doi.org/10.1007/s00217-013-2060-4>
- 745 Revilla, I., & Gonzalez-SanJose, M. L. (2003). Compositional changes during the storage of red
746 wines treated with pectolytic enzymes: Low molecular-weight phenols and flavan-3-ol derivative
747 levels. *Food Chemistry*, 80(2), 205–214. [https://doi.org/10.1016/S0308-8146\(02\)00255-8](https://doi.org/10.1016/S0308-8146(02)00255-8)
- 748 Río Segade, S., Torchio, F., Giacosa, S., Ricauda Aimonino, D., Gay, P., Lambri, M., Dordoni, R.,
749 Gerbi, V., & Rolle, L. (2014). Impact of several pre-treatments on the extraction of phenolic
750 compounds in winegrape varieties with different anthocyanin profiles and skin mechanical
751 properties. *Journal of Agricultural and Food Chemistry*, 62(33), 8437-8451.
752 <https://doi.org/10.1021/jf502656d>

- 753 Rizzini, F. M., Bonghi, C., & Tonutti, P. (2009). Postharvest water loss induces marked changes in
754 transcript profiling in skins of wine grape berries. *Postharvest Biology and Technology*, 52(3),
755 247-253. <https://doi.org/10.1016/j.postharvbio.2008.12.004>
- 756 Rolle, L., Torchio, F., Giacosa, S., & Gerbi, V. (2009). Modifications of mechanical characteristics
757 and phenolic composition in berry skins and seeds of Mondeuse winegrapes throughout the on-
758 vine drying process. *Journal of the Science of Food and Agriculture*, 89(11), 1973-1980.
759 <https://doi.org/10.1002/jsfa.3686>
- 760 Rolle, L., Giacosa, S., Río Segade, S., Ferrarini, R., Torchio, F., & Gerbi, V. (2013). Influence of
761 different thermohygro-metric conditions on changes in instrumental texture properties and phenolic
762 composition during postharvest withering of 'Corvina' winegrapes (*Vitis vinifera* L.). *Drying*
763 *Technology*, 31(5), 549-564. <https://doi.org/10.1080/07373937.2012.745092>
- 764 Rusjan, D., & Mikulic- Petkovsek, M. (2017). Double maturation raisonnée: the impact of on- vine
765 berry dehydration on the berry and wine composition of Merlot (*Vitis vinifera* L.). *Journal of the*
766 *Science of Food and Agriculture*, 97(14), 4835-4846. <https://doi.org/10.1002/jsfa.8354>
- 767 Sabel, A., Bredefeld, S., Schlander, M., & Claus, H. (2017). Wine phenolic compounds:
768 Antimicrobial properties against yeasts, lactic acid and acetic acid bacteria. *Beverages*, 3(3), 29.
769 <https://doi.org/10.3390/beverages3030029>
- 770 Salagoïty- Auguste, M. H., & Bertrand, A. (1984). Wine phenolics—analysis of low molecular
771 weight components by high performance liquid chromatography. *Journal of the Science of Food*
772 *and Agriculture*, 35(11), 1241-1247. <https://doi.org/10.1002/jsfa.2740351116>
- 773 Scienza, A. (2013). Sweet wines: The essence of European civilization. In Mencarelli, F., Tonutti, P.,
774 (Eds.) *Sweet, Reinforced and Fortified Wines* (pp. 5-25). Chichester: John Wiley & Sons Ltd.
775 <https://doi.org/10.1002/9781118569184>
- 776 Serratos, M. P., Lopez-Toledano, A., Merida, J., & Medina, M. (2008a). Changes in color and
777 phenolic compounds during the raisining of grape cv. Pedro Ximenez. *Journal of Agricultural and*
778 *Food Chemistry*, 56(8), 2810-2816. <https://doi.org/10.1021/jf073278k>
- 779 Serratos, M. P., Lopez-Toledano, A., Medina, M., & Merida, J. (2008b). Drying of Pedro Ximenez
780 grapes in chamber at controlled temperature and with dipping pretreatments. Changes in the color
781 fraction. *Journal of Agricultural and Food Chemistry*, 56(22), 10739-10746.
782 <https://doi.org/10.1021/jf8021767>

- 783 Serratos, M. P., Lopez-Toledano, A., Medina, M., & Merida, J. (2011). Characterisation of the
784 colour fraction of Pedro Ximenez Andalusian sweet wines. *South African Journal of Enology and*
785 *Viticulture*, 32(1), 155-163. <http://www.sawislibrary.co.za/dbtextimages/72572.pdf>
- 786 Somers, T. C., Vérette, E., & Pocock, K. F. (1987). Hydroxycinnamate esters of *Vitis vinifera*:
787 changes during white vinification, and effects of exogenous enzymic hydrolysis. *Journal of the*
788 *Science of Food and Agriculture*, 40(1), 67-78. <https://doi.org/10.1002/jsfa.2740400109>
- 789 Stivala, M. G., Vilecco, M. B., Enriz, D., & Fernández, P. A. (2017). Effect of phenolic compounds
790 on viability of wine spoilage lactic acid bacteria. A structure-activity relationship study. *American*
791 *Journal of Enology and Viticulture*, 68(2), 228-233. <https://doi.org/10.5344/ajev.2016.16084>
- 792 Tang, K., Li, J. M., Wang, B., Ma, L., & Xu, Y. (2013). Evaluation of nonvolatile flavor compounds
793 in Vidal icewine from China. *American journal of enology and viticulture*, 64(1), 110-117.
794 <https://doi.org/10.5344/ajev.2012.12037>
- 795 Tian, R. R., Li, G., Wan, S. B., Pan, Q. H., Zhan, J. C., Li, J. M., Zhang, Q. H. & Huang, W. D.
796 (2009a). Comparative study of 11 phenolic acids and five flavan-3-ols in cv. Vidal: impact of
797 natural icewine making versus concentration technology. *Australian Journal of Grape and Wine*
798 *Research*, 15(3), 216-222. <https://doi.org/10.1111/j.1755-0238.2009.00055.x>
- 799 Tian, R. R., Pan, Q. H., Zhan, J. C., Li, J. M., Wan, S. B., Zhang, Q. H., & Huang, W. D. (2009b).
800 Comparison of phenolic acids and flavan-3-ols during wine fermentation of grapes with different
801 harvest times. *Molecules*, 14(2), 827-838. <https://doi.org/10.3390/molecules14020827>
- 802 Toffali, K., Zamboni, A., Anesi, A., Stocchero, M., Pezzotti, M., Levi, M., & Guzzo, F. (2011). Novel
803 aspects of grape berry ripening and post-harvest withering revealed by untargeted LC-ESI-MS
804 metabolomics analysis. *Metabolomics*, 7(3), 424-436. <https://doi.org/10.1007/s11306-010-0259-y>
- 805 Torchio, F., Urcan, D. E., Lin, L., Gerbi, V., Giacosa, S., Rìo Segade, S., Pop, N., Lambri, M. &
806 Rolle, L. (2016). Influence of different withering conditions on phenolic composition of Avanà,
807 Chatus and Nebbiolo grapes for the production of 'Reinforced' wines. *Food Chemistry*, 194, 247-
808 256. <https://doi.org/10.1016/j.foodchem.2015.08.009>
- 809 Vanbeneden, N., Gils, F., Delvaux, F., & Delvaux, F. R. (2008). Formation of 4-vinyl and 4-ethyl
810 derivatives from hydroxycinnamic acids: Occurrence of volatile phenolic flavour compounds in
811 beer and distribution of Pad1-activity among brewing yeasts. *Food Chemistry*, 107(1), 221-230.
812 <https://doi.org/10.1016/j.foodchem.2007.08.008>

- 813 Vanzo, A., Cecotti, R., Vrhovsek, U., Torres, A. M., Mattivi, F., & Passamonti, S. (2007). The fate
814 of trans-caftaric acid administered into the rat stomach. *Journal of Agricultural and Food*
815 *Chemistry*, 55(4), 1604-1611. <https://doi.org/10.1021/jf0626819>
- 816 Vérette, E., Noble, A. C., & Somers, T. C. (1988). Hydroxycinnamates of *Vitis vinifera*: Sensory
817 assessment in relation to bitterness in white wines. *Journal of the Science of Food and*
818 *Agriculture*, 45(3), 267-272. <https://doi.org/10.1002/jsfa.2740450310>
- 819 Wang, X. J., Li, Y. K., Song, H. C., Tao, Y. S., & Russo, N. (2020). Phenolic matrix effect on aroma
820 formation of terpenes during simulated wine fermentation–Part I: Phenolic acids. *Food*
821 *Chemistry*, 341, 128288. <https://doi.org/10.1016/j.foodchem.2020.128288>
- 822 Waterhouse, A. L., & Teissedre, P. L. (1997). Levels of phenolics in California varietal wines. *ACS*
823 *Symposium Series*, 661, 12-23. <https://doi.org/10.1021/bk-1997-0661.ch003>
- 824 Waterhouse, A. L. (2002). Wine phenolics. *Annals of the New York Academy of Sciences*, 957(1), 21-
825 36. <https://doi.org/10.1111/j.1749-6632.2002.tb02903.x>
- 826 Zamboni, A., Minoia, L., Ferrarini, A., Tornielli, G. B., Zago, E., Delledonne, M., & Pezzotti, M.
827 (2008). Molecular analysis of post-harvest withering in grape by AFLP transcriptional
828 profiling. *Journal of Experimental Botany*, 59(15), 4145-4159. <https://doi.org/10.1093/jxb/ern256>
- 829 Zamboni, A., Di Carli, M., Guzzo, F., Stocchero, M., Zenoni, S., Ferrarini, A., Tononi, P., Toffali,
830 K., Desiderio, A., Lilley, K. S. & Pè, M. E. (2010). Identification of putative stage-specific
831 grapevine berry biomarkers and omics data integration into networks. *Plant Physiology*, 154(3),
832 1439-1459. <https://doi.org/10.1104/pp.110.160275>
- 833 Zenoni, S., Fasoli, M., Guzzo, F., Dal Santo, S., Amato, A., Anesi, A., Commisso, M., Herderich,
834 M., Ceoldo, S., Avesani, L., Pezzotti, M., & Tornielli, G.B. (2016). Disclosing the molecular basis
835 of the postharvest life of berry in different grapevine genotypes. *Plant Physiology*, 172(3), 1821-
836 1843. <https://doi.org/10.1104/pp.16.00865>
- 837 Zenoni, S., Amato, A., D'Inca, E., Guzzo, F., & Tornielli, G. B. (2020). Rapid dehydration of grape
838 berries dampens the post-ripening transcriptomic program and the metabolite profile
839 evolution. *Horticulture Research*, 7(1), 1-15. <https://doi.org/10.1038/s41438-020-00362-5>
- 840 Zoccatelli, G., Zenoni, S., Savoi, S., Dal Santo, S., Tononi, P., Zandonà, V., Dal Cin, A., Pezzotti,
841 M., & Tornielli, G. B. (2013). Skin pectin metabolism during the postharvest dehydration of

842 berries from three distinct grapevine cultivars. *Australian Journal of Grape and Wine*
843 *Research*, 19(2), 171-179. <https://doi.org/10.1111/ajgw.12014>

844 Zou, H., Kilmartin, P. A., Inglis, M. J., & Frost, A. (2002). Extraction of phenolic compounds during
845 vinification of Pinot Noir wine examined by HPLC and cyclic voltammetry. *Australian Journal*
846 *of Grape and Wine Research*, 8(3), 163-174. <https://doi.org/10.1111/j.1755-0238.2002.tb00252.x>

847

Journal Pre-proof

848 **FIGURE CAPTION**

849 **Figure 1** –Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) flow
850 diagram of the study selection process: evolution of phenolic acids in winegrapes undergoing a
851 withering process.

852

Journal Pre-proof

853 **Table 1** – Identification of the question key elements (PICO).

Key element	Response
P – population of interest	Phenolic acids of withered winegrape
I – intervention of interest	Withering process
C – comparator	Phenolic acids of fresh winegrape
O – outcome	Changes and evolution of phenolic acids

854

Journal Pre-proof

855 **Table 2** – List of keywords useful to construct strings with Boolean operators.

Keyword	Synonyms	Plural/singular forms	Alternative words/different spellings
Withering			Dehydration, drying, raisining
Withered grapes		Withered grape	Dehydrated grapes, dried grapes, raisin grapes
Grape	Winegrape	Grapes, winegrapes	
Sweet wines	Passito wines, withered wines, raisin wines,	Sweet wine	Straw wines, Icewines , Reinforced wines, Fortified wines, Dessert wines, Botrytized wines, Marsala, Porto, Sauternes, Tokaji, Sforzato, Sfusat, Amarone, Recioto, Aleatico, Vin Santo, Vin de Paille
Phenolic acids			HCA, HCAs, HCTAs, HBA, HBAs, hydroxycinnamic acids, hydroxycinnamics. hydroxybenzoic acids, hydroxybenzoics

856

857 **Table 3** – Search strings used in all the selected sources of information.

Search string	Databases		
	Scopus	ScienceDirect	Web of Science
(phenolic acids) AND (withered grapes)	36	94	3
"phenolic acids" AND "sweet wines"	98	88	2
("phenolic acids") AND ("ice wine" OR "ice wines" OR "icewine" OR "icewines")	47	33	4
("HCAs" OR "HCTAs" OR "hydroxycinnamics") AND ("withering" OR "dehydration" OR "drying" OR "raisining") AND ("grape" OR "winegrape")	54	2115	0
("HCA" OR "hydroxycinnamic" OR "hydroxybenzoic") AND ("withering" OR "withered" OR "raisin" OR "dehydration") AND ("grape" OR "winegrape")	370	717	14
("HBA" OR "HBAs" OR "hydroxybenzoics") AND ("withering" OR "withered" OR "raisin" OR "dehydration") AND ("grape" OR "winegrape")	6	430	0
("HCAs" OR "HCTAa" OR "hydroxyinnamics") AND ("withered" OR "dehydrated" OR "dried" OR "raisin") AND ("grape" OR "winegrape")	56	391	0
("HCA" OR "hydroxycinnamic" OR "hydroxybenzoic") AND ("withered" OR "dehydrated" OR "dried" OR "raisin") AND ("grape" OR "winegrape")	849	2721	14
("HBA" OR "HBAs" OR "hydroxybenzoics") AND ("withered" OR "dehydrated" OR "dried" OR "raisin") AND ("grape" OR "winegrape")	24	1541	0
("phenolic acids") AND ("reinforced" OR "fortified" OR "dessert" OR "straw" OR "botrytized" OR "passito") AND ("wine" OR "wines")	950	976	7
("phenolic acids") AND ("Sauternes" OR "Tokaji" OR "Sforzato" OR "Sfursat" OR "Amarone" OR "Recioto" OR "Aleatico" OR "Vin Santo")	53	44	1
("phenolic acids") AND ("Marsala" OR "Porto" OR "Vin de Paille" OR "sweet wine" OR "withered wine" OR "raisin wine")	1593	861	146
("hydroxycinnamic" OR "HCA" OR "HCTAs" OR "hydroxybenzoic" OR "HBA") AND ("reinforced" OR "fortified" OR "passito") AND "wines"	244	599	6
("hydroxycinnamic" OR "hydroxybenzoic") AND ("Sauternes" OR "Tokaji" OR "Sfursat" OR "Amarone" OR "Recioto" OR "Aleatico" OR "Vin Santo")	38	56	1
("hydroxycinnamic" OR "hydroxybenzoic") AND ("Marsala" OR "Porto" OR "Vin de Paille" OR "sweet wine" OR "withered wine" OR "raisin wine")	810	660	131

858

Table 4 – Extraction table of the main outcomes on the evolution of phenolic acids in winegrapes during the **On-vine** withering process.

Author, Year	Title/purpose	On-vine/Off-vine withering	Variety	Outcomes/main findings on phenolic acids of withered grapes
Avizcuri-Inac et al. (2018)	Chemical and sensory characterization of sweet wines obtained by different techniques	On-vine natural dehydration vs off-vine	Tempranillo, Grenache, Viura, Cagazal, Riesling Gewürztraminer, Verdil	Sweet wines obtained using different techniques were classified in three clusters based on phenolic compounds: the first one was characterized by the highest contents of HBAs and HCAs ; the second cluster was by intermediate amounts of phenolic compounds ; and the third, including four artificial icewines, was described by the lowest amounts of phenolic compounds, particularly HBAs .
Blanco-Ulate et al. (2015)	Developmental and metabolic plasticity of white-skinned grape berries in response to <i>Botrytis cinerea</i> during noble rot	On-vine: harvest of grapes with symptoms of noble rot caused by <i>Botrytis cinerea</i> strain BcDW1 and other strains present in the field in three different stages of infection	Sémillon	The increased levels of HCAs derivatives were probably due to the activation of the shikimate and phenylpropanoid metabolic pathways by noble rot infection, normally associated with the ripening of red-berry cultivars.
Figueiredo-González et al. (2013a)	Effects on color and phenolic composition of sugar concentration processes in dried-on-or dried-off-vine grapes and their aged or not natural sweet wines.	On-vine natural withering vs Off-vine withering	All varieties studied in reviewed articles	<i>Review on the effect of withering on the phenolic composition of grapes and wines</i>
Hong et al. (2012)	Influence of plant defense system and fungal growth in <i>Botrytis cinerea</i> -infected Chardonnay berries	On-vine: harvest of botrytized grapes	Chardonnay	Metabolomic investigation revealed a decrease in the concentration of HCTAs as <i>Botrytis cinerea</i> infection increased.
Ivanova et al. (2011)	Identification and changes of polyphenolic compounds in red and white grape varieties grown in R. Macedonia during ripening	On-vine: late harvest	Vranec, Smederevka, Merlot, Chardonnay	The decrease in the content of total phenols in the overripening and late-harvest phase could be the result of a change in the HCA ester concentration in skins, due to oxidation/polymerization reactions for white and red varieties and the participation in acetylated anthocyanin synthesis in red varieties.
Kallitsounakis and Catarino (2020)	An overview on botrytized wines	On-vine: botrytized wine	All varieties investigated in the reviewed studies	<i>Review on the different stages of botrytized wine production. The authors provided a section on HCAs and their esters.</i>
Kilmartin et al. (2007)	Polyphenol content and browning of Canadian icewines	On-vine: icewine production process vs cryogenic extraction method ("REAL" vs "FAUX" Icewines)	Riesling and Vidal	Differences in the amounts of HCAs were observed between "REAL" and "FAUX" icewines in 1999-2000 vintages. The overall concentration of HCAs in "FAUX" icewines was up to 11-times and 21-times higher than that of the "REAL" Riesling and Vidal icewines, respectively. In the 2002 vintage, instead, the authors found relatively high concentrations of HCAs both in "REAL" and "FAUX" icewines, probably owing to the matching of the earliest harvest dates in that year.

Author, Year	Title/purpose	On-vine/Off-vine withering	Variety	Outcomes/main findings on phenolic acids of withered grapes
Loizzo et al. (2013)	Phenolics, aroma profile, and in vitro antioxidant activity of Italian dessert Passito wine from Saracena (Italy)	On-vine: overripe grapes (Malvasia and Guarnaccia) + Off-vine: in the typical <i>cannizza</i> under the sun, covered during the night (Moscato, not macerated grapes collected few weeks before harvest)	Malvasia, Guarnaccia, and Moscato	HBAs were the most abundant class of marker phenolic compounds, particularly gallic acid , which showed the highest content (376.5 mg/L).
Lukić et al. (2016)	Changes in physicochemical and volatile aroma compound composition of Gewürztraminer wine as a result of late and ice harvest	On-vine: late harvest and icewine production process	Gewürztraminer	Higher amounts of HCAs ethyl esters were found in late harvest and icewines than those in regular harvest and of 4-vinylguaiacol in late harvest ones (but not in icewines), suggesting a concentration effect of HCAs in the special wines compared to the control.
Mikulic-Petrovsek et al. (2017)	Effects of partial dehydration techniques on the metabolite composition in 'Refosk' grape berries and wine	On-vine: double maturation raisonnée (DMR) vs off-vine: berry partial dehydration in the chamber (PDC: 22–23 °C, 60% relative humidity, and ventilation)	Refosk	Only one HCA compound was detected in grape skins of Refosk (<i>p</i>-coumaric acid hexose, 0.72-1.48 mg/100 g dry weight) and its content decreased over the withering process , particularly for PDC-treated berries. Seven HCA derivatives were detected in wines, caftaric acid being the most abundant (11.43-20.68 mg/L) . Wines obtained with PDC-withered grapes showed the highest amounts of HCAs and derivatives compared to those of the DMR wines and the control (1.8 and 1.4-times higher, respectively) .
Rusjan et al. (2017)	Double maturation raisonnée: the impact of on-vine dehydration on the berry and wine composition of Merlot	On-vine: double maturation raisonnée (DMR) , cutting 1-year old canes with fruitful shoots during berry ripening	Merlot	A reduction in the contents of HCA and derivatives in DMR grapes was observed compared to that of the control; instead, a high presence of these compounds was found in DMR wines , in some cases significantly higher than that in control wines.
Tang et al. (2013)	Evaluation of nonvolatile flavor compounds in Vidal icewines from China	On-vine: icewine production process	Vidal	HBAs were the major phenolic acids in icewines , representing more than 74% of the total amount of the class: protocatechuic acid (7.43-9.48 mg/L) showed the highest content and syringic acid (0.20-0.36 mg/L) the lowest . As regards HCAs, the main compounds were ferulic (1.07-2.14 mg/L) and <i>p</i>-coumaric (0.20-1.39 mg/L) acids . No significant differences were found in phenolic acid concentrations between wines characterized by different pressing levels .
Tian et al. (2009a)	Comparative study of 11 phenolic acids and five flavan-3-ols in cv. Vidal: impact of natural icewine making versus concentration technology	On vine: naturally frozen juice vs artificially produced (refrigerator-frozen and concentrated-juice)	Vidal	The presence and concentration of phenolic acids could differentiate between icewine production methods . At the end of experimental fermentations, the phenolic acid content increased by 20% in the wine produced from refrigerator-frozen juice and increased by 2.5-2.7 times in the wine from concentrated juice compared to that of the wine made from naturally frozen juice.
Tian et al. (2009b)	Comparison of phenolic acids and flavan-3-ols during wine fermentation of grapes with different harvest times	On-vine: dry wine, semi-sweet wine, icewine with low alcohol levels	Vidal	For the production of icewines, increased contents of HBAs and HCAs were observed when grape harvest time was delayed .

Table 5 –Extraction table of the main outcomes on the evolution of phenolic acids in winegrapes during the **Off-vine controlled** withering process.

Author, Year	Title/purpose	On-vine/Off-vine withering	Variety	Outcomes/main findings on phenolic acids of withered grapes
Bellincontro et al. (2009)	Study of postharvest water stress of winegrape using non-destructive techniques	Off-vine: in perforated plastic trays inside a ventilated small tunnel at different controlled conditions (10 °C with 1.5 m/s airflow; 10 °C with 2.5 m/s airflow; 20 °C with 1.5 m/s airflow. Relative humidity: 45%)	Cesanese	Among the HCAs detected, caftaric acid was very sensitive to atmosphere variation. Its content was 368.95, 513.40, and 393.28 mg/kg for grapes withered at 10 °C-1.5 m/s airflow, 10 °C-2.5 m/s airflow, and 20 °C-1.5 m/s airflow, respectively, and in all the cases it was significantly lower than that of the control (644.99 mg/kg).
Bonghi et al. (2012)	Phenol compound metabolism and gene expression in the skin of winegrape berries subjected to partial postharvest dehydration	Off-vine: in controlled conditions, slow and rapid rates of up to 10% and 30% weight loss	Raboso Piave	A slight increase in caffeic acid contents was observed in rapidly dehydrated samples at 10% weight loss and in <i>p</i> -coumaric acid content under slow dehydration at 30% weight loss. Reduced content of gallic acid was observed in rapidly dehydrated samples. The results obtained on gene expression patterns suggested a complex expression regulation of the PAL multigene family.
Budić-Leto et al. (2017)	Differentiation between Croatian dessert wine Prosek and dry wines based on phenolic composition	Off-vine with a system for temperature control and ventilation (maximum daily temperature of 40 °C)	Prosek, Plavac, and Posip	Prošek dessert wines resulted in significantly lower concentrations of HCAs than those of dry wines.
Frangipane et al. (2007)	Study of phenolic compounds in Aleatico grapes dried in a forced air tunnel	Off-vine: in a forced air tunnel with different temperatures (from 15 to 30 °C) throughout 30 days of withering	Aleatico	In the skin, <i>trans</i> -caftaric acid content increased at temperatures between 17 and 23 °C, but it declined when the temperature increased to 30 °C, evidencing no substantial differences compared to the initial value. In the juice, <i>trans</i> -caftaric acid content remained almost constant until midway through the process, after which its concentration tripled. Contents of syringic and gallic acids in grape juice, instead, decreased during the withering process.
Frangipane et al. (2012)	Effect of drying process in chamber at controlled temperature on the grape phenolic compounds	Off-vine, in a chamber in controlled conditions (18 –21 °C, 68% relative humidity, 1 m/s airspeed) for 16 days	Roscetto	A slight decrease in the content of HCTAs (expressed as mg/1000 berries) was observed in the juice, probably because of oxidation reactions that occurred during the withering process. HBA content showed the largest decrease.
Marquez et al. (2012)	Colour and phenolic compounds in sweet red wines from Merlot and Tempranillo grapes chamber-dried under controlled conditions	Off-vine: in chamber drying under controlled conditions at a constant temperature of 40 °C and initial relative humidity of 30%	Tempranillo and Merlot	A great increase in HBA and HCTA amounts was detected in withered Merlot and Tempranillo grapes. The concentration of HBAs increased during the first 24-48 h of maceration, and then no significant differences or slight decrease were found in Merlot and Tempranillo, respectively; moreover, contents of HCTA esters increased over the first 24 h of maceration and then decreased.

Author, Year	Title/purpose	On-vine/Off-vine withering	Variety	Outcomes/main findings on phenolic acids of withered grapes
Marquez et al. (2014)	Antioxidant activity and phenolic profile changes during the winemaking of Cabernet sauvignon sweet wines	Off-vine: in controlled condition at a constant temperature of 40 °C and an initial relative humidity of 20%.	Cabernet sauvignon	The overall amounts of phenolic acids increased during off-vine withering and subsequent experimental winemaking, namely, for HBAs . The observed increase peaked at 48 h of skin maceration . Additionally, contents of HCAs also increased for all the compounds detected except for cis-fertaric acid, which showed the opposite trend . Moreover, the evolution of the total amount of phenolic acids was correlated to the antioxidant activity .
Mencarelli et al. (2010)	Chemical and biochemical change of healthy phenolic fractions in winegrape by means of postharvest dehydration	Off-vine: in different controlled conditions (10, 20, or 30 °C and 30% relative humidity)	Aleatico	The highest increase in the overall content of phenolic acids (caftaric and coutaric acids) in withered winegrapes was observed at 20 °C and 20% weight loss , while at 10 °C, the increase was lower, and at 30 °C, it decreased for grapes at 20% and 30% weight loss. An upregulation in the expression of the PAL gene was detected in the berries exposed to temperatures of 10 and 20 °C at 10% weight loss. Nevertheless, transcriptomic abundance did not correlate well with metabolomic abundance .
Negri et al. (2017)	The induction of noble rot (<i>Botrytis cinerea</i>) infection during postharvest withering changes the metabolome of Garganega grapevine berries	Off-vine: in perforated plastic boxes in a ventilated withering facility under natural conditions (17-20 °C, 79% relative humidity) until the grapes reached about 30% weight loss ; then half number of plastic boxes were covered to increase relative humidity (88-94%) to induce noble rot development , whereas the remaining boxes were maintained at 68-75% relative humidity for 32 days. The length of the process was 61 days.	Garganega	The authors observed a decrease in HCTA concentrations (coutaric, caftaric, and fertaric acids) in Botrytis-induced grapes , probably owing to the degradation reactions by fungal metabolism .
Nicoletti et al. (2013)	Postharvest dehydration of Nebbiolo grapes grown at altitude for <i>Sfursat</i> wine production is affected by time of defoliation	Off-vine: in controlled-environment rooms (10, 20, or 30 °C; 60% relative humidity; 1 m/s air flow)	Nebbiolo	During withering, all samples lost phenolic acids (caftaric and coutaric acids) except for grapes harvested from not-defoliated vines and withered at 10 °C , in which the amounts of phenolic acids increased.
Panceri et al. (2013)	Effect of dehydration process on mineral content, phenolic compounds, and antioxidant activity of Cabernet sauvignon and Merlot grapes	Off-vine: under controlled conditions of 7 °C, 35% relative humidity, and volumetric airflow of 12 m ³ /s (dehydration until 30% and 40% weight loss)	Cabernet sauvignon and Merlot	Among HBAs detected, gallic acid was the predominant compound in grapes and wines. While contents of HBAs increased in Cabernet sauvignon, a decrease was observed in Merlot grapes , suggesting an effect induced by genotype. Increased contents of caffeic acid in free form were reported after 40% dehydration for both varieties .
Panceri et al. (2015)	Effect of grape dehydration under controlled conditions on chemical composition and sensory characteristics of Cabernet sauvignon and Merlot wines	Off-vine: under controlled conditions with a constant temperature of 7 °C, 35% relative humidity, and volumetric airflow of 12 m ³ /s (dehydration until 30% and 40% weight loss)	Cabernet sauvignon and Merlot	Total content of HCAs was generally higher only for Merlot wines made from grapes at 30% weight loss than that of the control wine. Higher contents of HBAs were found in the wines produced from Cabernet sauvignon and Merlot grapes at 30% and 40% weight loss .

Author, Year	Title/purpose	On-vine/Off-vine withering	Variety	Outcomes/main findings on phenolic acids of withered grapes
Panceri and Bordignon-Luiz (2017)	Impact of grape dehydration process on the phenolic composition of wines during bottle ageing	Off-vine: under controlled conditions with a constant temperature of 7 °C, 35% relative humidity, and airflow of 12 m ³ /s (dehydration until 30% and 40% weight loss)	Cabernet sauvignon and Merlot	The wines produced from dehydrated grapes evidenced higher amounts of gallic, protocatechuic, vanillic, and caffeic acids than that of the control. An increase in the concentration of phenolic acids (gallic, protocatechuic, and <i>p</i> -coumaric acids) was observed during 22 months of bottle ageing in all Cabernet sauvignon and Merlot wines made from withered grapes .
Piano et al. (2013)	Focusing on bioactive compounds in Uvalino grapes	Off-vine: in a room at a constant temperature of 24 °C and 30% humidity vs On-vine: not-harvested overripe grapes (control)	Uvalino	HCTA contents (caftaric, coumaric, ferulic acids) slightly increased after 7 days of withering/overripening , and this increase was maintained up to the 21st day (final point).
Serratos et al. (2008b)	Drying of Pedro Ximenez grapes in chamber at controlled temperature and with dipping pretreatments	Off-vine: at a controlled temperature of 40 and 50 °C vs. traditional sun-drying	Pedro Ximenez	An increase in the concentration of gallic acid in the juice from withered grapes was observed during the withering process, from 2.33 to 7.24 mg/L at a temperature of 40 °C and 10.2 mg/L at 50 °C. Similarly, also HCA concentrations increased by 50% or doubled after withering at 40 or 50 °C, respectively.
Toffali et al. (2011)	Novel aspects of grape berry ripening and post-harvest withering revealed by untargeted LC-ESI-MS metabolomics analysis	Off-vine: in controlled conditions (13-17 °C and about 65% relative humidity)	Corvina	Anthocyanins acylated with <i>p</i>-coumaric acid started to accumulate after veraison but continued to increase thereafter, during the withering process . Consequently, the ratio of non-acylated anthocyanins to coumaroyl anthocyanins fell from 6.4 in ripening and ripe berries to 3.7 in withering and withered berries .
Torchio et al. (2016)	Influence of different withering conditions on phenolic composition of Avanà, Chatus and Nebbiolo grapes for the production of 'Reinforced' wines	Off-vine: different controlled conditions such as slow withering (18 °C, 40% relative humidity, and 0.9 m/s airflow) and fast withering (28 °C, 40% relative humidity and 0.9 m/s airflow)	Avanà, Chatus, Nebbiolo	The concentration effect due to withering prevailed over the degradation reactions for grape pulp HCTAs in two varieties out of the three tested, particularly for Chatus . These amounts were not influenced by the withering rate.
Zenoni et al. (2016)	Disclosing the molecular basis of the postharvest life of berry in different grapevine genotypes	Off-vine: in controlled conditions with gradually decreasing temperature (from 16 ° to 7 °C) and gradually increasing relative humidity (from 55% to 80%)	Corvina, Sangiovese, Merlot, Oseleta, Syrah, and Cabernet sauvignon	The positive or negative final balance of HCAs/HBAs depended on genotype , and it was negative in Corvina .

Table 6 –Extraction table of the main outcomes on the evolution of phenolic acids in winegrapes during the **Off-vine natural** withering process.

Author, Year	Title/purpose	On-vine/Off-vine withering	Variety	Outcomes/main findings on phenolic acids of withered grapes
Constantinou et al. (2017)	Metabolic fingerprinting of must obtained from sun-dried grapes of two indigenous Cypriot cultivars destined for the production of 'Commandaria': a protected designation of origin product	Off-vine: sun-drying process, during which the grapes were manually turned over periodically. The sun-drying process lasted 10-12 days.	Xynisteri and Mavro	Among phenolic compounds, HBAs represented the most predominant group in Xynisteri grapes (62.1% of total phenolics) and were an important group in Mavro (42.3%). Total amounts of HBAs increased in musts of both varieties during sun-drying. As regards HCAs and derivatives , they were the second major group of phenolic compounds, contributing to 25.7% and 39.8% for Xynisteri and Mavro, respectively. During withering, a six-times increase in their content was observed , even if the concentration of some singular compounds such as caffeic acid dihexoside and fertaric acid isomer decreased .
Constantinou et al. (2018)	The impact of postharvest dehydration methods on qualitative attributes and chemical composition of 'Xynisteri' grape must	Off-vine: traditional sun-drying method vs. alternative dehydration methods (multiple horizontal wires, multiple vertical pallets, low greenhouse, hot-air dryer treatment)	Xynisteri	Total HBA contents in musts from grapes withered with multiple horizontal wires, low greenhouse, and hot-air treatments increased by 1.8-, 2.3- and 2.7-times that of fresh grapes, respectively. Traditional sun-drying revealed the lowest effect on HCAs (1.5-times) , while other withering methods showed a strong increase (from 1.7 to 6.0 times), which was higher than that of the concentration effect.
Figueiredo-González et al. (2013b)	Evolution of colour and phenolic compounds during Garnacha Tintorera grape raisining	Off-vine: in a room with natural ventilation over 83 days, grapes were placed in a single layer in plastic boxes, turned over, and checked weekly	Garnacha tintorera	The overall concentration of HCA esters decreased from 32 to 14 mg/kg of berry during 83 days of withering, and this decrease began after the 16 th day. A decreasing trend in HCA esters was observed during withering for the two isomers: cis-caftaric (from 0.38 to 0.07 mg/kg of berry), trans-caftaric (from 19 to 9.7 mg/kg of berry), cis-coutaric (from 4.0 to 1.1 mg/kg of berry) and trans-coutaric (from 9.0 to 3.2 mg/kg of berry).
Figueiredo-González et al. (2014a)	Garnacha-Tintorera based sweet wines: detailed phenolic composition by HPLC/DAD - ESI/MS analysis	Off-vine: for naturally sweet wine, in plastic boxes at natural conditions of temperature and humidity	Garnacha tintorera	The concentration of HBAs was lower in the two sweet wines than that of the dry wine (control), particularly for the fortified one, mainly driven by the steep decrease in gallic acid content. HCTAs and free HCAs were present at high concentrations (45 mg/L) in dry base wine, but their presence decreased probably owing to enzymatic browning reactions.
Figueiredo-González et al. (2014b)	The phenolic chemistry and spectrochemistry of red sweet winemaking and oak-aging	Off-vine: in plastic boxes at natural conditions of temperature and humidity vs. fortified sweet wine	Garnacha tintorera	An increase in phenolic acid amounts was observed for both natural and fortified sweet wines during the oak-ageing period of 6 and 12 months. The concentration of HBAs increased , starting from 32 and 40 mg/L at the beginning in natural and fortified sweet wines, respectively, and averaging about 50 mg/L in both sweet wines at the end of the ageing , probably because of the transfer from the oak wood. Moreover, a decreasing trend was found in the contents of HCA esters (caftaric and coutaric acids). The simultaneous increase of HCA free forms suggested a slow hydrolysis process, but this increase was not proportional to the decrease of the contents of esters.

Author, Year	Title/purpose	On-vine/Off-vine withering	Variety	Outcomes/main findings on phenolic acids of withered grapes
Nievierowski et al. (2021)	Role of partial dehydration in a naturally ventilated room on the mycobiota, ochratoxins, volatile and phenolic composition of Merlot grapes	Off-vine: in a naturally ventilated room for 21 days	Merlot	The main differences were observed for four HBAs (ethyl gallate, <i>p</i> -hydroxybenzoic acid, gallic acid-hexose, and gallic acid) and two HCAs (caffeic and caftaric acids). The amounts of these compounds increased more than two times during withering compared to that of fresh grapes.
Serratos et al. (2008a)	Changes in color and phenolic compounds during the raisining of grape cv. Pedro Ximenez (for the production of sweet wines)	Off-vine: sun-drying grape raisining on mats in slight hills facing south (40-50 °C) for 7 days	Pedro Ximenez	The withering process enhanced the concentration of HBAs in grape must. Particularly, <i>p</i> -hydroxybenzoic acid amount strongly increased by 188% , suggesting an additional increase by means other than concentration. Meanwhile the content of gallic acid increased by 33% , suggesting that this compound can participate in other reactions during the withering process. As regards HCTAs , the amounts of caftaric, coutaric, and fertaric acids increased more for cis- isomers than for trans-isomers. These increases were lower than those expected, indicating a simultaneous degradation of these compounds during withering that contribute to the typical browning of the sun- dried grapes.

Identification

Records identified through
database searching
(n=16883)

Additional records identified
through other sources
(n=0)

Screening

Records after duplicates removed
(n=8549)

Eligibility

Records screened
(n=8549)


Records excluded at title
or abstract review
(n=8455)

Included

Full-text articles assessed for
eligibility
(n=94)

Full-text articles excluded
(not completely meeting
inclusion criteria)
(n=55)

Studies included in qualitative synthesis
(n=39)



Highlights

- Phenolic acids are represented by hydroxybenzoic and hydroxycinnamic compounds
- Their evolution during withering involves a complex balance of different processes
- Concentration effect usually prevails in withered grapes increasing their contents
- Withering conditions and genetic regulation are relevant in determining the content
- Phenolic acids have been proposed as authenticity markers for icewines

Journal Pre-proof