

AperTO - Archivio Istituzionale Open Access dell'Università di Torino

Campylobacter spp. prevalence and mitigation strategies in the broiler production chain

This is the author's manuscript

Original Citation:

Availability:

This version is available <http://hdl.handle.net/2318/1839754> since 2022-02-10T10:53:35Z

Published version:

DOI:10.1016/j.fm.2022.103998

Terms of use:

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)

1 ***Campylobacter* spp. prevalence and mitigation strategies in the broiler production chain**

2 Jatziri Mota-Gutierrez^a, Lukas Lis^b, Amaia Lasagabaster^c, Ibai Nafarrate^c, Ilario Ferrocino^a, Luca
3 Cocolin^a, Kalliopi Rantsiou^{a*}

4
5
6 **Affiliations:**

7 ^a Department of Agricultural, Forest, and Food Sciences, University of Turin, Largo Paolo
8 Braccini 2, 10095, Grugliasco, Turin, Italy

9 ^b PTC Phage Technology Center GmbH, Im Kompetenzzentrum BioSecurity, Bönen, Germany

10 ^c AZTI, Food Research, Basque Research and Technology Alliance (BRTA), Parque Tecnológico
11 de Bizkaia, Astondo Bidea 609, 48160 Derio, Bizkaia, Spain.

12

13 **Corresponding author:**

14 * Kalliopi Rantsiou, Ph.D., Department of Agricultural, Forest and Food Sciences of the
15 University of Turin (DISAFA), University of Turin, Grugliasco, Turin, 10095, Italy

16 E-mail: kalliopi.rantsiou@unito.it

17 Telephone: 011 670 8870, fax: 011 670 8549

18

19 **Scientific section:** Microbiology and Food safety

20 **ABSTRACT**

21 This study aims to discuss the microbial ecology of the broiler gut environment,
22 *Campylobacter* prevalence across the broiler production chain with a follow-up focus on a
23 possible mitigation strategy based on the use of bacteriophages. Scientific literature published
24 from the last two decades was reviewed and data were collected to establish the ranges of
25 *Campylobacter* loads from different samples. Results showed that the pathogen load in the
26 sample is likely to increase from the different stages of the production chain. Contamination of
27 water and feed represents the most notable source of contamination during the primary
28 production, while cross-contamination of broiler carcasses, skin, and meat occurs during the
29 slaughter, dressing, and processing via machinery, work surfaces, water, and air partially due to
30 the leaking of contaminated feces from visceral rupture. Knowledge gaps were identified and
31 included: a lack of studies detecting *Campylobacter* in broilers in most of the European countries
32 over the last decade and a low number of studies determining the bacterial load in crates used to
33 transport broilers to the slaughterhouse. Determining the prevalence of *Campylobacter* in the
34 broiler industry will enable us to set critical control points to produce broiler flocks and meat
35 products with a low risk of *Campylobacter* contamination.

36

37 **Keywords:** Microbiota, chicken, bacteriophage, biocontrol, poultry, public health risk

38 1. INTRODUCTION

39 Zoonoses are defined as those infectious diseases naturally transmitted from vertebrate
40 animals to humans through direct or indirect contact (food or water contamination). Zoonotic
41 agents include a wide variety of bacteria, viruses, protozoa, insects, and helminths. According to
42 the World Health Organization (WHO), almost 600 million cases of food-borne zoonoses were
43 reported worldwide in 2015, of which 52% were caused by pathogenic bacteria (WHO, 2015).
44 *Campylobacter* is the most common pathogen responsible for food-borne zoonotic diseases in
45 humans and it is considered a serious public health issue in both developing and developed
46 countries. The most recent summary report of the European Food Safety Authority (EFSA)
47 stated that campylobacteriosis is the most frequently reported food-borne zoonoses in the
48 European Union (EU), with 220 682 confirmed cases in 2019, reported from 18 different
49 members states (EFSA and European Centre of Disease Prevention and Control (ECDC), 2019).
50 The most common source of infection in humans due to *Campylobacter* are broiler meat and
51 milk (EFSA and European Centre of Disease Prevention and Control (ECDC), 2019).

52 *Campylobacter* is a genus of microaerophilic and Gram-negative bacteria belonging to
53 the *Proteobacteria* phylum. Bacterial cells generally appear as slender, spirally curved, or
54 comma-shaped rods characterized by inability to form endospores and ability to change to
55 spherical or coccoid form under stress conditions (Lastovica et al., 2014). *Campylobacter* species
56 have an optimal growth temperature, O₂, and CO₂ concentration of 30- 42°C, 5-10%, and 3-5%,
57 respectively (Lastovica et al., 2014). In terms of biochemical characteristics of *Campylobacter*
58 species, they are oxidase-positive, with the capacity to reduce fumarate to succinate but are not
59 able to metabolize lipids, starch, gelatin, and casein (Lastovica et al., 2014). Currently, 33
60 species have been taxonomically described from this genus, but this number is increasing due to

61 the identification of new species (bacterio.net). The main natural reservoir of
62 thermophilic/thermotolerant *Campylobacter* species has been extensively reported in warm-
63 blood animals, including most mammals, birds, and food-producing animals (Silva et al., 2011).
64 The mechanism of colonization, adherence, and invasion of host by pathogenic *Campylobacter*
65 species such as *C. jejuni*, has been previously reviewed elsewhere (Elmi et al., 2021). In
66 summary, successful colonization of the bacteria depends on the ability to attach to the gut
67 mucosa, motility, chemotaxis, spiral shape of the cell, the functionality of the flagella, production
68 of toxins and secreted proteins (cytolethal distending toxin -CDT) and other virulence factors
69 (Elmi et al., 2021).

70 Interestingly, *Campylobacter* does not multiply outside a warm-blooded host due to the
71 absence of microaerobic conditions. However, they can survive when protected from dryness. In
72 this regard, it has been shown that *Campylobacter* can survive up to 3 months in slurries and
73 dirty water (Nicholson et al., 2005). The mechanism of survival of the *Campylobacter* species
74 when exposed to stress environments has been explained by its ability to form a biofilm on
75 abiotic surfaces, this biofilm ensures a supply of nutrients and mechanical protection to survive
76 (Johnson et al., 2017). Although the bacteria cannot multiply outside the animal hosts or in food
77 during storage, it has been shown that they are able to develop complex mechanisms of virulence
78 which remain poorly understood.

79 Recent studies have facilitated a greater appreciation of the complex mechanism of
80 virulence of the members of the *Campylobacter* genus. The full genome of *Campylobacter*
81 *jejuni*, has elucidated strain-specific genetic diversity with high genome plasticity (Bacon et al.,
82 2000). The ability to survive and adapt to stress environments indicates that *C. jejuni* harbors
83 complex virulence and fitness factors (Tegtmeyer et al., 2021). Virulence factors and

84 pathogenicity islands in *C. jejuni* have been identified and are reported elsewhere (Ali et al.,
85 2012; Bacon et al., 2000; Sierra-Arguello et al., 2021; Tang et al., 2020; Zhang et al., 2017).
86 Virulence-associated genes such as *flaA*, *cadF*, *cdtA*, *cdtB*, *cdtC*, *cheY*, *iamA*, and *virB11* were
87 recently identified in *C. jejuni* and *Campylobacter coli* isolates originating from broiler of 31-day
88 or 37-day age at the rearing period to slaughtering process (Tang et al., 2020). Besides the
89 identification of virulence-associated genes, invasion-associated genes, plasmid genes and CDT-
90 associated genes were also recognized in *C. jejuni* strains isolated from cloacal, broiler carcasses,
91 and broiler slaughterhouses (Sierra-Arguello et al., 2021). The aforementioned genes are
92 involved in the adhesion, invasion, chemotaxis, motility, toxin-activity, and host immune system
93 evasion. Interestingly, there is a difference of virulence factors between *C. jejuni* and *C. coli*,
94 where a higher number of virulence genes were retrieved in *C. jejuni* if compared with *C. coli*,
95 this difference might contribute to the higher colonization of *C. jejuni* in the broilers' intestines
96 (Tang et al., 2020; Zhang et al., 2017).

97 Antibiotic resistance in *Campylobacter* is also considered a global trend. In this regard,
98 *C. jejuni* and *C. coli* had shown a multi-drug resistance to several antibiotics such as
99 tetracyclines, macrolides, aminoglycosides, and β -lactams. However, a higher number of
100 antibiotic resistance genes were retrieved for *C. jejuni* if compared with *C. coli* (Tang et al.,
101 2020). On the increase of antibiotic resistance to more than one class of antibiotics, further
102 research is needed to understand the mechanism of antimicrobial resistance to improve not only
103 human but also animal health.

104 Besides the genetic makeup, the main factors that influence the occurrence of *Campylobacter*
105 in broilers are related to the host gut environment, production chain, or farm practices (Barker et
106 al., 2020; Djennad et al., 2017; McKenna et al., 2020; Perez-Arnedo and Gonzalez-Fandos,

107 2019; Sibanda et al., 2018; Tang et al., 2020). A conceptual framework of the factors increasing
108 the occurrence of *Campylobacter* and a prevention guideline to stipulate the best conditions and
109 food processing management to reduce the risk of *Campylobacter* contamination in the broiler
110 production chain has been developed (EFSA, Panel on Biological Hazards, 2011; Lyngstad,
111 Jonsson, Hofshagen, & Heier, 2008).

112 Several intervention methods have been developed in recent years, such as the combination
113 of strict biosecurity measures, good manufacturing practice (GMP), hazard analysis and critical
114 control points (HACCP), *Campylobacter* vaccines, antibiotic alternatives to control
115 *Campylobacter*, probiotics, and phytochemicals (Deng et al., 2020; European Food Safety
116 Authority Panel on Biological Hazards, 2011; Umar et al., 2016; Ushanov et al., 2020).

117 However, the problem has not been completely eradicated and the prevalence of this pathogen is
118 still high. Ante- and post-mortem veterinary inspections of broilers are routinely used at the
119 slaughterhouse level as a strategy to ensure that meat does not bear fecal or other contaminants.

120 However, the presence of *Campylobacter* in broiler carcasses cannot be detected visually. As an
121 attempt to mitigate this issue, the application of *Campylobacter*-specific bacteriophages has
122 emerged as one of the most promising approaches to be applied within the farm-to-fork poultry
123 process (Atterbury et al., 2003; Fischer et al., 2013; Hammerl et al., 2014; Kittler et al., 2013;
124 Richards et al., 2019). In this context, this review focuses on discussing the most updated
125 scientific achievements made on the microbial ecology of the gastrointestinal (GI) tract of
126 broilers and the interaction between chickens' gut microbiota and *Campylobacter*,
127 *Campylobacter* prevalence across the broiler production chain with a follow up of the application
128 of bacteriophage along the farm-to-fork process.

129

130 2. MICROBIAL ECOLOGY IN CHICKEN' ECOSYSTEMS

131 Chicken ecosystems harbor complex, diverse, and dynamic microbial communities
132 composed of bacteria, protozoa, fungi, yeasts, bacteriophages, and other viruses. The integrity of
133 the GI tract and the gut microbiota composition has a direct influence on chicken' health,
134 affecting the development of their digestive and immune systems (Clavijo and Flórez, 2018;
135 Khan et al., 2020). The avian gut microorganisms are mainly responsible for the continuous
136 generation of energy and nutrients, such as vitamins (K and B groups), amino acids, short-chain
137 fatty acids (SCFA like, butyric, propionic, and lactic acids), ammonia, antimicrobial compounds
138 (bacteriocins) and the decrease of triglyceride concentrations in the avian gut (Vispo and
139 Karasov, 1997). The positive effect of the production of SCFA on the host includes the inhibition
140 of pathogens, reduction of the pH levels in the colon, and stimulation of the cell proliferation in
141 the gut epithelium (Christl et al., 1997; Dibner and Richards, 2005; Ricke, 2003). In contrast,
142 chickens' growth can be depressed when competition for energy and protein occurs between the
143 commensal microbiota. As a result of this competition, toxic metabolites (amino acid catabolites)
144 can be produced and impact the growth of the animal (Shang et al., 2018).

145 As reported in recent years, the composition and function of chickens' microbiota vary
146 depending on biological changes within and between hosts (age, sex, maternal factors, and
147 breed), and environmental factors (biosecurity levels, housing, litter, feed access and antibiotic
148 administration, hygiene, location, and climate) (Kers et al., 2018). Regarding the biological
149 variation within and between hosts, it has been demonstrated that the chickens' microbiota
150 richness increases during the first week of life (Ballou et al., 2016; Crhanova et al., 2011), while
151 the number of different microbial taxa decreases with chicken age (Lu et al., 2003). The

152 microbial composition of chickens does not only change with chicken age but it is also
153 influenced by the location in the digestive tract and diet (Shang et al., 2018).

154 Significant progress has been made in understanding the microbial taxonomic
155 composition of the different sections of the chickens' GI tract. Overall, *Firmicutes* is the phylum
156 most abundant in both ilea and ceca of chickens (Kumar et al., 2018; Lu et al., 2003; Oakley et
157 al., 2014). Interestingly, in the ceca, the relative abundance of members of the *Clostridium* genus
158 increased by 10 fold between weeks 1 and 6 post-hatch (Oakley and Kogut, 2016). Concerning
159 the bacterial community within the small intestine, in this section lactobacilli mainly dominated
160 the microbial ecosystem (Gong et al., 2007; Lu et al., 2003). According to Lu *et al.*, the
161 microbial composition of the ceca and ileum of three days of age broilers (chickens for meat
162 production) fed with a vegetarian corn-soy diet was mainly composed of *Lactobacillus*
163 *delbrueckii* (13 and 42%, respectively), *Clostridium* spp. (31 and 1%, respectively) and
164 *Clostridium perfringens* (13 and 16%, respectively), however differences in the microbial
165 composition between these sections (ceca and ileum) were reported (Lu et al., 2003). Regarding
166 seven to 49 days old chickens, *Clostridium* spp. remained the most abundant bacteria in the ceca,
167 followed by *Ruminococcus*, while differences in the relative abundance of bacterial species
168 overtime were reported in the ileum (Lu et al., 2003). In detail, *Lactobacillus acidophilus* (50-
169 59%) was the most abundant bacteria in chickens of seven to 21 days of age, while a unique
170 community was reported in the ileum of three, 28, and 49 days of age broilers. Interestingly,
171 regardless of the absence of *Clostridium* spp. and *Ruminococcus* spp. in the ileum section at an
172 early age (between three to 14 days), significant differences of the microbiota between the
173 different sections of the GI tract (ileum and ceca) were only found after 14 days of age (Lu et al.,
174 2003).

175 Overall, it is still not clear whether *Proteobacteria* and *Bacteroidetes* are also abundant
176 phyla during the first six weeks in the chicken's ceca (Shang et al., 2018). These contradictory
177 results may be related to the different chicken breeds studied. According to Kers and colleagues
178 (2018) a variation at the phylum level from the ceca samples of broiler breeds (meat production)
179 and laying-type chickens (egg production) were observed at different time points using 16S
180 rRNA gene amplicon sequencing. Specifically, at zero hr, *Firmicutes* was the most abundant
181 phylum reported in meat-type chicks (Pedroso et al., 2016; Danzeisen et al., 2011), while
182 *Proteobacteria* was identified in laying-type chicks (Ballou et al., 2016). This variability may be
183 due to sample types (fecal vs cecal), feed intervention, and/or the technical aspects of the
184 microbial identification as reported elsewhere (Shang et al., 2018). However, from one to 42
185 days of age, *Firmicutes* was the most abundant phylum regardless of the type of broiler breeds.
186 Furthermore, *Candidatus arthromitus*, a desirable bacterium commonly associated with healthy
187 GI tracts in animals, has been also identified in the jejunum and ileum of chickens (Gong et al.,
188 2007, 2002).

189 ***2.1 Chicken diet and intestinal microbiota interaction***

190 The characteristics of the chickens, feed management, the use of medications or vaccines,
191 the environmental conditions of the poultry house, and the housing systems are known factors
192 that have a short- or long-term effect on the intestinal microbiota composition and immune
193 system development of chickens (Kers et al., 2018). The effect of feed management on the
194 intestinal microbiota has been extensively studied (Apajalahti, 2005; McKenna et al., 2020;
195 Singh et al., 2014; Stanley et al., 2012; Takeshita et al., 2021). Differences in the relative
196 abundance of bacterial species in fecal and cecal samples of broilers following high and low
197 growth diets were reported elsewhere (Singh et al., 2014; Stanley et al., 2012). Knarreborg and

198 colleagues (2002) demonstrated that the divergence in feed can also increase or decrease the
199 relative abundance of a specific bacterial group: the aforementioned study shows how pellet feed
200 increases the number of *Enterococcus* spp. and coliforms and decreases lactobacilli species and
201 *C. perfringens* in the ileum of broilers when compared with mash feed (Knarreborg et al., 2002).
202 In contrast, when chickens consumed corn, this diet favors a decrease in the number of clostridia,
203 enterococci, and lactobacilli, while when chickens consumed wheat, it favors the increase of
204 bifidobacteria (Apajalahti, 2005). Interestingly, the amount of protein in the chicken feed also
205 changed microbial composition, where high amount of protein showed a lower relative
206 abundance of lactobacilli species compared with chicken feed with a low amount of proteins diet
207 (Takeshita et al., 2021). The difference in microbial community structure between production
208 systems together with different management parameters such as stocking density has also shown
209 to alter the microbiota of broilers (McKenna et al., 2020). To date, broilers' microbiota studies
210 have focused on identifying bacterial composition while the identification of other components,
211 such as fungi, phages, or viruses, remain unclear. A better understanding of the role and
212 interactions between mycobiota, phagobiota, and virobiota with the broiler microbial ecosystem
213 may help to improve chicken productivity, health, and welfare and develop novel strategies for
214 controlling the prevalence of *Campylobacter* spp. in broilers (Silva et al., 2011).

215 ***2.2 Campylobacter and its interactions with gut microbiota in chickens***

216 *Campylobacter* typically occurs within two weeks onwards and increases during broilers
217 life cycle (Ijaz et al., 2018; Kalupahana et al., 2013; McKenna et al., 2020; Neill et al., 1984;
218 Thibodeau et al., 2015). It usually grows in the ceca and cloaca and can also colonize the spleen,
219 blood, and liver (Lin, 2009). *C. jejuni* and *C. coli* are the most commonly isolated
220 *Campylobacter* species in broiler samples at different production stages. Interestingly, recent

221 studies show contradictory results on how microbial ecology influences *Campylobacter*
222 colonization. According to Sofka and colleagues the presence of *Campylobacter* decreased the
223 diversity of intestinal microbiota when compared with non-colonized broilers (Sofka et al.,
224 2015). In alignment with the aforementioned study, a significant difference in the relative
225 abundance of the microbial operational taxonomic units detected in the ceca of chickens fed with
226 different diets, at different ages from three different commercial broiler farms were reported
227 between *Campylobacter*-positive and -negative chickens (Takeshita et al., 2021). Interestingly,
228 the decrease in lactobacilli abundance in chicken ceca was associated with high levels of
229 *Campylobacter*, while the growth of *Campylobacter* increased the levels of *Enterobacteriaceae*
230 (Sakaridis et al., 2018). However, according to McKenna and colleagues the presence of
231 *Campylobacter* in broilers reared under standard industrial growing systems increased the cecal
232 microbial community structure (McKenna et al., 2020). Whereas the transplantation of cecal
233 microbial in chickens shows no significant difference in the ceca microbial communities of
234 different inbred chickens (Chintoan-Uta et al., 2020). The factors affecting host-pathogen
235 ecology in terms of the microbiome and the microbial dynamics and *Campylobacter* presence
236 remain poorly studied at an industrial or small-scale farm level.

237

238 **3. CAMPYLOBACTER PREVALENCE IN THE BROILER PRODUCTION CHAIN**

239 In terms of *Campylobacter* diversity in the broiler chain production, a recent study has
240 demonstrated that *C. jejuni* predominates during the rearing broiler period while more isolates of
241 *C. coli* were detected during the slaughtering process (Chen et al., 2010; Tang et al., 2020; Zhang
242 et al., 2018).

243 *Campylobacter* from chicken reservoirs may reach humans through the environment or by
244 direct contact and mainly through the consumption of raw and undercooked contaminated broiler
245 meat and meat products. The Scientific Opinion of the Panel on Biological Hazards (BIOHAZ),
246 EFSA in 2010 estimated that the majority of human campylobacteriosis is attributed to the
247 chicken reservoir as a whole (50 – 80%), while the handling, preparation, and consumption of
248 broiler meat may account only for 20 to 30% (EFSA, 2010). Other food products such as
249 unpasteurized dairy products and contaminated water are also vehicles of transmission of this
250 pathogen, but *Campylobacter* infections are less common from these sources compared with
251 meat products (Josefsen et al., 2015). To provide an indication of the possible control points for
252 *Campylobacter* at the broiler meat production chain, this review describes the prevalence of
253 *Campylobacter* along the process and discusses the risk factors that influence the level of
254 contamination (Figure 1).

255 ***3.1 Primary production***

256 The increases of *Campylobacter* loads during the first weeks of age of commercial flocks
257 (Umar et al., 2016) indicate that vertical transmission of the pathogen does not commonly occur;
258 however, it still represents an important risk factor for the *Campylobacter* colonization in
259 broilers (Bull et al., 2008; European Food Safety Authority Panel on Biological Hazards, 2011;
260 Humphrey, 2006; Tang et al., 2020). The protection of young broilers' GI tract against
261 colonization of *Campylobacter* has been associated with *Campylobacter*-specific maternal
262 antibodies (MAB) (Sahin et al., 2001; Vandeplass et al., 2010). MAB, predominantly
263 immunoglobulin G (IgG) class, are transported from the egg yolk across the yolk sac membrane
264 into the embryonic circulation to protect young broiler chickens from *C. jejuni* infections
265 (Linden and Roth, 1978). However, MAB levels change with the increase in the broiler age. In

266 detail, the highest level of MAB is reported during the first two weeks after hatching, followed
267 by a decrease, reaching minimal values at the third and fourth weeks of age (Sahin et al., 2001;
268 Vandeplass et al., 2010). The increased risk of *Campylobacter* colonization in broilers from week
269 three to week six was demonstrated elsewhere, as shown in Table 1 (Ingesa-Capaccioni et al.,
270 2016; Perez-Arnedo and Gonzalez-Fandos, 2019; Tang et al., 2020; Tangkham et al., 2016).
271 Interestingly, the prevalence of *Campylobacter* spp. and *C. jejuni* at the end of the rearing period
272 is higher in open housing compared with environmentally controlled housing (Tangkham et al.,
273 2016). Noteworthy, a higher prevalence of *C. jejuni* isolates compared with *C. coli* was observed
274 from the broiler rearing period at the farm level in China (Tang et al., 2020), while in Spain, *C.*
275 *coli* has not been detected throughout chickens progeny (Ingesa-Capaccioni et al., 2016; Perez-
276 Arnedo and Gonzalez-Fandos, 2019).

277 Natural colonization of broilers by single or multiple *Campylobacter* species rapidly occurs
278 through horizontal transmission from vectors such as domestic pets, insects, rodents and
279 migratory birds, farm equipment, transport vehicles, farmworkers, drinking water, feed, litter,
280 air, or thinning (Bang et al., 2002; Carvalho et al., 2010; Gharib Naseri et al., 2012; Koolman et
281 al., 2014; Robyn et al., 2013; Schroeder et al., 2014; Stern et al., 2001). Based on the most recent
282 studies considered in this review, the highest prevalence of *Campylobacter* presence has been
283 detected on drinking water, feed, workers boots, and broilers' fecal and cecal samples after
284 thinning, while the lowest prevalence of this bacteria was found on samples from the litter, air
285 and the infrastructure of the farm (Table 1).

286 The colonization of the flocks with *Campylobacter* can also be introduced from external
287 visitors to the farm, maintenance personnel, bird catching crew, close livestock, when operators
288 visit more than one farm or are negligent regarding hygienic practices (Berndtson et al., 1996;

289 Cardinale et al., 2004; Hald et al., 2000; Johnsen et al., 2007; Kapperud et al., 1993; A. M.
290 Ridley et al., 2011). Besides human traffic, the proximity of the fresh litter, larger thinning
291 crews, increasing the number of broiler houses on-site, and the presence of dead broilers in the
292 house enhances the survival of *Campylobacter* and thus the risk of positive flocks is increased
293 (Cardinale et al., 2004; Koolman et al., 2014; Lyngstad et al., 2008). Inadequate biosecurity such
294 as broiler houses older than 15 years, absence of anterooms and barriers in each house, the use of
295 shared tools between houses, long downtime, and drinking systems with bells or cups represent a
296 common risk factor for *Campylobacter* colonization of broiler flocks (Sommer et al., 2016).
297 Partial thinning of broiler flocks has also been considered as a potential risk factor for
298 *Campylobacter* colonization of the remaining animals, due to the difficulty of keeping
299 biosecurity measures during the process (Hermans et al., 2011).

300 The dispersion of *Campylobacter* originating from broilers into the environment represents
301 an important factor leading to increased contamination. In broiler, *C. jejuni* is the most
302 predominant species colonizing the flocks, followed by *C. coli* and occasionally by other species
303 (Rossler et al., 2019; Umar et al., 2016). Once the first bird is infected, *Campylobacter* is
304 horizontally transmitted to most of the birds in a flock within only a few days, reaching between
305 10^6 and 10^8 CFU/g in their intestinal tract, and they remain colonized until slaughter (Marotta et
306 al., 2015).

307 The increased water consumption by chickens during summer months increases the risk of
308 drinking water contaminated with *Campylobacter* (Herman et al., 2002; Lyngstad et al., 2008).
309 This association has been observed in northern European countries, such as Sweden, Denmark,
310 Norway, and the Netherlands (Jore et al., 2010). The changes in the temperature throughout the
311 year can explain the increase in water consumption during summer. However, other factors such

312 as the increment in the abundance of transmission vectors of *Campylobacter* (flies) and the need
313 for ventilation in the poultry house of broilers can also increase the probabilities of the
314 introduction of *Campylobacter* into the environment (Hald et al., 2008; Jore et al., 2010). Also, it
315 must be underlined that, as expected, climate change is an important factor that might increase
316 the prevalence of *Campylobacter* in the future, placing public health at risk. It is worth noting
317 that colonized birds are predominantly asymptomatic, with no negative effect on their health
318 (Pielsticker et al., 2012).

319 ***3.2 Transportation before slaughter***

320 The transportation step consists of moving the broilers from farm facilities to the
321 slaughterhouse. During this process, the animals are confined in crowded crates or cages,
322 deprived of water and feed, and undergo continuous stress that affect not only the meat quality
323 but also the *Campylobacter* prevalence, which could increase due to cross-contamination. If a
324 single broiler is colonized, *Campylobacter* will be spread to the environment and will be
325 transmitted to the rest through fecal matter or feathers contact (Stern et al., 2001; Whyte et al.,
326 2001).

327 In comparison with the primary production, the pre- and post-transportation steps have
328 been less studied over the last two decades (Table 2). Overall, from the literature available we
329 observed that during transportation, the floor and transportation cage/crates account for the
330 highest risk of cross-contamination in broilers (Stern et al., 2001; Willis et al., 2002), while no
331 clear contamination level in fecal samples between pre-and post-transportation steps is observed
332 (Whyte et al., 2001). One likely explanation for this phenomenon is that catching and placing the
333 chickens in cages/crates to transport them to the processing plant increases the risk of
334 contamination, mostly due to cross-contamination during transportation (Slader et al., 2002).

335 However, recent literature has demonstrated the ineffectiveness of cleaning naturally
336 contaminated crates before using them to transport broilers and reduce *Campylobacter* infections
337 (A. Ridley et al., 2011). Further, transportation stress alters excretion rates of *Campylobacter* in
338 the fecal material of broilers (Whyte et al., 2001).

339 ***3.3 Slaughter, dressing, and processing***

340 Colonization of *Campylobacter* in broilers during rearing and transportation steps
341 contributes to the contamination of this bacteria in the slaughterhouse causing cross-
342 contamination. During the slaughter process, the transmission of *Campylobacter*-infected flocks
343 to non-infected (Shange et al., 2019; Umar et al., 2016) is mainly due to the leaking of
344 contaminated feces from visceral rupture to the chickens (García-Sánchez et al., 2019; Hermans
345 et al., 2011). In the European Union (EU), it was observed that batches of broilers whose
346 intestines were colonized with *Campylobacter* yielded carcasses with high numbers of
347 *Campylobacter* (EFSA, Panel on Biological Hazards, 2011). However, *Campylobacter* counts on
348 carcasses varied among slaughterhouses. These differences found on the levels of
349 *Campylobacter* loads among slaughterhouses might be related to the different hygiene practices
350 used between countries. Interestingly, *Campylobacter* strains in chickens are not necessarily the
351 same as those isolated from processed carcasses, which suggests that cross-contamination occurs
352 during processing (Slader et al., 2002).

353 This cross-contamination can occur during the entire slaughter, dressing, and processing
354 including the chilling room, plucking, evisceration and portioning areas, or via machinery, work
355 surfaces, process water and air (Allen et al., 2003; Arnold and Silvers, 2000; Corry and Atabay,
356 2003; Haas et al., 2005; Hue et al., 2010; Johnsen et al., 2007). Overall, based on the most recent
357 studies considered in this review, the highest prevalence of *Campylobacter* spp. was detected on

358 the defeathering, evisceration, operation tables, worker's gloves, shackles, and conveyor belt
359 equipment, while the lowest prevalence of *Campylobacter* spp. was detected on the sink, floor
360 and chopping boards and knife swabs during slaughter, dressing, and processing as shown in
361 Table 3 (García-Sánchez et al., 2017; Khan et al., 2018; Tang et al., 2020; Zhang et al., 2018). In
362 agreement with data from the primary production, where the surfaces and equipment in the
363 facility are the main sources for *Campylobacter*, a high prevalence of this bacteria in broiler
364 samples and carcasses has been also reported during the slaughter, dressing and processing
365 (Table 3) (Carrillo et al., 2014; Casagrande Proietti et al., 2018; García-Sánchez, Melero, Diez,
366 Jaime, & Rovira, 2018; Ingesa-Capaccioni et al., 2016; Khan et al., 2018; Korsak, Maćkiw,
367 Rożynek, & Żyłowska, 2015; Perez-Arnedo & Gonzalez-Fandos, 2019; Williams & Oyarzabal,
368 2012; Zhang et al., 2018).

369 The variation of the prevalence of *Campylobacter* spp. during the different processes'
370 steps can also be explained due to technical aspects of *Campylobacter* detection (sampling
371 procedures, storage, DNA extraction, selection of targeting region, and PCR primers and the
372 sequencing platforms used). Culture-based isolation approaches is considered as a standard
373 method for the detection and enumeration of the different *Campylobacter* spp. of products
374 intended for human consumption, animal feeding, environmental samples in the area of food and
375 feed production, and samples from the primary production stage (ISO, 2017). However, the
376 limitations of this technique rely on the difficulties to isolate *Campylobacter* from samples with
377 heavy contamination and the rapid loss in cultivability of isolates. To overcome the challenges in
378 traditional phenotype-based methods for the identification of *Campylobacter*, different DNA-
379 based approaches have become widely used due to the speed, and reproducibility to confirm
380 *Campylobacter* identification (Johannessen et al., 2020). Nevertheless, besides the great

381 advances made in establishing a less time-consuming sampling protocol and more amenable to
382 couple with DNA-base methods, currently, there is no standard procedure for a fast screening of
383 *Campylobacter* at the retail level. One also notes that pathogenesis or virulence factors that
384 certain *Campylobacter* sequence types may have is an important feature to consider for
385 predicting future *Campylobacter* outbreaks and accurate identification in the context of risk
386 assessment.

387

388 4. CONTROL STRATEGIES

389 A direct relationship between the reduction of *Campylobacter*'s load at the different broiler
390 production stages and the reduction of public health risk has been linked to an effective control
391 strategy. Reducing the numbers of *Campylobacter* on the carcasses by one log₁₀-unit, would
392 reduce the public health risk by between 50 and 90%, and, reducing counts by more than two
393 log₁₀-units would reduce the public health risk by more than 90% (European Food Safety
394 Authority Panel on Biological Hazards, 2011). To reduce *Campylobacter* loads, the EU has
395 recently developed the Commission Regulation (EU) 2017/1495, which sets microbiological
396 limits regarding *Campylobacter* spp. in carcasses of broilers (European Commission, 2017). In
397 recent years several *Campylobacter* control strategies have been developed; most of them
398 focused on the reduction of *Campylobacter* colonization at the farm level which consequently
399 decreased *Campylobacter* loads into the slaughterhouse, resulting in a low concentration or
400 absence of the pathogen on the final product (Wagenaar et al., 2006).

401 The main strategies to control *Campylobacter* spp. colonization at the farm level is based
402 on the reduction of environmental exposure (biosecurity and hygienic measures), the increase of

403 broiler resistance to colonization (competitive exclusion, vaccination, application of pre-and pro-
404 biotics, organic acids, or phytochemicals etc.), the use of alternative antimicrobials
405 (bacteriophage therapy and bacteriocin treatment), and/or selection of specific breeding to
406 increase the resistance of broiler chickens to colonization (European Food Safety Authority
407 Panel on Biological Hazards, 2011; Umar et al., 2016). Besides, the BIOHAZ sets sanitation
408 practices during thinning to prevent *Campylobacter* from entering broiler houses at primary
409 production, and the application and monitor system of the decontamination of carcasses (using
410 chemical or physical treatments) are recommended (EFSA, Panel on Biological Hazards, 2011).

411 At the transportation stage the improvement of hygienic measures by removing feed and
412 litter, cleaning and disinfecting transport crates, are the main strategies studied (Meunier et al.,
413 2016). However, besides the importance of the transportation step, at the moment the BIOHAZ
414 has not published any recommendation to prevent and/or reduce the contamination of
415 *Campylobacter* during this processing step. At slaughter, dressing, and processing the most
416 common and effective strategies used to reduce *Campylobacter* loads is the application of
417 specific food safety protocols and strict hygienic practices (HACCP), separating *Campylobacter*-
418 infected flocks from non-infected, physical treatments (scalding, chilling) and chemical
419 decontamination of carcasses using chlorine compounds or chlorine-based antimicrobials
420 (Osimani et al., 2017; Silva et al., 2011). One of the disadvantages of using physical treatments
421 is that it contributes to the change of organoleptic properties of the food products, which would
422 make them less desirable to the consumers. In addition, physical decontamination is allowed in
423 the United States but not in the EU.

424 In the EU, bacteriophages or bacteriocins in the feed are used to reduce the load of
425 *Campylobacter* in the GI tract of broilers before slaughtering, a reduction of the slaughter age of

426 broilers, implementation and improvement of the sanitation practices during slaughter (including
427 the design of adequate equipment with the prevention of fecal leakage), and training food
428 handlers with better hygienic practices to prevent or reduce the *Campylobacter* colonization in
429 the slaughter, dressing and processing steps are recommendations made by the EFSA to promote
430 good processing practices (EFSA, Panel on Biological Hazards, 2011).

431

432 **5. THE APPLICATION OF BACTERIOPHAGES TO COMBAT**

433 ***CAMPYLOBACTER* IN BROILER PRODUCTION**

434 Despite the extensive efforts from the broiler industry, food safety authorities, and
435 academia, there is no effective, reliable, and practical intervention control strategy able to reduce
436 the prevalence of *Campylobacter* from the farm-to-fork process. As the incidence of antibiotic-
437 resistant *Campylobacter* strains is increasing, the development of novel non-antibiotic anti-
438 *Campylobacter* treatments is becoming critical (Johnson et al., 2017). Treatment strategies that
439 have shown highly promising results for *Campylobacter* control in broiler chickens are currently
440 under development. Among them, the use of specific bacteriophages (phages) as biocontrol
441 agents is considered one of the most promising strategies to reduce the prevalence of
442 *Campylobacter* in the broiler production chain (Atterbury et al., 2003; Carvalho et al., 2010; El-
443 Shibiny et al., 2009; Fischer et al., 2013; Hammerl et al., 2014). Bacteriophages are viruses that
444 specifically infect and kill bacteria, widely distributed in the environment from the human GI
445 tract to the deep ocean, and often naturally present in animals such as, broilers (Dion et al., 2020;
446 Nafarrate et al., 2021). The interest in using phages as a safety strategy in food production relies
447 on its selectivity towards the pathogen of concern, it is harmless to humans, animals, and plants,
448 and does not affect the existing commensal microbiota or alter food properties.

449 *Campylobacter*-specific phage cocktail (phiCcolBB12, phiCcolBB35, and phiCcolBB37) has
450 been applied at pre-slaughter and post-slaughter stages to reduce bacterial loads (Carvalho et al.,
451 2012). In detail, the application of *Campylobacter*-specific bacteriophages in the broiler
452 production chain has been tested in several studies, focusing on the administration of
453 bacteriophages into the drinking water during the rearing cycle of broilers (Loc Carrillo et al.,
454 2005; El-Shibiny et al., 2009; Fischer et al., 2013; Hammerl et al., 2014; Kittler et al., 2013;
455 Richards, Connerton, & Connerton, 2019) or using phages on raw and processed meat or raw
456 liver (Atterbury et al., 2003; Firlieyanti et al., 2016; Goode et al., 2003).

457 During the rearing cycle of broilers, the reduction rates in the cecal content achieved by the
458 addition of single bacteriophages or bacteriophage cocktails (phage NCTC 12673, 12674, and
459 12678) showed promising potential reducing bacterial loads between 2.5 to 3.2 log CFU/g
460 (Fischer et al., 2013; Kittler et al., 2013). In addition, it was demonstrated that no adverse effects
461 on the broiler microbiota occur after administering a phage cocktail (CP20 and CP30A) in
462 contrast to administering broad-spectrum antibiotic treatments, which can yield dysbiosis in the
463 gut microbiota (Richards et al., 2019). Overall, the results from most of the studies in broilers
464 conclude that bacteriophages (NCTC 12672, 12673, 12674, 12678, 12669, 12671, 12684, CP8,
465 CP34, CP81, Cj6, phiCcolIBB35, phiCcolIBB37, and phiCcolIBB12) can effectively reduce
466 *Campylobacter* levels if they are administered 24-48 h prior to slaughter (Ushanov et al., 2020).

467 Besides the use of bacteriophages in the primary production, a mean reduction of
468 approximately one log CFU/g of *Campylobacter* on broiler products was obtained using single
469 bacteriophages (NCTC 12674 and 12673) during meat processing (Atterbury et al., 2003; Goode
470 et al., 2003). In this case, different authors suggest the application of bacteriophages
471 (phiCcolIBB35, phiCcolIBB37, and phiCcolIBB12) at high titers to achieve successful

472 reductions in *Campylobacter* counts (Carvalho et al., 2012). In addition, bacteriophages (CP8
473 and CP30) have also shown successful dispersal of *Campylobacter* biofilms and a reduction of
474 viable cells ranging from one to three log CFU/cm² (Siringan et al., 2011), indicating an
475 additional potential field for phage application to target biofilms in meat processing facilities.

476 From an epidemiological point of view, the bacteriophage treatment in the production chain
477 can contribute to a drastic reduction of the infection risk for the consumer. According to the
478 model prediction, phage therapy could lead to a reduction of *Campylobacter* in the fecal matter
479 and on the surface of broiler chicken meat (one log each), resulting on a 90% risk reduction for
480 consumer infection (Havelaar et al., 2007). Clinical data suggest the tolerability and/or
481 effectiveness of phage therapy to reduce antibiotic-resistant infections in humans, but also phage
482 resistance (El Haddad et al., 2019; Zhvania et al., 2017). Bacteria can promote phage attack
483 mainly through spontaneous chromosomal mutations governed by Darwinian dynamics, leading
484 to the emergence of phage resistance and consequently treatment failure (Luong et al., 2020).
485 Identifying new phages with different binding sites to improve efficacy may aid in the prevention
486 of problems related to phage resistance (Wright et al., 2019).

487 More research is needed to find routes of administration, phage selection, the order of phage
488 exposure frequency of administration, dosage, phage resistance, pharmacokinetic and
489 pharmacodynamic properties of the phages, and improve bacteriophage efficacy against
490 *Campylobacter* and broiler meat safety. It is worth noting that bacteriophages should not be
491 considered as a substitute for the control strategies developed so far, but rather seen as a
492 complementary strategy. Successful control of *Campylobacter* could probably be achieved by
493 implementing strict biosecurity and hygiene measures in combination with bacteriophage
494 treatments.

495

496 **6. CONCLUSION**

497 The high occurrence of *Campylobacter* along the broiler production chain is a serious threat
498 to public health. This review revealed that abiotic factors have the potential to contribute to
499 cross-contamination of *Campylobacter*. Furthermore, the transfer of contaminated content of the
500 bird intestine, or persisting biofilm on equipment/surfaces represents likely the source of cross-
501 contamination during the broiler production. Effective *Campylobacter* control measures along
502 the whole broiler production chain are, therefore, needed to improve broiler meat safety,
503 resulting in a reduction of the incidence of human campylobacteriosis. The use of
504 *Campylobacter*-specific bacteriophages at different points from farm-to-fork (in livestock,
505 slaughter, and/or processing facilities) has been proposed as an additional strategy of a
506 multistage bio-security measure to assure safer chicken products for the consumer. The use of
507 multi-omics approaches can help us to increase our understanding of the ability of this foodborne
508 pathogen to persist through the water and the food chain, its environmental niche, and how it
509 interacts with bacteriophages. Progress in this field will help us to better understand how to
510 assess the environmental conditions and nutritional requirements to reduce the risk of
511 *Campylobacter* contamination in the broiler production chain.

512

513 **AUTHORS CONTRIBUTIONS.** Conceptualization, J.MG, L.C, K.R; Data curation: J.MG,
514 I.N, L.L; Project administration, K.R; Investigation related to chapter 1, 2 and 3, J.MG;
515 Investigation related to chapter 4, J.MG, A.L, I.N; Investigation related to chapter 5, A.L, I.N,

516 L.L; Validation, Review and Editing, K.R, L.L, I.F, L.C, A.L, I.N, J.MG; Writing-original Draft
517 Preparation, J.MG; Supervision, K.R.

518

519 **CONFLICTS OF INTEREST.** The authors declare no conflict of interest.

520

521 **ACKNOWLEDGMENT.** The authors gratefully acknowledge Maria Rita Corvaglia for
522 helping in construction of the database. This research was supported by the European Knowledge
523 and Innovation Community (KIC), within the EIT Food program “C-SNIPER: *Campylobacter*-
524 Specific Nullification via Innovative Phage-mediated Enteropathogen Reduction” (Project ID
525 19141).

526

527 **FUNDING.** This research did not receive any specific grant from funding agencies in the
528 public, commercial, or not-for-profit sectors.

529

530 **7. REFERENCES**

531 Ali, A., Soares, S.C., Santos, A.R., Guimarães, L.C., Barbosa, E., Almeida, S.S., Abreu, V.A.C.,
532 Carneiro, A.R., Ramos, R.T.J., Bakhtiar, S.M., Hassan, S.S., Ussery, D.W., On, S., Silva,
533 A., Schneider, M.P., Lage, A.P., Miyoshi, A., Azevedo, V., 2012. *Campylobacter fetus*
534 subspecies: Comparative genomics and prediction of potential virulence targets. *Gene* 508,
535 145–156. <https://doi.org/10.1016/j.gene.2012.07.070>

536 Allen, V.M., Tinker, D.B., Hinton, M.H., Wathes, C.M., 2003. Dispersal of micro-organisms in
537 commercial defeathering systems. *Br. Poult. Sci.* 44, 53–59.
538 <https://doi.org/10.1080/0007166031000085436>

539 Apajalahti, J., 2005. Comparative gut microflora, metabolic challenges, and potential
540 opportunities of gastrointestinal. *J. Appl. Poult. Res.* 14, 444–453.
541 <https://doi.org/https://doi.org/10.1093/japr/14.2.444>

542 Arnold, J., Silvers, S., 2000. Comparison of poultry processing equipment surfaces for
543 susceptibility to bacterial attachment and biofilm formation. *Poult. Sci.* 79, 1215–1221.
544 <https://doi.org/https://doi.org/10.1093/ps/79.8.1215>

545 Atterbury, R.J., Connerton, P.L., Dodd, C.E.R., Rees, C.E.D., Connerton, I.F., 2003. Application
546 of host-specific bacteriophages to the surface of chicken skin leads to a reduction in
547 recovery of *Campylobacter jejuni*. *Appl. Environ. Microbiol.* 69, 6302–6306.
548 <https://doi.org/10.1128/AEM.69.10.6302-6306.2003>

549 Bacon, D.J., Alm, R., Burr, D.H., Hu, L., Kopecko, D.J., Ewing, C.P., Trust, T.J., Guerry, P.,
550 2000. Involvement of a plasmid in virulence of *Campylobacter jejuni* 81-176. *Infect.*
551 *Immun.* 68, 4384–4390. <https://doi.org/10.1128/IAI.68.8.4384-4390.2000>

552 bacterio.net, n.d. List of prokaryotic names with standing in nomenclature (LPSN): Genus,
553 *Campylobacter* [WWW Document]. URL <https://www.bacterio.net/genus/campylobacter>
554 (accessed 3.19.21).

555 Ballou, A.L., Ali, R.A., Mendoza, M.A., Ellis, J.C., Hassan, H.M., Croom, W.J., Koci, M.D.,
556 2016. Development of the chick microbiome: How early exposure influences future

557 microbial diversity. *Front. Vet. Sci.* 3, 1–12. <https://doi.org/10.3389/fvets.2016.00002>

558 Bang, D.D., Wedderkopp, A., Pedersen, K., Madsen, M., 2002. Rapid PCR using nested primers
559 of the 16s rRNA and the hippuricase (*hipO*) genes to detect *Campylobacter jejuni* and
560 *Campylobacter coli* in environmental samples. *Mol. Cell. Probes* 16, 359–369.
561 <https://doi.org/10.1006/mcpr.2002.0434>

562 Barker, C.R., Painset, A., Swift, C., Jenkins, C., Godbole, G., Maiden, M.C.J., Dallman, T.J.,
563 2020. Microevolution of *Campylobacter jejuni* during long-term infection in an
564 immunocompromised host. *Sci. Rep.* 10. <https://doi.org/10.1038/s41598-020-66771-7>

565 Berndtson, E., Danielsson-Tham, M.L., Engvall, A., 1996. *Campylobacter* incidence on a
566 chicken farm and the spread of *Campylobacter* during the slaughter process. *Int. J. Food*
567 *Microbiol.* 32, 35–47. [https://doi.org/10.1016/0168-1605\(96\)01102-6](https://doi.org/10.1016/0168-1605(96)01102-6)

568 Bull, S.A., Thomas, A., Humphrey, T., Ellis-Iversen, J., Cook, A.J., Lovell, R., Jorgensen, F.,
569 2008. Flock health indicators and *Campylobacter* spp. in commercial housed broilers reared
570 in Great Britain. *Appl. Environ. Microbiol.* 74, 5408–5413.
571 <https://doi.org/10.1128/AEM.00462-08>

572 Cardinale, E., Tall, F., Guèye, E.F., Cisse, M., Salvat, G., 2004. Risk factors for *Campylobacter*
573 spp. infection in Senegalese broiler-chicken flocks. *Prev. Vet. Med.* 64, 15–25.
574 <https://doi.org/10.1016/j.prevetmed.2004.03.006>

575 Carrillo, C.D., Plante, D., Iugovaz, I., Kenwell, R., Belanger, G., Boucher, F., Poulin, N.,
576 Trottier, Y.L., 2014. Method-dependent variability in determination of prevalence of
577 *Campylobacter jejuni* and *Campylobacter coli* in Canadian retail poultry. *J. Food Prot.* 77,

578 1682–1688. <https://doi.org/10.4315/0362-028x.jfp-14-133>

579 Carvalho, C.M., Gannon, B.W., Halfhide, D.E., Santos, S.B., Hayes, C.M., Roe, J.M., Azeredo,
580 J., 2010. The in vivo efficacy of two administration routes of a phage cocktail to reduce
581 numbers of *Campylobacter coli* and *Campylobacter jejuni* in chickens. *BMC Microbiol.* 10,
582 232. <https://doi.org/https://doi.org/10.1186/1471-2180-10-232>

583 Casagrande Proietti, P., Pergola, S., Bellucci, S., Menchetti, L., Miraglia, D., Franciosini, M.P.,
584 2018. Occurrence and antimicrobial susceptibility of *Campylobacter* spp. on fresh and
585 refrigerated chicken meat products in Central Italy. *Poult. Sci.* 97, 2895–2901.
586 <https://doi.org/10.3382/ps/pey147>

587 Chen, X., Naren, G.-W., Wu, C.-M., Wang, Y., Dai, L., Xia, L.-N., Luo, P.-J., Zhang, Q., Shen,
588 J.-Z., 2010. Prevalence and antimicrobial resistance of *Campylobacter* isolates in broilers
589 from China. *Vet. Microbiol.* 144, 133–139. <https://doi.org/10.1016/j.vetmic.2009.12.035>

590 Chintoan-Uta, C., Wisedchanwet, T., Glendinning, L., Bremner, A., Psifidi, A., Vervelde, L.,
591 Watson, K., Watson, M., Stevens, M.P., 2020. Role of cecal microbiota in the differential
592 resistance of inbred chicken lines to colonization by *campylobacter jejuni*. *Appl. Environ.*
593 *Microbiol.* 86. <https://doi.org/10.1128/AEM.02607-19>

594 Christl, S.U., Bartram, H.P., Paul, A., Kelber, E., Scheppach, W., Kasper, H., 1997. Bile acid
595 metabolism by colonic bacteria in continuous culture: Effects of starch and pH. *Ann. Nutr.*
596 *Metab.* 41, 45–51. <https://doi.org/10.1159/000177977>

597 Clavijo, V., Flórez, M.J. V, 2018. The gastrointestinal microbiome and its association with the
598 control of pathogens in broiler chicken production: A review. *Poult. Sci.* 97, 1006–1021.

599 <https://doi.org/10.3382/ps/pex359>

600 Corry, J.E.L., Atabay, H.I., 2003. Poultry as a source of *Campylobacter* and related organisms. *J.*
601 *Appl. Microbiol.* 90, 96S-114S. <https://doi.org/10.1046/j.1365-2672.2001.01358.x>

602 Crhanova, M., Hradecka, H., Faldynova, M., Matulova, M., Havlickova, H., Sisak, F., Rychlik,
603 I., 2011. Immune response of chicken gut to natural colonization by gut microflora and to
604 *Salmonella enterica* serovar enteritidis infection. *Infect. Immun.* 79, 2755–2763.
605 <https://doi.org/10.1128/iai.01375-10>

606 Danzeisen, J.L., Kim, H.B., Isaacson, R.E., Tu, Z.J., Johnson, T.J., 2011. Modulations of the
607 chicken cecal microbiome and metagenome in response to anticoccidial and growth
608 promoter treatment. *PLoS One* 6. <https://doi.org/10.1371/journal.pone.0027949>

609 Deng, W., Dittoe, D.K., Pavilidis, H.O., Chaney, W.E., Yang, Y., Ricke, S.C., 2020. Current
610 perspectives and potential of probiotics to limit foodborne *Campylobacter* in poultry. *Front.*
611 *Microbiol.* 11. <https://doi.org/10.3389/fmicb.2020.583429>

612 Dibner, J.J., Richards, J.D., 2005. Antibiotic growth promoters in agriculture: History and mode
613 of action actual usage of antimicrobials in Denmark voluntary and legislated bans. *Poult.*
614 *Sci.* 84, 634–643. <https://doi.org/https://doi.org/10.1093/ps/84.4.634>

615 Dion, M.B., Oechslin, F., Moineau, S., 2020. Phage diversity, genomics and phylogeny. *Nat.*
616 *Rev. Microbiol.* 18, 125–138. <https://doi.org/10.1038/s41579-019-0311-5>

617 Djennad, A., Nichols, G., Loiacono, G., Fleming, L., Kessel, A., Kovats, S., Lake, I., Sarran, C.,
618 Elson, R., Lane, C., Hoesser, C., Bailey, T., 2017. The seasonality and effects of temperature

619 and rainfall on *Campylobacter* infections. *Int. J. Popul. Data Sci.* 1, 1–10.
620 <https://doi.org/10.23889/ijpds.v1i1.51>

621 El-Shibiny, A., Scott, A., Timms, A., Metawea, Y., Connerton, P., Connerton, I., 2009.
622 Application of a group II campylobacter bacteriophage to reduce strains of campylobacter
623 jejuni and campylobacter coli colonizing broiler chickens. *J. Food Prot.* 72, 733–740.
624 <https://doi.org/10.4315/0362-028X-72.4.733>

625 El Haddad, L., Harb, C.P., Gebara, M.A., Stibich, M.A., Chemaly, R.F., 2019. A systematic and
626 critical review of bacteriophage therapy against multidrug-resistant ESKAPE organisms in
627 humans. *Clin. Infect. Dis.* 69, 167–178. <https://doi.org/10.1093/cid/ciy947>

628 Elmi, A., Nasher, F., Dorrell, N., Wren, B., Gundogdu, O., 2021. Revisiting *Campylobacter*
629 jejuni virulence and fitness factors: Role in sensing, adapting, and competing. *Front. Cell.*
630 *Infect. Microbiol.* 10. <https://doi.org/10.3389/fcimb.2020.607704>

631 European Commission, 2017. Commission Regulation (EU) 2017/1495 of 23 August 2017
632 amending Regulation (EC) No 2073/2005 as regards *Campylobacter* in broiler carcasses.
633 *Off. J. Eur. Union* L 218, 1–6.

634 European Food Safety Authority (EFSA), 2010. Analysis of the baseline survey on the
635 prevalence of *Campylobacter* in broiler batches and of *Campylobacter* and *Salmonella* on
636 broiler carcasses, in the EU, 2008 - Part B. *EFSA J.* 8, 1522.
637 <https://doi.org/10.2903/j.efsa.2010.1522>

638 European Food Safety Authority and European Centre of Disease Prevention and Control (EFSA
639 and ECDC), 2019. The European union one health 2018 zoonoses report. *EFSA J.* 17.

640 <https://doi.org/10.2903/j.efsa.2019.5926>

641 European Food Safety Authority Panel on Biological Hazards, (BIOHAZ), 2011. Scientific
642 opinion on *Campylobacter* in broiler meat production: control options and performance
643 objectives and/or targets at different stages of the food chain. *EFSA J.* 9, 2105.
644 <https://doi.org/10.2903/j.efsa.2011.2105>

645 Firleyanti, A.S., Connerton, P.L., Connerton, I.F., 2016. *Campylobacters* and their
646 bacteriophages from chicken liver: The prospect for phage biocontrol. *Int. J. Food*
647 *Microbiol.* 237, 121–127. <https://doi.org/10.1016/j.ijfoodmicro.2016.08.026>

648 Fischer, S., Kittler, S., Klein, G., Glünder, G., 2013. Impact of a single phage and a phage
649 cocktail application in broilers on reduction of *Campylobacter jejuni* and development of
650 resistance. *PLoS One* 8, 1–13. <https://doi.org/10.1371/journal.pone.0078543>

651 García-Sánchez, L., Melero, B., Diez, A.M., Jaime, I., Rovira, J., 2018. Characterization of
652 *Campylobacter* species in Spanish retail from different fresh chicken products and their
653 antimicrobial resistance. *Food Microbiol.* 76, 457–465.
654 <https://doi.org/10.1016/j.fm.2018.07.004>

655 García-Sánchez, L., Melero, B., Jaime, I., Hänninen, M.L., Rossi, M., Rovira, J., 2017.
656 *Campylobacter jejuni* survival in a poultry processing plant environment. *Food Microbiol.*
657 65, 185–192. <https://doi.org/10.1016/j.fm.2017.02.009>

658 García-Sánchez, L., Melero, B., Jaime, I., Rossi, M., Ortega, I., Rovira, J., 2019. Biofilm
659 formation, virulence and antimicrobial resistance of different *Campylobacter jejuni* isolates
660 from a poultry slaughterhouse. *Food Microbiol.* 83, 193–199.

661 <https://doi.org/10.1016/j.fm.2019.05.016>

662 Gharib Naseri, K., Rahimi, S., Khaki, P., 2012. Comparison of the effects of probiotic, organic
663 acid and medicinal plant on campylobacter jejuni challenged broiler chickens. *J. Agric. Sci.*
664 *Technol.* 14, 1485–1496.

665 Gong, J., Forster, R.J., Yu, H., Chambers, J.R., Wheatcroft, R., Sabour, P.M., Chen, S., 2002.
666 Molecular analysis of bacterial populations in the ileum of broiler chickens and comparison
667 with bacteria in the cecum. *FEMS Microbiol. Ecol.* 41, 171–179.
668 <https://doi.org/https://doi.org/10.1111/j.1574-6941.2002.tb00978.x>

669 Gong, J., Si, W., Forster, R.J., Huang, R., Yu, H., Yin, Y., Yang, C., Han, Y., 2007. 16S rRNA
670 gene-based analysis of mucosa-associated bacterial community and phylogeny in the
671 chicken gastrointestinal tracts: From crops to ceca. *FEMS Microbiol. Ecol.* 59, 147–157.
672 <https://doi.org/10.1111/j.1574-6941.2006.00193.x>

673 Goode, D., Allen, V.M., Barrow, P.A., 2003. Reduction of experimental Salmonella and
674 Campylobacter contamination of chicken skin by application of lytic bacteriophages. *Appl.*
675 *Environ. Microbiol.* 69, 5032–5036. <https://doi.org/10.1128/AEM.69.8.5032-5036.2003>

676 Haas, D., Posch, J., Schmidt, S., Wüst, G., Sixl, W., Feierl, G., Marth, E., Reinthaler, F.F., 2005.
677 A case study of airborne culturable microorganisms in a poultry slaughterhouse in Styria,
678 Austria. *Aerobiologia (Bologna)*. 21, 193–201. <https://doi.org/10.1007/s10453-005-9003-x>

679 Hald, B., Skovgård, H., Pedersen, K., Bunkenborg, H., 2008. Influxed insects as vectors for
680 Campylobacter jejuni and Campylobacter coli in Danish broiler houses. *Poult. Sci.* 87,
681 1428–1434. <https://doi.org/10.3382/ps.2007-00301>

682 Hald, B., Wedderkopp, A., Madsen, M., 2000. Thermophilic *Campylobacter* spp. in Danish
683 broiler production: A cross-sectional survey and a retrospective analysis of risk factors for
684 occurrence in broiler flocks. *Avian Pathol.* 29, 123–131.
685 <https://doi.org/10.1080/03079450094153>

686 Hammerl, J.A., Jäckel, C., Alter, T., Janzcyk, P., Stingl, K., Knüver, M.T., Hertwig, S., 2014.
687 Reduction of *campylobacter jejuni* in broiler chicken by successive application of group II
688 and group III phages. *PLoS One* 9, 1–17. <https://doi.org/10.1371/journal.pone.0114785>

689 Havelaar, A.H., Mangen, M.J.J., De Koeijer, A.A., Bogaardt, M.J., Evers, E.G., Jacobs-Reitsma,
690 W.F., Van Pelt, W., Wagenaar, J.A., De Wit, G.A., Van Der Zee, H., Nauta, M.J., 2007.
691 Effectiveness and efficiency of controlling *Campylobacter* on broiler chicken meat. *Risk*
692 *Anal.* 27, 831–844. <https://doi.org/10.1111/j.1539-6924.2007.00926.x>

693 Herman, L., Heyndrickx, M., Grijspeerdt, K., Vandekerchove, D., Rollier, I., De Zutter, L., 2002.
694 Routes for *Campylobacter* contamination of poultry meat: Epidemiological study study
695 from hatchery to slaughterhouse. *Epidemiol. Infect.* 131, 1169–1180.
696 <https://doi.org/https://doi.org/10.1017/S0950268803001183>

697 Hermans, D., Van Deun, K., Messens, W., Martel, A., Van Immerseel, F., Haesebrouck, F.,
698 Rasschaert, G., Heyndrickx, M., Pasmans, F., 2011. *Campylobacter* control in poultry by
699 current intervention measures ineffective: Urgent need for intensified fundamental research.
700 *Vet. Microbiol.* 152, 219–228. <https://doi.org/10.1016/j.vetmic.2011.03.010>

701 Hue, O., Le Bouquin, S., Laisney, M.J., Allain, V., Lalande, F., Petetin, I., Rouxel, S., Quesne,
702 S., Gloaguen, P.Y., Picherot, M., Santolini, J., Salvat, G., Bougeard, S., Chemaly, M., 2010.

703 Prevalence of and risk factors for *Campylobacter* spp. contamination of broiler chicken
704 carcasses at the slaughterhouse. *Food Microbiol.* 27, 992–999.
705 <https://doi.org/10.1016/j.fm.2010.06.004>

706 Humphrey, T., 2006. Are happy chickens safer chickens? Poultry welfare and disease
707 susceptibility. *Br. Poult. Sci.* 47, 379–391. <https://doi.org/10.1080/00071660600829084>

708 Ijaz, U.Z., Sivaloganathan, L., McKenna, A., Richmond, A., Kelly, C., Linton, M., Stratakos,
709 A.C., Lavery, U., Elmi, A., Wren, B.W., Dorrell, N., Corcionivoschi, N., Gundogdu, O.,
710 2018. Comprehensive longitudinal microbiome analysis of the chicken cecum reveals a shift
711 from competitive to environmental drivers and a window of opportunity for *Campylobacter*.
712 *Front. Microbiol.* 9. <https://doi.org/10.3389/fmicb.2018.02452>

713 Ingesa-Capaccioni, S., Jiménez-Trigos, E., Marco-Jiménez, F., Catalá, P., Vega, S., Marin, C.,
714 2016. *Campylobacter* epidemiology from breeders to their progeny in Eastern Spain. *Poult.*
715 *Sci.* 95, 676–683. <https://doi.org/10.3382/ps/pev338>

716 ISO 10272-2:2017, 2017. Microbiology of the food chain — Horizontal method for the detection
717 , and enumeration of *Campylobacter* spp.

718 Johannessen, G.S., Garofolo, G., Di Serafino, G., Koláčková, I., Karpíšková, R., Wiczorek, K.,
719 Osek, J., Christensen, J., Torp, M., Hoorfar, J., 2020. *Campylobacter* in chicken – Critical
720 parameters for international, multicentre evaluation of air sampling and detection methods.
721 *Food Microbiol.* 90, 103455. <https://doi.org/10.1016/j.fm.2020.103455>

722 Johnsen, G., Kruse, H., Hofshagen, M., 2007. Genotyping of thermotolerant *Campylobacter*
723 from poultry slaughterhouse by amplified fragment length polymorphism. *J. Appl.*

724 Microbiol. 103, 271–279. <https://doi.org/10.1111/j.1365-2672.2006.03238.x>

725 Johnson, T.J., Shank, J.M., Johnson, J.G., 2017. Current and potential treatments for reducing
726 *Campylobacter* colonization in animal hosts and disease in humans. *Front. Microbiol.* 8,
727 2019. <https://doi.org/10.3389/fmicb.2017.00487>

728 Jore, S., Viljugrein, H., Brun, E., Heier, B.T., Borck, B., Ethelberg, S., Hakkinen, M., Kuusi, M.,
729 Reiersen, J., Hansson, I., Olsson Engvall, E., Løfdahl, M., Wagenaar, J.A., van Pelt, W.,
730 Hofshagen, M., 2010. Trends in *Campylobacter* incidence in broilers and humans in six
731 European countries, 1997-2007. *Prev. Vet. Med.* 93, 33–41.
732 <https://doi.org/10.1016/j.prevetmed.2009.09.015>

733 Josefsen, M.H., Bhunia, A.K., Engvall, E.O., Fachmann, M.S.R., Hoorfar, J., 2015. Monitoring
734 *Campylobacter* in the poultry production chain - From culture to genes and beyond. *J.*
735 *Microbiol. Methods* 112, 118–125. <https://doi.org/10.1016/j.mimet.2015.03.007>

736 Kalupahana, R.S., Kottawatta, K.S.A., Kanankege, K.S.T., Van Bergen, M.A.P., Abeynayake,
737 P., Wagenaar, J.A., 2013. Colonization of *Campylobacter* spp. in broiler chickens and
738 laying hens reared in tropical climates with low-biosecurity housing. *Appl. Environ.*
739 *Microbiol.* 79, 393–395. <https://doi.org/10.1128/AEM.02269-12>

740 Kapperud, G., Skjerve, E., Vik, L., Hauge, K., Lysaker, A., Aalmen, I., Ostroff, S.M., Potter, M.,
741 1993. Epidemiological investigation of risk factors for *Campylobacter* colonization in
742 Norwegian broiler flocks. *Epidemiol. Infect.* 111, 245–256.
743 <https://doi.org/10.1017/S0950268800056958>

744 Kers, J.G., Velkers, F.C., Fischer, E.A.J., Hermes, G.D.A., Stegeman, J.A., Smidt, H., 2018.

745 Host and environmental factors affecting the intestinal microbiota in chickens. *Front.*
746 *Microbiol.* 9, 1–14. <https://doi.org/10.3389/fmicb.2018.00235>

747 Khan, J.A., Rathore, R.S., Abulreesh, H.H., Qais, F.A., Ahmad, I., 2018. Prevalence and
748 antibiotic resistance profiles of *Campylobacter jejuni* isolated from poultry meat and related
749 samples at retail shops in Northern India. *Foodborne Pathog. Dis.* 15, 218–225.
750 <https://doi.org/10.1089/fpd.2017.2344>

751 Khan, S., Moore, R.J., Stanley, D., Chousalkar, K.K., 2020. The gut microbiota of laying hens
752 and its manipulation with prebiotics and probiotics to enhance gut health and food safety.
753 *Appl. Environ. Microbiol.* 86. <https://doi.org/10.1128/AEM.00600-20>

754 Kittler, S., Fischer, S., Abdulmawjood, A., Glünder, G., Kleina, G., 2013. Effect of
755 bacteriophage application on *Campylobacter jejuni* loads in commercial broiler flocks.
756 *Appl. Environ. Microbiol.* 79, 7525–7533. <https://doi.org/10.1128/AEM.02703-13>

757 Knarreborg, A., Simon, M. A., Engberg, R. M., Jensen, B. B., & Tannock, G.W., 2002. Effects
758 of dietary fat source and subtherapeutic levels of antibiotic on the bacterial community in
759 the ileum of broiler chickens at various ages. *Appl. Environ. Microbiol.* 68, 1189–1195.
760 <https://doi.org/10.1128/AEM.68.12.5918>

761 Koolman, L., Whyte, P., Bolton, D.J., 2014. An investigation of broiler caecal *Campylobacter*
762 counts at first and second thinning. *J. Appl. Microbiol.* 117, 876–881.
763 <https://doi.org/10.1111/jam.12580>

764 Korsak, D., Maćkiw, E., Rożynek, E., Żyłowska, M., 2015. Prevalence of *Campylobacter* spp. in
765 retail chicken, turkey, pork, and beef meat in Poland between 2009 and 2013. *J. Food Prot.*

766 78, 1024–1028. <https://doi.org/10.4315/0362-028x.jfp-14-353>

767 Kumar, S., Chen, C., Indugu, N., Werlang, G.O., Singh, M., Kim, W.K., Thippareddi, H., 2018.
768 Effect of antibiotic withdrawal in feed on chicken gut microbial dynamics, immunity,
769 growth performance and prevalence of foodborne pathogens. *PLoS One* 13, 1–23.
770 <https://doi.org/10.1371/journal.pone.0192450>

771 Lastovica, A.J., On, S.L.W., Zhang, L., 2014. The family Campylobacteraceae, in: Rosenberg,
772 E., DeLong, E.F., Lory, S., Stackebrandt, E., Thompson, F. (Eds.), *The Prokariotes*.
773 Springer Berlin Heidelberg, Berlin, Heidelberg, pp. 307–335. [https://doi.org/10.1007/978-3-](https://doi.org/10.1007/978-3-642-39044-9_274)
774 [642-39044-9_274](https://doi.org/10.1007/978-3-642-39044-9_274)

775 Lin, J., 2009. Novel approaches for Campylobacter control in poultry. *Foodborne Pathog. Dis.* 6,
776 755–765. <https://doi.org/10.1089/fpd.2008.0247>

777 Linden, C.D., Roth, T.F., 1978. IgG receptors on foetal chick yolk sac. *J. Cell Sci.* 33, 317–328.

778 Loc Carrillo, C., Atterbury, R.J., Dillon, E., Scott, A., Connerton, I.F., Connerton, P.L., 2005.
779 Bacteriophage therapy to reduce Campylobacter jejuni colonization of broiler chickens.
780 *Appl. Environ. Microbiol.* 71, 6554–6563. <https://doi.org/10.1128/AEM.71.11.6554>

781 Lu, J., Idris, U., Harmon, B., Hofacre, C., Maurer, J., Lee, M., 2003. Diversity and succession of
782 the intestinal bacterial community of the maturing. *Appl. Environ. Microbiol.* 69, 6816–
783 6824. <https://doi.org/10.1128/AEM.69.11.6816>

784 Luong, T., Salabarria, A.-C., Roach, D.R., 2020. Phage therapy in the resistance era: Where do
785 we stand and where are we going? *Clin. Ther.* 42, 1659–1680.

786 <https://doi.org/10.1016/j.clinthera.2020.07.014>

787 Lyngstad, T.M., Jonsson, M.E., Hofshagen, M., Heier, B.T., 2008. Risk factors associated with
788 the presence of *Campylobacter* species in Norwegian broiler flocks. *Poult. Sci.* 87, 1987–
789 1994. <https://doi.org/10.3382/ps.2008-00132>

790 Marotta, F., Garofolo, G., Di Donato, G., Aprea, G., Platone, I., Cianciavichia, S., Alessiani, A.,
791 Di Giannatale, E., 2015. Population diversity of *Campylobacter jejuni* in poultry and its
792 dynamic of contamination in chicken meat. *Biomed Res. Int.* 2015.
793 <https://doi.org/10.1155/2015/859845>

794 McKenna, A., Ijaz, U.Z., Kelly, C., Linton, M., Sloan, W.T., Green, B.D., Lavery, U., Dorrell,
795 N., Wren, B.W., Richmond, A., Corcionivoschi, N., Gundogdu, O., 2020. Impact of
796 industrial production system parameters on chicken microbiomes: mechanisms to improve
797 performance and reduce *Campylobacter*. *Microbiome* 8, 128.
798 <https://doi.org/10.1186/s40168-020-00908-8>

799 Meunier, M., Guyard-Nicodème, M., Dory, D., Chemaly, M., 2016. Control strategies against
800 *Campylobacter* at the poultry production level: Biosecurity measures, feed additives and
801 vaccination. *J. Appl. Microbiol.* 120, 1139–1173. <https://doi.org/10.1111/jam.12986>

802 Nafarrate, I., Mateo, E., Miranda-Cadena, K., Lasagabaster, A., 2021. Isolation, host specificity
803 and genetic characterization of *Campylobacter* specific bacteriophages from poultry and
804 swine sources. *Food Microbiol.* 97. <https://doi.org/10.1016/j.fm.2021.103742>

805 Neill, S.D., Campbell, J.N., Greene, J.A., 1984. *Campylobacter* species in broiler chickens.
806 *Avian Pathol.* 13, 777–785. <https://doi.org/10.1080/03079458408418574>

807 Nicholson, F.A., Groves, S.J., Chambers, B.J., 2005. Pathogen survival during livestock manure
808 storage and following land application. *Bioresour. Technol.* 96, 135–143.
809 <https://doi.org/10.1016/j.biortech.2004.02.030>

810 Oakley, B.B., Kogut, M.H., 2016. Spatial and temporal changes in the broiler chicken cecal and
811 fecal microbiomes and correlations of bacterial taxa with cytokine gene expression. *Front.*
812 *Vet. Sci.* 3, 1–12. <https://doi.org/10.3389/fvets.2016.00011>

813 Oakley, B.B., Lillehoj, H.S., Kogut, M.H., Kim, W.K., Maurer, J.J., Pedroso, A., Lee, M.D.,
814 Collett, S.R., Johnson, T.J., Cox, N.A., 2014. The chicken gastrointestinal microbiome.
815 *FEMS Microbiol. Lett.* 360, 100–112. <https://doi.org/10.1111/1574-6968.12608>

816 Osimani, A., Aquilanti, L., Pasquini, M., Clementi, F., 2017. Prevalence and risk factors for
817 thermotolerant species of *Campylobacter* in poultry meat at retail in Europe. *Poult. Sci.* 96,
818 3382–3391. <https://doi.org/10.3382/ps/pex143>

819 Pedroso, A.A., Batal, A.B., Lee, M.D., 2016. Effect of in ovo administration of an adult-derived
820 microbiota on establishment of the intestinal microbiome in chickens. *Am. J. Vet. Res.* 77,
821 514–526. <https://doi.org/https://doi.org/10.2460/ajvr.77.5.514>

822 Perez-Arnedo, I., Gonzalez-Fandos, E., 2019. Prevalence of *Campylobacter* spp. in poultry in
823 three Spanish farms, a slaughterhouse and a further processing plant. *Foods* 8, 111.
824 <https://doi.org/10.3390/foods8030111>

825 Pielsticker, C., Glünder, G., Rautenschlein, S., 2012. Colonization properties of *Campylobacter*
826 *jejuni* in chickens. *Eur. J. Microbiol. Immunol.* 2, 61–65.
827 <https://doi.org/10.1556/eujmi.2.2012.1.9>

828 Richards, P.J., Connerton, P.L., Connerton, I.F., 2019. Phage biocontrol of campylobacter jejuni
829 in chickens does not produce collateral effects on the gut microbiota. *Front. Microbiol.* 10,
830 1–10. <https://doi.org/10.3389/fmicb.2019.00476>

831 Ricke, S.C., 2003. Perspectives in the use of organic acids and short chain fatty acids as
832 microbials. *Poult. Sci.* 82, 632–639. <https://doi.org/https://doi.org/10.1093/ps/82.4.632>

833 Ridley, A., Morris, V., Gittins, J., Cawthraw, S., Harris, J., Edge, S., Allen, V., 2011. Potential
834 sources of Campylobacter infection on chicken farms: Contamination and control of broiler-
835 harvesting equipment, vehicles and personnel. *J. Appl. Microbiol.* 111, 233–244.
836 <https://doi.org/10.1111/j.1365-2672.2011.05038.x>

837 Ridley, A.M., Morris, V.K., Cawthraw, S.A., Ellis-Iversen, J., Harris, J.A., Kennedy, E.M.,
838 Newell, D.G., Allen, V.M., 2011. Longitudinal molecular epidemiological study of
839 thermophilic campylobacters on one conventional broiler chicken farm. *Appl. Environ.*
840 *Microbiol.* 77, 98–107. <https://doi.org/10.1128/AEM.01388-10>

841 Robyn, J., Rasschaert, G., Hermans, D., Pasmans, F., Heyndrickx, M., 2013. Is allicin able to
842 reduce campylobacter jejuni colonization in broilers when added to drinking water? *Poult.*
843 *Sci.* 92, 1408–1418. <https://doi.org/10.3382/ps.2012-02863>

844 Rossler, E., Signorini, M.L., Romero-Scharpen, A., Soto, L.P., Berisvil, A., Zimmermann, J.A.,
845 Fusari, M.L., Olivero, C., Zbrun, M. V., Frizzo, L.S., 2019. Meta-analysis of the prevalence
846 of thermotolerant Campylobacter in food-producing animals worldwide. *Zoonoses Public*
847 *Health* 1–11. <https://doi.org/10.1111/zph.12558>

848 Sahin, O., Zhang, Q., Meitzler, J.C., Harr, B.S., Morishita, T.Y., Mohan, R., 2001. Prevalence,

849 antigenic specificity, and bactericidal activity of poultry anti-Campylobacter maternal
850 antibodies. *Appl. Environ. Microbiol.* 67, 3951–3957.
851 <https://doi.org/10.1128/AEM.67.9.3951-3957.2001>

852 Sakaridis, I., Ellis, R.J., Cawthraw, S.A., van Vliet, A.H., Stekel, D.J., Penell, J., Chambers, M.,
853 La Ragione, R.M., Cook, A.J., 2018. Investigating the association between the caecal
854 microbiomes of broilers and Campylobacter burden. *Front. Microbiol.* 9, 927.
855 <https://doi.org/10.3389/fmicb.2018.00927>

856 Schroeder, M.W., Eifert, J.D., Ponder, M.A., Schmale, D.G., 2014. Association of
857 Campylobacter spp. levels between chicken grow-out environmental samples and processed
858 carcasses. *Poult. Sci.* 93, 734–741. <https://doi.org/10.3382/ps.2013-03646>

859 Shang, Y., Kumar, S., Oakley, B., Kim, W.K., 2018. Chicken gut microbiota: Importance and
860 detection technology. *Front. Vet. Sci.* 5, 254. <https://doi.org/10.3389/fvets.2018.00254>

861 Shange, N., Gouws, P., Hoffman, L.C., 2019. Campylobacter and Arcobacter species in food-
862 producing animals: prevalence at primary production and during slaughter. *World J.*
863 *Microbiol. Biotechnol.* 35. <https://doi.org/10.1007/s11274-019-2722-x>

864 Sibanda, N., McKenna, A., Richmond, A., Ricke, S.C., Callaway, T., Stratakos, A.C.,
865 Gundogdu, O., Corcionivoschi, N., 2018. A review of the effect of management practices
866 on campylobacter prevalence in poultry farms. *Front. Microbiol.* 9, 1–9.
867 <https://doi.org/10.3389/fmicb.2018.02002>

868 Sierra-Arguello, Y.M., Perdoncini, G., Rodrigues, L.B., Ruschel dos Santos, L., Apellanis
869 Borges, K., Quedi Furian, T., Pippi Salle, C.T., de Souza Moraes, H.L., Pereira Gomes,

870 M.J., Pinheiro do Nascimento, V., 2021. Identification of pathogenic genes in
871 *Campylobacter jejuni* isolated from broiler carcasses and broiler slaughterhouses. *Sci. Rep.*
872 11. <https://doi.org/10.1038/s41598-021-84149-1>

873 Silva, J., Leite, D., Fernandes, M., Mena, C., Gibbs, P.A., Teixeira, P., 2011. *Campylobacter* spp.
874 As a foodborne pathogen: A review. *Front. Microbiol.* 2.
875 <https://doi.org/10.3389/fmicb.2011.00200>

876 Singh, K.M., Shah, T.M., Reddy, B., Deshpande, S., Rank, D.N., Joshi, C.G., 2014. Taxonomic
877 and gene-centric metagenomics of the fecal microbiome of low and high feed conversion
878 ratio (FCR) broilers. *J. Appl. Genet.* 55, 145–154. <https://doi.org/10.1007/s13353-013->
879 0179-4

880 Siringan, P., Connerton, P.L., Payne, R.J.H., Connerton, I.F., 2011. Bacteriophage-mediated
881 dispersal of *Campylobacter jejuni* biofilms. *Appl. Environ. Microbiol.* 77, 3320–3326.
882 <https://doi.org/10.1128/AEM.02704-10>

883 Slader, J., Domingue, G., Mcalpine, K., Owen, R.J., Bolton, F.J., Humphrey, T.J., 2002. Impact
884 of transport crate reuse and of catching and processing on *Campylobacter* and *Salmonella*
885 contamination of broiler chickens. *Society* 68, 713–719.
886 <https://doi.org/10.1128/AEM.68.2.713>

887 Sofka, D., Pfeifer, A., Gleiss, B., Paulsen, P., Hilbert, F., 2015. Changes within the intestinal
888 flora of broilers by colonisation with *Campylobacter jejuni*. *Berl. Munch. Tierarztl.*
889 *Wochenschr.* 128, 104–110.

890 Sommer, H. M., Høg, B. B., Larsen, L. S., Sørensen, A. I. V., Williams, N., Merga, J. Y., Osek,

891 J., 2016. Analysis of farm specific risk factors for *Campylobacter* colonization of broilers in
892 six European countries. *Microb. Risk Anal.* 2, 16–26.
893 <https://doi.org/https://doi.org/10.1016/j.mran.2016.06.002>

894 Stanley, D., Denman, S.E., Hughes, R.J., Geier, M.S., Crowley, T.M., Chen, H., Haring, V.R.,
895 Moore, R.J., 2012. Intestinal microbiota associated with differential feed conversion
896 efficiency in chickens. *Appl. Microbiol. Biotechnol.* 96, 1361–1369.
897 <https://doi.org/10.1007/s00253-011-3847-5>

898 Stern, N.J., Fedorka-Cray, P., Bailey, J.S., Cox, N.A., Craven, S.E., Hiatt, K.L., Mead, G.C.,
899 2001. Distribution of *Campylobacter* spp. in selected U.S. poultry production and
900 processing operations. *J. Food Prot.* 64, 1705–1710. [https://doi.org/10.4315/0362-028x-](https://doi.org/10.4315/0362-028x-64.11.1705)
901 [64.11.1705](https://doi.org/10.4315/0362-028x-64.11.1705)

902 Stern, N.J., Svetoch, E.A., Eruslanov, B. V., Pereygin, V. V., Mitsevich, E. V., Mitsevich, I.P.,
903 Pokhilenko, V.D., Levchuk, V.P., Svetoch, O.E., Seal, B.S., 2006. Isolation of a
904 *Lactobacillus salivarius* strain and purification of its bacteriocin, which is inhibitory to
905 *Campylobacter jejuni* in the chicken gastrointestinal system. *Antimicrob. Agents*
906 *Chemother.* 50, 3111–3116. <https://doi.org/10.1128/AAC.00259-06>

907 Takeshita, N., Watanabe, T., Ishida-Kuroki, K., Sekizaki, T., 2021. Transition of microbiota in
908 chicken cecal droppings from commercial broiler farms. *BMC Vet. Res.* 17.
909 <https://doi.org/10.1186/s12917-020-02688-7>

910 Tang, Y., Jiang, Q., Tang, H., Wang, Z., Yin, Y., Ren, F., Kong, L., Jiao, X., Huang, J., 2020.
911 Characterization and prevalence of *Campylobacter* spp. from broiler chicken rearing period

912 to the slaughtering process in eastern China. *Front. Vet. Sci.* 7.
913 <https://doi.org/10.3389/fvets.2020.00227>

914 Tangkham, W., Janes, M., Lemieux, F., 2016. Prevalence and distribution of *Campylobacter*
915 *jejuni* in small-scale broiler operations. *J. Food Prot.* 79, 75–81.
916 <https://doi.org/10.4315/0362-028x.jfp-15-331>

917 Tegtmeier, N., Sharafutdinov, I., Harrer, A., Soltan Esmaili, D., Linz, B., Backert, S., 2021.
918 *Campylobacter* virulence factors and molecular host-pathogen interactions. *Curr. Top.*
919 *Microbiol. Immunol.* 431, 169–202. https://doi.org/10.1007/978-3-030-65481-8_7

920 Thibodeau, A., Fravalo, P., Yergeau, E., Arsenault, J., Lahaye, L., Letellier, A., 2015. Chicken
921 caecal microbiome modifications induced by *Campylobacter jejuni* colonization and by a
922 non-antibiotic feed additive. *PLoS One* 10. <https://doi.org/10.1371/journal.pone.0131978>

923 Umar, S., Maiyah, A.T., Mushtaq, A., 2016. *Campylobacter* infections in poultry: Update on
924 challenges and potential immune interventions. *Worlds. Poult. Sci. J.* 72, 381–390.
925 <https://doi.org/10.1017/S0043933916000167>

926 Ushanov, L., Lasareishvili, B., Janashia, I., Zautner, A.E., 2020. Application of *Campylobacter*
927 *jejuni* phages: Challenges and perspectives. *Animals* 10.
928 <https://doi.org/10.3390/ani10020279>

929 Vandeplas, S., Beckers, Y., Théwis, A., Dubois-Dauphin, R., Thonart, P., Palm, R., 2010.
930 Prevalence and sources of *Campylobacter* spp. contamination in free-range broiler
931 production in the southern part of Belgium. *Biotechnol. Agron. Soc. Environ.* 14, 279–288.

932 Vispo, C., Karasov, W.H., 1997. The interaction of avian gut microbes and their host: An elusive
933 symbiosis. *Gastrointest. Microbiol.* 116–155. https://doi.org/10.1007/978-1-4615-4111-0_5

934 Wagenaar, J.A., Mevius, D.J., Havelaar, A.H., 2006. *Campylobacter* in primary animal
935 production and control strategies to reduce the burden of human campylobacteriosis. *Rev.*
936 *Sci. Tech.* 25, 581–94.

937 Whyte, P., Collins, J.D., McGill, K., Monahan, C., O’Mahony, H., 2001. The effect of
938 transportation stress on excretion rates of campylobacters in market-age broilers. *Poult. Sci.*
939 80, 817–820. <https://doi.org/10.1093/ps/80.6.817>

940 Williams, A., Oyarzabal, O.A., 2012. Prevalence of *Campylobacter* spp. in skinless, boneless
941 retail broiler meat from 2005 through 2011 in Alabama, USA. *BMC Microbiol.* 12, 184.
942 <https://doi.org/10.1186/1471-2180-12-184>

943 Willis, W.L., Murray, C., Talbott, C., 2002. *Campylobacter* isolation trends of cage versus floor
944 broiler chickens: A one-year study. *Poult. Sci.* 81, 629–631.
945 <https://doi.org/10.1093/ps/81.5.629>

946 World Health Organization (WHO), 2015. Who estimates of the global burden of foodborne
947 diseases. *Foodborne disase Burd. Epidemiol. refeence Gr.* 265.

948 Wright, R.C.T., Friman, V.-P., Smith, M.C.M., Brockhurst, M.A., 2019. Resistance evolution
949 against phage combinations depends on the timing and order of exposure. *MBio* 10.
950 <https://doi.org/10.1128/mBio.01652-19>

951 Zhang, X., Tang, M., Zhou, Q., Zhang, J., Yang, X., Gao, Y., 2018. Prevalence and

952 characteristics of *Campylobacter* throughout the slaughter process of different broiler
953 batches. *Front. Microbiol.* 9, 1–9. <https://doi.org/10.3389/fmicb.2018.02092>

954 Zhang, X., Yin, T., Du, X., Yang, W., Huang, J., Jiao, X., 2017. Occurrence and genotypes of
955 *Campylobacter* species in broilers during the rearing period. *Avian Pathol.* 46, 215–223.
956 <https://doi.org/10.1080/03079457.2016.1248374>

957 Zhvania, P., Hoyle, N.S., Nadareishvili, L., Nizharadze, D., Kutateladze, M., 2017. Phage
958 therapy in a 16-year-old boy with netherton syndrome. *Front. Med.* 4.
959 <https://doi.org/10.3389/fmed.2017.00094>

960

961 **Table legend**

962 **Table 1.** Prevalence of *Campylobacter* spp, *Campylobacter jejuni* and *Campylobacter coli*
963 from different sample types at different sampling points during the broiler primary production
964 using culture-based methods for the detection of *Campylobacter* expresses as percentage or log
965 CFU/g

966 **Table 2.** Prevalence of *Campylobacter* spp, *Campylobacter jejuni* and *Campylobacter coli*
967 from fecal samples and different equipment used to transport broilers during the pre-and post-
968 transportation of broilers to the slaughterhouse using culture-based methods for the detection of
969 *Campylobacter* expresses as percentage

970 **Table 3.** Prevalence of *Campylobacter* spp, *Campylobacter jejuni* and *Campylobacter coli*
971 from different sample types during slaughter, dressing and processing using different types of
972 polymerase chain reaction for the detection of *Campylobacter* expresses as percentage

973

974 **Figure legend**

975 **Figure 1.** Key steps identified and used to assess the prevalence of *Campylobacter* in the
976 broiler production chain. The location of studies investigating the effect of bacteriophage is
977 indicated with a bacteriophage icon. Color in blue is the prevalence of *Campylobacter* spp. and
978 color in yellow is the prevalence of *Campylobacter jejuni*, both expressed as percentage.

979 **Table 1**

FACTORS	DESCRIPTION	SAMPLE	<i>CAMPYLOBACTER</i>	<i>C. JEJUNI</i>	<i>C. COLI</i>	REFERENCE
Broiler age						
	14 d	Cloacal	5.00 %	100.00 %		(Ingesa-Capaccioni et al., 2016)
	42 d	Cloacal	62.00 %	67.00 %		(Ingesa-Capaccioni et al., 2016)
	14	Cloacal	0.00 %			(Perez-Arnedo and Gonzalez-Fandos, 2019)
	42 d	Cloacal	0.00 – 100.00 %			(Perez-Arnedo and Gonzalez-Fandos, 2019)
	31 d	Cloacal	39.30 %	46.00 %	26.00 %	(Tang et al., 2020)
	37 d	Cloacal	60.00 %	74.00 %	38.00 %	(Tang et al., 2020)
	7 d (control housing)	Fecal	5.30 %	5.30 %		(Tangkham et al., 2016)
	42 d (control housing)	Fecal	26.00 %	26.00 %		(Tangkham et al., 2016)
	21 d (control housing)	Fecal	83.30 %	18.70 %		(Tangkham et al., 2016)
	7 d (open housing)	Fecal	0.00 %	0.00 %		(Tangkham et al., 2016)
	21 d (open housing)	Fecal	93.30 %	70.70 %		(Tangkham et al., 2016)
	42 d (open housing)	Fecal	37.30 %	00 - 37.30 %		(Tangkham et al., 2016)
Water						
	Environmentally control		0.00 - 66.70 %	0.00 - 16.70 %		(Tangkham et al., 2016)
	Open		0.00 - 83.30 %	0.00 - 83.30 %		(Tangkham et al., 2016)
	Water of 31 d age		0.00 %			(Tang et al., 2020)
	Water of 37 d age		0.00 %			(Tang et al., 2020)
Feed						
	Environmentally control		0.00 - 83.30 %	0.00 - 33.30 %		(Tangkham et al., 2016)
	Open		0.00 - 33.30 %	0.00 - 33.30 %		(Tangkham et al., 2016)

	Feed of 31 d age	0.00 %		(Tang et al., 2020)
	Feed of 37 d age	0.00 %		(Tang et al., 2020)
Litter				
	Covering shoe	20.00 %		(Schroeder et al., 2014)
Air				
	Gelatin sample	15.00 %		(Schroeder et al., 2014)
	Air filter samples	0.00 - 10.00 %		(Johannessen et al., 2020)
Workers				
	Workers' boots swabs	0.00 - 60.00 %		(Johannessen et al., 2020)
Infrastructure of the farm				
		0.00 - 12.50 %		(Bang et al., 2002)
	Floor 31 d age	0.00 %		(Tang et al., 2020)
	Floor 37 d age	0.00 %		(Tang et al., 2020)
	Bedding 31 d age	10.00 %		(Tang et al., 2020)
	Bedding 37 d age	10.00 %	1.00 %	(Tang et al., 2020)
	Sole 31 d age	4.70 %	1.00 %	(Tang et al., 2020)
	Sole 37 d age	10.50 %	2.00 %	(Tang et al., 2020)
	Net 31 d age	0.00 %		(Tang et al., 2020)
	Net 37 d age	0.00 %		(Tang et al., 2020)
	Stool 31 d age	0.00 %		(Tang et al., 2020)
	Stool 37 d age	6.70 %	1.00 %	(Tang et al., 2020)
Management				
	First thinning	Cecal	27.00 - 100.00 %	(Koolman et al., 2014)
	Second thinning	Cecal	90.00 - 100.00 %	(Koolman et al., 2014)
	Drinking water + allicin	Cecal	5.38 log CFU/g	(Robyn et al., 2013)
	Feed			

	Cecal	4.2 - 7.5 log CFU/g	(Gharib Naseri et al., 2012)
	Fecal	6.3 - 7.2 log CFU/g	(Gharib Naseri et al., 2012)
Drink water + probiotics			
	Cecal	4.1 - 6.6 log CFU/g	(Gharib Naseri et al., 2012)
	Fecal	5.4 - 6.4 log CFU/g	(Gharib Naseri et al., 2012)
Feed + plant derivate			
	Cecal	4.2 - 6.3 log CFU/g	(Gharib Naseri et al., 2012)
	Fecal	5.5 - 6.5 log CFU/g	(Gharib Naseri et al., 2012)
Feed + organic acids			
	Cecal	4.0 - 6.2 log CFU/g	(Gharib Naseri et al., 2012)
	Fecal	4.1 - 5.6 log CFU/g	(Gharib Naseri et al., 2012)
Feed + bacteriocin	Fecal	ND	(Stern et al., 2006)
Feed + bacteriophages	Fecal	5.00 log CFU/g	(Carvalho et al., 2010)

980 Abbreviations: *C. jejuni*; *Campylobacter jejuni*, *C. coli*; *Campylobacter coli*, **ND**; *Not determined*

981

982 **Table 2**

FACTORS	DESCRIPTION	PREVALENCE (%)			REFERENCE
		<i>CAMPYLOBACTER</i>	<i>C. JEJUNI</i>	<i>C. COLI</i>	
Equipment					
	Cage		36.80	ND	(Willis et al., 2002)
	Floor		65.40	ND	(Willis et al., 2002)
	Pre-transportation cage	6.20 - 30.00		ND	(Stern et al., 2001)
	Post-transportation cage	42.50 - 85.00		ND	(Stern et al., 2001)
Fecal					
	Pre-transportation	57.10 - 80.00		ND	(Whyte et al., 2001)
	Post-transportation	60.00 - 80.00		ND	(Whyte et al., 2001)

983 **Abbreviations:** *C. jejuni*; *Campylobacter jejuni*, *C. coli*; *Campylobacter coli*, **ND**; *Not determined*

984

985 **Table 3**


PRODUCTION STAGE	FACTORS	DESCRIPTION	PREVALENCE (%)				REFERENCE	
			CAMPYLOBACTER	C. JEJUNI	C. COLI	AMPLIFICATION GENE		
Slaughter	Environment	Dirty defeathering swabs	60.00	80.00	20.00	<i>hipO-F, hipoO-R, hipO-P, ceuE-F, ceuE-R, ceuE-R, ceuE-P^Δ</i>	(García-Sánchez et al., 2017)	
		Clean defeathering swabs	54.50	90.00	10.00	<i>hipO-F, hipoO-R, hipO-P, ceuE-F, ceuE-R, ceuE-R, ceuE-P^Δ</i>	(García-Sánchez et al., 2017)	
		Dehairing swabs	31.80	3.00	4.00	16S rRNA, <i>mapA</i> , <i>ceuE^C</i>	(Tang et al., 2020)	
		Dirty evisceration swabs	78.00	100.00	0.00	<i>hipO-F, hipoO-R, hipO-P, ceuE-F, ceuE-R, ceuE-R, ceuE-P^Δ</i>	(García-Sánchez et al., 2017)	
		Clean evisceration swabs	56.40	100.00	0.00	<i>hipO-F, hipoO-R, hipO-P, ceuE-F, ceuE-R, ceuE-R, ceuE-P^Δ</i>	(García-Sánchez et al., 2017)	
		Evisceration	31.80	1.00	7.00	16S rRNA, <i>mapA</i> , <i>ceuE^C</i>	(Tang et al., 2020)	
		Dirty sink swabs	100.00	100.00	0.00	<i>hipO-F, hipoO-R, hipO-P, ceuE-F, ceuE-R, ceuE-R, ceuE-P^Δ</i>	(García-Sánchez et al., 2017)	
		Clean sink swabs	20.00	50.00	50.00	<i>hipO-F, hipoO-R, hipO-P, ceuE-F, ceuE-R, ceuE-R, ceuE-P^Δ</i>	(García-Sánchez et al., 2017)	
		Dirty floor swabs	22.70	100.00	0.00	<i>hipO-F, hipoO-R, hipO-P, ceuE-F, ceuE-R, ceuE-R, ceuE-P^Δ</i>	(García-Sánchez et al., 2017)	
		Clean floor swabs	30.00	83.30	16.70	<i>hipO-F, hipoO-R, hipO-P, ceuE-F, ceuE-R, ceuE-R, ceuE-P^Δ</i>	(García-Sánchez et al., 2017)	
		Operation table swabs			10.00 - 80.00	0.00 - 100.00	16S rRNA, <i>mapA</i> , <i>ceuE^C</i>	(Zhang et al., 2018)
		Partition		9.10	1.00	1.00	16S rRNA, <i>mapA</i> , <i>ceuE^C</i>	(Tang et al., 2020)
		Workers' gloves swabs			20.00 - 60.00	40.00 - 80.00	16S rRNA, <i>mapA</i> , <i>ceuE^C</i>	(Zhang et al., 2018)

	Cooling	36.40	5.00	5.00	16S rRNA, <i>mapA</i> , <i>ceuE^C</i>	(Tang et al., 2020)	
Broiler slaughterhouse	Cloacal swabs		0.00 - 63.00	0.00 - 13.60	16S rRNA, <i>mapA</i> , <i>ceuE^C</i>	(Zhang et al., 2018)	
	Carcass after plucking		0.00 - 73.30	0.00 - 85.70	16S rRNA, <i>mapA</i> , <i>ceuE^C</i>	(Zhang et al., 2018)	
	Carcass after evisceration		0.00 - 80.00	0.00 - 80.00	16S rRNA, <i>mapA</i> , <i>ceuE^C</i>	(Zhang et al., 2018)	
	Carcass after washing		0.00 - 76.70	0.00 - 95.20	16S rRNA, <i>mapA</i> , <i>ceuE^C</i>	(Zhang et al., 2018)	
	Carcass after chilling		0.00 - 87.50	0.00 - 100.00	16S rRNA, <i>mapA</i> , <i>ceuE^C</i>	(Zhang et al., 2018)	
	Carcass entrance (41 - 44 days age)	41.40	38.00	22.00	16S rRNA, <i>mapA</i> , <i>ceuE^C</i>	(Tang et al., 2020)	
	Carcass after dehairing	12.90	3.00	23.00	16S rRNA, <i>mapA</i> , <i>ceuE^C</i>	(Tang et al., 2020)	
	Carcass after evisceration	53.40	37.00	69.00	16S rRNA, <i>mapA</i> , <i>ceuE^C</i>	(Tang et al., 2020)	
	Carcass after cooling	14.80	12.00	16.00	16S rRNA, <i>mapA</i> , <i>ceuE^C</i>	(Tang et al., 2020)	
	Carcass after partition	13.60	8.00	18.00	16S rRNA, <i>mapA</i> , <i>ceuE^C</i>	(Tang et al., 2020)	
Dressing	Environment	Dirty shackles swabs	41.80	100.00	0.00	<i>hipO-F</i> , <i>hipoO-R</i> , <i>hipO-P</i> , <i>ceuE-F</i> , <i>ceuE-R</i> , <i>ceuE-P^Δ</i>	(García-Sánchez et al., 2017)
		Clean shackles swabs	38.00	94.70	5.30	<i>hipO-F</i> , <i>hipoO-R</i> , <i>hipO-P</i> , <i>ceuE-F</i> , <i>ceuE-R</i> , <i>ceuE-P^Δ</i>	(García-Sánchez et al., 2017)
		Dirty conveyor belt swabs	87.90	96.60	3.40	<i>hipO-F</i> , <i>hipoO-R</i> , <i>hipO-P</i> , <i>ceuE-F</i> , <i>ceuE-R</i> , <i>ceuE-P^Δ</i>	(García-Sánchez et al., 2017)
		Clean conveyor belt swabs	3.30	100.00	0.00	<i>hipO-F</i> , <i>hipoO-R</i> , <i>hipO-P</i> , <i>ceuE-F</i> , <i>ceuE-R</i> , <i>ceuE-P^Δ</i>	(García-Sánchez et al., 2017)
		Dirty sink swabs	0.00	0.00	0.00	<i>hipO-F</i> , <i>hipoO-R</i> , <i>hipO-P</i> , <i>ceuE-F</i> , <i>ceuE-R</i> , <i>ceuE-P^Δ</i>	(García-Sánchez et al., 2017)
		Clean sink swabs	0.00	0.00	0.00	<i>hipO-F</i> , <i>hipoO-R</i> , <i>hipO-P</i> , <i>ceuE-F</i> , <i>ceuE-R</i> , <i>ceuE-P^Δ</i>	(García-Sánchez et al., 2017)

		Dirty floor swabs	9.10	100.00	0.00	<i>hipO-F, hipO-R, hipO-P, ceuE-F, ceuE-R, ceuE-R, ceuE-P⁴</i>	(García-Sánchez et al., 2017)	
		Clean floor swabs	10.00	100.00	0.00	<i>hipO-F, hipO-R, hipO-P, ceuE-F, ceuE-R, ceuE-R, ceuE-P⁴</i>	(García-Sánchez et al., 2017)	
Broiler		Carcass processing plant	91.00			Not mentioned ^B	(Perez-Arnedo and Gonzalez-Fandos, 2019)	
		Skin on thighs	43.50			<i>glyA, hipO^C</i>	(Casagrande Proietti et al., 2018)	
		Skin off breast	44.40			<i>glyA, hipO^C</i>	(Casagrande Proietti et al., 2018)	
		Legs	54.30			Not mentioned ^B	(Perez-Arnedo and Gonzalez-Fandos, 2019)	
		Breast	46.00			Not mentioned ^B	(Perez-Arnedo and Gonzalez-Fandos, 2019)	
		Wings	87.00			Not mentioned ^B	(Perez-Arnedo and Gonzalez-Fandos, 2019)	
Processing	Environment	Chopping board and knives swabs		14.00		<i>hipO^B</i>	(Khan et al., 2018)	
	Partition		13.60	8.00	18.00	16S rRNA, <i>mapA, ceuE^C</i>	(Tang et al., 2020)	
	Broiler parts		Skin on thighs	51.20			<i>glyA, hipO^C</i>	(Casagrande Proietti et al., 2018)
			Skin off breast	2.70			<i>glyA, hipO^C</i>	(Casagrande Proietti et al., 2018)
			Unpacked thighs	51.60			23S rRNA, <i>glyA, hipO, sapB2^C</i>	(García-Sánchez et al., 2018)
			Unpacked breast	51.60			23S rRNA, <i>glyA, hipO, sapB2^C</i>	(García-Sánchez et al., 2018)
			Unpacked minced	21.90			23S rRNA, <i>glyA, hipO, sapB2^C</i>	(García-Sánchez et al., 2018)
			Unpacked marinated	56.30			23S rRNA, <i>glyA, hipO, sapB2^C</i>	(García-Sánchez et al., 2018)
			Mean unpacked	45.30			23S rRNA, <i>glyA, hipO, sapB2^C</i>	(García-Sánchez et al., 2018)
			Packed thighs	56.30			23S rRNA, <i>glyA, hipO, sapB2^C</i>	(García-Sánchez et al., 2018)
	Packed breast	45.30			23S rRNA, <i>glyA, hipO, sapB2^C</i>	(García-Sánchez et al., 2018)		

Packed minced	14.00			23S rRNA, <i>glyA</i> , <i>hipO</i> , <i>sapB2</i> ^C	(García-Sánchez et al., 2018)
Packed marinated	18.70			23S rRNA, <i>glyA</i> , <i>hipO</i> , <i>sapB2</i> ^C	(García-Sánchez et al., 2018)
Mean packed	33.60			23S rRNA, <i>glyA</i> , <i>hipO</i> , <i>sapB2</i> ^C	(García-Sánchez et al., 2018)
Raw broiler meat		36.00		<i>hipO</i> ^B	(Khan et al., 2018)
Broiler intestine		24.00		<i>hipO</i> ^B	(Khan et al., 2018)
Feathers		8.00		<i>hipO</i> ^B	(Khan et al., 2018)
Gizzard	59.75			23S rRNA, <i>mapA</i> , <i>ceuE</i> , <i>hipO</i> ^B	(Korsak et al., 2015)
Heart	49.66			23S rRNA, <i>mapA</i> , <i>ceuE</i> , <i>hipO</i> ^B	(Korsak et al., 2015)
Livers	44.08			23S rRNA, <i>mapA</i> , <i>ceuE</i> , <i>hipO</i> ^B	(Korsak et al., 2015)
Fillet	61.00			23S rRNA, <i>mapA</i> , <i>ceuE</i> , <i>hipO</i> ^B	(Korsak et al., 2015)
Breast	39.00			<i>glyA</i> , <i>hipO</i> , <i>ask</i> ^C	(Williams and Oyarzabal, 2012)
Tenderloins	26.00			<i>glyA</i> , <i>hipO</i> , <i>ask</i> ^C	(Williams and Oyarzabal, 2012)
Thighs	53.00 - 90.00	94.50	5.50	<i>glyA</i> , <i>hipO</i> , <i>ask</i> ^{AC}	(García-Sánchez et al., 2017; Williams and Oyarzabal, 2012)

987 Figure 1



988

