

AperTO - Archivio Istituzionale Open Access dell'Università di Torino

Long-term effect of asthma on the development of obesity among adults: an international cohort study, ECRHS

This is the author's manuscript

Original Citation:

Availability:

This version is available <http://hdl.handle.net/2318/1856044> since 2023-08-25T14:45:14Z

Published version:

DOI:10.1136/thoraxjnl-2021-217867

Terms of use:

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)

Thorax

Long-term effect of asthma on the development of obesity among adults: an international cohort study, ECRHS

Journal:	<i>Thorax</i>
Manuscript ID	thoraxjnl-2021-217867.R1
Article Type:	Original research
Date Submitted by the Author:	15-Feb-2022
Complete List of Authors:	<p>Moitra, Subhabrata; University of Alberta Faculty of Medicine & Dentistry, Department of Medicine Carsin, Anne-Elie ; ISGlobal Abramson, Michael; Monash University, School of Public Health and Preventive Medicine ACCORDINI, Simone; University of Verona, Unit of Epidemiology and Medical Statistics, Dept. Public Health and Community Medicine Amaral, Andre; Imperial College London, NHLI Anto, Josep; ISGlobal Bono, Roberto; University of Turin, Department of Public Health and Pediatrics Casas Ruiz, Lidia; University of Antwerp; KU Leuven, Department of Public Health and Primary Care Cerveri, Isa; University of Pavia, Division of Respiratory Diseases Chatzi, Leda; University of Crete, Department of Social Medicine; USC Keck School of Medicine, Department of Preventive Medicine; Maastricht University, Department of Genetics & Cell Biology Demoly, Pascal; Département de Pneumologie et Addictologie, University Hospital of Montpellier Arenas, Sandra; Hospital Galdakao-Usansolo, Pulmonology Department Forsberg, Bertil; Division of Occupational and Environmental Medicine, Umeå University Gilliland, Frank; USC Keck School of Medicine Gislason, Thorarinn; Landspítali University Hospital Gullón, Jose; University Hospital San Agustín, Pneumology Heinrich, Joachim; 1 Ludwig Maximilians University Munich, University Hospital Munich, , Institute and Outpatient Clinic for Occupational, Social and Environmental Medicine; 2 Helmholtz Zentrum München, German Research Center for Environmental Health,, Institut of Epidemiology Holm, Mathias; University of Gothenburg, Department of Occupational and Environmental Medicine Janson, Christer; Uppsala University, Department of Medical Sciences: Respiratory Medicine Jogi, Rain; Tartu University Hospital, Lung Clinic Gómez Real, Francisco; Haukeland University Hospital Jarvis, Debbie; Imperial College London Leynaert, Bénédicte; INSERM, U1152 Epidemiology Nowak, Dennis; Ludwig-Maximilians-Universität München Medizinische Fakultät</p>

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

	Probst-Hensch, Nicole; Swiss Tropical and Public Health Institute, Department Epidemiology and Public Health Sánchez-Ramos, José; University of Huelva Raheison-Semjen, Chantal; CHU de Bordeaux, Respiratory Diseases Siroux, Valerie; Inserm U1209, Univ Grenoble Alpes, Team of Environmental epidemiology Guerra, Stefano; University of Arizona Medical Center - University Campus, Arizona Respiratory Center Kogevinas, Manolis; ISGlobal Garcia-Aymerich, Judith; Centre de Recerca en Epidemiologia Ambiental
Keywords:	Asthma, Asthma Epidemiology, Clinical Epidemiology





I, the Submitting Author has the right to grant and does grant on behalf of all authors of the Work (as defined in the below author licence), an exclusive licence and/or a non-exclusive licence for contributions from authors who are: i) UK Crown employees; ii) where BMJ has agreed a CC-BY licence shall apply, and/or iii) in accordance with the terms applicable for US Federal Government officers or employees acting as part of their official duties; on a worldwide, perpetual, irrevocable, royalty-free basis to BMJ Publishing Group Ltd ("BMJ") its licensees and where the relevant Journal is co-owned by BMJ to the co-owners of the Journal, to publish the Work in this journal and any other BMJ products and to exploit all rights, as set out in our [licence](#).

The Submitting Author accepts and understands that any supply made under these terms is made by BMJ to the Submitting Author unless you are acting as an employee on behalf of your employer or a postgraduate student of an affiliated institution which is paying any applicable article publishing charge ("APC") for Open Access articles. Where the Submitting Author wishes to make the Work available on an Open Access basis (and intends to pay the relevant APC), the terms of reuse of such Open Access shall be governed by a Creative Commons licence – details of these licences and which [Creative Commons](#) licence will apply to this Work are set out in our licence referred to above.

Other than as permitted in any relevant BMJ Author's Self Archiving Policies, I confirm this Work has not been accepted for publication elsewhere, is not being considered for publication elsewhere and does not duplicate material already published. I confirm all authors consent to publication of this Work and authorise the granting of this licence.

1
2
3 **Long-term effect of asthma on the development of obesity among adults: an international**
4 **cohort study, ECRHS**
5
6
7

8 Subhabrata Moitra^{1,2,3*†}, Anne-Elie Carsin^{1,2,3†}, Michael J Abramson⁴, Simone Accordini⁵, André
9 FS Amaral⁶, Josep M Antó^{1,2,3}, Roberto Bono⁷, Lidia Casas Ruiz^{8,9}, Isa Cerveri¹⁰, Leda
10 Chatzi^{11,12,13}, Pascal Demoly^{14,15}, Sandra Dorado-Arenas¹⁶, Bertil Forsberg¹⁷, Frank D Gilliland¹²,
11 Thóraninn Gíslason^{18,19}, José A. Gullón²⁰, Joachim Heinrich²¹, Mathias Holm²², Christer Janson²³,
12 Rain Jõgi²⁴, Francisco Gomez Real^{25,26}, Deborah Jarvis^{6,27}, Bénédicte Leynaert^{28,29}, Dennis
13 Nowak²¹, Nicole Probst-Hensch^{30,31}, José Luis Sánchez-Ramos³², Chantal Raheison Semjen³³,
14 Valérie Siroux³⁴, Stefano Guerra^{1,2,3,35}, Manolis Kogevinas^{1,2,3}, Judith Garcia-Aymerich^{1,2,3}
15
16
17
18
19
20
21
22
23

24 ¹ISGlobal, Barcelona, Spain

25 ²Universitat Pompeu Fabra (UPF), Barcelona, Spain

26 ³CIBER Epidemiología y Salud Pública (CIBERESP), Barcelona, Spain

27 ⁴School of Public Health and Preventive Medicine, Monash University, Melbourne, Australia

28 ⁵Unit of Epidemiology and Medical Statistics, Department of Diagnostics and Public Health,
29 University of Verona, Verona, Italy

30 ⁶National Heart and Lung Institute, Imperial College London, London, UK

31 ⁷Department of Public Health and Paediatrics, University of Turin, Turin, Italy

32 ⁸Epidemiology and Social Medicine, University of Antwerp, Antwerp, Belgium

33 ⁹Centre for Environment and Health, Department of Public Health and Primary Care, KU Leuven,
34 Leuven, Belgium

35 ¹⁰Unit of Respiratory Diseases, IRCCS Policlinico San Matteo, University of Pavia, Pavia, Italy

36 ¹¹Department of Social Medicine, University of Crete, Greece

37 ¹²Department of Preventive Medicine, Keck School of Medicine, University of Southern California,
38 Los Angeles, CA, USA

39 ¹³Department of Genetics & Cell Biology, Maastricht University, Maastricht, the Netherlands

40 ¹⁴Department of Pulmonology, Division of Allergy, Hôpital Arnaud de Vileneuve, University
41 Hospital of Montpellier, Montpellier, France

42 ¹⁵Inserm, Sorbonne Université, Equipe, EPAR - IPLESP, Paris, France
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 ¹⁶Osakidetza Basque Health Service, Galdakao University Hospital, Department of Respiratory
4
5 Medicine, Galdakao, Spain

6
7 ¹⁷Section of Sustainable Health, Department of Public Health and Clinical Medicine, Umeå
8
9 University, Umeå, Sweden

10
11 ¹⁸Department of Sleep, Landspítali - The National University Hospital of Iceland, Reykjavik,
12
13 Iceland

14
15 ¹⁹Faculty of Medicine, University of Iceland, Reykjavik, Iceland

16
17 ²⁰Pneumology Department. University Hospital San Agustín. Avilés. Spain

18
19 ²¹Institute and Outpatient Clinic for Occupational and Environmental Medicine, Clinic Center,
20
21 Ludwig Maximilian University, Comprehensive Pneumology Centre Munich, member DZL,
22
23 German Centre for Lung Research, Munich, Germany.

24
25 ²²Department of Public Health and Community Medicine, Institute of Medicine, University of
26
27 Gothenburg, Gothenburg, Sweden

28
29 ²³Department of Medical Sciences, Respiratory, Allergy and Sleep research, Uppsala University,
30
31 Uppsala, Sweden

32
33 ²⁴Lung Clinic, Tartu University Hospital, Tartu, Estonia

34
35 ²⁵Department of Clinical Science, University of Bergen, Bergen, Norway

36
37 ²⁶Department of Gynecology and Obstetrics, Haukeland University Hospital, Bergen, Norway

38
39 ²⁷MRC Centre for Environment and Health, Imperial College London, London, United Kingdom

40
41 ²⁸Inserm- U1168, VIMA (Aging and Chronic Diseases. Epidemiological and Public Health
42
43 Approaches), Villejuif, France

44
45 ²⁹UMR-S 1168, UVSQ, Univ Versailles St-Quentin-en-Yvelines, St-Quentin-en-Yvelines, France

46
47 ³⁰Swiss Tropical and Public Health Institute, Basel, Switzerland

48
49 ³¹University of Basel, Basel, Switzerland

50
51 ³²Department of Nursing, University of Huelva, Huelva, Spain

52
53 ³³INSERM U1219, Institute of Public health and Epidemiology, Bordeaux University, Bordeaux,
54
55 France

56
57 ³⁴University Grenoble-Alpes, CNRS, Team of Environmental Epidemiology applied to
58
59 Reproduction and Respiratory Health, IAB- Institute for Advanced Biosciences, Grenoble, France

60
³⁵Asthma and Airway Disease Research Center, University of Arizona, Tucson, AZ, USA

1
2
3
4
5 ***Current affiliation:** Alberta Respiratory Centre, Department of Medicine, University of Alberta,
6
7 Edmonton, Canada
8
9

10 †Equal contribution
11
12
13

14 **Correspondence to:**

15
16 Dr Subhabrata Moitra, Alberta Respiratory Centre, Division of Pulmonary Medicine, Department
17
18 of Medicine, 559, HMRC, University of Alberta, Edmonton, AB T6G 2R3, Canada
19

20 Tel: +1-780-604-0263
21

22 Email: moitra@ualberta.ca
23
24
25

26 **Word Count:** 3,055 (excluding abstract, references, tables, and figure legends)
27

28 **No. of Tables:** 1
29

30 **No. of Figures:** 3
31
32

33 This manuscript has an online supplement.
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

KEY MESSAGES

- **What is the key question?**

Is asthma a risk factor for the development of obesity in adults?

- **What is the bottom line?**

Asthmatics, particularly those who were non-atopic, had longer disease duration or were on oral corticosteroids, were at a higher risk of developing obesity later in life. This association was not mediated by physical activity.

- **Why read on?**

Although an association between obesity and asthma has been well established, whether asthma can relate to increased risk of future obesity has been only studied in children. Understanding the link between asthma and obesity is key for a proper asthma management.

ABSTRACT

Introduction: Obesity is a known risk factor for asthma. Although some evidence showed asthma causing obesity in children, the link between asthma and obesity has not been investigated in adults.

Methods: We used data from the European Community Respiratory Health Survey (ECRHS), a cohort study in 11 European countries and Australia in 3 waves between 1990 and 2014, at intervals of approximately 10 years. We considered two study periods: from ECRHS I (t) to ECRHS II ($t+1$), and from ECRHS II (t) to ECRHS III ($t+1$). We excluded obese (BMI ≥ 30 kg/m²) individuals at visit t . The relative risk (RR) of obesity at $t+1$ associated with asthma at t was estimated by multivariable modified Poisson regression (lag) with repeated measurements. Additionally, we examined the association of atopy and asthma medication on the development of obesity.

Results: We included 7,576 participants in the period ECRHS I-II (51.5% female, mean (SD) age of 34 (7) years), and 4,976 in ECRHS II-III (51.3% female, 42 (8) years). 9% of participants became obese in ECRHS I-II and 15% in ECRHS II—III. The risk of developing obesity was higher among asthmatics than non-asthmatics (RR: 1.22, 95%CI: 1.07 to 1.38), and particularly higher among non-atopic than atopic (1.47; 1.17 to 1.86 vs 1.04; 0.86 to 1.27), those with longer disease duration (1.32; 1.10 to 1.59 in >20 years vs 1.12; 0.87 to 1.43 in ≤ 20 years), and those on oral corticosteroids (1.99; 1.26 to 3.15 vs 1.15; 1.03 to 1.28). Physical activity was not a mediator of this association.

Conclusion: This is the first study showing that adult asthmatics have a higher risk of developing obesity than non-asthmatics, particularly those non-atopic, of longer disease duration or on oral corticosteroids.

Keywords: Asthma, ECRHS, Obesity, Oral corticosteroids, Physical activity

INTRODUCTION

Several studies demonstrating the co-existence of obesity and asthma across the world suggest the presence of common aetiological factors between these conditions.[1-5] Asthma and obesity share some common socioeconomic, behavioural, and environmental risk factors, all of which could trigger the expression of genes, leading to the development of either of these diseases. Previous studies focussed mainly on the mechanisms by which obesity could lead to asthma. Several obesity-associated biological phenomena have been proposed, such as by altering the lung compliance (causing airflow limitation), increasing the synthesis of immune-modulators, affecting the sympathetic nervous system, or by modulating gene function.[1, 2] The inverse relationship, i.e., whether asthma is a risk factor for later obesity had not received much attention until recently.

In a longitudinal cohort study of kindergarten and first-grade children, Chen and colleagues (2017) observed that the non-obese children who were diagnosed with asthma during the recruitment phase were at increased risk of developing obesity during the 10-years follow up.[6] This was confirmed in a pooled analysis of 18 European cohorts associating physician diagnosed asthma at 3-4 years with incident obesity at 8 years, the risk being higher among children with active asthma symptoms.[7] Another recent study demonstrated a link between high peak flow variability in childhood and a steeper increase of body mass index (BMI) in adulthood.[8] Despite studies on children and young adolescents, whether and to what extent asthma increased the risk for subsequent obesity among adults remained unclear. Asthma in adults may be causally and aetiologically more complex than childhood asthma (which is predominantly allergic asthma), therefore, the magnitude of the asthma-obesity risk is likely to be different in adults.

In line with the previously reported observations, we hypothesized that asthma could be a risk factor for obesity at a later stage of life. Therefore, we aimed to investigate the occurrence of obesity attributable to asthma in participants of the European Community Respiratory Health Survey (ECRHS) over two period of 10 years.

METHODS

Design and participants

ECRHS is a population-based cohort study initiated in 1990-94 (ECRHS-I) with over 18,000 participants aged between 20 and 44 years from 30 centres in 14 countries, which had two follow-up rounds at approximately 10-year intervals (ECRHS-II in 1999-2003 and ECRHS-III in 2010-2014). In the recruitment phase, a postal questionnaire was sent containing items about asthma symptoms and exacerbation history in the last 12 months, current medication, and allergic conditions such as nasal symptoms and hay fever. From those who responded to the postal questionnaire, participants were either selected from a random sample or a symptomatic sample and were invited for more detailed investigation that included an interviewer-administered questionnaire and lung function testing.[9, 10] The questionnaires and clinical investigations were repeated in the next follow-ups (ECRHS-II and III).

The design and selection of participants of the present study are presented in **Figure 1**. We considered two study periods: from ECRHS I (t) to ECRHS II ($t+1$), and from ECRHS II (t) to ECRHS III ($t+1$). At each period we classified the participants as “no asthma” or “current asthma” (see definitions below). To study only incident obesity, we excluded individuals who were obese (BMI $\geq 30\text{kg/m}^2$) at visit t . To avoid misclassification in the definition of the ‘current asthma’ and ‘no asthma’ groups, we excluded participants with inactive asthma (i.e. self-reported doctor diagnosed asthma, with no respiratory symptoms within the last 12 months). We finally excluded subjects with missing asthma or obesity information at either t or $t+1$, giving a total of 8,716 participants from 26 centres of 11 European countries and Australia. Ethical approval of the study was obtained from the ethical review boards or committees of the respective centres. All participants provided signed informed consent. Detailed information about the ethical approval, informed consent and data management were described elsewhere.[10]

- Figure 1

Definition of asthma-related variables

Asthma-related information was collected at each survey. We defined current asthma as giving a positive answer to the questions “Have you ever had asthma?” and “Was this confirmed by a

1
2
3 doctor?”, and having reported at least one of the following in the last 12 months: waking up by an
4 attack of shortness of breath, any asthma attack, or use of asthma medications.[11-13] Asthma
5 duration at t was derived as the number of years between age at t and reported age at first asthma
6 onset. Atopy was defined as a serum specific IgE antibody concentration of ≥ 0.35 kU_A/L to at
7 least one of four common allergens (house dust mite, cat, timothy grass or *Cladosporium*
8 *herbarum*), as defined previously.[14] We also obtained information on asthma drug treatment
9 from the questions “Have you used any pills, capsules, tablets or medicines, other than inhaled
10 medicines, to help your breathing at any time in the last 12 months?”, and defined whether each
11 patient took inhaled (ICS) oral corticosteroids (OCS) or both during the last 12 months. Only at
12 ECRHS II, information was collected on whether OCS were used continuously, for a short-course,
13 or as rescue medication. The modified Medical Research Council (mMRC) Dyspnoea Scale was
14 used to define dyspnoea (considered if they had mMRC grade ≥ 2).[15]
15
16
17
18
19
20
21
22
23
24
25
26
27

28 **Obesity**

29 Height and weight of all participants were measured by physical examination during the clinical
30 visit at each survey. BMI was calculated as weight in kilograms divided by the square of height in
31 metres. Obesity was defined as BMI ≥ 30 kg/m². [16]
32
33
34
35
36
37

38 **Other relevant information**

39 At each study wave, we collected data on socio demographic and other clinical factors using
40 questionnaires. These included sex, age, education (age at which the participants finished their
41 formal education) and smoking status (never, former and current) and intensity (pack-years).
42
43
44
45

46 Information about physical activity of the participants was obtained at ECRHS-II and III from a
47 self-completed questionnaire in which they were asked about the weekly frequency and duration
48 of vigorous physical activity.[17-19] Those who exercised at least two times a week for at least 1
49 hour were considered physically active.
50
51
52
53
54

55 **Statistical analyses**

56 Relative risks (RR) and 95% confidence intervals (CIs) for the association of current asthma,
57 asthma with atopy, asthma duration, and asthma with medication (ICS, OCS or both), with new
58
59
60

1
2
3 obesity onset were estimated using modified Poisson regression for repeated measures with time
4 of follow-up as offset.[20] Exposure was the asthma status at the survey time t (lag1) and outcome
5 was obesity at subsequent time $t+1$. Therefore, participants could contribute with one or two
6 observations, i.e., from ECRHS-I to ECRHS-II and/or from ECRHS-II to ECRHS-III (Figure 1).
7
8 Multivariate models were fitted after testing the following potential confounders: education, sex,
9 age, smoking status, and physical activity (between ECRHS-II and III only). Only those variables
10 associated (i.e., p-value <0.05) both with asthma and obesity were retained. Hence, the final
11 models were adjusted for age, sex, and smoking status at time t . Subjects nested in centre were
12 included as random effects to account for the repeated data design.
13
14
15
16
17
18
19
20
21

22 Models were presented overall and separately by sex. We performed meta-analysis to determine
23 if there was any heterogeneity between the participating countries and between centres.
24
25
26
27

28 Two secondary analyses were performed: (1) to determine if physical activity could be a mediator
29 in the asthma-obesity association, we tested 3 paths separately: physical activity at ECRHS-II
30 mediating the relationship between asthma at ECRHS-I and obesity at ECRHS-II (path 1),
31 physical activity at ECRHS-II mediating the relationship between asthma at ECRHS-I and obesity
32 at ECRHS-II (path 2), and physical activity at ECRHS-III mediating the relationship between
33 asthma at ECRHS-II and obesity at ECRHS-III (path 3); and (2) to estimate the impact of smoking
34 cessation on the association between asthma and new obesity onset, we stratified our models by
35 active smokers (at time t), smokers who quit during the study (between ECRHS-I and II, and
36 ECRHS-II and III), and never smokers.
37
38
39
40
41
42
43
44
45
46

47 Some sensitivity analyses were performed to check the robustness of our results. We repeated
48 our analyses: (1) excluding those who did not take part in all three ECRHS waves; (2) excluding
49 participants who reported avoiding physical activity because of respiratory symptoms; (3)
50 excluding those who reported dyspnoea at time t ; (4) defining current asthma as the a self-report
51 of doctor's diagnosis and at least one of the following within the last 12 months: waking up by an
52 attack of shortness of breath, any asthma attack, use of asthma medications, wheeze, shortness
53 of breath, waking up with chest tightness or attack of cough; and (5) using BMI change between
54
55
56
57
58
59
60

1
2
3 t and $t+1$ as the outcome variable in linear mixed regression. All analyses were performed in
4
5 STATA (version 12.1, STATA Corp. College Station, Texas, USA).
6
7

8 9 **RESULTS**

10 Distribution of the participants included in this analysis is presented in Table 1. We studied 7,576
11 participants in the period ECRHS I-II (51.5% female, mean (SD) age of 34 (7) years at ECRHS
12 I), and 4,976 participants in the period ECRHS II-III (51.3% female, 42 (8) years at ECRHS II).
13
14 Current smokers decreased during follow-up (30% in ECRHS I-II vs. 19% in ECRHS II-III). The
15
16 relative number of participants with current asthma remained the same across the study waves.
17
18 Overall, 9% of participants became obese from ECRHS-I to ECRHS-II and this proportion rose to
19
20 15% from ECRHS II to ECRHS-III. We observed that 37% of the study population was physically
21
22 active at ECRHS-II which increased slightly to 42% at ECRHS-III.
23
24
25
26
27

- 28 • Table 1

29
30
31 The association between asthma and obesity is presented in **Supplementary Table E1** and
32
33 **Figure 2**. Compared to 8.4% and 14.6% of non-asthmatics who developed obesity at ECRHS-II
34
35 and ECRHS-III respectively, 10.7% and 16.9% asthmatics developed obesity at ECRHS-II and
36
37 III. The absolute risk difference in asthmatics vs no asthmatics was 2.3% (95% CI 0.8 to 3.9%)
38
39 and the relative risk (RR) was 1.21 (95%CI: 1.07 to 1.37). After adjusting for potential confounders
40
41 (age, sex and smoking status at time t), the association between asthma and the development of
42
43 obesity remained significant (1.22; 1.07 to 1.38). The association was stronger among asthmatics
44
45 without atopy (1.47; 1.17 to 1.86), asthmatics with longer disease duration (1.32; 1.10 to 1.59),
46
47 asthmatics who used oral corticosteroids (1.99; 1.26 to 3.15) and those who used both oral and
48
49 inhaled corticosteroids (2.07; 1.19 to 3.59).
50
51

- 52 • Figure 2

53
54
55 We conducted post hoc analyses to understand better the association between oral
56
57 corticosteroids intake and incident obesity. The comparison between patients on continuous,
58
59
60

1
2
3 short-course or use as rescue medication of OCS suggested a higher risk of obesity among those
4 using continuously, but very small sample size precluded any kind of statistical testing
5
6
7 **(Supplementary Table E2)**.

8
9
10 The association of asthma and obesity was similar in males (1.27; 1.06 to 1.51) and females
11 (1.20; 1.01 to 1.42, **Supplementary Table E3**). The association of asthma with new onset obesity
12 was homogeneous across the participating countries (**Figure 3**) and centres (**Supplementary**
13 **Figure E1**). The overall estimates from the meta-analyses were similar to that reported in the
14 main analysis and there was no statistical evidence of heterogeneity ($I^2=0$, p-values for
15 heterogeneity were 0.70 and 0.96 for country and centre, respectively).
16
17
18
19
20
21
22
23

- 24 • Figure 3

25
26
27 No evidence of physical activity as a mediator in the asthma-obesity association was observed
28 (**Supplementary Table E4**). When stratified by smoking status, the risk of becoming obese in
29 asthmatics was higher among current smokers (1.46, 1.12 to 1.90) and among never smokers
30 (1.27, 1.08 to 1.49), while no association between asthma and obesity was observed among those
31 who stopped smoking (0.95; 0.65 to 1.37) (**Supplementary Table E5**). In the sensitivity analyses,
32 the estimates were comparable to the main analyses (**Supplementary Tables E6, E7, E8, E9**
33 **and E10**).
34
35
36
37
38
39
40
41
42

43 DISCUSSION

44
45 In this prospective cohort study, we found that individuals with active asthma were at higher risk
46 of developing obesity compared to individuals without asthma, that this association was higher
47 among non-atopic asthmatics, those with long disease duration or those using oral
48 corticosteroids, and that the association was not mediated by physical activity levels.
49
50
51
52
53

54
55 Our observation of accelerated weight gain among asthmatics is supported by a previous
56 prospective study that demonstrated asthma to be associated with increased weight gain at a
57 later stage in women,[21] although unlike our results, they did not find any such association in
58
59
60

1
2
3 men. Two recent longitudinal studies in children, one of them including more than 21,000 children
4 from 16 European cohorts, showed that childhood asthma was a risk factor for obesity at a later
5 age.[6, 7] Another recent report also demonstrated that children with high peak flow variability
6 were on a steeper increase of BMI up to young adult life that was not associated with asthma.[8]
7
8 This evidence suggests that asthma and obesity share several common physiological pathways
9 and a bi-directional association between asthma and obesity is plausible.
10
11
12
13
14
15

16 It has been debated whether there is any association between atopy and increased body
17 weight.[22] While several reports support this association, [23, 24] other studies have shown that
18 the risk of being overweight is higher among the non-atopic than those with marked allergic
19 sensitization.[25, 26] Our observation of a greater risk of obesity among non-atopic asthmatics
20 indicates that allergic sensitization is unlikely to explain the association between asthma and
21 obesity.
22
23
24
25
26
27
28
29

30 We observed that the risk of developing obesity was more pronounced among asthmatics who
31 were on oral corticosteroids than those who were not, in agreement with previous reports, in
32 adults and children with asthma, relating the intake of oral corticosteroids with abnormal weight
33 gain.[27-30] Unfortunately we could not test the role of corticosteroids dose or duration due to
34 small sample size of OCS users. The effects of long term use of oral corticosteroids on weight
35 gain are well established in asthma and general population,[27, 31] and supported by several
36 mechanisms, including but not limited to increased lipid uptake from the gut and storing in the
37 peripheral tissues,[30] or increased insulin resistance.[32] However, we also found an increased
38 and statistically significant risk of obesity among asthmatics who reported not having taken OCS.
39 One could argue that, if asthma really causes weight gain, this effect would be higher among
40 those with severe asthma, so the higher risk observed among OCS users is actually the result of
41 confounding by indication and not a result of the treatment itself.[33] This is supported by our
42 findings of higher obesity risk among asthmatics of longer disease duration..
43
44
45
46
47
48
49
50
51
52
53
54
55

56 A potential explanation for asthma causing weight gain is through the reduction of physical activity
57 levels. Patients with chronic respiratory diseases tend to be less physically active due to recurrent
58
59
60

1
2
3 episodes of breathlessness or persistent wheezing.[34] However, our analyses do not support
4 this hypothesis, as physical activity was not mediating the asthma-obesity association and this
5 association remained after excluding those participants who did not perform physical activity due
6 to breathlessness, in line with the previously mentioned study among asthmatic children who
7 developed obesity on follow-up independent of physical activity.[6]
8
9
10
11
12

13
14 Another potential explanation would be via a role of smoking. Smoking cessation has been
15 observed to augment weight gain at least in the short term.[35, 36] However, we found no
16 association between asthma and incident obesity among participants quitting smoking. On the
17 contrary, we observed greater relative risks among current smokers and non-smokers. Smoking
18 has been observed to reduce the efficacy of steroid treatment in asthma [37-39], so it could be
19 argued that asthmatics who smoke are at a higher risk of the effects of asthma (such as obesity)
20 because of worse disease control. Altogether suggests that there may be mechanisms other than
21 OCS use, reducing physical activity and smoking cessation, linking asthma to obesity, but these
22 are yet unknown.
23
24
25
26
27
28
29
30
31
32

33
34 Our study outcomes are of significant clinical importance. First, our observation of higher obesity
35 risk in asthmatics who took oral corticosteroids, advocates the minimal and precise use of oral
36 corticosteroids in asthma.[40] Albeit current recommendations of oral corticosteroids for asthma
37 management, several reports also indicate that oral steroids are still prescribed indiscriminately
38 in asthma, sometimes, even without diagnosing the severity of the disease properly.[41-43] This
39 could result in an increasing burden of adverse health effects in asthma, obesity being one of the
40 major consequences. Secondly, the observation of a greater risk of obesity among non-atopic
41 asthmatics directs our attention towards adult-onset asthma and supports increasing attention to
42 the reduction of adult risk factors, such as occupational exposures that account for up to 25% of
43 adult-onset asthma.[44] Thirdly, we also observed a higher risk of obesity among asthmatics who
44 were smokers. Although smoking cessation programmes have now become a part of the clinical
45 management of respiratory and cardiovascular diseases, evidence shows that not all patients
46 benefit from them, particularly in primary care.[45, 46] Advice on smoking cessation along with its
47 plausible consequences (such as temporary weight gain) should also be discussed with the
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 patients. Finally, although we did not find any evidence of physical activity mediating the asthma-
4 obesity interaction, it must be noted that lifestyle factors also significantly influence weight gain.
5
6 Thus, advice on healthy diet, proper physical activity and lifestyle modification should be
7
8 warranted as part of asthma management.
9

10
11
12 The strengths of our study included its longitudinal design, allowing the exclusion of obese
13 subjects at baseline, with a large cohort that was followed over a long period of time. Furthermore,
14
15 we performed several sensitivity analyses to test the robustness of our results against
16
17 assumptions about asthma or obesity miss-classification, residual confounding or model miss-
18
19 specification. Lastly, the results were homogeneous across the participating countries which
20
21 indicate that study findings can be extrapolated to other populations.
22
23

24
25
26 A limitation in this study is the potential bias in defining asthma and obesity status, given the lack
27
28 of information on when events happened during the 10 years follow-up period. However, our
29
30 results were robust to several sensitivity analyses. Although we were able to consider plausible
31
32 known confounders, the effect of other lifestyle factors (such as diet, only available in a subsample
33
34 of ECRHS at one single time point), psychosocial attributes, or genetic predisposition could not
35
36 be analysed. Finally, we were limited to explore in detail the role of OCS on obesity risk because
37
38 of the lack of detailed information on dosage, duration or their use for other diseases than asthma.
39
40

41 **CONCLUSION**

42
43 In summary, we demonstrated that participants with asthma were at a higher risk of developing
44
45 obesity later in life and the risk was more pronounced among those asthmatics who were non-
46
47 atopic, those with asthma of longer duration and those on oral corticosteroids. These results
48
49 support earlier results in children and warrant further clinico-epidemiological and experimental
50
51 research, in order to determine the mechanisms through which asthma could trigger obesity risk
52
53 among adults.
54
55
56
57
58
59
60

AUTHOR CONTRIBUTIONS

S Moitra, A-E Carsin, M Kogevinas, and J Garcia-Aymerich conceived the study design, analysed the results, interpreted the data, and drafted the manuscript. S Guerra interpreted the data; drafted the article or revised it; provided intellectual content of critical importance to the work described. All other authors revised the manuscript and provided critical comments. The joint first authors and the last author take the responsibility for the data integrity.

FUNDING AND ACKNOWLEDGEMENT

ECRHS I

Co-ordinating Centre (London): P Burney, S Chinn, C Luczynska†, D Jarvis, E Lai.

Project Management Group: P Burney (Project leader-UK), S Chinn (UK), C Luczynska† (UK), D Jarvis (UK), P Vermeire† (Antwerp), H Kesteloot (Leuven), J Bousquet (Montpellier), D Nowak (Hamburg), J Prichard† (Dublin), R de Marco† (Verona), B Rijcken (Groningen), JM Anto (Barcelona), J Alves (Oporto), G Boman (Uppsala), N Nielsen (Copenhagen), P Paoletti (Pisa).

Financial support: The following grants helped to fund the local studies. Australia: Asthma Foundation of Victoria, Allen and Hanbury's, Belgium: Belgian Science Policy Office, National Fund for Scientific Research, Estonia: Estonian Science Foundation, grant no 1088, France: Ministère de la Santé, Glaxo France, Insitut Pneumologique d'Aquitaine, Contrat de Plan Etat-Région Languedoc-Rousillon, CNMATS, CNMRT (90MR/10, 91AF/6), Ministre delegué de la santé, RNSP, France; GSF, Germany: Bundesminister für Forschung und Technologie, Italy: Ministero dell'Università e della Ricerca Scientifica e Tecnologica, CNR, Regione Veneto grant RSF n. 381/05.93, Norway: Norwegian Research Council project no. 101422/310, Spain: Fondo de Investigación Sanitaria (#91/0016-060-05/E, 92/0319 and #93/0393), Hospital General de Albacete, Hospital General Juan Ramón Jiménez, Dirección Regional de Salud Pública (Consejería de Sanidad del Principado de Asturias), CIRIT (1997 SGR 00079) and Servicio Andaluz de Salud, Sweden: The Swedish Medical Research Council, the Swedish Heart Lung Foundation, the Swedish Association against Asthma and Allergy, Switzerland: Swiss national

1
2
3 Science Foundation grant 4026-28099, UK: National Asthma Campaign, British Lung Foundation,
4 Department of Health, South Thames Regional Health Authority.
5
6
7

8
9 Coordination: The co-ordination of this work was supported by the European Commission and
10 the authors and participants are grateful to the late C. Baya and M. Hallen for their help during
11 the study and K. Vuylsteek and the members of the COMAC for their support.
12
13
14

15 16 **ECRHS II**

17
18 Steering Committee: U. Ackermann-Lieblich (University of Basel, Switzerland); N. Kuenzli
19 (University of Basel, and University of Southern California, Los Angeles, USA); J.M. Antó and J.
20 Sunyer (Institut Municipal d' Investigació Mèdica (IMIM-IMAS), Universitat Pompeu Fabra (UPF),
21 Spain); P. Burney (project leader), S Chinn, D. Jarvis, J. Knox and C. Luczynska (King's College
22 London, UK); I. Cerveri (University of Pavia, Italy); R. de Marco† (University of Verona, Italy); T.
23 Gislason (Iceland University Hospital, Iceland); J. Heinrich and M. Wjst (GSF-Institute of
24 Epidemiology, Germany); C. Janson (Uppsala University, Sweden); B. Leynaert and F. Neukirch
25 (Institut National de la Santé et de la Recherche Médicale (INSERM), France); J. Schouten
26 (University of Groningen, The Netherlands); C. Svanes (University of Bergen, Norway); P.
27 Vermeire† (University of Antwerp, Belgium).
28
29

30
31 Principal Investigators and senior scientific teams: Australia: (M. Abramson, E.H Walters, J.
32 Raven); Belgium: South Antwerp and Antwerp City (P. Vermeire, J. Weyler, M. van Sprundel, V.
33 Nelen); Estonia: Tartu (R. Jõgi, A. Soon); France: Paris (F. Neukirch, B. Leynaert, R. Liard, M.
34 Zureik), Grenoble (I. Pin, J. Ferran-Quentin), Bordeaux (A. Taytard, C. Raheison), Montpellier
35 (J. Bousquet, P. J. Bousquet); Germany: Erfurt (J. Heinrich, M. Wjst, C. Frye, I. Meyer); Iceland:
36 Reykjavik (T. Gislason, E. Bjornsson, D. Gislason, K.B Jörundsdóttir); Italy: Turin (R. Bono, M.
37 Bugiani, P. Piccioni, E. Caria, A. Carosso, E. Migliore, G. Castiglioni), Verona (R. de Marco†, G.
38 Verlato, E. Zanolin, S. Accordini, A. Poli, V. Lo Cascio, M. Ferrari, I. Cazzoletti), Pavia (A.
39 Marinoni, S. Villani, M. Ponzio, F. Frigerio, M. Comelli, M. Grassi, I. Cerveri, A. Corsico); Norway:
40 Bergen (A. Gulsvik, E. Omenaas, C. Svanes, B. Laerum); Spain: Albacete (J. Martinez-Moratalla
41 Rovira, E. Almar, M. Arévalo, C. Boix, G. González, J.M. Ignacio García, J. Solera, J. Damián),
42 Galdakao (N. Muñozguren, J. Ramos, I. Urrutia, U. Aguirre), Barcelona (J. M. Antó, J. Sunyer,
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 M. Kogevinas, J. P. Zock, X. Basagaña, A. Jaen, F. Burgos, C. Acosta), Huelva (J. Maldonado,
4 A. Pereira, J.L. Sanchez), Oviedo (F. Payo, I. Huerta, A. de la Vega, L. Palenciano, J. Azofra, A.
5 Cañada); Sweden: Göteborg (K. Toren, L. Lillienberg, A. C. Olin, B. Balder, A. Pfeifer-Nilsson, R.
6 Sundberg), Umea (E. Norrman, M. Soderberg, K.A. Franklin, B. Lundback, B. Forsberg, L.
7 Nystrom), Uppsala (C. Janson, G. Boman, D. Norback, G. Wieslander, M. Gunnbjornsdottir);
8 Switzerland: Basel (N. Küenzli, B. Dibbert, M. Hazenkamp, M. Brutsche, U. Ackermann-Liebrich);
9 United Kingdom: Ipswich (D. Jarvis, R. Hall, D. Seaton), Norwich (D. Jarvis, B. Harrison).

10
11
12
13
14
15
16
17
18 Financial Support: Australia: National Health and Medical Research Council; Belgium: Antwerp:
19 Fund for Scientific Research (grant code, G.0402.00), University of Antwerp, Flemish Health
20 Ministry; Estonia: Tartu Estonian Science Foundation grant no 4350; France: (All) Programme
21 Hospitalier de Recherche Clinique—Direction de la Recherche Clinique (DRC) de Grenoble 2000
22 number 2610, Ministry of Health, Ministère de l'Emploi et de la Solidarité, Direction Générale de
23 la Santé, Centre Hospitalier Universitaire (CHU) de Grenoble, Bordeaux: Institut Pneumologique
24 d'Aquitaine, Grenoble: Comité des Maladies Respiratoires de l'Isère, Montpellier: Aventis
25 (France), Direction Regionale des Affaires Sanitaires et Sociales Languedoc-Roussillon, Paris:
26 Union Chimique Belge- Pharma (France), Aventis (France), Glaxo France; Germany: Erfurt
27 GSF—National Research Centre for Environment and Health, Deutsche
28 Forschungsgemeinschaft (grant code, FR1526/1-1), Hamburg: GSF—National Research Centre
29 for Environment and Health, Deutsche Forschungsgemeinschaft (grant code, MA 711/4-1);
30 Iceland: Reykjavik, Icelandic Research Council, Icelandic University Hospital Fund; Italy: Pavia
31 GlaxoSmithKline Italy, Italian Ministry of University and Scientific and Technological Research
32 (MURST), Local University Funding for Research 1998 and 1999, Turin: Azienda Sanitaria Locale
33 4 Regione Piemonte (Italy), Azienda Ospedaliera Centro Traumatologico Ospedaliero/Centro
34 Traumatologico Ortopedico—Istituto Clinico Ortopedico Regina Maria Adelaide Regione
35 Piemonte, Verona: Ministero dell'Università e della Ricerca Scientifica (MURST), Glaxo
36 Wellcome spa; Norway: Bergen: Norwegian Research Council, Norwegian Asthma and Allergy
37 Association, Glaxo Wellcome AS, Norway Research Fund; Spain: Fondo de Investigación
38 Sanitarias (grant codes, 97/0035-01, 99/0034-01 and 99/0034-02), Hospital Universitario de
39 Albacete, Consejería de Sanidad, Barcelona: Sociedad Española de Neumología y Cirugía
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 Toracica, Public Health Service(grant code, R01 HL62633-01), Fondo de Investigaciones
4 Santarias (grant codes, 97/0035-01, 99/0034-01, and 99/0034-02), Consell Interdepartamentalde
5 Recerca i Innovacio´ Tecnolo`gica (grant code, 1999SGR 00241) Instituto de Salud Carlos III;
6 Red deCentros de Epidemiologi´a y Salud Pu´blica, C03/09,Redde Basesmoleculares y
7 fisiolo´gicas de lasEnfermedadesRespiratorias,C03/011and Red de Grupos Infancia y Medio
8 Ambiente G03/176, Huelva: Fondo de Investigaciones Santarias (grant codes, 97/0035-01,
9 99/0034-01, and 99/0034-02), Galdakao: Basque Health Department, Oviedo: Fondo de
10 Investigaciones Sanitaria (97/0035-02, 97/0035, 99/0034-01, 99/0034-02, 99/0034-04, 99/0034-
11 06, 99/350, 99/0034--07), European Commission (EU-PEAL PL01237), Generalitat de Catalunya
12 (CIRIT 1999 SGR 00214), Hospital Universitario de Albacete, Sociedad Espa˜ola de Neumologıa
13 y Cirugıa Tor´cica (SEPAR R01 HL62633-01) Red de Centros de Epidemiologıa y Salud P´blica
14 (C03/09), Red de Bases moleculares y fisiol´gicas de las Enfermedades Respiratorias (C03/011)
15 and Red de Grupos Infancia y Medio Ambiente (G03/176);97/0035-01, 99/0034-01, and99/0034-
16 02); Sweden: Gteborg, Umea, Uppsala: Swedish Heart Lung Foundation, Swedish Foundation
17 for Health Care Sciences and Allergy Research, Swedish Asthma and Allergy Foundation,
18 Swedish Cancer and Allergy Foundation, Swedish Council for Working Life and Social Research
19 (FAS); Switzerland: Basel Swiss National Science Foundation, Swiss Federal Office for Education
20 and Science, Swiss National Accident Insurance Fund; UK: Ipswich and Norwich: Asthma UK
21 (formerly known as National Asthma Campaign).

22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Coordination: The coordination of this work was supported by the European Commission, as part
of their Quality of Life programme, (Grant code: QLK4-CT-1999-01237).

ECRHS III

Principal Investigators and senior scientific teams: Australia: Melbourne (M.Abramson, G. Benke,
S. Dharmage, B. Thompson, S. Kaushik, M Matheson). Belgium: South Antwerp & Antwerp City
(J. Weyler, H.Bentouhami, V. Nelen). Estonia: Tartu (R. Jgi, H.Orru). France: Bordeaux (C.
Raherison, P.O Girodet) Grenoble (I. Pin, V. Siroux, J.Ferran, J.L Cracowski) Montpellier (P.
Demoly, A.Bourdin, I. Vachier) Paris (B. Leynaert, D. Soussan, D. Courbon, C. Neukirch, L.
Alavoine, X. Duval, I. Poirier). Germany: Erfurt (J. Heinrich, E. Becker, G. Woelke, O. Manuwald)

1
2
3 Hamburg (H. Magnussen, D. Nowak, A-M Kirsten). Iceland: Reykjavik (T. Gislason, B.
4 Benediktsdottir, D. Gislason, E.S Arnardottir, M. Clausen, G. Gudmundsson, L. Gudmundsdottir,
5 H. Palsdottir, K. Olafsdottir, S. Sigmundsdottir, K. Bara-Jörundsdottir). Italy: Pavia (I .Cerveri,
6 A.Corsico, A. Grosso, F. Albicini, E. Gini, E.M Di Vincenzo, V. Ronzoni, S. Villani, F. Campanella,
7 M. Gnesi, F. Manzoni, L. Rossi, O. Ferraro) Turin: (M. Bugiani, R. Bono, P. Piccioni, R. Tassinari,
8 V. Bellisario, G. Trucco) Verona: (R de Marco†, S. Accordini, L. Calciano, L. Cazzoletti, M. Ferrari,
9 A.M Fratta Pasini, F. Locatelli, P. Marchetti, A. Marcon, E. Montoli, G. Nguyen, M. Olivieri, C.
10 Papadopoulou, C.Posenato, G. Pesce, P. Vallerio, G. Verlato, E. Zanolin). Norway: (C. Svanes,
11 E. Omenaas, A. Johannessen, T. Skorge, F. Gomez Real). Spain: Albacete (J. Martinez-Moratalla
12 Rovira, E. Almar, A. Mateos, S. García, A. Núñez, P.López, R. Sánchez, E Mancebo), Barcelona:
13 (J-M. Antó, J.P Zock, J Garcia-Aymerich, M Kogevinas, X. Basagaña, A.E. Carsin, F. Burgos, C.
14 Sanjuas, S Guerra, B. Jacquemin, P. Davdand) Galdakao: (N. Muñozguren, I. Urrutia, U. Aguirre,
15 S. Pascual), Huelva:(J Antonio Maldonado, A. Pereira, J. Luis Sánchez, L. Palacios, Oviedo: (F.
16 Payo, I. Huerta, N. Sánchez, M. Fernández, B. Robles). Sweden: Göteborg (K. Torén, M. Holm,
17 J-L Kim, A-C. Olin, A. Dahlman-Höglund), Umea (B. Forsberg, L. Braback, L Modig, B Järholm,
18 H Bertilsson, K.A Franklin, C Wahlgreen), Uppsala: (B Andersson, D Norback, U Spetz Nystrom,
19 G Wieslander, G.M Bodinaa Lund, K Nisser); Switzerland: Basel (N.M. Probst-Hensch, N. Künzli,
20 D. Stolz, C. Schindler, T. Rochat, J.M. Gaspoz, E. Zemp Stutz, M.Adam, C. Autenrieth, I. Curjuric,
21 J. Dratva, A. Di Pasquale, R. Ducret-Stich, E. Fischer, L. Grize, A. Hensel, D. Keidel, A. Kumar,
22 M. Imboden, N. Maire, A. Mehta, H. Phuleria, M. Ragettli, M. Ritter, E. Schaffner, G.A Thun, A.
23 Ineichen, T. Schikowski, M. Tarantino, M. Tsai. UK: London (P. Burney, D. Jarvis, S. Kapur, R.
24 Newson, J. Potts), Ipswich: (N. Innes), Norwich: (A. Wilson).

25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47 Financial Support: Australia: National Health & Medical Research Council. Belgium: Antwerp
48 South, Antwerp City: Research Foundation Flanders (FWO), grant code G.0.410.08.N.10 (both
49 sites). Estonia: Tartu- SF0180060s09 from the Estonian Ministry of Education. France: (All)
50 Ministère de la Santé. Programme Hospitalier de Recherche Clinique (PHRC) national 2010.
51 Bordeaux: INSERM U897 Université Bordeaux segalen, Grenoble: Comite Scientifique
52 AGIRadom 2011. Paris: Agence Nationale de la Santé, Région Ile de France, domaine d'intérêt
53 majeur (DIM). Germany: Erfurt: German Research Foundation HE 3294/10-1 Hamburg: German
54
55
56
57
58
59
60

1
2
3 Research Foundation MA 711/6-1, NO 262/7-1. Iceland: Reykjavik, The Landspítali University
4 Hospital Research Fund, University of Iceland Research Fund, ResMed Foundation, California,
5 USA, Orkuveita Reykjavíkur (Geothermal plant), Vegagerðin (The Icelandic Road Administration
6 (ICERA). Italy: All Italian centres were funded by the Italian Ministry of Health, Chiesi Farmaceutici
7 SpA, in addition Verona was funded by Cariverona foundation, Education Ministry (MIUR).
8 Norway: Norwegian Research council grant no 214123, Western Norway Regional Health
9 Authorities grant no 911631, Bergen Medical Research Foundation. Spain: Fondo de
10 Investigación Sanitaria (PS09/02457, PS09/00716, PS09/01511, PS09/02185, PS09/03190),
11 Servicio Andaluz de Salud, Sociedad Española de Neumología y Cirugía Torácica (SEPAR
12 1001/2010). Fondo de Investigación Sanitaria (PS09/02457), Barcelona: Fondo de Investigación
13 Sanitaria (FIS PS09/00716), Galdakao: Fondo de Investigación Sanitaria (FIS 09/01511) Huelva:
14 Fondo de Investigación Sanitaria (FIS PS09/02185) and Servicio Andaluz de Salud Oviedo:
15 Fondo de Investigación Sanitaria (FIS PS09/03190). Sweden: All centres were funded by The
16 Swedish Heart and Lung Foundation, The Swedish Asthma and Allergy Association, The Swedish
17 Association against Lung and Heart Disease, Swedish Research Council for health, working life
18 and welfare (FORTE) Göteborg: Also received further funding from the Swedish Council for
19 Working life and Social Research. Umea also received funding from Vasterbotten Country Council
20 ALF grant. Switzerland: The Swiss National Science Foundation (grants no 33CSCO-134276/1,
21 33CSCO-108796, 3247BO-104283, 3247BO-104288, 3247BO-104284, 3247-065896, 3100-
22 059302, 3200-052720, 3200-042532, 4026-028099) The Federal office for forest, environment
23 and landscape, The Federal Office of Public Health, The Federal Office of Roads and Transport,
24 the canton's government of Aargau, Basel-Stadt, Basel-Land, Geneva, Luzern, Ticino, Valais and
25 Zürich, the Swiss Lung League, the canton's Lung League of Basel Stadt/ Basel, Landschaft,
26 Geneva, Ticino, Valais and Zurich, SUVA, Freiwillige Akademische Gesellschaft, UBS Wealth
27 Foundation, Talecris Biotherapeutics GmbH, Abbott Diagnostics, European Commission 018996
28 (GABRIEL), Wellcome Trust WT 084703MA. UK: Medical Research Council (Grant Number
29 92091). Support also provided by the National Institute for Health Research through the Primary
30 Care Research Network.
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 Coordination: The coordination was funded through the Medical Research Council (Grant Number
4
5 92091).
6
7

8 †Deceased
9

10 11 12 **CONFLICT OF INTEREST**

13
14 *S. Moitra* reports Long-Term Research fellowship and Young Scientist Sponsorship from the
15 European Respiratory Society, outside the submitted work. *M. Abramson* holds investigator-
16 initiated grants for unrelated research from Pfizer and Boehringer-Ingelheim. He has undertaken
17 an unrelated consultancy for and received assistance with conference attendance from Sanofi
18 and has also received a speaker's fee from GSK, outside the submitted work. *P. Demoly* reports
19 personal fees from ALK, Stallergenes Greer, IQVIA, Chiesi, AstraZeneca, Thermo Fisher
20 Scientific, Ménarini, Bausch & Lomb, Mylan, ASIT Biotech, Novartis, Sanofi, Regeneron, outside
21 the submitted work. *R. Jõgi* reports grants from Estonian Research Council Personal Research
22 Grant no 562, during the conduct of the study; personal fees from Consultancy, grants from
23 Grants/grants pending, personal fees from Payment for lectures, personal fees from
24 Travel/accommodations/meeting expenses, outside the submitted work. *C. Raheison* reports
25 personal fees from ALK, Astra Zeneca, GSK, Novartis, outside the submitted work. Other authors
26 do not have any conflict of interest to declare.
27
28
29
30
31
32
33
34
35
36
37
38
39
40

41 **DATA SHARING POLICY**

42 The datasets used and analysed during the current study are available from the authors upon
43 reasonable request.
44
45
46
47
48

49 **REFERENCES**

- 50
51 1. Ford ES, Mannino DM, Redd SC, et al. Body mass index and asthma incidence among
52 USA adults. *Eur. Respir. J.* 2004;24(5):740-4.
53
54 2. Ali Z, Ulrik CS. Obesity and asthma: a coincidence or a causal relationship? A systematic
55 review. *Respir. Med.* 2013;107(9):1287-300.
56
57
58
59
60

- 1
2
3 3. Schatz M, Zeiger RS, Zhang F, et al. Overweight/obesity and risk of seasonal asthma
4 exacerbations. *J Allergy Clin Immunol Pract* 2013;1(6):618-22.
5
- 6
7 4. Newson RB, Jones M, Forsberg B, et al. The association of asthma, nasal allergies, and
8 positive skin prick tests with obesity, leptin, and adiponectin. *Clin. Exp. Allergy* 2014;44(2):250-
9
10 60.
11
- 12 5. Peters U, Dixon AE, Forno E. Obesity and asthma. *J. Allergy Clin. Immunol.*
13 2018;141(4):1169-79.
14
- 15 6. Chen Z, Salam MT, Alderete TL, et al. Effects of Childhood Asthma on the Development
16 of Obesity among School-aged Children. *Am. J. Respir. Crit. Care Med.* 2017;195(9):1181-88.
17
18 7. Contreras ZA, Chen Z, Roumeliotaki T, et al. Does early onset asthma increase childhood
19 obesity risk? A pooled analysis of 16 European cohorts. *Eur. Respir. J.* 2018;52(3):pii: 1800504.
20
21 8. Lombardi E, Stern DA, Sherrill D, et al. Peak flow variability in childhood and body mass
22 index in adult life. *J. Allergy Clin. Immunol.* 2019;143(3):1224-26 e9.
23
24 9. Burney PG, Luczynska C, Chinn S, et al. The European Community Respiratory Health
25 Survey. *Eur. Respir. J.* 1994;7(5):954-60.
26
27 10. Jarvis D, Newson R, Janson C, et al. Prevalence of asthma-like symptoms with ageing.
28 *Thorax* 2018;73(1):37-48.
29
30 11. de Marco R, Marcon A, Jarvis D, et al. Prognostic factors of asthma severity: a 9-year
31 international prospective cohort study. *J. Allergy Clin. Immunol.* 2006;117(6):1249-56.
32
33 12. Cazzoletti L, Marcon A, Corsico A, et al. Asthma severity according to Global Initiative for
34 Asthma and its determinants: an international study. *Int. Arch. Allergy Immunol.* 2010;151(1):70-
35 9.
36
37 13. Janson C, Accordini S, Cazzoletti L, et al. Pharmacological treatment of asthma in a
38 cohort of adults during a 20-year period: results from the European Community Respiratory Health
39 Survey I, II and III. *ERJ Open Res* 2019;5(1).
40
41 14. Le Moual N, Carsin AE, Siroux V, et al. Occupational exposures and uncontrolled adult-
42 onset asthma in the European Community Respiratory Health Survey II. *Eur. Respir. J.*
43 2014;43(2):374-86.
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

- 1
2
3 15. Bestall JC, Paul EA, Garrod R, et al. Usefulness of the Medical Research Council (MRC)
4 dyspnoea scale as a measure of disability in patients with chronic obstructive pulmonary disease.
5
6 *Thorax* 1999;54(7):581-6.
7
8
9 16. Chinn S, Jarvis D, Melotti R, et al. Smoking cessation, lung function, and weight gain: a
10 follow-up study. *Lancet* 2005;365(9471):1629-35; discussion 00-1.
11
12 17. Washburn RA, Goldfield SR, Smith KW, et al. The validity of self-reported exercise-
13 induced sweating as a measure of physical activity. *Am. J. Epidemiol.* 1990;132(1):107-13.
14
15 18. Rovio S, Kareholt I, Helkala EL, et al. Leisure-time physical activity at midlife and the risk
16 of dementia and Alzheimer's disease. *Lancet Neurol.* 2005;4(11):705-11.
17
18 19. Fuertes E, Markevych I, Jarvis D, et al. Residential air pollution does not modify the
19 positive association between physical activity and lung function in current smokers in the ECRHS
20 study. *Environ. Int.* 2018;120:364-72.
21
22 20. Zou G. A modified poisson regression approach to prospective studies with binary data.
23 *Am. J. Epidemiol.* 2004;159(7):702-6.
24
25 21. Hasler G, Gergen PJ, Ajdacic V, et al. Asthma and body weight change: a 20-year
26 prospective community study of young adults. *Int. J. Obes. (Lond.)* 2006;30(7):1111-8.
27
28 22. Boulet LP. Obesity and atopy. *Clin. Exp. Allergy* 2015;45(1):75-86.
29
30 23. Chen Y, Rennie D, Cormier Y, et al. Association between obesity and atopy in adults. *Int.*
31 *Arch. Allergy Immunol.* 2010;153(4):372-7.
32
33 24. Luo X, Xiang J, Dong X, et al. Association between obesity and atopic disorders in
34 Chinese adults: an individually matched case-control study. *BMC Public Health* 2013;13:12.
35
36 25. von Mutius E, Schwartz J, Neas LM, et al. Relation of body mass index to asthma and
37 atopy in children: the National Health and Nutrition Examination Study III. *Thorax*
38 2001;56(11):835-8.
39
40 26. Visness CM, London SJ, Daniels JL, et al. Association of childhood obesity with atopic
41 and nonatopic asthma: results from the National Health and Nutrition Examination Survey 1999-
42 2006. *J. Asthma* 2010;47(7):822-9.
43
44 27. Sullivan PW, Ghushchyan VH, Globe G, et al. Oral corticosteroid exposure and adverse
45 effects in asthmatic patients. *J. Allergy Clin. Immunol.* 2018;141(1):110-16 e7.
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

- 1
2
3 28. Cooper V, Metcalf L, Versnel J, et al. Patient-reported side effects, concerns and
4 adherence to corticosteroid treatment for asthma, and comparison with physician estimates of
5 side-effect prevalence: a UK-wide, cross-sectional study. *NPJ Prim Care Respir Med*
6 2015;25:15026.
7
8
9
10 29. Sweeney J, Patterson CC, Menzies-Gow A, et al. Comorbidity in severe asthma requiring
11 systemic corticosteroid therapy: cross-sectional data from the Optimum Patient Care Research
12 Database and the British Thoracic Difficult Asthma Registry. *Thorax* 2016;71(4):339-46.
13
14
15 30. Umlawska W. Adipose tissue content and distribution in children and adolescents with
16 bronchial asthma. *Respir. Med.* 2015;109(2):200-7.
17
18
19 31. Berthon BS, MacDonald-Wicks LK, Wood LG. A systematic review of the effect of oral
20 glucocorticoids on energy intake, appetite, and body weight in humans. *Nutr. Res.*
21 2014;34(3):179-90.
22
23
24 32. Carpaj OA, van den Berge M. The asthma-obesity relationship: underlying mechanisms
25 and treatment implications. *Curr. Opin. Pulm. Med.* 2018;24(1):42-49.
26
27
28 33. Garcia-Aymerich J, Marrades RM, Monso E, et al. Paradoxical results in the study of risk
29 factors of chronic obstructive pulmonary disease (COPD) re-admission. *Respir. Med.*
30 2004;98(9):851-7.
31
32
33 34. van 't Hul AJ, Frouws S, van den Akker E, et al. Decreased physical activity in adults with
34 bronchial asthma. *Respir. Med.* 2016;114:72-7.
35
36
37 35. Bush T, Lovejoy JC, Deprey M, et al. The effect of tobacco cessation on weight gain,
38 obesity, and diabetes risk. *Obesity (Silver Spring)* 2016;24(9):1834-41.
39
40
41 36. Jain P, Danaei G, Robins JM, et al. Smoking cessation and long-term weight gain in the
42 Framingham Heart Study: an application of the parametric g-formula for a continuous outcome.
43 *Eur. J. Epidemiol.* 2016;31(12):1223-29.
44
45
46 37. Chaudhuri R, Livingston E, McMahon AD, et al. Cigarette smoking impairs the therapeutic
47 response to oral corticosteroids in chronic asthma. *Am. J. Respir. Crit. Care Med.*
48 2003;168(11):1308-11.
49
50
51 38. Lazarus SC, Chinchilli VM, Rollings NJ, et al. Smoking affects response to inhaled
52 corticosteroids or leukotriene receptor antagonists in asthma. *Am. J. Respir. Crit. Care Med.*
53 2007;175(8):783-90.
54
55
56
57
58
59
60

- 1
2
3 39. Kane B, Kolsum U, Southworth T, et al. The effects of smoking on the lipopolysaccharide
4 response and glucocorticoid sensitivity of alveolar macrophages of patients with asthma. *Chest*
5 2009;136(1):163-70.
6
7
8 40. Bourdin A, Adcock I, Berger P, et al. How can we minimise the use of regular oral
9 corticosteroids in asthma? *Eur. Respir. Rev.* 2020;29(155):190085.
10
11 41. Bhulani N, Lalani S, Ahmed A, et al. Knowledge of asthma management by general
12 practitioners in Karachi, Pakistan: comparison with international guidelines. *Prim. Care Respir. J.*
13 2011;20(4):448-51.
14
15 42. Lommatzsch M, Sauerbeck IS, Wilmer C, et al. Oral corticosteroid prescription for asthma
16 by general practitioners: A three-year analysis in Germany. *Respir. Med.* 2021;176:106242.
17
18 43. Blakey J, Chung LP, McDonald VM, et al. Oral corticosteroids stewardship for asthma in
19 adults and adolescents: A position paper from the Thoracic Society of Australia and New Zealand.
20 *Respirology* 2021;26(12):1112-30.
21
22 44. MacKinnon M, To T, Ramsey C, et al. Improving detection of work-related asthma: a
23 review of gaps in awareness, reporting and knowledge translation. *Allergy Asthma Clin. Immunol.*
24 2020;16(1):73.
25
26 45. Coleman T, Murphy E, Cheater F. Factors influencing discussion of smoking between
27 general practitioners and patients who smoke: a qualitative study. *Br. J. Gen. Pract.*
28 2000;50(452):207-10.
29
30 46. Houston TK, Scarinci IC, Person SD, et al. Patient smoking cessation advice by health
31 care providers: the role of ethnicity, socioeconomic status, and health. *Am. J. Public Health*
32 2005;95(6):1056-61.
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Table 1: Demographic profile of the study participants in ECRHS I-II and in ECRHS II-III

	ECRHS I-II n=7576	ECRHS II-III n=4976
Sex (female), n (%)	3905 (51.5%)	2555 (51.3%)
Age <i>t</i> (years), mean (SD)	34.0 (7.1)	41.5 (7.8)
Age <i>t</i> +1 (years), mean (SD)	42.7 (7.2)	53.9 (7.1)
Education, n (%)		
Low	813 (10.8%)	409 (8.5%)
Medium	2715 (36.0%)	1675 (34.7%)
High	4024 (53.3%)	2744 (56.8%)
Country, n (%)		
Australia	387 (5.1%)	212 (4.3%)
Belgium	528 (7.0%)	287 (5.8%)
Estonia	250 (3.3%)	132 (2.7%)
France	862 (11.4%)	737 (14.8%)
Germany	534 (7.0%)	545 (11.0%)
Iceland	434 (5.7%)	320 (6.4%)
Italy	512 (6.8%)	210 (4.2%)
Norway	500 (6.6%)	298 (6.0%)
Spain	1353 (17.9%)	806 (16.2%)
Sweden	1231 (16.2%)	749 (15.1%)
Switzerland	483 (6.4%)	403 (8.1%)
UK	502 (6.6%)	277 (5.6%)
Smoking status <i>t</i> , n (%)		
Never	3272 (43.3%)	2201 (44.4%)
Former	2043 (27.0%)	1814 (36.6%)
Current	2247 (29.7%)	939 (19.0%)
Current asthma <i>t</i> , n (%)		
No asthma	6859 (90.5%)	4495 (90.2%)
Current asthma, all	717 (9.5%)	490 (9.8%)
Current asthma, <i>without atopy</i>	271 (3.6%)	190 (3.8%)
Current asthma, <i>with atopy</i>	446 (5.9%)	300 (6.0%)
Current asthma, <i>≤20 years since onset</i>	406 (5.4%)	218 (4.4%)
Current asthma, <i>>20 years since onset</i>	308 (4.1%)	266 (5.3%)
Current asthma, <i>no ICS</i>	494 (6.5%)	260 (5.2%)
Current asthma, <i>with ICS</i>	223 (2.9%)	230 (4.6%)
Current asthma, <i>no OCS</i>	664 (8.8%)	454 (9.1%)
Current asthma, <i>with OCS</i>	53 (0.7%)	36 (0.7%)
Current asthma, <i>only ICS</i>	183 (2.4%)	199 (4.0%)
Current asthma, <i>ICS & OCS</i>	40 (0.5%)	31 (0.6%)
BMI <i>t</i> (kg/m ²), mean (SD)	23.3 (2.8)	24.1 (2.9)
BMI <i>t</i> +1 (kg/m ²), mean (SD)	25.0 (3.6)	26.0 (3.8)
Obesity (yes) <i>t</i> +1, n (%)	655 (8.6%)	736 (14.8%)
Dyspnoea (yes) <i>t</i> , n (%)	256 (3.4%)	182 (3.7%)
Physical Activity (yes) <i>t</i> +1, n (%)	2627 (37.0%)	2029 (41.8%)

1
2
3 ECRHS: European Community Respiratory Health Survey; ICS: inhaled corticosteroids; OCS: oral
4 corticosteroids; BMI: body mass index.

5
6
7 Current asthma defined as a self-report of a doctor's asthma diagnosis and at least one of the following,
8
9 within the last 12 months: waking up by an attack of shortness of breath, any asthma attack or use of
10
11 asthma medications. Individuals with inactive asthma (reporting asthma diagnosis without any current
12
13 symptoms) were excluded from the study (see Methods).
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Confidential: For Review Only

1
2
3 **Figure Legends:**
4
5
6

7 **Figure 1: Schematic representation of the study design**
8
9

10 **Figure 2: Association between asthma, asthma with atopy, asthma duration, and asthma**
11 **with medication at t and incident obesity at $t+1$.** Data presented as relative risk (RR) and 95%
12 confidence interval (95%CI). RR was calculated from modified Poisson regression with robust
13 standard errors and subjects nested in centre as random effects. All models (for all participants)
14 were adjusted for age, sex and smoking status at time t . See full figures in Supplementary Table
15 E1.
16
17
18
19
20
21
22
23

24 **Figure 3: Meta-analysis results of the association between asthma at t and obesity at $t+1$**
25 **by country**
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60




Figure 1: Schematic representation of the study design

79x45mm (300 x 300 DPI)




Figure 2: Association between asthma, asthma with atopy, asthma duration, and asthma with medication at t and incident obesity at t+1.

Data presented as relative risk (RR) and 95% confidence interval (95%CI). RR was calculated from modified Poisson regression with robust standard errors and subjects nested in centre as random effects. All models (for all participants) were adjusted for age, sex and smoking status at time t. See full figures in Supplementary Table E1.

173x140mm (300 x 300 DPI)




Figure 3: Meta-analysis results of the association between asthma at t and obesity at t+1 by country

1082x838mm (72 x 72 DPI)

1
2
3 **Long-term effect of asthma on the development of obesity among adults: an international cohort**
4 **study, ECRHS**
5
6

7 Subhabrata Moitra, Anne-Elie Carsin, Michael J Abramson, Simone Accordini, André FS Amaral, Josep
8 M Antó, Roberto Bono, Lidia Casas Ruiz, Isa Cerveri, Leda Chatzi, Pascal Demoly, Sandra Dorado-
9 Arenas, Bertil Forsberg, Frank D Gilliland, Thóraninn Gíslason, José A. Gullón, Joachim Heinrich,
10 Mathias Holm, Christer Janson, Rain Jõgi, Francisco Gomez Real, Deborah Jarvis, Bénédicte Leynaert,
11 Dennis Nowak, Nicole Probst-Hensch, José Luis Sánchez-Ramos, Chantal Raheison Semjen, Valérie
12 Siroux, Stefano Guerra, Manolis Kogevinas, Judith Garcia-Aymerich
13
14
15
16

17 **List of Supplementary Tables and Figures**
18
19
20

- 21 • Supplementary Table E1: Association between asthma, asthma with atopy, asthma duration, and
22 asthma with medication at t and incident obesity at $t+1$
23
- 24 • Supplementary Table E2: Association between OCS dosage at t and obesity at $t+1$ in asthmatics
25
- 26 • Supplementary Table E3: Association between asthma, asthma with atopy, asthma duration, and
27 asthma with medication at t and incident obesity at $t+1$, stratified by sex
28
- 29 • Supplementary Table E4: Mediation analyses testing the mediating role of physical activity on the
30 association between asthma and incident obesity
31
- 32 • Supplementary Table E5: Association between asthma, asthma with atopy, asthma duration, and
33 asthma with medication at t and incident obesity at $t+1$, stratified by smoking pattern.
34
- 35 • Supplementary Table E6: Association between asthma, asthma with atopy, asthma duration, and
36 asthma with medication at t and incident obesity at $t+1$, including only participants who took part in
37 all the three waves of ECRHS I, II and III.
38
- 39 • Supplementary Table E7: Association between asthma, asthma with atopy, asthma duration, and
40 asthma with medication at t and incident obesity at $t+1$, excluding the participants who reported
41 avoiding physical activity because of wheezing/asthma.
42
- 43 • Supplementary Table E8: Association between asthma, asthma with atopy, asthma duration, and
44 asthma with medication at t and incident obesity at $t+1$, excluding the participants with dyspnoea
45 at t .
46
- 47 • Supplementary Table E9: Association between asthma, asthma with atopy, asthma duration, and
48 asthma with medication at t and incident obesity at $t+1$, using a broader asthma definition.
49
- 50 • Supplementary Table E10: Association between asthma, asthma with atopy, asthma duration,
51 and asthma with medication at t and BMI changes between t and $t+1$.
52
- 53 • Supplementary Figure E1: Meta-analysis of the association between asthma at t and obesity at
54 $t+1$ by ECRHS centre.
55
56
57
58
59
60

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46

Supplementary Table E1: Association between asthma, asthma with atopy, asthma duration, and asthma with medication at *t* and incident obesity at *t+1*

	Obesity at visit, % (n/N)		RR (95%CI)	
	ECRHS II	ECRHS III	Unadjusted	Adjusted
No asthma	8.4% (578/6859)	14.6% (653/4486)	1	1
Asthma	10.7% (77/717)	16.9% (83/490)	1.21 (1.07 to 1.37)	1.22 (1.07 to 1.38)
No asthma	8.4% (578/6859)	14.6% (653/4486)	1	1
Asthma, without atopy	12.9% (35/271)	21.0% (40/190)	1.46 (1.16 to 1.85)	1.47 (1.17 to 1.86)
Asthma, with atopy	9.4% (42/446)	14.3% (43/300)	1.05 (0.86 to 1.28)	1.04 (0.86 to 1.27)
No asthma	8.4% (578/6859)	14.6% (653/4486)	1	1
Current asthma, ≤20 years since onset	10.3% (42/406)	16.7% (31/218)	1.17 (0.96-1.42)	1.12 (0.87-1.43)
Current asthma, >20 years since onset	11.4% (35/308)	19.2% (51/266)	1.36 (1.11-1.65)	1.32 (1.10-1.59)
No asthma	8.4% (578/6859)	14.6% (653/4486)	1	1
Asthma, no ICS	9.1% (45/494)	18.8% (49/260)	1.22 (1.09 to 1.35)	1.21 (1.09 to 1.34)
Asthma, on ICS	14.4% (32/223)	14.8% (34/230)	1.20 (0.93 to 1.55)	1.21 (0.94 to 1.56)
No asthma	8.4% (578/6859)	14.6% (653/4486)	1	1
Asthma, no OCS	10.2% (68/664)	16.1% (73/454)	1.15 (1.03 to 1.28)	1.15 (1.03 to 1.28)
Asthma, on OCS	17.0% (9/53)	27.8% (10/36)	1.98 (1.26 to 3.11)	1.99 (1.26 to 3.15)
No asthma	8.4% (578/6859)	14.6% (653/4486)	1	1
Asthma, ICS only	13.1% (24/183)	13.1% (26/199)	1.06 (0.83 to 1.35)	1.07 (0.84 to 1.36)
Asthma, on OCS & ICS	20.0% (8/40)	25.8% (8/31)	2.05 (1.18 to 3.56)	2.07 (1.19 to 3.59)

ECRHS: European Community Respiratory Health Survey; ICS: inhaled corticosteroids; OCS: oral corticosteroids; RR: relative risk.

Current asthma defined as a self-report of a doctor's asthma diagnosis and at least one of the following, within the last 12 months: waking up by an attack of shortness of breath, any asthma attack or use of asthma medications. Individuals with inactive asthma (reporting asthma diagnosis without any current symptoms) were excluded from the study (see Methods).

RR was calculated from modified Poisson regression with robust standard errors and subjects nested in centre as random effects. All models (for all participants) were adjusted for age, sex and smoking status at time *t*.

Supplementary Table E2. Association between OCS dosage at *t* and obesity at *t+1* in asthmatics

	n* at ECRHS II	New obesity at ECRHS III	RR (95%CI)
Asthma - no OCS	405	15%	
Asthma – OCS as rescue medication	9	11%	Not computable
Asthma – OCS for a short course	14	14%	
Asthma – OCS continuously	4	100%	

* Numbers do not add to the total due to missing or inconsistent data in asthma treatment

Supplementary Table E3: Association between asthma, asthma with atopy, asthma duration, and asthma with medication at *t* and incident obesity at *t*+1, stratified by sex

	Male			Female		
	Obesity at visit, % (n/N)		RR (95% CI)	Obesity at visit, % (n/N)		RR (95% CI)
	ECRHS II	ECRHS III	Adjusted	ECRHS II	ECRHS III	Adjusted
No asthma	8.7 (292/3347)	14.6 (320/2196)	1	8.1 (286/3512)	14.6 (334/2293)	1
Asthma	9.9 (32/324)	18.4 (42/228)	1.27 (1.06 to 1.51)	11.4 (45/393)	15.6 (41/262)	1.20 (1.01 to 1.42)
No asthma	8.7 (292/3347)	14.6 (320/2196)	1	8.1 (286/3512)	14.6 (334/2293)	1
Asthma, no atopy	12.9 (13/101)	24.2 (16/66)	1.62 (1.06 to 2.49)	12.9 (22/170)	19.4 (24/124)	1.38 (1.06 to 1.79)
Asthma, with atopy	8.5 (19/223)	16.0 (26/162)	1.11 (0.83 to 1.46)	10.3 (23/223)	12.3 (17/138)	1.04 (0.78 to 1.39)
No asthma	8.7 (292/3347)	14.6 (320/2196)	1	8.1 (286/3512)	14.6 (334/2293)	1
Current asthma, ≤20 years since onset	8.9 (14/158)	15.9 (14/88)	1.16 (0.80-1.68)	11.3 (28/248)	13.1 (17/130)	1.22 (0.98-1.51)
Current asthma, >20 years since onset	11.0 (18/164)	19.6 (27/138)	1.36 (1.10-1.66)	11.8 (17/144)	18.8 (24/128)	1.32 (1.01-1.72)
No asthma	8.7 (292/3347)	14.6 (320/2196)	1	8.1 (286/3512)	14.6 (334/2293)	1
Asthma, no ICS	7.9 (19/242)	21.1 (28/133)	1.26 (1.04 to 1.52)	10.3 (26/252)	16.5 (21/127)	1.22 (0.99 to 1.52)
Asthma, on ICS	15.8 (13/82)	14.7 (14/95)	1.28 (0.87 to 1.86)	13.5 (19/141)	14.8 (20/135)	1.17 (0.86 to 1.59)
No asthma	8.7 (292/3347)	14.6 (320/2196)	1	8.1 (286/3512)	14.6 (334/2293)	1
Asthma, no OCS	9.6 (30/311)	17.3 (37/214)	1.20 (1.00 to 1.45)	10.8 (38/353)	15.0 (36/240)	1.14 (0.97 to 1.33)
Asthma, on OCS	15.4 (2/13)	35.7 (5/14)	2.45 (1.45 to 4.13)	17.5 (7/40)	22.7 (5/22)	1.81 (1.01 to 3.24)
No asthma	8.7 (292/3347)	14.6 (320/2196)	1	8.1 (286/3512)	14.6 (334/2293)	1
Asthma, ICS only	15.3 (11/72)	11.9 (10/84)	1.11 (0.71 to 1.73)	11.7 (13/111)	13.9 (16/115)	1.04 (0.79 to 1.37)
Asthma, on OCS & ICS	20.0 (2/10)	36.4 (4/11)	2.62 (1.62 to 4.24)	20.0 (6/30)	20.0 (4/20)	1.81 (0.91 to 3.59)

ECRHS: European Community Respiratory Health Survey; ICS: inhaled corticosteroids; OCS: oral corticosteroids; RR: relative risk.

Current asthma defined as a self-report of a doctor’s asthma diagnosis and at least one of the following, within the last 12 months: waking up by an attack of shortness of breath, any asthma attack or use of asthma medications. Individuals with inactive asthma (reporting asthma diagnosis without any current symptoms) were excluded from the study (see Methods).

RR was calculated from modified Poisson regression with robust standard errors and subjects nested in centre as random effects. All models (for all participants) were adjusted for age, sex and smoking status at time *t*.

Supplementary Table E4: Mediation analyses testing the mediating role of physical activity on the association between asthma and incident obesity

Mediator tested	Mediation path tested	n	Direct Effect	Average mediation	Total Effect	% mediated
Active/inactive at ECRHS II	Asthma I → PA II → OBE II	7098	*0.02111	-0.000185	0.020926	-0.82%
Active/inactive at ECRHS II	Asthma I → PA II → OBE III	4449	0.01856	0.000496	0.01906	1.67%
Active/inactive at ECRHS III	Asthma II → PA III → OBE III	4863	0.01852	-0.00037	0.0181	-1.50%

PA: Physical Activity. OBE: new obesity onset. I, II and III: ECRHS I, ECRHS II and ECRHS III

Mediator and outcome modelled as binary.

* p<0.05.

Supplementary Table E5: Association between asthma, asthma with atopy, asthma duration, and asthma with medication at t and incident obesity at t+1, stratified by smoking pattern.

	Current smoker (N=2214)			Quitters (N=1303)			Never smoker (N=3846)		
	Obesity at visit, %			Obesity at visit, %			Obesity at visit, %		
	ECRHS II	ECRHS III	RR (95% CI)	ECRHS II	ECRHS III	RR (95% CI)	ECRHS II	ECRHS III	RR (95% CI)
No asthma	7.6	12.7	1	11.5	19.6	1	8.0	13.2	1
Asthma	13.1	13.9	1.46 (1.12 to 1.90)	12.2	17.1	0.95 (0.65 to 1.37)	11.1	15.6	1.27 (1.08 to 1.49)
No asthma	7.6	12.7	1	11.5	19.6	1	8.0	13.2	1
Asthma, without atopy	14.1	13.9	1.53 (0.93 to 2.52)	20.0	19.4	1.14 (0.69 to 1.91)	12.7	20.7	1.52 (1.13 to 2.04)
Asthma, with atopy	12.2	13.9	1.38 (0.96 to 1.99)	8.8	15.0	0.79 (0.42 to 1.50)	10.3	12.8	1.12 (0.91 to 1.39)
No asthma	7.6	12.7	1	11.5	19.6	1	8.0	13.2	1
Current asthma, ≤20 y since onset	14.0	8.6	1.42 (0.84-2.39)	11.5	17.1	0.92 (0.48-1.76)	9.2	13.0	1.07 (0.80-1.41)
Current asthma, >20 y since onset	11.8	19.4	1.55 (1.03-2.33)	13.0	17.5	0.99 (0.58-1.67)	14.0	17.5	1.48 (1.15-1.91)
No asthma	7.6	12.7	1	11.5	19.6	1	8.0	13.2	1
Asthma, no ICS	11.7	15.9	1.46 (1.03 to 2.05)	8.3	18.9	0.91 (0.59 to 1.39)	9.8	16.9	1.27 (1.06 to 1.54)
Asthma, on ICS	17.6	10.7	1.46 (0.73 to 2.93)	23.1	15.4	0.99 (0.55 to 1.80)	14.0	14.0	1.25 (0.92 to 1.71)
No asthma	7.6	12.7	1	11.5	19.6	1	8.0	13.2	1
Asthma, no OCS	13.3	12.1	1.45 (1.08 to 1.96)	12.8	17.1	0.96 (0.66 to 1.39)	10.8	15.3	1.24 (1.06 to 1.45)
Asthma, on OCS	10.0	33.3	2.17 (0.88 to 5.36)	0.0	16.7	0.77 (0.11 to 5.14)	15.4	18.8	1.62 (0.72 to 3.64)
No asthma	7.6	12.7	1	11.5	19.6	1	8.0	13.2	1
Asthma, no ICS	11.7	15.9	1.47 (1.07 to 2.02)	8.3	18.9	0.91 (0.59 to 1.39)	9.8	16.9	1.27 (1.05 to 1.54)
Asthma, on ICS only	17.9	12.5	1.59 (0.87 to 2.92)	25.0	14.7	1.00 (0.53 to 1.86)	12.9	13.0	1.14 (0.87 to 1.51)
Asthma, on ICS & OCS	16.7	0.0	1.69 (0.39 to 7.25)	0.0	20.0	0.96 (0.15 to 6.01)	19.0	21.4	1.91 (0.84 to 4.31)

ECRHS: European Community Respiratory Health Survey; ICS: inhaled corticosteroids; OCS: oral corticosteroids; RR: relative risk.

1
2
3 Current asthma defined as a self-report of a doctor's asthma diagnosis and at least one of the following, within the last 12 months: waking up by an attack of shortness of
4 breath, any asthma attack or use of asthma medications. Individuals with inactive asthma (reporting asthma diagnosis without any current symptoms) were excluded from
5 the study (see Methods).
6

7 RR was calculated from modified Poisson regression with robust standard errors and subjects nested in centre as random effects. All models (for all participants) were
8 adjusted for age, sex and smoking status at time t .
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46

Supplementary Table E6: Association between asthma, asthma with atopy, asthma duration, and asthma with medication at t and incident obesity at $t+1$, including only participants who took part in all the three waves of ECRHS I, II and III.

	Obesity at visit, %		RR (95% CI)
	ECRHS II	ECRHS III	
No asthma	8.1	13.7	1
Asthma	11.9	15.6	1.25 (1.04 to 1.51)
No asthma	8.1	13.7	1
Asthma, without atopy	14.5	19.0	1.52 (1.12 to 2.07)
Asthma, with atopy	10.4	13.5	1.09 (0.87 to 2.07)
No asthma	8.1	13.7	1
Current asthma, ≤ 20 years since onset	10.0	12.3	1.05 (0.75-1.47)
Current asthma, >20 years since onset	14.4	18.0	1.43 (1.15-1.79)
No asthma	8.1	13.7	1
Asthma, no ICS	8.7	17.6	1.22 (1.00 to 1.50)
Asthma, on ICS	19.0	13.3	1.29 (0.97 to 1.72)
No asthma	8.1	13.7	1
Asthma, no OCS	11.2	15.1	1.19 (1.00 to 1.42)
Asthma, on OCS	19.4	21.9	1.97 (1.14 to 3.39)
No asthma	8.1	13.7	1
Asthma, no treatment	8.7	17.6	1.22 (1.00 to 1.49)
Asthma, on ICS only	17.2	12.1	1.13 (0.85 to 1.51)
Asthma, on ICS & OCS	25.9	21.4	2.21 (1.22 to 4.00)

ECRHS: European Community Respiratory Health Survey; ICS: inhaled corticosteroids; OCS: oral corticosteroids; RR: relative risk.

Current asthma defined as a self-report of a doctor's asthma diagnosis and at least one of the following, within the last 12 months: waking up by an attack of shortness of breath, any asthma attack or use of asthma medications. Individuals with inactive asthma (reporting asthma diagnosis without any current symptoms) were excluded from the study (see Methods).

RR was calculated from modified Poisson regression with robust standard errors and subjects nested in centre as random effects. All models (for all participants) were adjusted for age, sex and smoking status at time t .

Supplementary Table E7: Association between asthma, asthma with atopy, asthma duration, and asthma with medication at *t* and incident obesity at *t+1*, excluding the participants who reported avoiding physical activity because of wheezing/asthma.

	Obesity at visit, %		
	ECRHS II	ECRHS III	RR (95% CI)
No asthma	8.2	14.2	1
Asthma	9.9	15.3	1.14 (0.96 to 1.34)
No asthma	8.2	14.2	1
Asthma, without atopy	12.0	19.6	1.40 (1.02 to 1.92)
Asthma, with atopy	8.9	12.6	0.98 (0.79 to 1.22)
No asthma	8.2	14.2	1
Current asthma, ≤ 20 years since onset	9.2	13.2	1.03 (0.77-1.37)
Current asthma, >20 years since onset	11.1	17.0	1.25 (1.00-1.56)
No asthma	8.2	14.2	1
Asthma, no ICS	7.5	17.6	1.10 (0.97 to 1.24)
Asthma, on ICS	15.7	12.2	1.20 (0.88 to 1.63)
No asthma	8.2	14.2	1
Asthma, no OCS	9.6	14.6	1.09 (0.93 to 1.27)
Asthma, on OCS	15.6	27.3	2.02 (1.00 to 4.10)
No asthma	8.2	14.2	1
Asthma, no treatment	7.5	17.6	1.10 (0.97 to 1.24)
Asthma, on ICS only	14.7	10.4	1.06 (0.79 to 1.42)
Asthma, on ICS & OCS	21.7	27.8	2.37 (1.12 to 4.99)

ECRHS: European Community Respiratory Health Survey; ICS: inhaled corticosteroids; OCS: oral corticosteroids; RR: relative risk.

Current asthma defined as a self-report of a doctor's asthma diagnosis and at least one of the following, within the last 12 months: waking up by an attack of shortness of breath, any asthma attack or use of asthma medications. Individuals with inactive asthma (reporting asthma diagnosis without any current symptoms) were excluded from the study (see Methods).

RR was calculated from modified Poisson regression with robust standard errors and subjects nested in centre as random effects. All models (for all participants) were adjusted for age, sex and smoking status at time *t*.

Supplementary Table E8: Association between asthma, asthma with atopy, asthma duration, and asthma with medication at t and incident obesity at $t+1$, excluding the participants with dyspnoea at t .

	Obesity at visit, %		
	ECRHS II	ECRHS III	RR (95% CI)
No asthma	8.2	14.4	1
Asthma	9.7	15.9	1.14 (1.00 to 1.30)
No asthma	8.2	14.4	1
Asthma, without atopy	12.0	19.8	1.41 (1.15 to 1.72)
Asthma, with atopy	8.4	13.6	0.98 (0.78 to 1.24)
No asthma	8.2	14.4	1
Current asthma, ≤ 20 years since onset	8.8	13.2	1.01 (0.78-1.31)
Current asthma, >20 years since onset	11.1	17.9	1.27 (1.02-1.59)
No asthma	8.2	14.4	1
Asthma, no ICS	8.1	18.7	1.18 (1.07 to 1.31)
Asthma, on ICS	13.7	12.3	1.08 (0.80 to 1.46)
No asthma	8.2	14.4	1
Asthma, no OCS	9.4	15.6	1.11 (0.97 to 1.26)
Asthma, on OCS	14.3	20.0	1.58 (1.05 to 2.37)
No asthma	8.2	14.4	1
Asthma, no ICS	8.1	18.8	1.18 (1.07 to 1.31)
Asthma, ICS only	13.1	11.8	1.01 (0.74 to 1.37)
Asthma, on OCS & ICS	16.7	16.0	1.52 (0.89 to 2.59)

ECRHS: European Community Respiratory Health Survey; ICS: inhaled corticosteroids; OCS: oral corticosteroids; RR: relative risk.

Current asthma defined as a self-report of a doctor's asthma diagnosis and at least one of the following, within the last 12 months: waking up by an attack of shortness of breath, any asthma attack or use of asthma medications. Individuals with inactive asthma (reporting asthma diagnosis without any current symptoms) were excluded from the study (see Methods).

RR was calculated from modified Poisson regression with robust standard errors and subjects nested in centre as random effects. All models (for all participants) were adjusted for age, sex and smoking status at time t .

Supplementary Table E9: Association between asthma, asthma with atopy, asthma duration, and asthma with medication at t and incident obesity at $t+1$, using a broader asthma definition.

	Obesity at visit, %		
	ECRHS II	ECRHS III	RR (95% CI)
No asthma	7.9%	14.3%	1
Asthma	10.1%	15.5%	1.16 (1.04-1.30)
No asthma	7.9%	14.3%	1
Asthma, without atopy	12.3%	18.0%	1.38 (1.10-1.71)
Asthma, with atopy	8.8%	14.0%	1.03 (0.86-1.24)
No asthma	7.9%	14.3%	1
Current asthma, ≤ 20 years since onset	9.8%	13.5%	1.09 (0.92-1.28)
Current asthma, > 20 years since onset	10.2%	17.2%	1.24 (1.03-1.48)
No asthma	7.9%	14.3%	1
Asthma, no ICS	8.5%	16.5%	1.13 (1.01-1.27)
Asthma, on ICS	15.0%	13.5%	1.24 (0.99-1.54)
No asthma	7.9%	14.3%	1
Asthma, no OCS	9.4%	14.7%	1.09 (0.97-1.23)
Asthma, on OCS	22.0%	27.8%	2.41 (1.61-3.61)
No asthma	7.9%	14.3%	1
Asthma, no ICS	8.5%	16.5%	1.13 (1.01-1.26)
Asthma, ICS only	13.3%	11.6%	1.06 (0.85-1.32)
Asthma, on OCS+ICS	24.2%	26.7%	2.41 (1.43-4.01)

ECRHS: European Community Respiratory Health Survey; ICS: inhaled corticosteroids; OCS: oral corticosteroids; RR: relative risk.

Current asthma defined as a self-report of a doctor's asthma diagnosis as having reported an asthma diagnosis and at least one of the following, within the last 12 months: waking up by an attack of shortness of breath, any asthma attack, use of asthma medications, wheeze, shortness of breath, waking up with chest tightness or attack of cough. Individuals with inactive asthma (reporting asthma diagnosis without any current symptoms) were excluded from the study (see Methods).

RR was calculated from modified Poisson regression with robust standard errors and subjects nested in centre as random effects. All models (for all participants) were adjusted for age, sex and smoking status at time t .

Supplementary Table E10: Association between asthma, asthma with atopy, asthma duration, and asthma with medication at t and BMI changes between t and $t+1$.


	BMI change (kg/m ²), m (SD)		Beta (95% CI)*
	ECRHS I-II	ECRHS II-III	
No asthma	1.55 (2.2)	1.86 (2.6)	
Asthma	1.78 (2.2)	1.66 (2.7)	0.20 (0.01 – 0.38)
No asthma	1.55 (2.2)	1.86 (2.6)	
Asthma, without atopy	2.05 (2.3)	1.85 (2.9)	0.48 (0.17-0.78)
Asthma, with atopy	1.64 (2.1)	1.56 (2.5)	0.04 (-0.19-0.27)
No asthma	1.55 (2.2)	1.86 (2.6)	
Current asthma, ≤ 20 years since onset	1.91 (2.2)	1.61 (2.8)	0.25 (0.01-0.49)
Current asthma, > 20 years since onset	1.61 (2.2)	1.71 (2.6)	0.12 (-0.15-0.39)
No asthma	1.55 (2.2)	1.86 (2.6)	
Asthma, no ICS	1.70 (2.1)	1.81 (2.84)	0.11 (-0.10 – 0.33)
Asthma, on ICS	1.97 (2.3)	1.49 (2.41)	0.39 (0.07 – 0.71)
No asthma	1.55 (2.2)	1.86 (2.6)	
Asthma, no OCS	1.76 (2.2)	1.62 (2.6)	0.17 (-0.02 – 0.36)
Asthma, on OCS	2.12 (2.7)	2.10 (2.8)	0.56 (-0.10 – 1.22)
No asthma	1.55 (2.2)	1.86 (2.6)	
Asthma, no treatment	1.70 (2.1)	1.81 (2.8)	0.11 (-0.10 – 0.32)
Asthma, on ICS only	1.95 (2.2)	1.44 (2.4)	0.37 (0.02 – 0.72)
Asthma, on ICS & OCS	2.09 (2.9)	1.77 (2.4)	0.53 (-0.26 – 1.31)

ECRHS: European Community Respiratory Health Survey; ICS: inhaled corticosteroids; OCS: oral corticosteroids; RR: relative risk.

Current asthma defined as a self-report of a doctor's asthma diagnosis and at least one of the following, within the last 12 months: waking up by an attack of shortness of breath, any asthma attack or use of asthma medications. Individuals with inactive asthma (reporting asthma diagnosis without any current symptoms) were excluded from the study (see Methods).

Coefficients estimated from linear mixed regression with subjects nested in centre as random effects. All models (for all participants) were adjusted for age, sex, smoking status and BMI at time t , and an interaction term for period.

Supplementary Figure E1: Meta-analysis of the association between asthma at t and obesity at $t+1$ by ECRHS centre.



RR was calculated from modified Poisson regression with robust standard errors and subjects nested in centre as random effects. All models were adjusted for age, sex and smoking status at time t .

I-squared: variation in estimated effect attributable to heterogeneity. ES: effect estimate.