

AperTO - Archivio Istituzionale Open Access dell'Università di Torino

**Hermetia illucens meal inclusion in low-fishmeal diets for rainbow trout (*Oncorhynchus mykiss*): Effects on the growth performance, nutrient digestibility coefficients, selected gut health traits, and health status indices**

**This is a pre print version of the following article:**

*Original Citation:*

*Availability:*

This version is available <http://hdl.handle.net/2318/1863881> since 2024-02-16T11:58:28Z

*Published version:*

DOI:10.1016/j.anifeedsci.2022.115341

*Terms of use:*

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.



(Article begins on next page)

# Animal Feed Science and Technology

## Dietary *Hermetia illucens* meal inclusion in commercial diets for rainbow trout (*Oncorhynchus mykiss*): effects on nutrient digestibility, growth performance, and fish health

--Manuscript Draft--

<b>Manuscript Number:</b>	
<b>Article Type:</b>	Research Paper
<b>Section/Category:</b>	Aquatic animal species
<b>Keywords:</b>	Black soldier fly, commercial feed, fish, growth performance, gut microbiota, insect meal, nutrient digestibility.
<b>Corresponding Author:</b>	Francesco Gai CNR ISPA Grugliasco, Italy
<b>First Author:</b>	Ilaria Biasato
<b>Order of Authors:</b>	Ilaria Biasato Giulia Chemello Sara Bellezza Oddon Ilario Ferrocino Maria Rita Corvaglia Christian Caimi Andrea Resconi Aman Paul Michel van Spankeren Maria Teresa Capucchio Elena Colombino Luca Cocolin Francesco Gai Achille Schiavone Laura Gasco
<b>Abstract:</b>	<p>The effects of including <i>Hermetia illucens</i> (HI) meal in diets for rainbow trout have already been widely characterized, but data related to its utilization in commercial diets (especially when gut microbiota is considered) are quite scarce. This study aimed to investigate the impact of HI meal inclusion in commercial diets for rainbow trout by assessing fish growth performance, nutrient digestibility, histomorphological traits of intestine and main organs, and intestinal microbiota. In the 133-days growth trial, 600 rainbow trout were randomly distributed to 4 dietary treatments (3 replicate tanks/diet, 50 fish/tank): a low fishmeal-based diet as control (HI0), and three experimental diets including 80, 160 e 320 g/kg of HI meal as fed as replacement of 25, 50 and 100% of fishmeal (HI25, HI50 and HI100, respectively). At the end of the trial, growth parameters, condition factor and somatic indices were assessed, and gut, stomach, liver and spleen samples (12 fish/diet) were collected for histomorphological analyses. Feed and posterior intestine content were also sampled to characterize the feed and gut microbiota respectively. In the digestibility trial, 216 trout (3 tanks/treatment, 18 fish/tank) were used to evaluate the <i>in vivo</i> apparent digestibility coefficients (ADC) of the same diets. Unaffected growth performance, condition factor, somatic indices, nutrient digestibility, and histomorphological features were observed in the HI-fed rainbow trout (<math>P &gt; 0.05</math>). Increasing levels of HI meal inclusion in the feeds determined a progressive increase in the relative abundance of Firmicutes and Actinobacteria</p>

	<p>phyla, and Staphylococcus , Enterococcus , Oceanobacillus and Actinomyces genera, whereas Proteobacteria – as well as Lactobacillus and Listeria – displayed a gradual reduction. Dietary HI meal inclusion increased the Chao1 index of the fish gut microbiota, but, at the same time, reduced the Shannon index (<math>P &lt; 0.05</math>). The HI25 and HI50 fish also displayed higher relative abundance of Actinobacteria when compared to the other dietary treatments, as well as decreased Bacteroidetes (False Discovery Rate [FDR] <math>&lt; 0.05</math>). Furthermore, Actinomyces , Bacillus , Enterococcus , Staphylococcus , and Oceanobacillus resulted to be enriched in the posterior gut microbiota of the HI-fed fish (FDR <math>&lt; 0.05</math>). Differently, dietary HI meal inclusion determined a reduction of Campylobacter and Listeria , as well as Clostridium , Lactobacillus , Leuconostoc , Pediococcus , unclassified members (U.m.) of Peptostreptococceae, Weissella , Vagococcus , and Lactococcus . In conclusion, HI meal can be used in commercial diets for rainbow trout up to high inclusion levels (32%) without negatively affecting the growth performance, nutrient digestibility, somatic indices and histomorphological features of the animals. Furthermore, a positive modulation of the gut microbiota in terms of selection of short chain fatty acids (SCFAs)-producing bacteria and reduction of foodborne disease-causing pathogens was herein observed.</p>
<b>Suggested Reviewers:</b>	<p>Ines Guerreiro  msguerreiro@gmail.com  She is an expert in fish nutrition.</p> <p>Giuliana Parisi  giuliana.parisi@unifi.it  She is an expert in fish nutrition and insect meal utilization in fish.</p>
<b>Opposed Reviewers:</b>	

**From:** Dr. Francesco Gai  
Institute of Sciences of Food Production  
National Research Council  
Largo Paolo Braccini 2  
Grugliasco (TO), 10095  
0116709232  
francesco.gai@ispa.cnr.it

**To:** Dr. Kumar  
Co-Editor  
*Animal Feed Science and Technology*

January 08<sup>th</sup>, 2022

Dear Dr. Kumar,

I am pleased to submit an original research article entitled “Dietary *Hermetia illucens* meal inclusion in commercial diets for rainbow trout (*Oncorhynchus mykiss*): effects on nutrient digestibility, growth performance, and fish health” for consideration for publication in *Animal Feed Science and Technology*.

In this manuscript, we investigated the effects of insect meal inclusion in commercial diets for rainbow trout. Despite several studies about the impact of insect meal utilization having already been performed in different fish species (rainbow trout included), this manuscript represents the first scientific evidence not only about the insect-related gut microbiota modulation when a commercial diet is administered, but also the first scientific evidence ever related to the characterization of the microbiota of an insect-based feed.

We believe that this manuscript is appropriate for publication by *Animal Feed Science and Technology* because it provides novel and useful information about the role of taurine in rainbow trout.

This manuscript has not been published and is not under consideration for publication elsewhere. We have no conflicts of interest to disclose.

Thank you for your consideration.

Best regards,

Dr. Francesco Gai

## Highlights

- *Hermetia illucens* meal does not alter the growth performance of rainbow trout
- *Hermetia illucens* meal does not affect nutrient digestibility of rainbow trout
- *Hermetia illucens* meal does not impair the health status of rainbow trout
- *Hermetia illucens* meal does positively modulate the gut microbiota of rainbow trout

### **Author statement**

**Ilaria Biasato:** conduct the experiment, sampling, statistical analysis, and writing the initial draft,

**Giulia Chemello:** conduct the experiment, sampling, statistical analysis, and writing the initial draft,

**Sara Bellezza Oddon:** fish feeding, sampling, and reviewing the final draft,

**Ilario Ferrocino:** feed and gut microbiota analyses, and reviewing the final draft,

**Christian Caimi:** fish feeding, sampling, and reviewing the final draft,

**Andrea Resconi:** fish feeding, sampling, and reviewing the final draft,

**Aman Paul:** feed production and reviewing the final draft,

**Michel van Spankeren:** feed production and reviewing the final draft,

**Maria Teresa Capucchio:** histomorphological analysis and reviewing the final draft,

**Elena Colombino:** histomorphological analysis and reviewing the final draft,

**Luca Cocolin:** feed and gut microbiota analyses, and reviewing the final draft,

**Francesco Gai:** planning the research activity and reviewing the final draft,

**Achille Schiavone:** planning the research activity and reviewing the final draft,

**Laura Gasco:** coordination, funding acquisition, planning the research activity, and reviewing the final draft.

### **Declaration of Competing Interest**

There are no competing financial, professional, or personal interests that might have influenced the presentation of the work described in this manuscript.

**Dietary *Hermetia illucens* meal inclusion in commercial diets for rainbow trout (*Oncorhynchus mykiss*): effects on nutrient digestibility, growth performance, and fish health**

I. Biasato<sup>a,\*</sup>, G. Chemello<sup>b,\*</sup>, S. Bellezza Oddon<sup>a</sup>, I. Ferrocino<sup>a</sup>, M.R. Corvaglia<sup>a</sup>, C. Caimi<sup>a</sup>, A. Resconi<sup>a</sup>, A. Paul<sup>c</sup>, M. van Spankeren<sup>c</sup>, M.T. Capucchio<sup>d,e</sup>, E. Colombino<sup>d</sup>, L. Cocolin<sup>a</sup>, F. Gai<sup>e,§</sup>, A. Schiavone<sup>d,e</sup>, and L. Gasco<sup>a</sup>.

<sup>a</sup>Department of Agricultural, Forest and Food Sciences, University of Turin, Grugliasco (TO), Italy

<sup>b</sup>Department of Life and Environmental Sciences, Marche Polytechnic University, Ancona, Italy

<sup>c</sup>Protix BV, Dongen, The Netherlands

<sup>d</sup>Department of Veterinary Sciences, University of Turin, Grugliasco (TO), Italy

<sup>e</sup>Institute of Sciences of Food Production, National Research Council, Grugliasco (TO), Italy

\*B.I. and G.C. contributed equally to this paper.

§Corresponding author: Dr. Francesco Gai, Institute of Sciences of Food Production, National Research Council, Grugliasco (TO), 10095, Italy. Email address: francesco.gai@ispa.cnr.it



## Abstract

The effects of including *Hermetia illucens* (HI) meal in diets for rainbow trout have already been widely characterized, but data related to its utilization in commercial diets (especially when gut microbiota is considered) are quite scarce. This study aimed to investigate the impact of HI meal inclusion in commercial diets for rainbow trout by assessing fish growth performance, nutrient digestibility, histomorphological traits of intestine and main organs, and intestinal microbiota. In the 133-days growth trial, 600 rainbow trout were randomly distributed to 4 dietary treatments (3 replicate tanks/diet, 50 fish/tank): a low fishmeal-based diet as control (HI0), and three experimental diets including 80, 160 e 320 g/kg of HI meal as fed as replacement of 25, 50 and 100% of fishmeal (HI25, HI50 and HI100, respectively). At the end of the trial, growth parameters, condition factor and somatic indices were assessed, and gut, stomach, liver and spleen samples (12 fish/diet) were collected for histomorphological analyses. Feed and posterior intestine content were also sampled to characterize the feed and gut microbiota respectively. In the digestibility trial, 216 trout (3 tanks/treatment, 18 fish/tank) were used to evaluate the *in vivo* apparent digestibility coefficients (ADC) of the same diets. Unaffected growth performance, condition factor, somatic indices, nutrient digestibility, and histomorphological features were observed in the HI-fed rainbow trout ( $P > 0.05$ ). Increasing levels of HI meal inclusion in the feeds determined a progressive increase in the relative abundance of Firmicutes and Actinobacteria phyla, and *Staphylococcus*, *Enterococcus*, *Oceanobacillus* and *Actinomyces* genera, whereas Proteobacteria – as well as *Lactobacillus* and *Listeria* – displayed a gradual reduction. Dietary HI meal inclusion increased the Chao1 index of the fish gut microbiota, but, at the same time, reduced the Shannon index ( $P < 0.05$ ). The HI25 and HI50 fish also displayed higher relative abundance of Actinobacteria when compared to the other dietary treatments, as well as decreased Bacteroidetes (False Discovery Rate [FDR]  $< 0.05$ ). Furthermore, *Actinomyces*, *Bacillus*, *Enterococcus*, *Staphylococcus*, and *Oceanobacillus* resulted to be enriched in the posterior gut microbiota of the HI-fed fish (FDR  $< 0.05$ ). Differently, dietary HI meal inclusion determined a reduction of *Campylobacter* and *Listeria*, as well as *Clostridium*, *Lactobacillus*,

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

*Leuconostoc*, *Pediococcus*, unclassified members (U.m.) of Peptostreptococceae, *Weissella*, *Vagococcus*, and *Lactococcus*. In conclusion, HI meal can be used in commercial diets for rainbow trout up to high inclusion levels (32%) without negatively affecting the growth performance, nutrient digestibility, somatic indices and histomorphological features of the animals. Furthermore, a positive modulation of the gut microbiota in terms of selection of short chain fatty acids (SCFAs)-producing bacteria and reduction of foodborne disease-causing pathogens was herein observed.

### Keywords

Black soldier fly, commercial feed, fish, growth performance, gut microbiota, insect meal, nutrient digestibility.

### Abbreviations

AA, amino acid; ADF, acid detergent fiber; ADC, apparent digestibility coefficient; CF, coefficient of fatness; CY, carcass yield; CP, crude protein; DM, dry matter; EE, ether extract; FA, fatty acid; FAME, fatty acid methyl esters; FCR, feed conversion ratio; FDR, false discovery rate; FM, fishmeal; HE, Haematoxylin & Eosin; HI0, control diet; HI25, *Hermetia illucens* meal as replacement of 25% of fishmeal; HI50, *Hermetia illucens* meal as replacement of 50% of fishmeal; HI100, *Hermetia illucens* meal as replacement of 100% of fishmeal; HSI, hepatosomatic index; iFBW, individual final body weight; iIBW, individual initial body weight; iWG, individual weight gain; NDF, neutral detergent fiber; OTUs, Operational Taxonomic Units; PER, protein efficiency ratio; SCFAs, short chain fatty acids; SGR, specific growth rate; TFA, total fatty acids; Vh, villus height; VSI, viscerosomatic index.

### Introduction

The commercial rearing of insects for feed production represents a market that has grown rapidly in the recent years, being also ready to scale up production (All About Feed, 2020). Recent economic

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

projections highlighted that the global edible insects market is expected to reach around USD 8 billion and a volume of 730,000 tonnes by 2030, with a CAGR of 24.4% and 27.8%, respectively, during the forecast period from 2019 to 2030 (Meticulous Research®, 2020). Since the use of insect proteins was firstly authorized in aquafeed by the EU (Annexe II of Regulation 2017/893 of the 24<sup>th</sup> of May, 2017), the aquaculture segment dominated the insect market, with a consumption of more than 50% (around 5,000 tonnes) of the European animal feed produced from insects (IPIFF, 2019). The rapid development of the insect sector is related to the strong ability of insects to transform the nutrients losses (food waste) back into the food chain in forms of protein-rich animal feed, thus allowing them to fully embrace the concept of “circular economy” (Ojha et al., 2020). Among the farmed insect species, the black soldier fly (*Hermetia illucens*, HI) represents the most popular choice for mass production, because of its short life cycle, better feed conversion ratio (FCR), and the efficiency in bioconversion (50–60%) and recovery of nutrients from a wide spectrum of organic materials (Sheppard et al., 1994). This scenario has stimulated the insect producers in the EU to invest more than € 600 million in scaling up their production in 2019, with more than € 2.5 billion being even invested in 2020 (All About Feed, 2020). However, this growth is strictly connected with two important challenges, such as the meet of consumer’s expectations (in terms of consumption of safe, nutritious, and high-quality products) and the update of the regulatory framework (as no animal-based foodstuff can be used to feed insects, with the exception of the ones listed in the Reg. (EU) 2021/1372). In order to overcome these barriers (and, accelerate the scale up process), the insect producers need to currently test their products in the experimental setup.

In order to assess if a novel feed ingredient (such as insect-based products) can be suitable for fish feeding, a two-way approach is commonly adopted. First, the nutritional profile of the feed source needs to be fully characterized, as well as the feed acceptance, the growth performance and the nutrient digestibility by the fish (Rawski et al., 2020). Secondly, the implications for animal health must be investigated, with the attention being mainly directed towards the role of the gut. Indeed, the health status of the intestine (in terms of morphological development, mucin production, and

1 microbiota/microbiome) is fundamental to guarantee a proper health and growth of the fish (Józefiak  
2 et al., 2019; Caimi et al., 2020). So far, the administration of fishmeal (FM)-based diets containing  
3  
4 high levels of HI meal up to 40% has been reported to not influence (Renna et al., 2017; Cappelozza  
5  
6 et al., 2019; Cardinaletti et al., 2019) or worsen (St-Hilarie et al., 2007; Sealey et al., 2011; Dumas et  
7  
8 al., 2018) the growth performance of rainbow trout (*Oncorhynchus mykiss*), with some authors also  
9  
10 reporting unaffected (Renna et al., 2017) or reduced (Dumas et al., 2018; Cardinaletti et al., 2019)  
11  
12 length of the intestinal villi. In parallel, the gut mucin production has been described as unaltered  
13  
14 (Elia et al., 2018), while a positive modulation of the intestinal microbiota in terms of increased  
15  
16 microbial diversity, selection of potentially beneficial bacteria, and reduction of potential pathogens  
17  
18 has been identified in HI-fed fish (Bruni et al., 2018; Huyben et al., 2019; Rimoldi et al., 2019; Terova  
19  
20 et al., 2019). However, the potential of using HI-based products in commercial diets – which are low-  
21  
22 FM feeds with plant-derived proteins as additional protein sources – has recently started being  
23  
24 explored at low inclusion levels only (3-15%; Caimi et al., 2021). Furthermore, no data about gut  
25  
26 microbiota modulation in rainbow trout fed HI-based commercial diets are available yet.  
27  
28  
29  
30  
31  
32

33  
34 Therefore, the present study aims to investigate the effects of including increasing levels of a partially  
35  
36 defatted HI meal in commercial diets for rainbow trout as partial or total replacement of FM. In  
37  
38 particular, the attention was herein focused on the fish growth performance, nutrient digestibility, and  
39  
40 gut health parameters.  
41  
42  
43  
44  
45

## 46 **Materials and Methods**

47  
48 Two experimental trials (a digestibility and a growth trial, respectively) were conducted at the  
49  
50 Experimental Facility of the Department of Agricultural, Forest, and Food Sciences (DISAFA) of the  
51  
52 University of Turin (Italy). The experimental protocol was designed according to the guidelines of  
53  
54 the European and Italian regulations on the care and use of experimental animals (European directive  
55  
56 86 609/EEC, put into law in Italy with D.L. 116/92). The experimental protocol was approved by the  
57  
58 Ethical Committee of the University of Turin (protocol n° 143811).  
59  
60  
61  
62  
63  
64  
65

1  
2 *Experimental diets*  
3

4 Two diets containing FM (206 g/kg as fed; HI0) or a partially defatted HI meal produced in the  
5 experimental facility of a Dutch insect producer (Protix BV, Dongen, The Netherlands – 320 g/kg as  
6 fed; HI100) in substitution of 100% of FM were formulated by Research Diet Services BV (Utrecht,  
7 The Netherlands) and DISAFA. For nutrient digestibility evaluation, 10 g/kg as fed of Diamol (an  
8 acid insoluble ash) was added as inert marker. The two diets were formulated to be isonitrogenous,  
9 isolipidic, and isoenergetic. After that, two additional experimental diets were prepared by mixing:  
10 1) 750 g/kg as fed of HI0 and 250 g/kg as fed of HI100 (HI25), and 2) 500 g/kg as fed of HI0 and  
11 500 g/kg as fed of HI100 (HI50). The control diet (HI0) was formulated to mimic a commercial diet  
12 for rainbow trout, while the four experimental diets included increasing levels of HI meal in  
13 substitution of 0% (HI0), 25% (HI25), 50% (HI50) and 100% (HI100) of FM (corresponding to  
14 dietary HI meal inclusion levels of 0, 80, 160 and 320 g/kg as fed, respectively). The four diets (shown  
15 in Table 1) were prepared as extruded feed by Research Diet Services BV and shipped to the  
16 Experimental Facility of DISAFA. The diets were stored at 0-4°C and 85-90% RH in dark room  
17 before feeding to the fish.  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40

41 *Chemical analyses of feed*  
42

43 Feed samples were ground using a cutting mill (MLI 204; Bühler AG, Uzwil, Switzerland) and  
44 analysed for dry matter (DM; AOAC #934.01), crude protein (CP; AOAC #984.13), acid detergent  
45 fiber (ADF; AOAC# 973.18) and ash (AOAC #942.05) contents according to AOAC International  
46 (2000). Feed samples were also analysed for ether extract (EE; AOAC #2003.05) content according  
47 to AOAC International (2003), while the neutral detergent fiber (NDF) was analysed according to  
48 Van Soest et al. (1991);  $\alpha$ -amylase (Sigma Aldrich, Saint Louis, MO, USA) was added, but no sodium  
49 sulphite, and the results were corrected for the residual ash content. The GE content was determined  
50 using an adiabatic calorimetric bomb (C7000; IKA, Staufen, Germany). The fatty acid (FA)  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

1 composition of the experimental diets was assessed using the method described by Schmid et al.  
2 (2009). Fatty acid methyl esters (FAME) were separated, identified and quantified on the basis of the  
3 chromatographic conditions reported by Renna et al. (2014). The results were expressed as g/100 g  
4 of DM of total detected fatty acids (TFA). All the chemical analyses of the feeds were performed in  
5 duplicate (proximate composition) and triplicate (FA composition). The proximate composition and  
6 the FA profile of the experimental diets are shown in Table 1 and 2, respectively. Feed were also  
7 sampled for the microbiota assessment (please, see “DNA extraction and 16S rRNA amplicon target  
8 sequencing” subsection).

### 20 *Digestibility trial*

21 An *in vivo* digestibility trial was performed in order to determine the apparent digestibility  
22 coefficients (ADC) of the diets. A total of two hundred and sixteen trout (purchased from a private  
23 fish hatchery [“Troticoltura Bassignana”, Cuneo, Italy], with a weight of  $160.25 \pm 8.24$  g) were  
24 divided into twelve 250-L cylindroconical tanks (3 replicate tanks/diet, 18 fish/tank) connected to a  
25 flow-through open system where artesian well water (constant T of  $13 \pm 1$  °C) was supplied (tank  
26 water inflow: 8 L/min), and the dissolved oxygen levels were measured every two weeks (range: 7.6-  
27 8.7 mg/L). After 14 days of acclimatization with the experimental diets, the fish were fed by hand to  
28 visual satiety twice a day (8:00 am and 3:00 pm). The ADC were measured using the indirect acid-  
29 insoluble ash method, with 1% Diamol being used as inert marker. The faeces were collected daily  
30 from each tank for four consecutive week, using a continuous automatic device, as described by  
31 Chemello et al. (2020). The faeces were frozen ( $-20$  °C) and successively freeze-dried and stored  
32 until chemical analyses. The ADC of DM (ADCDM), crude protein (ADCCP), ether extract  
33 (ADCEE) and gross energy (ADCGE) were calculated according to Chemello et al. (2020).

### 57 *Growth trial*

1 A total of six hundred rainbow trout were purchased from a private fish hatchery (“Troticoltura  
2 Bassignana”, Cuneo, Italy). After a four-week period of acclimation (during which the fish were fed  
3 a commercial diet [42% CP and 22% EE; Skretting Italia Spa, Mozzecane, Verona, Italy]), the  
4 rainbow trout were submitted to a light anaesthesia (MS-222; PHARMAQ Ltd., Sandleheath, UK; 60  
5 mg/L), individually weighed ( $112.86 \pm 8.41$  g) using electronic scales (KERN PLE-N v. 2.2; KERN  
6 & Sohn GmbH, Balingen-Frommern, Germany; d: 0.1), and randomly allotted to twelve 300-L,  
7 rectangular-shaped tanks (three replicate tanks per diet, fifty fish per tank) connected to the same  
8 flow-through open water system of the digestibility trial. The fish were fed 1.4% of the tank biomass  
9 for the first 123 days of trial, while the feeding rate was reduced to 1.1% for the remaining 20 days.  
10 In particular, the fish were fed by hand, twice a day (08:00 and 15:00) and six days per week. Feed  
11 intake was checked at each administration, and feed distribution was immediately interrupted if fish  
12 stopped eating. In order to update the daily feeding rate, the biomass tanks were weighed in bulk  
13 every 14 days. Mortality was daily checked. The experimental trial lasted 133 days.  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33

### 34 *Growth performance*

35  
36 At the end of the growth trial, the fish were left unfed for one day, submitted to a light anaesthesia  
37 (MS-222; PHARMAQ Ltd., Sandleheath, UK; 60 mg/L) and individually weighed (KERN PLE-N  
38 v.2.2; KERN and SOHN GmbH, Balingen-Frommern, Germany; d: 0.1). The following performance  
39 indices were calculated per each tank:  
40  
41  
42  
43  
44

- 45 1. Survival (%) =  $100 - [(\text{number of dead fish} / \text{number of fish at start}) \times 100]$
  - 46 2. Individual weight gain (iWG, g) = average individual final body weight (iFBW, g) – average  
47 individual initial body weight (iIBW, g)
  - 48 3. Feed conversion ratio (FCR) = total feed supplied (g, DM) / WG (g)
  - 49 4. Protein efficiency ratio (PER) = WG (g) / total protein fed (g, DM)
  - 50 5. Specific growth rate (SGR, % day<sup>-1</sup>) =  $[(\ln \text{FBW} - \ln \text{IBW}) / \text{number of feeding days}] \times 100$
- 51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

### *Condition factor and somatic indices*

At the end of the growth trial, twenty-one fish per dietary treatment (seven fish/tank) were killed by over anaesthesia (MS-222; PHARMAQ Ltd., Sandleheath, UK; 500 mg/L). The fish were individually weighted (KERN PLE-N v.2.2; KERN and SOHN GmbH, Balingen-Frommern, Germany; d: 0.1), and fish total length was manually measured to determine the Fulton's condition factor (K). The fish were then dissected in order to calculate the carcass yield (CY). Liver, gut and perivisceral fat were successively weighted (KERN PLE-N 420-3N; KERN and SOHN GmbH, Balingen-Frommern, Germany; d: 0.001) to calculate the hepatosomatic index (HSI), the viscerosomatic index (VSI), and the coefficient of fatness (CF) as follows:

- $K = [\text{fish weight (g)} / (\text{body length})^3 \text{ (cm)}] \times 100;$
- $CY (\%) = [\text{total weight without gut and gonad (g)} / \text{fish weight (g)}] \times 100;$
- $HSI (\%) = [\text{liver weight (g)} / \text{fish weight (g)}] \times 100;$
- $VSI (\%) = [\text{gut weight (g)} / \text{fish weight (g)}] \times 100;$
- $CF (\%) = [\text{perivisceral fat weight (g)} / \text{fish weight (g)}] \times 100.$

### *Sampling and processing*

At the end of the growth trial, twelve fish per dietary treatment (four fish per tank) were also killed by over anaesthesia (MS-222; PHARMAQ Ltd., Sandleheath, UK; 500 mg/L) and submitted to morphometric and histopathological investigations. Anterior (the tract immediately after the pyloric caeca) and posterior (the tract 1 cm before the anus) gut segment samples (approximately 2 cm in length) were excised, flushed with 0.9% saline to remove all the content, and fixed in 10% buffered formalin solution for histomorphological investigations, as well as liver, spleen and stomach. All the tissues were routinely embedded in paraffin wax blocks, sectioned at 5  $\mu\text{m}$  thickness, mounted on glass slides, and stained with Haematoxylin & Eosin (HE) for morphometric (gut) and histopathological (gut, liver, spleen and stomach) investigations. The posterior intestine content was



1  
2 also collected into sterile plastic tubes with appropriate squeezing, cooled at 4 °C (for a maximum of  
3 2 hours) and frozen at -80°C until DNA extraction.  
4  
5  
6

### 7 *Histomorphological investigations*

8  
9 One slide per each intestinal segment was examined by means of light microscopy, and one randomly  
10 selected high power field per each slide was captured with a Nikon DS-Fi1 digital camera (Nikon  
11 Corporation, Minato, Tokyo, Japan) coupled to a Zeiss Axiophot microscope (Carl Zeiss,  
12 Oberkochen, Germania) using a 2.5× objective lens. The NIS-Elements F software (Nikon  
13 Corporation, Minato, Tokyo, Japan) was then used for image capturing. All the morphometric  
14 measurements were performed by Image-Pro Plus 6.0 software (Media Cybernetics Inc., Bethesda,  
15 Rockville, MD, USA) on 10 well-oriented and intact villi in order to evaluate the villus height (Vh,  
16 from the villus tip to submucosa) (Renna et al., 2017). The observed histopathological findings were  
17 evaluated in all the organs using the semi-quantitative scoring system previously established (Elia et  
18 al., 2018): absent (score = 0), focal to multifocal, mild (score = 1), multifocal, moderate (score = 2),  
19 and multifocal to diffuse, severe (score = 3). Gut histopathological findings were separately assessed  
20 for mucosa (inflammatory infiltrates) and submucosa (inflammatory infiltrates and Gut-Associated  
21 Lymphoid Tissue [GALT] activation) for each segment, according to Biasato et al. (2019). The total  
22 score of each gut segment was obtained by adding up the mucosa and submucosa scores. All the  
23 slides were blind assessed by two independent observers and the discordant cases were reviewed,  
24 using a multi-head microscope, until unanimous consensus was reached.  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50

### 51 *DNA extraction and 16S rRNA amplicon target sequencing*

52 The total genomic DNA (gDNA) was extracted using the RNeasy Power Microbiome KIT (Qiagen,  
53 Milan, Italy), according to the manufacturer's instructions. One µL of RNase (Illumina Inc., San  
54 Diego, CA, USA) was added to digest RNA in the DNA samples with an incubation of 1 h at 37°C.  
55 The DNA was then quantified using the NanoDrop and standardized at 5 ng/µL. The gDNA was used  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

1  
2 to assess the microbiota composition by the amplification of the V3-V4 region of the 16S rRNA gene  
3 (Klindworth et al., 2013). The PCR products were purified according to the Illumina metagenomic  
4 standard procedure (Illumina Inc., San Diego, CA, USA). The sequencing was performed by a MiSeq  
5 Illumina instrument with V3 chemistry and generated 250 bp paired-end reads, according to the  
6 manufacturer's instructions.  
7  
8  
9

### 10 11 12 13 14 *Bioinformatics and statistical analysis* 15

16 The experimental unit was the tank for growth performance and nutrient digestibility, and the fish for  
17 somatic indices, histomorphological findings and 16S rRNA sequences.  
18  
19

20 Paired-end reads were first merged using FLASH software with default parameters (Magoc and  
21 Salzberg, 2011). Joint reads were further quality filtered (at Phred < Q20) using QIIME 1.9.0 software  
22 (Caporaso et al., 2010) and the pipeline recently described (Biasato et al., 2018). The Operational  
23 Taxonomic Units (OTUs) clustering was obtained at 97% of similarity and taxonomy assignment was  
24 assessed by Greengenes 16S rRNA gene database v. 2013. The OTUs table was rarefied at the lowest  
25 number of sequence and displayed the highest taxonomy resolution. The vegan package of R (Dixon,  
26 2003) was used to calculate the alpha diversity. The diversity indices were further analyzed by  
27 pairwise comparisons using Wilcoxon rank sum test to assess differences among the dietary  
28 treatments. The OTUs table filtered for relative abundance (>0.2% in at least five samples) was used  
29 to perform Anosim statistical test in R environment, and Pairwise Kruskal-Wallis tests allowed to  
30 find significant differences in microbial taxa abundance according to the dietary treatment. P-values  
31 were adjusted for multiple testing and a false discovery rate (FDR) < 0.05 considered as significant.  
32  
33

34 The statistical analysis of growth performance, nutrient digestibility, somatic indices and  
35 histomorphological findings was performed using IBM SPSS Statistics v, 26,0 (IBM, Armonk, NY,  
36 USA). Growth performance, nutrient digestibility and somatic indices data were analysed by one-  
37 way ANOVA. The Shapiro–Wilk test was used to check dependent variables for normality. The  
38 assumption of equal variances was assessed by Levene's homogeneity of variance test. If such an  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

1 assumption did not hold, the Brown-Forsythe statistic was performed to test for the equality of group  
2 means instead of the F one. Pairwise multiple comparisons were performed to test the difference  
3  
4 between each pair of means (Tukey's test and Tamhane's T2 in the cases of equal variances assumed  
5 or not assumed, respectively). The morphometric indices were analysed by fitting a general linear  
6  
7 model that allowed the morphometric indices (Vh) to depend on three fixed factors (diet, intestinal  
8  
9 segment, and interaction between diet and intestinal segment). The interactions between the levels of  
10  
11 the fixed factors were evaluated by pairwise contrasts. Histopathological scores were analysed by  
12  
13 Chi-square test. The results obtained from normally distributed data were expressed as mean (growth  
14  
15 performance, nutrient digestibility and somatic indices) or least square mean (morphometric indices)  
16  
17 and pooled standard error of the mean (SEM), while those obtained from not normally distributed  
18  
19 data (histopathological findings) as n (%). P values  $\leq 0.05$  were considered statistically significant.  
20  
21  
22  
23  
24  
25  
26  
27  
28

## 29 **Results**

### 30 *Chemical analyses of the feed*

31  
32  
33 The experimental diets were not fully comparable in terms of macronutrients (Table 1). In particular,  
34  
35 the HI100 diet showed numerically higher DM and CP, and lower EE when compared to the HI0 diet  
36  
37 (+2.21%, +4.86%, and -8.13%, respectively). However, the proximate composition of the HI25 and  
38  
39 HI50 diets was overall similar to that of the HI0 (Table 1). As far as the FA profile is concerned  
40  
41 (Table 2), the lauric (C12:0), myristic (C14:0), palmitic (C16:0), stearic (C18:0), oleic (C18:1 c9) and  
42  
43 linoleic (C18:2 n6) acids were the most represented FA in all the experimental diets. In particular, the  
44  
45 lauric, myristic, and palmitic acids increased with increasing HI meal inclusion levels, while the oleic  
46  
47 and linoleic acids displayed the opposite trend (Table 2). Subsequently, the total saturated fatty acids  
48  
49 (SFA) increased following the increased inclusion of the insect meal, whereas the total  
50  
51 monounsaturated and polyunsaturated fatty acids (MUFA and PUFA, respectively) decreased (Table  
52  
53  
54 2). The decrease in the PUFA was determined by the decrease in the arachidonic (C20:4 n6),  
55  
56 eicosapentaenoic (EPA, C20:5 n3), docosapentaenoic (DPA, C22:5 n3) and docosahexaenoic (DHA,  
57  
58  
59  
60  
61  
62  
63  
64  
65

1 C22:6 n3) acids, thus furtherly explaining the progressive reduction in the n3 and n6 FA (and the  
2 n6/n3 as well) identified in the experimental diets (Table 2).  
3  
4  
5  
6

### 7 *Digestibility trial*

8  
9 Dietary HI meal inclusion did not influence the nutrient digestibility of the rainbow trout ( $P > 0.05$ ,  
10 Table 3).  
11  
12  
13  
14  
15

### 16 *Growth trial*

#### 17 *Growth performance*

18  
19 Growth performance of the rainbow trout are summarized in Table 4. The fish readily accepted all  
20 the experimental diets, with all the supplied feed being consumed and no feed refusals being recorded  
21 during the experimental trial. Survival was high for all the dietary treatments (range: 95.33-97.33),  
22 being also unaffected by dietary HI meal inclusion ( $P > 0.05$ , Table 4). Similarly, all the other growth  
23 parameters were not affected by insect meal utilization ( $P > 0.05$ , Table 4).  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35

#### 36 *Condition factor and somatic indices*

37  
38 Dietary HI meal inclusion did not influence either the condition factor or the somatic indices of the  
39 rainbow trout ( $P > 0.05$ , Table 5).  
40  
41  
42  
43  
44

#### 45 *Histomorphological investigations*

46  
47 Data regarding the morphometric measurements of the Vh in the anterior and posterior gut are  
48 reported in Table 6. The Vh was not influenced by the diet and the interaction between the diet and  
49 the intestinal segment ( $P > 0.05$ , Table 6), but it only depended on the intestinal segment ( $P < 0.001$ ,  
50 Table 6). Independently of the dietary HI meal inclusion, the Vh showed a proximo-distal increasing  
51 gradient from the anterior to the posterior gut ( $P < 0.001$ , Table 6).  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

1 The histopathological alterations observed in liver, spleen, stomach, anterior and posterior gut are  
2 summarized in Table 7. In liver, absent to mild, focal to multifocal lymphoplasmacytic infiltrates, as  
3 well as absent to moderate, multifocal to diffuse fatty changes of the hepatocytes were observed in  
4 all the dietary treatments. Mild, focal to multifocal hemosiderosis, along with moderate, focal to  
5 multifocal white pulp hyperplasia were also recorded in all the experimental groups. No signs of  
6 immune cell infiltration were observed in the stomach, except for the HI100 group (8.3%). All the  
7 fish displayed mild, focal to multifocal lymphoplasmacytic infiltrates in both the anterior and the  
8 posterior intestine. However, dietary HI meal inclusion did not influence either the severity or the  
9 distribution of the observed histopathological alterations ( $P > 0.05$ , Table 7).  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23

#### 24 *Feed 16S rRNA amplicon target sequencing*

25 The feed samples were overall characterized by a simple microbiota, with Firmicutes, Cyanobacteria  
26 and Proteobacteria representing the main bacterial phyla, and *Lactobacillus*, *Listeria*, *Leuconostoc*,  
27 *Streptococcus* and *Photobacterium* the most abundant genera (Figure 1). Increasing levels of HI meal  
28 inclusion in the feeds determined a progressive increase in the relative abundance of Firmicutes and  
29 Actinobacteria phyla, whereas Proteobacteria displayed a gradual reduction (Figure 1A).  
30 Furthermore, the relative abundance of *Lactobacillus* and *Listeria* decreased with increasing levels  
31 of dietary HI meal inclusion, while *Staphylococcus*, *Enterococcus*, *Oceanobacillus* and *Actinomyces*  
32 showed the opposite trend (Figure 1B).  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47

#### 48 *Posterior gut 16S rRNA amplicon target sequencing*

49 After sequencing and quality filtering, 895,348 reads were used for the downstream analysis (with a  
50 median value of  $18,371 \pm 10,395$  reads/sample). The rarefaction analysis and the estimated sample  
51 coverage indicated that there was a satisfactory coverage of all the samples (ESC median value of  
52 96%). The alpha diversity analysis also revealed a significant increase in the Chao1 index of the  
53 posterior gut microbiota from the HI-fed rainbow trout, whereas the Shannon index displayed the  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

opposite trend ( $P < 0.05$ , Figure 2). By plotting the Principal Component Analysis (PCA), a clear separation between the fish fed the control and the HI-based diets was also observed, with a higher beta diversity being furtherly identified in the posterior gut microbiota from the HI25 rainbow trout when compared to the HI50 and HI100 groups ( $P < 0.001$ , Figure 3).

The characterization of the posterior gut microbiota of the rainbow trout overall revealed Firmicutes, Actinobacteria and Proteobacteria as predominant phyla (Figure 4A), while *Staphylococcus*, *Lactobacillus*, *Enterococcus*, *Oceanobacillus*, *Actinomyces*, *Streptococcus* and *Weissella* resulted to be the most abundant genera (Figure 4B). At phylum level (Figure 5), the HI25 and the HI50 fish showed a significant increase in the relative abundance of Actinobacteria in comparison with the HI0 group ( $FDR < 0.05$ ). On the contrary, Bacteroidetes phylum was significantly less abundant in the rainbow trout fed the HI25 and the HI50 diets when compared to the HI0 one ( $FDR < 0.05$ ). As far as genus level is concerned (Figure 6), the HI-fed fish showed a significant increase in the relative abundance of *Actinomyces*, *Bacillus*, *Enterococcus*, *Oceanobacillus*, and *Staphylococcus* ( $FDR < 0.05$ ). Differently, the relative abundance of *Clostridium*, *Campylobacter*, *Listeria*, *Lactobacillus*, *Lactococcus*, *Leuconostoc*, *Pediococcus*, unclassified members (U.m.) of Peptostreptococceae, *Vagococcus*, and *Weissella* genera was significantly decreased in the rainbow trout fed the HI-based diets in comparison with the HI0 group. No changes related to the different HI meal inclusion levels were, however, identified for both the phyla and the genera ( $FDR > 0.05$ ).

## Discussion

### *Digestibility trial*

The apparent digestibility of the nutrients and the energy of the HI-based diets was analogous to that recorded for the C diet, as already underlined by previous research (Renna et al., 2017; Caimi et al., 2021). This is indicative of a good, proper nutrient availability, which reasonably explains the unaffected growth performance highlighted in the HI-fed rainbow trout.

## *Growth trial*

### *Growth performance*

The growth performance of the rainbow trout of the present study were not affected by dietary HI meal inclusion, as already underlined by previous research (Renna et al., 2017; Cappellozza et al., 2019; Cardinaletti et al., 2019; Caimi et al., 2021). This represents a positive finding, since increasing levels of HI larva (26.4% [Dumas et al., 2018]) and prepupa (29.8% [St-Hilarie et al., 2007] or 32.80% [Sealey et al., 2011]) meals in the diets for rainbow trout have also been reported to worsen either the weight gain (St-Hilarie et al., 2007; Sealey et al., 2011) or the feed efficiency (St-Hilarie et al., 2007; Dumas et al., 2018) of the fish. Those different outcomes could be attributed to a potential, reduced nutrient availability of the HI diets (Sealey et al., 2011), which, in turn, is partially related to the use of full-fat HI meals (St-Hilarie et al., 2007; Sealey et al., 2011). Indeed, lower lipid (St-Hilarie et al., 2007; Sealey et al., 2011), GE (St-Hilarie et al., 2007), and CP (Dumas et al., 2018) contents were identified in the whole body (St-Hilarie et al., 2007; Dumas et al., 2018) and the muscle (Sealey et al., 2011) of rainbow trout fed the HI-based diets than the control. Despite no whole-body composition analysis having been performed in the current research, the unaffected nutrient digestibility herein observed in the HI-fed fish reasonably suggests no alterations in the nutrient availability as well. Apart from the nutritional composition, the quality of the HI meal in terms of rearing substrates on which the HI larvae were reared may exert a significant influence as well. Indeed, the use of manure from swine (St-Hilarie et al., 2007) and dairy cows (Sealey et al., 2011) may not represent an optimal rearing substrate when compared to the vegetable waste (Cappellozza et al., 2019; Cardinaletti et al., 2019; Caimi et al., 2021).

### *Condition factor and somatic indices*

Both the condition factor and the somatic indices of the rainbow trout of the present study were not significantly influenced by dietary HI meal inclusion. This is in agreement with previous research studies assessing the effects of HI meal utilization in rainbow trout, which also reported analogous K

1 (Renna et al., 2017; Cardinaletti et al., 2019), HSI (Sealey et al., 2011) and VSI (Bruni et al., 2018)  
2 values. All the dietary treatments showed K values higher than 1, thus implying that fish are in good  
3  
4 physiological state of well-being and, in turn, that dietary HI meal inclusion does not alter the  
5  
6 condition, fatness, or wellbeing of fish (Muddasir and Imtiaz, 2016; Renna et al., 2017). The  
7  
8 unaffected HSI and VSI are indicative of the absence of significant diseases in either the liver or the  
9  
10 gastrointestinal tract of the HI-fed rainbow trout, as altered values of HSI have previously been  
11  
12 ascribed to metabolic problems or liver deficiencies (Dernekbaşı, 2012), and no HI-related hepatic or  
13  
14 gastrointestinal histopathological alterations were herein identified. The unaltered values of CF in the  
15  
16 fish fed the HI-based diets is also indicative of a proper nutrient availability (Sealey et al., 2011).  
17  
18  
19  
20  
21  
22  
23

#### 24 *Histomorphological features*

25  
26 Dietary HI meal inclusion did not significantly affect the gut morphology of the rainbow trout of the  
27  
28 current research, as already reported by Renna et al. (2017). This is in agreement with the unaffected  
29  
30 growth performance herein observed in the HI-fed fish, thus suggesting no negative repercussions on  
31  
32 either the digestion or the absorption of the nutrients by the intestine. A shortening of the gut villi  
33  
34 (Dumas et al., 2018) and fold (Cardinaletti et al., 2019) has also previously been reported in rainbow  
35  
36 trout fed diets containing HI meal, with the growth performance of the fish being, however, impaired  
37  
38 with the highest inclusion level only (26.4% [Dumas et al., 2018]). Independently of HI utilization,  
39  
40 the posterior intestine of the rainbow trout of the present study showed longer villi than the anterior  
41  
42 segment. This appears to be in contrast with Khojasteh et al. (2009), which reported progressively  
43  
44 shorter villi toward the posterior intestine. However, the simultaneous presence of short and long villi  
45  
46 in both the gut segments – as well as the villi length changes throughout the fish cycle – has recently  
47  
48 been reported in rainbow trout (Verdile et al., 2020), thus making further investigations needed.  
49  
50  
51  
52  
53

54  
55 The histopathological alterations observed in the fish of the current research were also not  
56  
57 significantly influenced by HI meal utilization, thus suggesting no negative effects of HI on fish  
58  
59 health. Elia et al. (2018) previously described similar findings in the liver, spleen and anterior  
60  
61  
62  
63  
64  
65



1 intestine of rainbow trout, with no HI-related alterations being analogously identified. The fatty and  
2 inflammatory changes in liver and gastrointestinal tract, respectively, are the common result of the  
3 high-energy diet administered to salmonids, while the spleen reactivity appears to be aspecific.  
4 Furthermore, the histopathological alterations were identified in both the control- and the HI-fed fish,  
5 also resulting to be predominantly mild to moderate.  
6  
7  
8  
9

#### 10 11 12 13 14 *Feed and gut microbiota*

15 Firmicutes, Cyanobacteria and Proteobacteria phyla dominated the microbiota of the feed used in the  
16 present study. This is partially in agreement with Terova et al. (2019), which identified a  
17 predominance of Firmicutes, Proteobacteria and Actinobacteria in FM-based diets for rainbow trout.  
18 However, the detection of high percentages of Cyanobacteria represents an unexpected finding.  
19 Cyanobacteria has recently been found in the gut microbiota of marine (Salas-Leiva et al., 2020) and  
20 freshwater (Jiang et al., 2020; Zeng et al., 2020) species, being also one of the most abundant  
21 prokaryotes in sea (Korlević et al., 2016; Quéméneur et al., 2020) and anthropogenic-induced  
22 eutrophied freshwaters (Zhang et al., 2021). Considering that the biomass which supplies the FM  
23 industry is mainly composed of small pelagic species (Péron et al., 2010), it seems reasonable that  
24 the feed microbiota herein characterized reflect the gut microbiota of the fish species (and their  
25 rearing environment as well) used to produce the FM. The identification of high relative abundances  
26 of Firmicutes and Proteobacteria – which are two of the dominant bacterial phyla of the fish gut  
27 microbiota (Butt and Volkoff, 2019) – further supports such hypothesis. A similar consideration can  
28 also be made for the most represented bacterial genera detected in feed microbiota, as *Lactobacillus*  
29 (Tarnecki et al., 2017; Huyben et al., 2020; Yu et al., 2021), *Leuconostoc*, *Streptococcus* (Tarnecki  
30 et al., 2017) and *Photobacterium* (Huyben et al., 2020) constitute the core microbiota of several  
31 marine species, with the latter OTU being particularly characteristic of piscivores such the pelagic  
32 species (Huang et al., 2020). Differently, the detection of high percentages of *Listeria* may rise  
33 worrying concerns in terms of food safety, as some species (especially *L. monocytogenes*) are  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

1 involved in foodborne outbreaks of listeriosis (Buchanan et al., 2017). Since the consumption of raw  
2 and smoked seafood is one of the most common predisposing factor to develop such disease and  
3  
4 *Listeria* has frequently been isolated in marine finfish (Basha et al., 2019), the fish species herein  
5  
6 used to produce the FM could have potentially carried *Listeria* to the feeds.  
7  
8

9 The HI-based diets used in the current research were characterized by a progressive increase in the  
10  
11 relative abundance of Firmicutes and Actinobacteria phyla, whereas Proteobacteria displayed a  
12  
13 gradual reduction. This is in agreement with Terova et al. (2019), which described the same scenario  
14  
15 in feeds containing increasing levels of HI prepupa meal as FM replacement. This represents the  
16  
17 logical consequence of substituting the FM (which is obtained by carnivorous fish) with the insect  
18  
19 meal (which is obtained by larvae reared on vegetable substrates). Indeed, plant ingredients in the  
20  
21 diet are commonly associated with a higher Firmicutes:Proteobacteria ratio when compared to animal  
22  
23 protein-based diet, which, on the contrary, stimulates the proliferation of Proteobacteria (Rimoldi et  
24  
25 al., 2018). A clear increase in the relative abundance of *Staphylococcus*, *Enterococcus*,  
26  
27 *Oceanobacillus* and *Actinomyces* was also identified in the HI-based diets, thus partially agreeing  
28  
29 with the findings reported by Terova et al. (2019). The detection of increasing percentages of  
30  
31 *Oceanobacillus* represents, however, a novel, difficult-to-explain result, as this taxon has been  
32  
33 reported to dominate the gut microbiota of healthy shrimp, crab and clam (Sun et al., 2019).  
34  
35 Furthermore, despite Rimoldi et al. (2021) having recently discovered *Oceanobacillus* in HI-based  
36  
37 feed only, its relationship with insects remains to be fully elucidated. High amounts of *Lactobacillus*  
38  
39 in diets containing HI meal are also common (Terova et al., 2019; Rimoldi et al., 2021), while the  
40  
41 HI-based feeds used in the present study displayed a progressive reduction of this genus. This finding  
42  
43 – as well as the decrease of *Listeria* – is reasonably related to the FM replacement by insect meal, as  
44  
45 these OTUs are herein hypothesized to depend on the fish species used to produce the FM.  
46  
47

48 Dietary HI meal inclusion increased the gut microbial richness in the fish of the current research, but,  
49  
50 at the same time, reduced its diversity. This partially contrasts with the majority of the previous  
51  
52 findings in rainbow trout, which identified unaffected or higher Chao1 and Shannon indices in the  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

1 HI-fed fish when compared to those fed the control diet (Bruni et al., 2018; Huyben et al., 2019;  
2 Rimoldi et al., 2019; Terova et al., 2019; Rimoldi et al., 2021). This represents a challenging scenario,  
3  
4 as reduced bacterial diversity may determine less competition for incoming pathogens, thus favouring  
5  
6 their colonization of the gastrointestinal tract of fish and the development of several diseases  
7  
8 frequently related to several diseases (Terova et al., 2019). However, the rainbow trout fed the HI-  
9  
10 based diets of the present study remained healthy throughout the experimental trial, also showing no  
11  
12 significant histopathological alterations.  
13  
14  
15

16 Firmicutes, Actinobacteria and Proteobacteria represented the dominant bacterial phyla in both the  
17  
18 control- and HI-fed fish of the current research. These findings overall agree with the previous  
19  
20 research carried out in rainbow trout (Desai et al., 2012; Wong et al., 2013; Ingerslev et al., 2014;  
21  
22 Bruni et al., 2018; Rimoldi et al., 2018; Huyben et al., 2019; Terova et al., 2019; Pelusio et al., 2020).  
23  
24

25 In relation to the genera composition, *Staphylococcus*, *Lactobacillus* and *Streptococcus* mainly  
26  
27 colonized the posterior gut microbiota of the fish fed either the control or the HI-based diets in the  
28  
29 present study. These findings are also in agreement with the previous studies, which observed  
30  
31 *Lactobacillus* (Wong et al., 2013; Ingerslev et al., 2014; Rimoldi et al., 2018; Huyben et al., 2019;  
32  
33 Terova et al., 2019; Pelusio et al., 2020), *Streptococcus* (Ingerslev et al., 2014; Rimoldi et al., 2018;  
34  
35 Pelusio et al., 2020) and *Staphylococcus* (Bruni et al., 2018; Terova et al., 2019) as main bacterial  
36  
37 genera in the cecal microbiota of rainbow trout.  
38  
39  
40  
41  
42

43 In the current research, the utilization of HI meal at 25% and 50% inclusion levels determined higher  
44  
45 relative abundance of Actinobacteria phylum in the fish posterior gut microbiota when compared to  
46  
47 the HI0 group. A significant increase in Actinobacteria has also previously been reported in HI-fed  
48  
49 rainbow trout (Huyben et al., 2019; Terova et al., 2019), as well as the increment in Firmicutes (Bruni  
50  
51 et al., 2018; Huyben et al., 2019; Terova et al., 2019) and the reduction of Proteobacteria (Huyben et  
52  
53 al., 2019; Terova et al., 2019). On one hand, the increase in Actinobacteria herein observed partially  
54  
55 reflects the high relative abundance of this bacterial phylum detected in the HI-based diets; on the  
56  
57 other, some genera belonging to Actinobacteria (such as *Actinomyces*) are often identified as chitin  
58  
59  
60  
61  
62  
63  
64  
65

1 degraders (Beier and Bertilsson, 2013), thus partially explaining its high abundance in the HI-fed  
2 rainbow trout. Despite Firmicutes and Proteobacteria percentages being similar among the  
3 experimental treatments, the HI25 and the HI50 fish of the present study also displayed lower relative  
4 abundance of Bacteroidetes in their posterior gut microbiota in comparison with the HI0 group.  
5 Bacteroidetes members are well-known to be involved in the fermentation of dietary non-starch  
6 polysaccharides (NSP; den Besten et al., 2013). Since the HI-based diets were characterized by a  
7 progressive reduction of wheat meal content (which has considerable quantity of NSP), the decrease  
8 in Bacteroidetes may represent a reasonable consequence. Chitin is another NSP, but the chitinolytic  
9 bacteria mainly belong to Firmicutes (Cody, 1989) and Actinobacteria (Beier and Bertilsson, 2013)  
10 phyla, thus furtherly explaining the reduction of Bacteroidetes herein observed.

11 *Actinomyces*, *Bacillus*, *Enterococcus*, *Oceanobacillus*, and *Staphylococcus* resulted to be enriched in  
12 the posterior gut microbiota of the HI-fed rainbow trout of the current research. On the one hand, this  
13 partially reflects the microbiota of the HI-based feeds (characterized by high percentages of  
14 *Actinomyces*, *Enterococcus*, *Oceanobacillus*, and *Staphylococcus*); on the other, these changes can  
15 be attributable to chitin. Indeed, apart from the already mentioned chitin degrading activity of  
16 *Actinomyces* (Beier and Bertilsson, 2013), many *Bacillus* species are chitinolytic (Cody, 1989). As  
17 lactic acid bacteria (LAB), *Enterococcus* is also capable of using chitin as prebiotic (Terova et al.,  
18 2019), while novel chitinolytic *Staphylococcus* species have recently been characterized (Gürkök and  
19 Görmez, 2016). In agreement with the findings herein observed, a significant increase in *Actinomyces*,  
20 *Enterococcus* (Terova et al., 2019), *Staphylococcus* (Bruni et al., 2018) and *Bacillus* (Rimoldi et al.,  
21 2021) has also been reported in rainbow trout fed diets containing HI meal. These changes can be  
22 beneficial for the health status of the fish gut, as bacterial fermentation of chitin leads to short-chain  
23 fatty acids (SCFAs) production (Borrelli et al., 2017; Yu et al., 2019). Indeed, SCFAs (such as butyric,  
24 propionic and acetic acids) act as energy source, promote the proliferation of intestinal epithelial cells,  
25 exert the antimicrobial activity by lowering intestinal pH, modulate the composition of intestinal  
26 microbiota, and enhance the immune response of the fish (Li et al., 2019). In the present study, dietary  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

1 HI meal inclusion also determined a significant reduction of *Clostridium*, *Campylobacter*, *Listeria*,  
2 *Lactobacillus*, *Lactococcus*, *Leuconostoc*, *Pediococcus*, unclassified members (U.m.) of  
3  
4 Peptostreptococceae, *Vagococcus*, and *Weissella* in the fish gut microbiota. The decrease in LAB  
5  
6 such as *Lactobacillus*, *Leuconostoc* and *Pediococcus* – which have been reported to proliferate in HI-  
7  
8 fed rainbow trout (Huyben et al., 2019; Terova et al., 2019; Rimoldi et al., 2021) – seems difficult to  
9  
10 explain, especially because *Enterococcus* (previously described as LAB) was, however, significantly  
11  
12 enriched. This discrepancy may be caused by the different HI meal adopted (prepupae [Terova et al.,  
13  
14 2019] vs larvae), but the capability of insects to stimulate the growth of some LAB at the expense of  
15  
16 others deserves future investigations. The reduction of *Clostridium* could not represent a relevant  
17  
18 finding, since this taxon is characteristic of the intestinal microbiota from endotherms (Eckburg et  
19  
20 al., 2005) and is involved in the degradation of the cellulolytic fibers (which are not predominant in  
21  
22 diets for carnivorous fish) (Chapagain et al., 2019). A similar consideration can also be made for  
23  
24 Peptostreptococcaceae family, whose members exert the generic function of utilizing proteinaceous  
25  
26 substrates and carbohydrates (Fu et al., 2019). On the contrary, the decrease in *Weissella* may  
27  
28 represent a potential challenging outcome, as this genus includes probiotic bacteria (Kühlwein et al.,  
29  
30 2013) and displays antimicrobial activity against a wide range of microorganisms (Patterson et al.,  
31  
32 2010). However, such reduction could have successfully been compensated by the chitin and the  
33  
34 lauric acid contained in the HI meal, which have been reported to exert antimicrobial activity against  
35  
36 both the Gram-negative (Marono et al., 2017) and the Gram-positive (Skrivanova et al., 2006)  
37  
38 bacteria. As a reasonable consequence, the HI antimicrobial properties may have determined the  
39  
40 decrease in *Lactococcus*, *Vagococcus*, *Campylobacter* and *Listeria*. Indeed, the reduction of  
41  
42 *Lactococcus* and *Vagococcus* – whose distinct species have been related to the development of a  
43  
44 growing number of diseases (Ringø and Gatesoupe, 1998) – can be considered a positive finding, but  
45  
46 the most remarkable HI-related outcome is represented by the decreased proliferation of  
47  
48 *Campylobacter* and *Listeria*. Similarly, to what was already pointed out for *Listeria*, *Campylobacter*  
49  
50 is one of the most common agents of food-borne diseases (Kreling et al., 2020), thus making their  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

1 reduction particularly interesting within a food safety scenario. The reduced percentage of *Listeria*  
2 identified in the HI-based diets could also partially explain its reduction in the gut, but the difference  
3  
4 in the corresponding relative abundances (about 7% vs 0.4%) reasonably suggests an active role of  
5  
6  
7 HI meal as well.  
8  
9

## 10 11 **Conclusions**

12 In conclusion, HI meal can be used in commercial diets for rainbow trout up to high inclusion levels  
13  
14 (320 g/kg as fed) without negatively affecting the growth performance, nutrient digestibility, somatic  
15  
16 indices and histomorphological features of the animals. Therefore, considering that the low FM-diets  
17  
18 are nowadays the most adopted fish feeds from a sustainability perspective, the possibility of  
19  
20 including either low or high inclusion levels of HI meal without incurring in adverse outcomes  
21  
22 represents a promising scenario. Furthermore, a positive modulation of the gut microbiota in terms  
23  
24 of selection of SCFAs-producing bacteria and reduction of foodborne disease-causing pathogens was  
25  
26 herein observed for the first time when rainbow trout were administered with low FM-diets containing  
27  
28 HI meal. In the light of such positive findings, future investigations also assessing the gut  
29  
30 metagenome and metabolome are mandatory in order to fully characterize the HI way of action in the  
31  
32 fish gut.  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42

## 43 **Author statement**

44 **Ilaria Biasato:** conduct the experiment, sampling, statistical analysis, and writing the initial draft,  
45  
46

47 **Giulia Chemello:** conduct the experiment, sampling, statistical analysis, and writing the initial draft,  
48  
49

50 **Sara Bellezza Oddon:** fish feeding, sampling, and reviewing the final draft,  
51  
52

53 **Ilario Ferrocino:** feed and gut microbiota analyses, and reviewing the final draft,  
54  
55

56 **Christian Caimi:** fish feeding, sampling, and reviewing the final draft,  
57  
58

59 **Andrea Resconi:** fish feeding, sampling, and reviewing the final draft,  
60  
61

62 **Aman Paul:** feed production and reviewing the final draft,  
63  
64  
65

1 **Michel van Spankeren:** feed production and reviewing the final draft,

2 **Maria Teresa Capucchio:** histomorphological analysis and reviewing the final draft,

3  
4 **Elena Colombino:** histomorphological analysis and reviewing the final draft,

5  
6  
7 **Luca Cocolin:** feed and gut microbiota analyses, and reviewing the final draft,

8  
9 **Francesco Gai:** planning the research activity and reviewing the final draft,

10  
11 **Achille Schiavone:** planning the research activity and reviewing the final draft,

12  
13  
14 **Laura Gasco:** coordination, funding acquisition, planning the research activity, and reviewing the  
15  
16 final draft.

## 17 18 19 20 21 **Funding**

22  
23  
24 The research was supported by the University of Turin (2019) and by The Protix BV, Dongen, The  
25  
26 Netherlands.

## 27 28 29 30 31 **Acknowledgements**

32  
33  
34 The authors are grateful to Mr. Dario Sola for the fish care and the technical support.

## 35 36 37 38 39 **Declaration of Competing Interest**

40  
41 There are no competing financial, professional, or personal interests that might have influenced the  
42  
43 presentation of the work described in this manuscript.

## 44 45 46 47 48 **References**

- 49  
50  
51 1) All About Feed, 2021. Insects in feed: Sector ready to upscale.  
52  
53 <https://www.allaboutfeed.net/all-about/new-proteins/insects-in-feed-sector-ready-to-upscale/>  
54  
55 (accessed 15 October 2021).  
56  
57  
58 2) Meticulous Market Research Pvt. Ltd, 2019. Edible Insects Market by Product Type (Whole  
59  
60 Insect, Insect Powder, Insect Meal, Insect Type (Cricket, Black Soldier Fly, Mealworms),  
61  
62  
63  
64  
65

1 Application (Animal Feed, Protein Bar and Shakes, Bakery, Confectionery, Beverages) -  
2 Global Forecast to 2030.  
3

- 4  
5 3) European Commission, 2017. Commission Regulation (EU) 2017/893 of 24 May 2017  
6 amending Annexes I and IV to Regulation (EC) No 999/2001 of the European Parliament and  
7 of the Council and Annexes X, XIV and XV to Commission Regulation (EU) No 142/2011  
8 as regards the provisions on processed animal protein. Official J. L138:92–116.  
9  
10  
11  
12  
13  
14 4) European Commission, 2021. Commission Regulation (EU) 2021/1372 of 17 August 2021  
15 amending Annexes IV to Regulation (EC) No 999/2001 of the European Parliament and of  
16 the Council as regards the prohibition to feed non-ruminant farmed animals, other than fur  
17 animals, with protein derived from animals. Official J. L295: 1-17.  
18  
19  
20  
21  
22  
23  
24 5) Ojha, S., Bußler, S., Schlüter, OK., 2020. Food waste valorisation and circular economy  
25 concepts in insect production and processing. Waste Manag. 118, 600-609. doi:  
26 10.1016/j.wasman.2020.09.010.  
27  
28  
29  
30  
31 6) Sheppard, D.C., Newton, G.L., Thompson, S.A., Savage, S., 1994. A value added manure  
32 management system using the black soldier fly. Bioresour. Technol. 50, 275-279. doi:  
33 10.1016/0960-8524(94)90102-3.  
34  
35  
36  
37  
38 7) Rawski, M., Mazurkiewicz, J., Kierończyk, B., Józefiak, D., 2020. Black Soldier Fly Full-Fat  
39 Larvae Meal as an Alternative to Fish Meal and Fish Oil in Siberian Sturgeon Nutrition: The  
40 Effects on Physical Properties of the Feed, Animal Growth Performance, and Feed  
41 Acceptance and Utilization. Animals. 10, 2119. doi: 10.3390/ani10112119.  
42  
43  
44  
45  
46  
47  
48 8) Józefiak, A., Nogales-Mérida, S., Rawski, M., Kierończyk, B., Mazurkiewicz, J., 2019.  
49 Effects of insect diets on the gastrointestinal tract health and growth performance of Siberian  
50 sturgeon (*Acipenser baerii* Brandt, 1869). BMC Vet. Res. 15, 348. doi: 10.1186/s12917-019-  
51 2070-y.  
52  
53  
54  
55  
56  
57  
58 9) Caimi, C., Gasco, L., Biasato, I., Malfatto, V., Varello, K., Prearo, M., Pastorino, P., Bona,  
59 M. C., Francese, D. F., Schiavone, A., Elia, A.C., Dörr, A. J. M., Gai, F., 2020. Could dietary  
60  
61  
62  
63  
64  
65



- 1 black soldier fly meal inclusion affect the liver and intestinal histological traits and the  
2 oxidative stress biomarkers of Siberian sturgeon (*Acipenser baerii*) juveniles? *Animals*. 10,  
3 155. doi: 10.3390/ani10010155.
- 4  
5  
6  
7 10) Renna, M., Schiavone, A., Gai, F., Dabbou, S., Lussiana, C., Malfatto, V., Prearo, M.,  
8 Capucchio, M.T., Biasato, I., Biasibetti, E., De Marco, M., Brugiapaglia, A., Zoccarato, I.,  
9 Gasco, L., 2017. Evaluation of the suitability of a partially defatted black soldier fly (*Hermetia*  
10 *illucens* L.) larvae meal as ingredient for rainbow trout (*Oncorhynchus mykiss* Walbaum)  
11 diets. *J. Anim. Sci. Biotechnol.* 8, 57. doi: 10.1186/s40104-017-0191-3.  
12  
13  
14  
15  
16  
17  
18  
19 11) Cappellozza, S., Leonardi, M.G., Savoldelli, S., Carminati, D., Rizzolo, A., Cortellino, G.,  
20 Terova, G., Moretto, E., Badaile, A., Concheri, G., Saviane, A., Bruno, D., Bonelli, M, Caccia,  
21 S., Casartelli, M., Tettamanti, G., 2019. A First Attempt to Produce Proteins from Insects by  
22 Means of a Circular Economy. *Animals*. 9, 278. doi: 10.3390/ani9050278.  
23  
24  
25  
26  
27  
28  
29 12) Cardinaletti, G., Randazzo, B., Messina, M., Zarantoniello, M., Giorgini, E., Zimbelli, A.,  
30 Bruni, L., Parisi, G., Olivotto, I., and Tulli, F. (2019). Effects of Graded Dietary Inclusion  
31 Level of Full-Fat *Hermetia illucens* Prepupae Meal in Practical Diets for Rainbow Trout  
32 (*Oncorhynchus mykiss*). *Animals*. 9, 1-15. doi: 10.3390/ani9050251.  
33  
34  
35  
36  
37  
38  
39 13) St-Hilaire, S., Sheppard, C., Tomberlin, J.K., Irving, S., Newton, L., McGuire, M.A., Mosley,  
40 E.E., Hady, R., Sealey, W., 2007. Fly prepupae as a feedstuff for rainbow trout, *Oncorhynchus*  
41 *mykiss*. *J. World. Aquacult. Soc.* 38, 59–67. doi: 10.1111/j.1749-7345.2006.00073.x.  
42  
43  
44  
45  
46 14) Sealey, W.M., Gaylord, T.G., Barrows, F.T., Tomberlin, J.K., McGuire, M.A., Ross, C., St-  
47 Hilaire, S., 2011. Sensory analysis of rainbow trout, *Oncorhynchus mykiss*, fed enriched black  
48 soldier fly prepupae, *Hermetia illucens*. *J. World. Aquacult. Soc.* 42, 34–45. doi:  
49 10.1111/j.1749-7345.2010.00441.x.  
50  
51  
52  
53  
54  
55 15) Dumas, A., Raggi, T., Barkhouse, J., Lewis, E., and Weltzien, E, 2018. The oil fraction and  
56 partially defatted meal of black soldier fly larvae (*Hermetia illucens*) affect differently growth  
57 performance, feed efficiency, nutrient deposition, blood glucose and lipid digestibility of  
58  
59  
60  
61  
62  
63  
64  
65

- rainbow trout (*Oncorhynchus mykiss*). *Aquaculture*. 492, 24-34. doi:  
10.1016/j.aquaculture.2018.03.038.
- 16) Bruni, L., Pastorelli, R., Viti, C., Gasco, L., Parisi, G., 2018. Characterisation of the intestinal microbial communities of rainbow trout (*Oncorhynchus mykiss*) fed with *Hermetia illucens* (black soldier fly) partially defatted larva meal as partial dietary protein source. *Aquaculture*. 487, 56-63. doi: 10.1016/j.aquaculture.2018.01.006.
- 17) Huyben, D., Vidaković, A., Werner Hallgren, S., Langeland, M., 2019. High-throughput sequencing of gut microbiota in rainbow trout (*Oncorhynchus mykiss*) fed larval and pre-pupae stages of black soldier fly (*Hermetia illucens*). *Aquaculture*. 500, 485-491. doi: 10.1016/j.aquaculture.2018.10.034.
- 18) Rimoldi, S., Gini, E., Iannini, F., Gasco, L., Terova, G., 2019. The Effects of Dietary Insect Meal from *Hermetia illucens* Prepupae on Autochthonous Gut Microbiota of Rainbow Trout (*Oncorhynchus mykiss*). *Animals*. 9, 143. doi: 10.3390/ani9040143.
- 19) Terova, G., Rimoldi, S., Ascione, C., Gini, E., Ceccotti, C. Gasco, L., 2019. Rainbow trout (*Oncorhynchus mykiss*) gut microbiota is modulated by insect meal from *Hermetia illucens* prepupae in the diet. *Rev. Fish. Biol. Fish.* 29, 465–486. doi: 10.1007/s11160-019-09558-y.
- 20) Caimi, C., Biasato, I., Chemello, G., Bellezza Oddon, S., Lussiana, C., Malfatto, V.M., Capucchio, M.T., Colombino, E., Schiavone, A., Gai, F., Trocino, A., Brugiapaglia, A., Renna, M., Gasco, L., 2021. Dietary inclusion of a partially defatted black soldier fly (*Hermetia illucens*) larva meal in low fishmeal-based diets for rainbow trout (*Oncorhynchus mykiss*). *J. Anim. Sci. Biotechnol.* 12, 50. doi: 10.1186/s40104-021-00575-1.
- 21) AOAC International. 2000. Official methods of analysis of AOAC International. 16<sup>th</sup> ed. Gaithersburg: Association of Official Analytical Chemists.
- 22) AOAC International, 2003. Official methods of analysis of AOAC International. 17<sup>th</sup> ed. Gaithersburg: Association of Official Analytical Chemists.

- 1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65
- 23) Van Soest, P.J., Robertson, J.B., Lewis, B.A., 1991. Methods for dietary fiber, neutral detergent fiber, and nonstarch polysaccharides in relation to animal nutrition. *J. Dairy. Sci.* 74, 3583–97. Doi: 10.3168/jds.S0022-0302(91)78551-2.
- 24) Schmid, A., Collomb, M., Hadorn, R.. 2009. Fatty acid composition of Swiss cooked sausages. *Fleischwirtschaft.* 89, 101–4.
- 25) Renna, M., Gasmi-Boubaker, A., Lussiana, C., Battaglini, L.M., Belfayez, K., Fortina, R., 2014. Fatty acid composition of the seed oils of selected *Vicia L.* taxa from Tunisia. *Ital. J. Anim. Sci.* 13, 308–16. doi: 10.4081/ijas.2014.3193.
- 26) Chemello, G., Renna, M., Caimi, C., Guerreiro, I., Oliva-Teles, A., Enes, P., Biasato, I., Schiavone, A., Gai, F., Gasco, L., 2020. Partially Defatted *Tenebrio molitor* Larva Meal in Diets for Grow-Out Rainbow Trout, *Oncorhynchus mykiss* (Walbaum): Effects on Growth Performance, Diet Digestibility and Metabolic Responses. *Animals.* 10, 229. doi: 10.3390/ani10020229.
- 27) Biasato, I., Renna, M., Gai, F., Dabbou, S., Meneguz, M., Perona, G., Martinez, S., Lajusticia, A.C.B., Bergagna, S., Sardi, L., Capucchio, M.T., Bressan, E., Dama, A., Schiavone, A., Gasco, L., 2019. Partially defatted black soldier fly larva meal inclusion in piglet diets: effects on the growth performance, nutrient digestibility, blood profile, gut morphology and histological features. *J. Anim. Sci. Biotechnol.* 10, 12. doi: 10.1186/s40104-019-0325-x.
- 28) Magoč, T. Salzberg, S.L., 2011. FLASH: Fast length adjustment of short reads to improve genome assemblies. *Bioinformatics.* 27, 2957–2963. doi: 10.1093/bioinformatics/btr507.
- 29) Caporaso, J.G., Kuczynski, J., Stombaugh, J., Bittinger, K., Bushman, F.D, Costello, E.K., Fierer, N., Peña, A.G., Goodrich, J.K., Gordon, J.I., et al. QIIME allows analysis of high-throughput community sequencing data. *Nat. Methods.* 7, 335–336. doi: 10.1038/nmeth.f.303.
- 30) Biasato, I., Ferrocino, I., Biasibetti, E., Grego, E., Dabbou, S., Sereno, A., Gai, F., Gasco, L., Schiavone, A., Cocolin, L., Capucchio, M.T., 2018. Modulation of intestinal microbiota,

- morphology and mucin composition by dietary insect meal inclusion in free-range chickens. BMC Vet. Res. 14, 383. doi: 10.1186/s12917-018-1690-y.
- 31) Dixon, P., 2003. VEGAN, a package of R functions for community ecology. J. Veg. Sci. 14, 927–930. doi: 10.1111/j.1654-1103.2003.tb02228.x.
- 32) Henry, M., Gasco, L., Piccolo, G., Fountoulaki, E., 2015. Review on the use of insects in the diet of farmed fish: past and future. Anim. Feed. Sci. Technol. 203, 1–22. doi: 10.1016/j.anifeedsci.2015.03.001.
- 33) Muddasir, J., Imtiaz, A., 2016. Length weight relationship and condition factor of snow trout, *Schizothorax plagiostomus* (Heckel, 1838) from Lidder River. Kashmir. Int. J. Fish. Aquat. Stud. 4, 131–6.
- 34) Dernekbaşı, S., 2021. Digestibility and liver fatty acid composition of rainbow trout (*Oncorhynchus mykiss*) fed by graded levels of canola oil. Turk. J. Fish. Aquat. Sci. 12, 105–13. doi: 10.4194/1303-2712-v12\_1\_13.
- 35) Khojasteh, S.M.B., Sheikhzadeh, F., Mohammadnejad, D., Azami, A., 2009. Histological, histochemical and ultrastructural study of the intestine of rainbow trout (*Oncorhynchus mykiss*). World. Appl. Sci. J. 6, 1525–1531.
- 36) Verdile. N., Pasquariello, R., Scolari, M., Sciré, G., Brevini, T.A.L., Gandolfi, F., 2020. A detailed study of rainbow trout (*Oncorhynchus mykiss*) intestine revealed that digestive and absorptive functions are not linearly distributed along its length. Animals. 10, 745. doi: 10.3390/ani10040745.
- 37) Elia, A.C., Capucchio, M.T., Caldaroni, B., Magara, G., Dörr, A.J.M., Biasato, I., Biasibetti, E., Righetti, M., Pastorino, P., Prearo, M., Gai, F., Schiavone, A., Gasco, L., 2018. Influence of *Hermetia illucens* meal dietary inclusion on the histological traits, gut mucin composition and the oxidative stress biomarkers in rainbow trout (*Oncorhynchus mykiss*). Aquaculture, 496, 50-57. 10.1016/j.aquaculture.2018.07.009.

- 1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65
- 38) Salas-Leiva, J., Mazón-Suástegui, J.M., Teles, A., Tovar-Ramírez, D., 2020. Structure and predictive metabolic contribution of intestinal microbiota of Longfin yellowtail (*Seriola rivoliana*) juveniles in aquaculture systems. *Mol. Biol. Rep.* 47, 9627-9636. doi: 10.1007/s11033-020-05970-x.
- 39) Jiang, M., Xu, M., Ying, C., Yin, D., Dai, P., Yang, Y., Ye, K., Liu, K., 2019. The intestinal microbiota of lake anchovy varies according to sex, body size, and local habitat in Taihu Lake, China. *Microbiologyopen*. 9, e00955. doi: 10.1002/mbo3.955.
- 40) Zeng, A., Tan, K., Gong, P., Lei, P., Guo, Z., Wang, S., Gao, S., Zhou, Y., Shu, Y., Zhou, X., Miao, D., Zeng, F., Liu, H., 2020. Correlation of microbiota in the gut of fish species and water. *3 Biotech*. 10, 472. doi: 10.1007/s13205-020-02461-5.
- 41) Korlević, M., Šupraha, L., Ljubešić, Z., Henderiks, J., Ciglencčki, I., Dautović, J., Orlić, S., 2016. Bacterial diversity across a highly stratified ecosystem: A salt-wedge Mediterranean estuary. *Syst. Appl. Microbiol.* 39, 398-408. doi: 10.1016/j.syapm.2016.06.006.
- 42) Quéméneur, M., Bel Hassen, M., Armougom, F., Khammeri, Y., Lajnef, R., Bellaaj-Zouari, A., 2020. Prokaryotic Diversity and Distribution Along Physical and Nutrient Gradients in the Tunisian Coastal Waters (South Mediterranean Sea). *Front. Microbiol.* 11, 593540. doi: 10.3389/fmicb.2020.593540.
- 43) Zhang, Z., Fan, X., Peijnenburg, W.J.G.M., Zhang, M., Sun, L., Zhai, Y., Yu, Q., Wu, J., Lu, T., Qian, H., 2021. Alteration of dominant cyanobacteria in different bloom periods caused by abiotic factors and species interactions. *J. Environ. Sci.* 99, 1-9. doi: 10.1016/j.jes.2020.06.001.
- 44) Péron, G., Mittaine, J-F., Le Gallic, B., 2010. Where do fishmeal and fish oil products come from? An analysis of the conversion ratios in the global fishmeal industry. *Marine Policy* 34, 815–820.
- 45) Butt, R.L., Volkoff, H., 2019. Gut Microbiota and Energy Homeostasis in Fish. *Front. Endocrinol. (Lausanne)*. 10, 9. doi: 10.3389/fendo.2019.00009.

- 1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65
- 46) Tarnecki, A.M., Burgos, F.A., Ray, C.L., Arias, C.R., 2017. Fish intestinal microbiome: diversity and symbiosis unravelled by metagenomics. *J. Appl. Microbiol.* 123, 2-17. doi: 10.1111/jam.13415.
- 47) Huyben, D., Rimoldi, S., Ceccotti, C., Montero, D., Betancor, M., Iannini, F., Terova, G., 2020. Effect of dietary oil from *Camelina sativa* on the growth performance, fillet fatty acid profile and gut microbiome of gilthead Sea bream (*Sparus aurata*). *PeerJ.* 8, e10430. doi: 10.7717/peerj.10430.
- 48) Yu, J., Wang, Y., Xiao, Y., Li, X., Xu, X., Zhao, H., Wu, L., Li, J., 2021. Effects of chronic nitrate exposure on the intestinal morphology, immune status, barrier function, and microbiota of juvenile turbot (*Scophthalmus maximus*). *Ecotoxicol. Environ. Saf.* 207, 111287. doi: 10.1016/j.ecoenv.2020.111287.
- 49) Huang, Q., Sham, R.C., Deng, Y., Mao, Y., Wang, C., Zhang, T., Leung, K.M.Y., 2020. Diversity of gut microbiomes in marine fishes is shaped by host-related factors. *Mol. Ecol.* 29, 5019-5034. doi: 10.1111/mec.15699.
- 50) Buchanan, R.L., Gorris, L.G.M., Hayman, M.M., Jackson, T.C., Whiting, R.C., 2017. A review of *Listeria monocytogenes*: An update on outbreaks, virulence, dose-response, ecology, and risk assessments. *Food. Control.* 75, 1-13. doi: 10.1016/j.foodcont.2016.12.016.
- 51) Basha, K.A., Kumar, N.R., Das, V., Reshmi, K., Rao, B.M., Lalitha, K.V., Joseph, T.C., 2019. Prevalence, molecular characterization, genetic heterogeneity and antimicrobial resistance of *Listeria monocytogenes* associated with fish and fishery environment in Kerala, India. *Lett. Appl. Microbiol.* 69, 286-293. doi: 10.1111/lam.13205.
- 52) Rimoldi, S., Terova, G., Ascione, C., Giannico, R., Brambilla, F., 2018. Next generation sequencing for gut microbiome characterization in rainbow trout (*Oncorhynchus mykiss*) fed animal by-product meals as an alternative to fishmeal protein sources. *PLoS One.* 13, e0193652. doi: 10.1371/journal.pone.0193652.

- 1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65
- 53) Rimoldi, S., Antonini, M., Gasco, L., Moroni, F., Terova, G., 2021. Intestinal microbial communities of rainbow trout (*Oncorhynchus mykiss*) may be improved by feeding a *Hermetia illucens* meal/low-fishmeal diet. *Fish. Physiol. Biochem.* 47, 365-380. doi: 10.1007/s10695-020-00918-1.
- 54) Desai, A.R., Links, M.G., Collins, S.A., Mansfield, G.S., Drew, M.D., Van Kessel, A.G., Hill, J.E., 2012. Effects of plant-based diets on the distal gut microbiome of rainbow trout (*Oncorhynchus mykiss*). *Aquaculture.* 350–353, 134–142. doi: 10.1016/j.aquaculture.2012.04.005.
- 55) Wong, S., Waldrop, T., Summerfelt, S., Davidson, J., Barrows, F., Kenney, P.B., Welch, T., Wiens, G.D., Snekvik, K., Rawls, J.F., Good, C., 2013. Aquacultured rainbow trout (*Oncorhynchus mykiss*) possess a large core intestinal microbiota that is resistant to variation in diet and rearing density. *Appl. Environ. Microbiol.* 79, 4974–4984. <https://doi.org/10.1128/AEM.00924-13>.
- 56) Ingerslev, H-C., von Gersdorff Jørgensen, L., Strube, M.L., Larsen, N., Dalsgaard, I., Boye, M., Madsen, L., 2014. The development of the gut microbiota in rainbow trout (*Oncorhynchus mykiss*) is affected by first feeding and diet type. *Aquaculture.* 424–425, 24–34. doi: 10.1016/j.aquaculture.2013.12.032.
- 57) Pelusio, N.F., Rossi, B., Parma, L., Volpe, E., Ciulli, S., Piva, A., D'Amico, F., Scicchitano, D., Candela, M., Gatta, P.P., Bonaldo, A., Grilli, E., 2020. Effects of increasing dietary level of organic acids and nature-identical compounds on growth, intestinal cytokine gene expression and gut microbiota of rainbow trout (*Oncorhynchus mykiss*) reared at normal and high temperature. *Fish. Shellfish. Immunol.* 107, 324-335. doi: 10.1016/j.fsi.2020.10.021.
- 58) Beier, S., Bertilsson, S., 2013. Bacterial chitin degradation-mechanisms and ecophysiological strategies. *Front. Microbiol.* 4, 149. doi: 10.3389/fmicb.2013.00149.

- 1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65
- 59) den Besten, G., van Eunen, K., Groen, A.K., Venema, K., Reijngoud, D.J., Bakker, B.M., 2013. The role of short-chain fatty acids in the interplay between diet, gut microbiota, and host energy metabolism. *J. Lipid. Res.* 54, 2325-40. doi: 10.1194/jlr.R036012.
- 60) Cody, R.M., 1989. Distribution of chitinase and chitobiase in bacillus. *Current. Microbiology* 19, 201–205. <https://doi.org/10.1007/BF01570162>.
- 61) Gürkök, S., Görmez. A., 2016. Isolation and characterization of novel chitinolytic bacteria. *AIP Conference Proceedings* 1726, 020017. <https://doi.org/10.1063/1.4945843>.
- 62) Borrelli, L., Coretti, L., Dipineto, L., Bovera, F., Menna, F., Chiariotti, L., Nizza, A., Lembo, F., Fioretti, A., 2017. Insect-based diet, a promising nutritional source, modulates gut microbiota composition and SCFAs production in laying hens. *Sci. Rep.* 7, 16269. doi: 10.1038/s41598-017-16560-6.
- 63) Yu, M., Li, Z., Chen, W., Rong, T., Wang, G., Ma, X.J., 2019. *Hermetia illucens* larvae as a potential dietary protein source altered the microbiota and modulated mucosal immune status in the colon of finishing pigs. *J. Anim. Sci Biotechnol.* 10, 50. doi: 10.1186/s40104-019-0358-1.
- 64) Li, Z., Tran, N.T., Ji, P., Sun, Z., Wen, X., Li, S., 2019. Effects of prebiotic mixtures on growth performance, intestinal microbiota and immune response in juvenile chu's croaker, *Nibea coibor*. *Fish. Shellfish. Immunol.* 89, 564-573. doi: 10.1016/j.fsi.2019.04.025.
- 65) Eckburg, P.B., Bik, E.M., Bernstein, C.N., Purdom, E., Dethlefsen, L., Sargent, M., Gill, S.R., Nelson, K.E., Relman, D.A., 2005. Diversity of the human intestinal microbial flora. *Science.* 308, 1635-8. doi: 10.1126/science.1110591.
- 66) Chapagain, P., Arivett, B., Cleveland, B.M., Walker, D.M., Salem, M., 2019. Analysis of the fecal microbiota of fast- and slow-growing rainbow trout (*Oncorhynchus mykiss*). *BMC Genomics.* 20, 788. doi: 10.1186/s12864-019-6175-2.



- 1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65
- 67) Fu, P.P., Xiong, F., Feng, W.W., Zou, H., Wu, S.G., Li, M., Wang, G.T., Li, W.X., 2019. Effect of intestinal tapeworms on the gut microbiota of the common carp, *Cyprinus carpio*. Parasit Vectors. 12, 252. doi: 10.1186/s13071-019-3510-z.
- 68) Kühlwein, H., Emery, M.J., Rawling, M.D., Harper, G.M., Merrifield, D.L., Davies, S.J., 2013. Effects of a dietary  $\beta$ -(1,3)(1,6)-D-glucan supplementation on intestinal microbial communities and intestinal ultrastructure of mirror carp (*Cyprinus carpio* L.). J. Appl. Microbiol. 115, 1091-106. doi: 10.1111/jam.12313.
- 69) Patterson, M.F., McKay, A.M., Connolly, M., Linton, M., 2010. Effect of high pressure on the microbiological quality of cooked chicken during storage at normal and abuse refrigeration temperatures. Food. Microbiol. 27, 266-73. doi: 10.1016/j.fm.2009.10.007.
- 70) Marono, S., Loponte, R., Lombardi, P., Vassalotti, G., Pero, M.E., Russo, F., Gasco, L., Parisi, G., Piccolo, G., Nizza, S., Di Meo, C., Attia, Y.A., Bovera, F., 2017. Productive performance and blood profiles of laying hens fed *Hermetia illucens* larvae meal as total replacement of soybean meal from 24 to 45 weeks of age. Poult. Sci. 96, 1783-1790. doi: 10.3382/ps/pew461.
- 71) Skrivanova, E., Marounek, M., Benda, V., Brezina, P., 2006. Susceptibility of *Escherichia coli*, *Salmonella* sp., and *Clostridium perfringens* to organic acids and monolaurin. Veterinarni Medicina 51, 81–88.
- 72) Ringø, E., Gatesoupe, F.-J., 1998. Lactic acid bacteria in fish: a review, Aquaculture. 160, 177-203. doi: 10.1016/S0044-8486(97)00299-8.
- 73) Kreling, V., Falcone, F.H., Kehrenberg, C., Hensel, A., 2020. *Campylobacter* sp.: Pathogenicity factors and prevention methods-new molecular targets for innovative antivirulence drugs? Appl. Microbiol. Biotechnol. 104, 10409-10436. doi: 10.1007/s00253-020-10974-5.

**Table 1.** Feed ingredients and proximate composition of the experimental diets.

	HI meal	HI0	HI25	HI50	HI100
Ingredients, g/kg as fed					
Fish meal		206	154.50	103	0
Soybean protein concentrate		150	150	150	150
Wheat gluten meal		100	100	100	100
Corn gluten meal		70	70	70	70
Soybean meal		40	40	40	40
Wheat meal		240.50	218.23	195.95	151.40
HI meal		0	80	160	320
Fish oil		50	50	50	50
Soybean oil		123.50	111.38	99.25	75
Vit. min. premix (1%)		10	10	10	10
DL methionine		0	0.28	0.55	1.10
L-lysine HCL		0	0.60	1.20	2.40
Diamol		10	10	10	10
Lime fine		0	1.62	3.25	6.50
Monocalcium phosphate		0	2	4	8
Salt		0	1.63	3.25	5
Magnesium oxide		0	0.15	0.30	0.60
Proximate composition <sup>a</sup>					
DM, g/100g	96.24	93.93	94.81	94.41	96.01
CP, g/100g DM <sup>b</sup>	51.71	43.75	44.01	44.84	45.88
EE, g/100g DM	20.43	19.43	19.23	18.04	17.85
Ash, g/100g DM	5.65	6.85	6.65	6.47	6.71
NDF, g/100g DM	N.A.	20.89	25.08	8.26	9.91
ADF, g/100g DM	N.A.	1.63	2.21	3.08	3.88
NFE, g/100g DM <sup>c</sup>	N.A.	23.90	24.92	25.06	25.58
GE, MJ/Kg <sup>c</sup>	22.04	22.23	22.33	22.08	22.47

Abbreviations: HI0, control diet; HI25, 25% of FM replaced by *Hermetia illucens* meal; HI50, 50% of FM replaced by *Hermetia illucens* meal; HI100, 100% of FM replaced by *Hermetia illucens* meal; DM, dry matter; CP, crude protein; EE, ether extract; ADF, acid detergent fiber; ADF<sub>n</sub>, acid detergent fiber nitrogen; NDF, neutral detergent fiber; NFE, nitrogen-free extract. <sup>a</sup>Values are reported as mean of duplicate analyses; <sup>b</sup>Conversion factors of 5.62 for the HI meal (Janssen et al., 2017) and 6.25 for the experimental diets; <sup>c</sup>Calculated as  $100 - [(100 - DM) + CP + EE + Ash]$ ; <sup>d</sup>Determined by calorimetric bomb.

**Table 2.** Fatty acid (FA) composition of the experimental diets.

	<b>HI0</b>	<b>HI25</b>	<b>HI50</b>	<b>HI100</b>
Fatty acids, g/100 g DM of TFA				
C10:0	0.00	5.90	12.32	14.87
C12:0	7.54	363.27	756.46	1298.35
C14:0	198.88	273.55	352.25	468.21
C15:0 iso	7.37	6.87	6.75	5.61
C15:0 anteiso	9.15	8.47	10.01	8.36
C14:1 c + C15:0	24.06	25.71	26.49	27.19
C16:0 iso	4.41	4.66	4.21	4.07
C16:0	1480.80	1570.42	1627.03	1663.50
C17:0 iso	20.61	19.82	19.09	15.73
C17:0 anteiso	19.35	17.37	20.76	18.41
C16:1 c	226.49	244.14	269.18	289.74
C17:1 c9	19.94	20.02	20.43	19.80
C18:0	486.70	491.09	476.63	443.73
C18:1 t	35.64	34.58	30.05	27.40
C18:1 c9	5186.36	5028.82	4851.09	4163.97
C18:1 c11	248.71	238.81	231.19	194.62
C18:1 c12	5.71	4.15	5.62	2.26
C18:1 c14 + t 16	18.77	13.95	15.35	9.80
C18:2 n6	6284.27	6029.43	5726.58	4698.82
C20:0	37.68	41.45	38.80	37.03
C18:3 n6	9.76	8.59	7.73	8.39
C20:1 c9	41.31	40.71	35.08	28.43
C20:1 c11	308.66	303.57	282.93	245.98
C18:3 n3	245.42	259.18	271.83	280.86
C20:2 n6	72.29	69.68	66.22	56.73
C18:4 n3	75.04	75.04	66.59	56.24
C22:0	9.60	10.09	10.44	9.19
C22:1 n9	322.80	290.28	270.81	227.25
C20:3 n6	41.46	39.34	35.67	30.64
C20:4 n6	24.29	22.58	19.59	12.26
C20:5 n3	295.22	279.60	251.54	193.51
C22:5 n3	62.39	60.51	59.32	52.80
C22:6 n3	235.30	192.50	175.99	132.08
Σ SFA	2371.59	2903.64	3417.38	4061.31
Σ MUFA	6414.39	6219.03	6011.73	5209.25
Σ PUFA	7279.99	6971.50	6624.90	5475.28
Σ n3	879.80	831.12	794.35	689.89
Σ n6	6405.91	6144.52	5836.18	4787.65
Σ n6/ Σ n3	7.28	7.39	7.35	6.94
TFA	16065.97	16094.16	16054.02	14745.84

Abbreviations: HI0, control diet; HI25, 25% of FM replaced by *Hermetia illucens* meal; HI50, 50% of FM replaced by *Hermetia illucens* meal; HI100, 100% of FM replaced by *Hermetia illucens* meal; c, cis; t, trans; SFA, saturated fatty acids; MUFA, monounsaturated fatty acids; PUFA, polyunsaturated fatty acids; TFA, total fatty acids.

**Table 3.** Apparent digestibility coefficients of dry matter, protein, ether extract and gross energy of the rainbow trout (n=4)

	HI0	HI25	HI50	HI100	SEM	P-value
<b>ADC DM (%)</b>	84.54	87.50	84.71	84.67	0.67	0.346
<b>ADC CP (%)</b>	95.07	95.85	94.44	94.32	0.25	0.086
<b>ADC EE (%)</b>	98.43	98.73	98.37	98.33	0.08	0.298
<b>ADC GE (%)</b>	92.26	93.10	91.25	90.84	0.41	0.192

Abbreviations: HI0, control diet; HI25, 25% of FM replaced by *Hermetia illucens* meal; HI50, 50% of FM replaced by *Hermetia illucens* meal; HI100, 100% of FM replaced by *Hermetia illucens* meal; SEM, standard error of the mean; P, probability; ADC, apparent digestibility coefficient; DM, dry matter; CP, crude protein; EE, ether extract; GE, gross energy.

**Table 4.** Survival and growth performance of the rainbow trout (n = 3).

	HI0	HI25	HI50	HI100	SEM	P-value
<b>Survival (%)</b>	96.00	95.33	97.33	96.67	0.48	0.557
<b>IBW (g)</b>	112.73	113.13	112.70	112.93	0.08	0.142
<b>FBW (g)</b>	467.53	463.60	469.37	474.70	3.33	0.756
<b>iWG (g)</b>	354.87	350.49	356.66	361.77	3.33	0.748
<b>FCR</b>	1.72	1.73	1.77	1.75	0.02	0.856
<b>PER</b>	1.33	1.32	1.26	1.25	0.02	0.299
<b>SGR (% day<sup>-1</sup>)</b>	0.90	0.90	0.89	0.91	0.01	0.762

Abbreviations: HI0, control diet; HI25, 25% of FM replaced by *Hermetia illucens* meal; HI50, 50% of FM replaced by *Hermetia illucens* meal; HI100, 100% of FM replaced by *Hermetia illucens* meal; SEM, standard error of the mean; P, probability; iIBW, individual initial body weight; iFBW, individual final body weight; iWG, individual weight gain; SGR, specific growth rate; FCR, feed conversion ratio; PER, protein efficiency ratio.

**Table 5.** Condition factor and somatic indices of the rainbow trout (n = 21).

	<b>HI0</b>	<b>HI25</b>	<b>HI50</b>	<b>HI100</b>	<b>SEM</b>	<b>p-value</b>
<b>K</b>	1.19	1.12	1.16	1.16	0.02	0.548
<b>CY</b>	89.98	88.72	89.17	89.37	0.26	0.392
<b>HSI</b>	1.12	1.08	1.07	1.08	0.03	0.938
<b>VSI</b>	8.42	8.47	8.16	7.88	0.15	0.488
<b>CF</b>	3.52	3.86	3.62	3.26	0.14	0.465

Abbreviations: HI0, control diet; HI25, 25% of FM replaced by *Hermetia illucens* meal; HI50, 50% of FM replaced by *Hermetia illucens* meal; HI100, 100% of FM replaced by *Hermetia illucens* meal; SEM, standard error of the mean; P, probability; K, condition factor; CY, carcass yield; HSI, hepatosomatic index; VSI, viscerosomatic index; CF, coefficient of fatness.

**Table 6.** Intestinal morphometric indices of the rainbow trout (n = 12).

	<b>Diet (D)</b>				<b>Intestinal segment (IS)</b>		<b>SEM</b>		<b>P-value</b>		
	<b>HI0</b>	<b>HI25</b>	<b>HI50</b>	<b>HI100</b>	<b>Anterior</b>	<b>Posterior</b>	<b>D</b>	<b>IS</b>	<b>D</b>	<b>IS</b>	<b>D x IS</b>
<b>Vh (mm)</b>	0.87	0.83	0.80	0.79	0.68 <sup>a</sup>	0.96 <sup>b</sup>	0.33	0.02	0.392	<0.001	0.982

Abbreviations: HI0, control diet; HI25, 25% of FM replaced by *Hermetia illucens* meal; HI50, 50% of FM replaced by *Hermetia illucens* meal; HI100, 100% of FM replaced by *Hermetia illucens* meal; SEM, standard error of the mean; P, probability; Vh, villus height.

Means with different superscript letters (a, b) indicate significant differences.

**Table 7.** Histopathological alterations of the rainbow trout (n = 12).

Variables	Dietary treatments				P-value
	HI0	HI25	HI50	HI100	
<b>Liver n (%)</b>					
Inflammation					0.110
Absent	12 (100)	12 (100)	9 (75)	10 (83.3)	
Mild	0 (0)	0 (0)	3 (25)	2 (16.7)	
Degeneration					0.088
Absent	0 (0)	1 (8.3)	5 (42)	3 (25)	
Mild	5 (41.7)	7 (58.3)	5 (42)	8 (66.7)	
Moderate	6 (50)	4 (33.4)	2 (16.7)	0 (0)	
Severe	1 (8.3)	0 (0)	0 (0)	1 (8.3)	
<b>Spleen n (%)</b>					
White pulp hyperplasia					0.495
Absent	9 (81.8)	10 (90.9)	11 (91.7)	12 (100)	
Mild	2 (18.2)	1 (9.1)	1 (8.3)	0 (0)	
Hemosiderosis					0.347
Absent	3 (27.3)	6 (54.5)	4 (33.3)	7 (58.3)	
Mild	8 (72.7)	5 (45.5)	8 (66.7)	5 (41.7)	
<b>Stomach inflammation n (%)</b>					0.395
Absent	12 (100)	12(100)	12 (100)	10 (83.4)	
Mild	0 (0)	0 (0)	0 (0)	1( 8.3)	
<b>Anterior gut inflammation n (%)</b>					1.00
Absent	9 (75)	9 (75)	9 (75)	9 (75)	
Mild	3 (25)	3 (25)	3 (25)	3 (25)	
<b>Posterior gut inflammation n (%)</b>					0.681
Absent	11 (91.7)	9 (75)	9 (75)	10 (83.3)	
Mild	1(8.3)	3 (25)	3 (25)	2 (16.7)	

Abbreviations: HI0, control diet; HI25, 25% of FM replaced by *Hermetia illucens* meal; HI50, 50% of FM replaced by *Hermetia illucens* meal; HI100, 100% of FM replaced by *Hermetia illucens* meal; SEM, standard error of the mean; P, probability.

## Figure captions

**Figure 1.** Relative abundance of the main bacterial phyla (A) and genera (B) in samples of commercial feeds containing low content of fishmeal (HI0), *Hermetia illucens* meal as replacement of 25% of fishmeal (HI50), *Hermetia illucens* meal as replacement of 50% of fishmeal (HI50), and *Hermetia illucens* meal as replacement of 100% of fishmeal (HI100).

**Figure 2.** Bacterial community alpha diversity in posterior gut samples of rainbow trout fed control (HI10), *Hermetia illucens* meal as replacement of 25% of fishmeal (HI50), *Hermetia illucens* meal as replacement of 50% of fishmeal (HI50), and *Hermetia illucens* meal as replacement of 100% of fishmeal (HI100) diets.

**Figure 3.** Bacterial community composition (weighted UniFrac beta diversity, PCA plots) in posterior gut samples of rainbow trout fed control (HI10), *Hermetia illucens* meal as replacement of 25% of fishmeal (HI50), *Hermetia illucens* meal as replacement of 50% of fishmeal (HI50), and *Hermetia illucens* meal as replacement of 100% of fishmeal (HI100) diets.

**Figure 4.** Relative abundance of the main bacterial phyla (A) and genera (B) in posterior gut samples of rainbow trout fed control (HI10), *Hermetia illucens* meal as replacement of 25% of fishmeal (HI50), *Hermetia illucens* meal as replacement of 50% of fishmeal (HI50), and *Hermetia illucens* meal as replacement of 100% of fishmeal (HI100) diets.

**Figure 5.** Relative abundance at phylum level of differentially abundant OTUs in in posterior gut samples of rainbow trout fed control (HI10), *Hermetia illucens* meal as replacement of 25% of fishmeal (HI50), *Hermetia illucens* meal as replacement of 50% of fishmeal (HI50), and *Hermetia*


1 *illucens* meal as replacement of 100% of fishmeal (HI100) diets. Pairwise Kruskal-Wallis test, FDR  
2 < 0.05.  
3  
4  
5  
6

7 **Figure 6.** Relative abundance at genus level of differentially abundant OTUs in in posterior gut  
8 samples of rainbow trout fed control (HI10), *Hermetia illucens* meal as replacement of 25% of  
9 fishmeal (HI50), *Hermetia illucens* meal as replacement of 50% of fishmeal (HI50), and *Hermetia*  
10 *illucens* meal as replacement of 100% of fishmeal (HI100) diets. Pairwise Kruskal-Wallis test, FDR  
11 < 0.05.  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65




Figure 1A

[Click here to access/download;Figure;Figure 1A.tif](#)










Figure 4A






Figure 5


[Click here to access/download;Figure;Figure 5.tiff](#)

Figure 6

