

AperTO - Archivio Istituzionale Open Access dell'Università di Torino

Trichoderma enriched compost, BCAs and potassium phosphite control Fusarium wilt of lettuce without affecting soil microbiome at genus level

This is a pre print version of the following article:

Original Citation:

Availability:

This version is available <http://hdl.handle.net/2318/1891861> since 2023-02-13T07:19:19Z

Published version:

DOI:10.1016/j.apsoil.2022.104678

Terms of use:

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)

1 **Title**

2 *Trichoderma* enriched compost, BCAs and potassium phosphite control Fusarium wilt of lettuce without
3 affecting soil microbiome at genus level

4

5 **Authors:**

6 Alessio Bellini^{1,2}, Giovanna Gilardi¹, Mohamed Idbella^{3,4}, Maurizio Zotti³, Massimo Pugliese^{1,2*},
7 Giuliano Bonanomi³, and Maria Lodovica Gullino¹

8

9 **Affiliations**

10 ¹*AGROINNOVA – Centre of Competence for the Innovation in the Agro-Environmental Sector,*
11 *University of Turin, Turin, Italy*

12 ²*Agricultural, Forestry and Food Sciences Department (DISAFA), University of Turin, Turin, Italy*

13 ³*Department of Agricultural Sciences, University of Naples Federico II, Portici, Italy*

14 ⁴*Faculty of Sciences and techniques, Hassan II University, Casablanca, Morocco*

15 *Corresponding author: M. Pugliese, E-mail: massimo.pugliese@unito.it

16

17

18

19

20 **Abstract**

21

22 *Fusarium oxysporum* f. sp. *lactucae* (Fol) is the causal agent of Fusarium wilt of lettuce, one of the most
23 troublesome diseases affecting lettuce worldwide. Chemical control strategies are inadequate due to
24 limited fungicide availability and consumer interest in organic vegetable production. Alternative control
25 strategies, such as biological control agents (BCAs), suppressive compost, and resistance inducers, have
26 been intensively studied to test their ability to reduce pathogen attacks. Research has been recently
27 focused on the influence of BCAs on the rhizosphere microbiota, which plays a critical role in soil
28 suppressiveness. In this work, three strategies of integrated pest management (IPM) were tested against
29 Fol attacks in two fields for two consecutive years: (i) a compost enriched with *Trichoderma*, (ii) a
30 combination of *T. gamsii* + *T. asperellum*, *Bacillus amyloliquefaciens* and potassium posphite and (iii) a
31 combination of *T. polysporum* + *T. atroviride*. The rhizosphere microbiota was characterized by high-
32 throughput sequencing of bacterial and eukaryotic rRNA gene markers. Obtained results indicated IPM
33 strategies statistically reduced disease severity, in both fields and years, from 50% to 70% compared to
34 untreated controls. An increased crop yield compared to untreated controls was also observed.
35 Predominant phyla were Proteobacteria, Firmicutes and Actinobacteria for bacteria, and Ascomycota for
36 fungi. However, microbiota populations were not affected by any of the treatments, nor were significant
37 differences observed when the soil microbial community was compared to that of untreated controls.
38 Conversely, large differences were observed when comparing the two fields and years, indicating an
39 important microbial buffering effect triggered by the soil.

40

41 **Keywords:** microbiota, rhizosphere, biocontrol agents, resistance inducers, seed born pathogen,
42 *Fusarium oxysporum* f. sp. *lactucae*.

43

44

45 **1. Introduction**

46 Lettuce is cultivated on 20,000 ha of agricultural land in Italy (ISTAT 2020). As an intensive crop, lettuce
47 yields are threatened by the presence of *Fusarium oxysporum* f. sp. *lactucae* (Fol), the causal agent of
48 Fusarium lettuce wilt. It is one of the most dangerous diseases affecting this crop worldwide (Matheron
49 and Gullino 2012; Gullino *et al.*, 2019). Characteristic symptoms of Fol infection are yellowing of leaves,
50 slow growth accompanied by brown or red streaks of the vascular system, and wilting to death.
51 Significant economic losses can occur if the disease is not properly managed (Matheron and Gullino
52 2012). In any case, Fol management is difficult for two reasons: (i) it spreads rapidly and easily as it is
53 seed-borne (Garibaldi *et al.*, 2004a) and (ii) has the ability to distinguish new races, in fact four are
54 already known and reported (Fujinaga *et al.*, 2001; 2003; Matheron and Gullino 2012; Gilardi *et al.*,
55 2017). Chemical control is not considered a sufficient strategy due to the reduced availability of soil
56 fumigants, the need to reduce chemical treatments to avoid environment and lettuce contamination, and
57 the increasing consumer demand for organic vegetable production. For these reasons, it is important to
58 develop and test new alternative control strategies against this pathogen. The use of biological control
59 agents (BCAs), organic amendments and resistance inducers are among the most studied (Bonanomi *et*
60 *al.*, 2007, 2010; Gilardi *et al.*, 2019, 2020). BCAs can act directly against pathogens via antibiosis,
61 parasitism or predation, or indirectly by colonizing the rhizosphere environment and using resources
62 more efficiently than pathogens (Pal and Gardener 2006). Several BCAs are registered in Europe against
63 soil-borne disease (Bardin and Pugliese, 2020). *Bacillus amyloliquefaciens* (former *subtilis*) QST713,
64 which is registered in Europe and Italy, is used to control *Pythium ultimum* and *Rhizoctonia solani*, but
65 also *Botrytis cinerea* and powdery mildews, via antibiosis and induced resistance (Bardin and Pugliese,
66 2020). *Trichoderma* strains are good soil and rhizosphere colonizers, and *Trichoderma gamsii* ICC080,
67 *T. asperellum asperellum* ICC012, *Trichoderma polysporum* IMI206039, and *Trichoderma atroviride*

68 IMI206040 are applied towards a broad spectrum of plant pathogens (EFSA, 2013; Bardin and Pugliese,
69 2020). *Trichoderma gamsii* ICC080 and *T. asperellum asperellum* ICC012 are registered in Europe and
70 Italy, while *Trichoderma polysporum* IMI206039 and *Trichoderma atroviride* IMI206040 are registered
71 in Europe.

72 Organic amendments, and composts, have been described for decades as suppressive against various
73 vascular pathogens, including Fol (Pugliese *et al.*, 2015; Gilardi *et al.*, 2016; De Corato *et al.*, 2018a,
74 2018b; Bonanomi *et al.*, 2018). Many studies have demonstrated that their microbiota plays an important
75 role in suppressive action (Reuveni *et al.*, 2002; Tilston *et al.*, 2002; Papatiriou *et al.*, 2013; De Corato
76 *et al.*, 2019). Their mechanisms of action are similar to those of BCAs, but the complexity and richness
77 of the compost microbiota suggest that those different mechanisms could be used and act synergistically.
78 Composts can also be enriched with BCAs to enhance their suppressive activity, as it has been indicated
79 that this technique could be the most promising to achieve long-term suppressiveness against vascular
80 pathogens (Pugliese *et al.*, 2011; Bonanomi *et al.*, 2018; Gilardi *et al.*, 2019). A compost enriched with
81 *Trichoderma virens* TW2 demonstrated to be effective in controlling Fusarium wilt on vegetable crops
82 (Gilardi *et al.*, 2019; Cucu *et al.*, 2019 and 2020b).

83 Resistance inducers are compounds of various origins that have been shown to activate the plant immune
84 system (Walters *et al.*, 2009; Akram and Anjum 2011; Alexandersson *et al.*, 2016). BCAs, and by
85 extension composts, have been shown to activate the induction of resistance in several pathosystems
86 (Compant *et al.*, 2005; Ongena *et al.*, 2007; Pieterse *et al.*, 2014; Akram *et al.*, 2015; Bellini *et al.*, 2021).

87 Integrated pest management (IPM) strategies are combinations of different measures based on the
88 principle of synergy between them to control pathogen attacks (Barzman *et al.*, 2015).

89 Soil microbial activities are responsible for nutrient transformation, decomposition of organic matter,
90 protection against pathogens and contribute to soil structure (Bowles *et al.*, 2014). The complex of
91 microorganisms inhabiting the rhizosphere can strengthen plants and protect them from both biotic and

92 abiotic stresses (Nihorimbere *et al.*, 2010; Chaudary *et al.*, 2021a), in fact, the rhizosphere microbiota
93 population is one of the greatest influences on plant and soil health (Berendsen, 2012; Kumari *et al.*,
94 2020; Chaudary *et al.*, 2022). The role of rhizosphere microbiota population is even more important for
95 the protective effect against vascular pathogens that invade plant tissues through the root system, as is
96 the case with Fol (Hubband and Gerik 1993; Gordon, 2017). The application of organic amendments and
97 BCAs may cause a shift in the composition and diversity of the rhizosphere microbiota, leading to a
98 change in soil suppressiveness. The combination of plant growth promoting rhizobacteria such as
99 *Bacillus* sp. and nanocompounds was demonstrated to have a strong influence on the composition of
100 rhizospheric microbiota on maize and to increase bacterial diversity and richness (Chaudary *et al.*,
101 2021b). The role of the rhizosphere microbiota in plant health is well established but it is not clear how
102 microorganisms applied as treatments interact with those already present in the soil and their ability to
103 establish permanent colonization. Studying the rhizosphere microbiota with amplicon-based sequencing
104 is a powerful tool (Simmons *et al.*, 2018; Elsayed *et al.*, 2020; Chaudary *et al.*, 2021b) to understand the
105 effects of IPM strategies on microbial populations at the end of the crop cycle and to determine whether
106 the protective effect guaranteed by the strategies is related to the shift in these populations. The key
107 difference between the conventional methods, i.e., Sanger sequencing and high-throughput sequencing
108 (HTS), is the sequencing volume. While the Sanger method sequences a single DNA fragment at a time,
109 HTS is massively parallel and sequences millions of fragments simultaneously per run. This process
110 enables the simultaneous sequencing of hundreds to thousands of genes. HTS also provides greater
111 discovery power to detect novel or rare variants with deep sequencing. Currently, HTS is used not only
112 for universal gene analysis but also for functional microbes (Hou *et al.*, 2018; Sultana *et al.*, 2019)
113 because it provides considerable information, is fast and easy to use, and is relatively inexpensive.
114 Furthermore, HTS was already applied and compared to conventional methods for studying the impact

115 of BCAs and biofertilizers on soil and rhizosphere microbial community (Cucu *et al.*, 2020a; Bellini *et*
116 *al.*, 2021).

117 In the present work, and based on previous studies (Gilardi *et al.*, 2016; 2019), three IPM strategies were
118 developed by using compost enriched with *T. virens* TW2 and the combinations of different BCAs
119 (*Trichoderma* spp. and *B. amyloliquefaciens*) alone or with potassium phosphite. Lettuce rhizosphere
120 composition and diversity were investigated for both bacteria (16S rRNA) and fungi (ITS) using Illumina
121 amplicon-based sequencing. The objectives of the work were: i) to evaluate the efficacy of the three
122 selected IPM strategies against lettuce Fusarium wilt under commercial and experimental fields with
123 different soils and ii) to study the rhizosphere microbiota of treated and untreated plants at genus/class
124 level.

125

126 **2. Material and Methods**

127

128 *2.1. Plants material and experimental design*

129 Field trials were conducted on two farms: a commercial farm in Moretta (CN, Italy) and an experimental
130 farm in Carmagnola (TO, Italy). The experiments were conducted under 360 m² and 64 m² plastic
131 tunnels, respectively, in Moretta and Carmagnola. The farm in Moretta (sand:silt:clay 56:19:25, pH 7.12
132 and 1.37% organic matter) had a natural infestation of Fol race 1, which causes significant crop losses in
133 susceptible lettuce cultivars (Gilardi *et al.*, 2019). The soil in Carmagnola (sand:silt:clay 68.16:10.7:21.1,
134 pH 8.2 and 0.94% organic matter) has no history of intensive lettuce cultivation nor evidence of natural
135 occurrence of the pathogen. For these reasons, the soil in Carmagnola was artificially infested with a
136 virulent strain of Fol race 1 coded as MYA-3040 (ATCC), previously isolated from lettuce wilted plants
137 in Italy, from the Agroinnova collection (Garibaldi *et al.*, 2002). Soil infestation was carried out as
138 follows: i) the pathogen was cultured in sterilized wheat kernels left at 23°C for two weeks, then ii) 100

139 g/m² of the colonized wheat kernels were incorporated into the soil at a depth of 10-15 cm using a rake.
140 The experiments were repeated in two consecutive years (2019 and 2020, see Table S1) with the lettuce
141 cultivar 'Voluski', classified as moderately susceptible to Fol (Gilardi *et al.*, 2017). The trial Carmagnola
142 2020 was done about 1 month later compared to the previous year, due to weather conditions and intense
143 rainfall. Trials lasted 33-41 days according to weather conditions. In both fields (Carmagnola and
144 Moretta), two-week-old lettuce plants were transplanted in a mulched soil at a density of 16 plants/m²,
145 with a randomized design of treatments and untreated controls, with four replicates to avoid the side
146 effect. Plants were irrigated with a drip irrigation system and grown in accordance with standard
147 cultivation practices in the region.

148

149 2.2. Integrated pest managements treatments

150 The three IPM strategies tested in this experiment were selected from those described in previous works
151 (Gilardi *et al.* 2016; 2019), choosing the best combination for the control of Fol in greenhouse and field
152 conditions. The strategies were designed and applied starting at the nursery level as follows: (i) a compost
153 enriched with *T. virens* TW2 (ANT's compost M; AgriNewTech s.r.l., Italy), (ii) a combination of
154 commercial BCAs *B. amyloliquefaciens* (former *subtilis*) QST713 (accession number CP025079,
155 Serenade Max, 15.6%, Bayer Crop Science, Italy), *T. gamsii* ICC080 (accession number IMI 392151) +
156 *T. asperellum asperellum* ICC012 (accession number IMI 392716) (Remedier, 2+2%, Isagro Ricerca,
157 Milan, Italy), and potassium phosphite (Alexin, 95PS, P₂O₅ 52%, K₂O 42%, Massò, Spain), and (iii) a
158 commercial mixture of *T. polysporum* IMI206039 (accession number IMI 206039, ATCC 20475) and *T.*
159 *atroviride* IMI206040 (accession number IMI 206040, ATCC 20476) (Binab solution 1+1%, BINAB
160 Bio-innovation AB, Helsingborg, Sweden). *T. virens* TW2 (accession number MZ222411) was isolated
161 from compost (Gilardi *et al.*, 2019; Cucu *et al.*, 2019, 2020a and 2020b; Bellini *et al.*, 2021). To simplify,
162 the tested IPM strategies are coded here as: i) Comp_Tricho, ii) Bac_Tricho and iii) Tricho. The products

163 Bac_Tricho and Tricho were applied as soil drench in the nursery. Comp_Tricho was used at sowing
164 (T0) at a dosage of 400 g/100 seedlings and applied in the field immediately before transplanting at a
165 dosage of 1 kg/m² for Moretta. In Carmagnola, Comp_Tricho was applied at a dosage of 0.5 kg/m²,
166 considering the soil mixture, the possibility of applying the treatment with the rototiller (in Moretta field)
167 or manually (in Carmagnola field), and because of the smaller size of the tunnel, which allows less deep
168 diffusion. The type, dosage and timing of application of the three strategies are shown in Table 1.

169

170 2.3. Lettuce yield and disease assessment

171 Yield and disease were evaluated at the end of each experiment. Sixteen plants per replicate were visually
172 evaluated, and dissected to assign rating scale 0 to 4 (Garibaldi *et al.*, 2004b), for a total of 64 plants for
173 each experimental thesis. Disease severity data were calculated as follows:

$$174 DS_{0-100} = \frac{\left(\frac{\sum N_{plants} \cdot Rating\ scale_{0-4} \cdot 100}{Total\ N\ recorded\ plants} \right)}{4}.$$

175 The same 64 plants were weighed at the end of the trials to measure their fresh weight. The data were
176 then analysed using SPSS 26 software by performing one-way analysis of variance (ANOVA) and
177 *Tukey's post hoc* tests to determine the statistical values of differences at the value of $\alpha = 0.05$.

178

179 2.4. Soil collection and DNA extraction

180 Rhizosphere samples were collected at the end of the experiment as follows: roots were shaken to avoid
181 any excess soil, and the remaining particles adhering root surface were collected in sterile vials for an
182 amount of 100 g from each plant. Three biological samples were collected from each treatment in both
183 fields and years. Each sample was made unifying the rhizosphere soil of five plants randomly chosen
184 inside each replicate. In total 48 samples were collected.

185 Microbial DNA extraction was performed using the “EZNA soil DNA kit” (Omega Bio-Tek, Norcross,
186 GA), following manufacturer’s instructions. DNA concentration was assessed by using a NanoDrop 2000
187 spectrophotometer (Thermo Fisher Scientific, Waltham MA). Amplicon based sequencing was
188 performed using V3 – V4 region (16S rRNA) for bacterial community (primers: 341F -
189 CCTAYGGGRBGCASCAG, 806R – GGACTACNNGGGTATCTAAT) and ITS2 region for fungal
190 community (primers: ITS3 - GCATCGATGAAGAACGCAGC, ITS4-
191 TCCTCCGCTTATTGATATGC). The analysis was done by Novogene using Illumina NovaSeq 6000
192 platform (Cambridge Science Park, Cambridge, CB4 0FW, United Kingdom).

193

194 2.5. Sequence data analysis

195 Demultiplexed fastq files were processed using the DADA2 pipeline (version 1.16.0) (Callahan *et al.*,
196 2016) in R software (4.0.4) (Team 2016). The resulting taxonomic units are referred to as amplicon
197 sequence variants (ASVs) and the identified Operational Taxonomic Units (OTUs) were 50. For bacterial
198 sequences, forward and reverse reads were trimmed to 250 bp and primer sequences were removed using
199 the following filter parameters: maxN = 0, maxEE for both reads = 2, truncQ = 2 (MaxEE corresponded
200 to the maximum expected errors, TruncQ represented the parameter that truncates reads on the first
201 occurrence of a quality score less than or equal to two, and MaxN was the maximum number of 'N' bases
202 accepted). Nearly 4 million reads were used to estimate the error rates by learnErrors function. Sequences
203 were dereplicated using derepFastq with default parameters and exact sequence variants were resolved
204 using the dada algorithm. The RemoveBimeraDenovo function was then used to remove chimeric
205 sequences. Fungal sequences were preliminary trimmed using Cutadapt software (Martin 2011) to
206 remove adapter sequences and low-quality ends (<Q20). For both the bacterial and fungal datasets, reads
207 with more than three errors in the forward reads and five errors in the reverse reads were removed.
208 Taxonomy was then assigned using assignTaxonomy based on the SILVA (v132) and UNITE (v7)

209 databases for bacterial and fungal communities, respectively (Quast *et al.*, 2013; Nilsson *et al.*, 2019).
210 Raw sequences are available at the National Centre for Biotechnology Information (NCBI), under
211 accession number PRJNA781120 with the title “BCAs application for soil microbiome”.

212

213 2.6. Statistical analysis and data visualization

214 Plotting was performed using PRIMER 7 software (Primer-E Ltd, Plymouth; UK). Alpha diversity
215 metrics were calculated. Heatmaps were created to assess variation in community composition at lowest
216 taxonomic levels including the most frequent 50 ASVs of both bacterial and fungal communities for all
217 the samples. In heatmaps, to assess the behaviour of group of variables according to IPM strategies,
218 clustering was made according index of association. Differently, samples were ordered according to
219 non-metric multidimensional scaling (NMDS) based on Bray-Curtis dissimilarity matrices. The
220 significance of changes in composition of the two communities analysed were tested by PERMANOVA
221 (999 permutations, Table S2), using the treatments, field, and year as fixed factors. The significance of
222 variation in the alpha diversity metrics of the two communities was assessed using the ANOVA test, and
223 specific changes in IPM strategies assessed by *post-hoc* Tukey test. The level of significant differences
224 was assessed at $\alpha < 0.05$. All statistical analyses were performed using R software. Furthermore,
225 functional group variation for the fungal community were analysed, identifying putative fungal
226 functional groups as well as their trophic modes using FUNGuild (Nguyen *et al.*, 2016). Co-occurrence
227 networks incorporating communities containing bacteria and fungi were based on single ASV and
228 generated using only the 50 most frequent ASVs for each bacteria and fungi. The pairwise correlations
229 between the ASVs were calculated using the Spearman correlation in R (version 3.3.2 and Hmisc package
230 4.0–1). Based on the statistical analysis, only strong and significant correlations were considered
231 (Spearman’s $r > 0.6$ or $r < -0.6$ and $p < 0.05$). The network visualization was made using Cytoscape
232 (version 3.8.2). Each edge represents correlation, and each node represents an ASV. A set of integrative

233 metrics were calculated and compared to describe the network topology. For example, the average
234 number of neighbours explains the complex pairwise connections and the average path length describes
235 node distribution. Pearson's correlation was used to identify the correlations between disease severity
236 and yield production and the first 50 most abundant ASVs for bacteria and fungi.

237 **3. Results**

238

239 *3.1. Disease assessment and yield*

240 In both years, the disease severity (DS) recorded in the control plots always showed a statistically higher
241 infection rate compared to the treated ones in Carmagnola (DS 34.4-43.3) and Moretta (DS 43.3-40.0),
242 with the only exception for the Comp_Tricho and Tricho strategies in the trial conducted in 2020 at the
243 Carmagnola experimental farm (Fig. 1a). At the Carmagnola site, Comp_Tricho, Bac_Tricho, Tricho and
244 the control provided DS of 17, 12.5, 14.3 and 34.4 in 2019 and 21.7, 10, 20.8 and 43.3 in 2020,
245 respectively; while at the Moretta site DS of 11, 13, 13.1 and 43.3 in 2019 and 24.4, 14.4, 18.1 and 40 in
246 2020 were recorded. Except for the case in Carmagnola in 2020, which had a problem with water runoff
247 due to the intense summer rains, all treatments showed a fresh weight between 3.7 and 5.3 kg/m², while
248 the controls ranged between 1.4 and 4.4 kg/m² (Fig. 1b). In Moretta, all treated plots had statistically
249 higher yield compared to the untreated control in 2020. The same trend was observed in Carmagnola in
250 2020 with the treatments Comp_Tricho and Bac_Tricho IPM compared to the control. The efficacies of
251 both treatments and yield production are shown in Figure 1.

252

253 *3.2. Microbial diversity*

254 Illumina amplicon-based sequencing (on 16S rRNA and ITS fragments) revealed that the composition
255 of the microbiota in the rhizosphere, expressed as diversity indices (number of species, number of reads,
256 and Shannon index), showed no statistical differences between samples for both bacterial and fungal
257 communities (Fig. 2-3).

258

259 3.3. *Rhizosphere associated bacteria*

260 The bacterial community showed a dominance of five phyla: Proteobacteria, Chloroflexi, Firmicutes,
261 Acidobacteria and Actinobacteria (Fig. 4a; Table S3) in all samples. There were observable differences
262 only between different trials, indicating some kind of site- and year-specific pattern. No similarities were
263 observed between the same treatments in different years or fields, nor were there any significant
264 differences between treated plants and controls. Moretta 2020 had the most similar pattern for the
265 bacterial community. In Moretta 2019 and in Carmagnola 2019, Comp_Tricho and control plants had a
266 similar rhizosphere bacterial community. Overall, no major differences were observed between controls
267 and treatments. Plotting the 50 most abundant bacterial community ASVs at genus level (Fig. S3) showed
268 that the greatest differences in community composition existed between the two fields. Specifically, most
269 bacterial ASVs were found in similar abundance in both Carmagnola and Moretta, while some ASVs
270 showed some specificity for one or the other field, such as *Rodopseudomonas*, *Enterobacteriaceae*,
271 *Paenisporosarcina*, which were found more frequently in the Carmagnola field, and *Blautia*,
272 *Faecalibacterium*, *Escherichia* and *Ralstonia*, which were more frequent in Moretta. Bacteria of the
273 genus *Bacillus*, which were slightly more abundant in Carmagnola, were the most abundant ASV, which
274 underlines the analysis. This genus was not more abundant in the plants treated with Bac_Tricho.

275

276 3.4. *Rhizosphere associated fungi*

277 A similar situation was observed for the fungal community (Fig. 4b; Table S3), where Ascomycota
278 occupied 75 to 90% of the observed phyla, followed by Basidiomycota, Mucoromycota, and
279 Aphelidiomycota, with variable proportions. There was also no clear separation between the rhizosphere
280 of the treated plants and that of the control among the fungi. More in detail, Carmagnola 2019 had a clear
281 pattern of similarity between samples, in Carmagnola 2020 there was a higher proportion of
282 Mucoromycota and Aphelidiomycota in Bac_Tricho, in Tricho and in the control samples. In Moretta

283 2019, there was the highest proportion of Basidiomycota (about 20%) in the Bac_Tricho treatment, while
284 Moretta 2020 had the highest proportion of unassigned sequences in the Tricho treated plants. The
285 heatmap of the 50 most abundant ASVs at genus level in the fungal community (Fig. S4) showed that
286 few of them dominated over the others. Ascomycota, for example, was the most abundant, with a greater
287 presence in the Carmagnola field compared to the Moretta field. In terms of fungi, there was also a
288 common core of ASVs between the two sites, but also some differences: *Fusarium* and *Chaetomium*
289 were more abundant in Moretta, while *Rhizopus* was slightly more abundant in Carmagnola.
290 *Trichoderma* did not show a greater predominance in the rhizosphere of the treated plants compared to
291 the control plants.

292

293 3.5. Bacterial and fungal distribution and co-occurrence

294 Considered at the whole community level with nMDS analysis, both bacteria and fungi showed no spatial
295 segregation as a function of treatments. The ordination of the bacterial community (Fig. 5a) clearly
296 separated by field and year, while that of the fungi (Fig. 5b) followed the site. Pearson correlations (Fig.
297 S2) between bacterial or fungal communities and disease severity or yield production for the Carmagnola
298 and Moretta fields generally showed no strong correlations for any individual ASV, except for
299 Carmagnola and yield in the bacterial community. The correlation basis network (Fig. 6) showed that the
300 Carmagnola and Moretta fields had different interactive structure when the community of bacteria and
301 fungi were analysed together. In the Moretta field, the correlations can be explained as more open, fungi
302 and bacteria seemed to co-occur and co-exclude each other, while in Carmagnola the trend was more
303 closed, as fungi seemed to interact only with each other and bacteria did the same. Figures (6a and 6b)
304 highlighted some genera that were considered more important for the experiments. *Fusarium* and
305 *Trichoderma* did not correlate in a strong way either in Carmagnola or in Moretta. *Bacillus*, which

306 dominated the population of both fields, showed few correlations in the Moretta field and had none in
307 the Carmagnola field.

308 3.6. Fungal functional guilds

309 The fungal community was analysed to identify the different guilds (Fig. 7). Twenty-one main guilds
310 were found, with a predominance of plant pathogens, animal pathogens, fungal parasites, endophytes and
311 wood saprophytes. In terms of phyla composition, there was no clear pattern identifying the treatments
312 in the different trials, nor was there a consistent difference between treated and control plants. The tricho
313 treatment of Carmagnola 2019 and the Bac_Tricho treatment of Moretta 2019 showed a greater number
314 of guilds compared to all other treatments. Seven trophic modes were found (Fig. S1) with dominance
315 for pathotroph-saprotroph-symbiotroph in Moretta (both 2019 and 2020) and increased abundance of
316 pathotroph-saprotroph and saprotroph-symbiotroph for Carmagnola field in both years.

317

318 4. Discussion

319 4.1. *Fusarium wilt control by IPM strategies*

320 The introduction of environmentally friendly strategies to control lettuce *Fusarium wilt* is extremely
321 important, as is a better understanding of how they behave under real farm conditions. In this work, the
322 effectiveness of three IPM strategies against Fol was tested under commercial and experimental
323 conditions in tunnels. The three IPM strategies tested showed a great ability to statistically reduce disease
324 severity in both fields and years, from 50% to 70% compared to untreated controls. This is in agreement
325 with preliminary experiments conducted in both greenhouse and field conditions (Cucu *et al.*, 2019;
326 Gilardi *et al.*, 2019). Also yield of treated lettuce plants significantly increased in 2020 trials, while in
327 2019 yields were not different from untreated control (Figure 1). This is in accordance with previous

328 publication (Gilardi *et al.*, 2016), in which it was clearly demonstrated that the fresh weight reduction of
329 the lettuce grown in the inoculated and treated soil was not significantly influenced by disease severity.

330

331 4.2. Impact on rhizosphere microbiota

332 At the phylum level, the microbiota data showed that there was no detectable treatment effect at the end
333 of the experiment. In fact, there was no clear differentiation between treated plants and the corresponding
334 controls, in contrast to a previous work where the same compost applied against *Phytophthora capsici*
335 protected *Cucurbita pepo* by altering its rhizosphere composition (Bellini *et al.*, 2020). In this case, the
336 experiment was conducted under greenhouse/pot conditions, using steam-sterilized peat as substrate,
337 which was a system less complex than soil in terms of microbial diversity. Some work reported a change
338 in rhizosphere microbiota when different BCAs were used, but only in pot systems (Liu *et al.*, 2021) or
339 when plant growth promoting bacteria were applied along with nanocomposites (Kumari *et al.*, 2020;
340 Chaudary *et al.*, 2022). While, in agreement with our results, Cucu *et al.*, (2020a) reported that field
341 application of BCAs did not alter the rhizosphere microbiome of *Cucurbita pepo* grown in agricultural
342 soil.

343

344 4.3. Impact on bacterial microbial diversity

345 Most ASVs were similar, but some differences were observed between the two fields. Proteobacteria,
346 Chloroflexi, Firmicutes, Acidobacteria and Actinobacteria were predominant phyla, and *Bacillus* was the
347 most abundant ASV for bacteria, which is consistent with the literature (Amin *et al.*, 2015). All
348 predominant phyla play an important role in ecological and metabolic functioning of the soil.
349 Proteobacteria are mainly involved in decomposition, nitrogen fixation, and humus formation
350 (Chaudhary *et al.*, 2021b). Chloroflexi and Acidobacteria are considered green chlorophototrophic
351 bacteria that can use chlorosomes for light-harvesting (Thweatt *et al.*, 2019). Abundance of Chloroflexi

352 was reported in wheat planted soil (Liu *et al.*, 2017), and increased with the application of nanozeolite
353 under wheat cultivation (Khali *et al.*, 2019) and of nanocompounds under maize cultivation (Chaudhary
354 *et al.*, 2021b). Firmicutes and *Bacillus* are considered important in plant growth promotion, facilitating
355 plant nutrient acquisition, in biocontrol of plant pathogens, as well as in the phytoremediation of heavy
356 metals (Amaresan *et al.*, 2020). Actinobacteria populations of soil participate in the decomposition of
357 organic matter, promote plant growth and are producers of secondary metabolites, which can be exploited
358 for disease suppression (Amaresan *et al.*, 2020). The Moretta field had a greater abundance of
359 *Gemmatimonadaceae*, *Blautia* and *Bifidobacterium* than the Carmagnola field. These bacterial ASVs
360 have been reported to be involved in chitin degradation and could be affected by fertilization and the use
361 of sludge amendments in the soil (Vo *et al.*, 2017; Hui *et al.*, 2020), which is consistent with the fact that
362 Moretta is a commercial farm subject to more intensive management and therefore conventional
363 fertilization. Moreover, the abundances of *Blautia* and *Bifidobacterium* could also be explained by the
364 difference in pH between the two fields, as the two bacterial genera have a neutral pH optimum (Cui *et*
365 *al.*, 2021; Liu *et al.*, 2021). Some of the ASVs found in greater presence in Carmagnola soil (such as
366 Enterobacteriaceae, *Pantaea* and *Lactococcus*) have been associated with the ability to degrade simple
367 sugar molecules (Degelmann *et al.*, 2009; Teuber *et al.*, 2006) and this may indicate a richer presence of
368 these molecules in this field compared to Moretta.

369

370 4.4. Impact on fungal microbial diversity

371 The 50 most abundant ASVs of fungi showed that the greatest dominance was in Ascomycota, which
372 was expected since this phylum is predominant in agricultural soils (Ma *et al.*, 2013; Egidi *et al.*, 2019).
373 *Fusarium* was very abundant in the Moretta field, as expected based on the history of Fol infection at
374 this site. *Chaetomium* was also more abundant in Moretta, again a genus containing species that degrade
375 cellulose-rich substrates, such as components in soil, straw, or wood (Aru *et al.*, 1997). Neither

376 *Trichoderma* nor *Bacillus* was found in greater abundance in the rhizosphere of plants treated with these
377 BCAs, even if BCAs and compost were applied together. Accordingly, very low Pearson correlation
378 coefficients were found between the 50 most abundant ASVs with yield production and disease severity.
379 Two-dimensional nMDS analyses showed that there was no treatment effect on the spatial distribution
380 of bacterial and fungal communities. Bacteria segregated by site and year, while for the fungal
381 community segregation was only observed for site. Network analyses of the two fields confirmed that
382 genera such as *Fusarium*, *Trichoderma*, and *Bacillus* were not involved in large co-occurrences or co-
383 exclusions; however, some differences were highlighted between the two fields, again suggesting that
384 the microbiota compositions of the rhizosphere samples at genus/class levels were mainly influenced by
385 soil type than by experimental treatments. Both in the present work and in Cucu *et al.* (2020) a standard
386 microbiome analysis was done at the genus / class level. Therefore, there might be shifts at the species
387 or strain level and natural strains may have replaced the applied strains. This aspect should be further
388 investigated in future research.

389 Nevertheless, a clear protective effect of the treatments against *Fusarium* wilt in lettuce was observed.
390 Apart from changing the rhizosphere microbial community, several mechanisms have been proposed to
391 explain the observed disease suppression, including the release of fungitoxic compounds (Blok *et al.*,
392 2000; Larkin & Griffin, 2006). Moreover, the induction of resistance may also have played a central role
393 in these experiments by activating the molecular pathways that can protect the plant from pathogen
394 attacks. Many studies have reported the ability of *Trichoderma* spp. to activate the immune system of the
395 plants they encounter (Fontenelle *et al.*, 2011; Ramírez-Cariño *et al.*, 2020; Sawant *et al.*, 2020). Indeed,
396 in a previous work with the *Capsicum annuum* - *Phytophthora capsici* pathosystem (Bellini *et al.*, 2021),
397 it was shown that the same *Trichoderma* TW2-enriched compost used in the first IPM strategy and
398 potassium phosphite (used in the second IPM strategy) activate systemic acquired resistance.

399

400 **5. Conclusion**

401 This study showed that the three IPM strategies tested here were able to reduce disease severity caused
402 by Fol in two different soils under commercial tunnel conditions in two consecutive years. The
403 rhizospheric microbiota at genus/class levels was not driven by the treatments done in nursery, but it was
404 shaped by the autochthonous soil microbial populations. Induction of resistance may have also been
405 involved in the protective effect of the treatments. Further studies should be considered to evaluate
406 whether lettuce resistance pathways can be activated by these IPM strategies, the effects on rhizospheric
407 microbiota at species/strain levels, and possible correlation of soil type and soil pH on microbial
408 community and disease suppression by BCAs.

409

410 **Acknowledgements**

411 The authors thank pathology department of University of Napoli Federico II for the collaboration and
412 hospitality, in particular professors Felice Scala, Matteo Lorito and Sheridan Woo.

413

414 **Funding**

415 This manuscript has received funding to Agroinnova, University of Turin, from the European Union's
416 Horizon 2020 Research and Innovation Program under grant agreements no. 817946 (EXCALIBUR) and
417 no. 633999 (EUCLID).

418

419 **References**

420 Akram, W., Anjum, T., 2011. Use of bioagents and synthetic chemicals for induction of systemic
421 resistance in tomato against diseases. *Int. Res. J. Agric. Sci. Soil Sci.* 1, 286-292.
422 Akram, W., Anjum, T., Ali, B., 2015. Searching ISR determinant/s from *Bacillus subtilis* IAGS174
423 against Fusarium wilt of tomato. *BioControl* 60, 271-280.

- 424 Alexandersson, E., Mulugeta, T., Lankinen, Å., Liljeroth, E., Andreasson, E., 2016. Plant resistance
425 inducers against pathogens in *Solanaceae* species—From molecular mechanisms to field application. *Int.*
426 *J. Mol. Sci* 17, 1673.
- 427 Amaresan, N., Kumar, M. S., Annapurna, K., Kumar, K., Sankaranarayanan, A. (eds), 2020. Beneficial
428 Microbes in Agro-Ecology: Bacteria and Fungi. Academic Press, Florida, USA, 932 pp.
- 429 Amin, M., Rakhisi, Z., Ahmady, A. Z., 2015. Isolation and identification of *Bacillus* species from soil
430 and evaluation of their antibacterial properties. *Avicenna J. Clin. Microbiol. Infect.* 2, 23233.
- 431 Aru, A., Munk-Nielsen, L., Federspiel, B.H., 1997. The soil fungus *Chaetomium* in the human paranasal
432 sinuses. *Eur. Arch. Otorhinolaryngol.* 254, 350–352. <https://doi.org/10.1007/BF02630728>
- 433 Bardin, M., Pugliese, M., 2020. Biocontrol Agents Against Diseases. In: Gullino, M., Albajes, R., Nicot,
434 P. (eds) *Integrated Pest and Disease Management in Greenhouse Crops. Plant Pathology in the 21st*
435 *Century*, vol 9. Springer, Cham., pp 385–407.
- 436 Barzman, M., Bärberi, P., Birch, A. N. E., Boonekamp, P., Dachbrodt-Saaydeh, S., Graf, B., Hommel,
437 B., Jensen, J. E., Kiss, J., Kudsk, P., Lamichhane, J. R., Messéan, A., Moonen, A.-C., Ratnadass, A.,
438 Ricci, P., Sarah, J.-L., Sattin, M., 2015. Eight principles of Integrated Pest Management. *Agron. Sustain.*
439 *Dev.* 35, 1199-1215. <https://doi.org/10.1007/s13593-015-0327-9>.
- 440 Bellini, A., Ferrocino, I., Cucu, M. A., Pugliese, M., Garibaldi, A., Gullino, M. L., 2020. A compost
441 treatment acts as a suppressive agent in *Phytophthora capsici*–*Cucurbita pepo* Pathosystem by modifying
442 the rhizosphere microbiota. *Front. Plant Sci.* 11, 885.
- 443 Bellini, A., Pugliese, M., Guarnaccia, V., Meloni, G. R., Gullino, L. M., 2021. Calcium oxide, potassium
444 phosphite and a *Trichoderma* enriched compost water suspension protect *Capsicum annuum* against
445 *Phytophthora capsici* by priming the immune system. *Pest Manag. Sci.* 77, 3484-3490.
- 446 Berendsen, R. L., Pieterse, C.M., Bakker, P.A., 2012. The rhizosphere microbiome and plant health.
447 *Trends Plant Sci.* 17, 478–486. <https://doi.org/10.1016/j.tplants.2012.04.001>

- 448 Blok, W.J., Lamers, J.G., Termorshuizen, A.J., Bollen, G.J., 2000. Control of soilborne plant pathogens
449 by incorporating fresh organic amendments followed by tarping. *Phytopath.* 90, 253–259.
- 450 Bonanomi, G., Antignani, V., Capodilupo, M., Scala, F., 2010. Identifying the characteristics of organic
451 soil amendments that suppress soilborne plant diseases. *Soil Biol. Biochem.* 42, 136–144. doi:
452 10.1016/j.soilbio.2009.10.012
- 453 Bonanomi, G., Antignani, V., Pane, C., Scala, F., 2007. Suppression of soilborne fungal diseases with
454 organic amendments. *J. Plant Pathol.* 89, 311–324.
- 455 Bonanomi, G., Lorito, M., Vinale, F., Woo, S. L., 2018. Organic amendments, beneficial microbes, and
456 soil microbiota: toward a unified framework for disease suppression. *Annu. Rev. Phytopathol.* 56, 1–20.
457 doi: 10.1146/annurev-phyto-080615-100046
- 458 Bowles, T.M., Acosta-Martínez, V., Calderón, F., Jackson, L.E., 2014. Soil enzyme activities, microbial
459 communities, and carbon and nitrogen availability in organic agroecosystems across an intensively-
460 managed agricultural landscape. *Soil Biol. Biochem.* 68, 252–262.
- 461 Callahan, B.J., McMurdie, P.J., Rosen, M.J., Han, A.W., Johnson A.J.A., Holmes, S.P., 2016. DADA2:
462 High-resolution sample inference from Illumina amplicon data. *Nat. Methods* 13, 581–583.
- 463 Chaudhary, P., Khati, P., Chaudhary, A., Maithani, D., Kumar, G., Sharma, A., 2021a. Cultivable and
464 metagenomic approach to study the combined impact of nanogypsum and *Pseudomonas taiwanensis* on
465 maize plant health and its rhizospheric microbiome. *PLoS ONE* 16 (4), e0250574.
- 466 Chaudhary, P., Sharma, A., Chaudhary, A., Khati, P., Gangola, S., Maithani, D., 2021b. Illumina base
467 high throughput analysis of microbial diversity of maize rhizosphere treated with nanocompounds and
468 *Bacillus* sp. *Appl. Soil Ecol.* 159, 103836.
- 469 Chaudhary, P., Chaudhary, A., Bhatt, P., Kumar, G., Khatoon, H., Rani, A., Kumar, S., Sharma, A., 2022.
470 Assessment of soil health indicators under the influence of nanocompounds and *Bacillus* spp. in field
471 condition. *Front. Environ. Sci.* 9, 769871.

- 472 Compant, S., Duffy, B., Nowak, J., Clément, C., Barka, E. A., 2005. Use of plant growth-promoting
473 bacteria for biocontrol of plant diseases: principles, mechanisms of action, and future prospects. Appl.
474 Environ. Microb. 71, 4951-4959.
- 475 Cucu, M. A., Gilardi, G., Pugliese, M., Matic, S., Gisi, U., Gullino, M. L., Garibaldi, A., 2019. Influence
476 of different biological control agents and compost on total and nitrification-driven microbial
477 communities at rhizosphere and soil level in a lettuce-*Fusarium oxysporum* f. sp. *lactucae*
478 pathosystem. J. Appl. Microbiol. 126, 905-918. Cucu, M. A., Gilardi, G., Pugliese, M., Ferrocino, I.,
479 Gullino, M. L., 2020a. Effects of biocontrol agents and compost against the *Phytophthora capsici* of
480 zucchini and their impact on the rhizosphere microbiota. Appl. Soil Ecol. 154, 103659.
- 481 Cucu, M.A., Gilardi, G., Pugliese, M., Gullino, M.L., Garibaldi, A., 2020b. An assessment of the
482 modulation of the population dynamics of pathogenic *Fusarium oxysporum* f. sp. *lycopersici* in the
483 tomato rhizosphere by means of the application of *Bacillus subtilis* QST713, *Trichoderma* sp. TW2 and
484 two composts. Biol. Contr. 142, 104158.
- 485 Cui, S., Zhu, D., Mao, B., Ma, F., Zhao, J., Zhang, H., Chen, W., 2021. Rapid evaluation of optimal
486 growth substrates and improvement of industrial production of *Bifidobacterium adolescentis* based on
487 the automatic feedback feeding method. LWT- Food Sci. Technol. 143, 110960.
- 488 De Corato, U., Patrino, L., Avella, N., Lacolla, G., Cucci, G., 2019. Composts from green sources show
489 an increased suppressiveness to soilborne plant pathogenic fungi: relationships between physicochemical
490 properties, disease suppression, and the microbiome. Crop Protect. 124, 104870. doi:
491 10.1016/j.cropro.2019.104870
- 492 De Corato, U., Salimbeni, R., De Pretis, A., 2018a. Suppression of soilborne pathogens in container
493 media amended with on-farm composted agrobioenergy wastes and residues under glasshouse condition.
494 J. Plant Dis. Protect. 125, 213–226. doi: 10.1007/s41348-017-0133-5

- 495 De Corato, U., Salimbeni, R., De Pretis, A., Patruno, L., Avella, N., Lacolla, G., Cucci, G., 2018b.
496 Microbiota from 'next-generation green compost' improves suppressiveness of composted Municipal-
497 Solid-Waste to soil-borne plant pathogens. *Biol. Control* 124, 1–17. doi:
498 10.1016/j.biocontrol.2018.05.020
- 499 Degelmann, D. M., Kolb, S., Dumont, M., Murrell, J. C., Drake, H. L., 2009. *Enterobacteriaceae*
500 facilitate the anaerobic degradation of glucose by a forest soil. *FEMS Microbiol. Ecol.* 68, 312-319.
- 501 EFSA, 2013. Conclusion on the peer review of the pesticide risk assessment of the active substance
502 *Trichoderma polysporum* strain IMI 206039. *EFSA J.* 11, 3035.
503 <https://doi.org/10.2903/j.efsa.2013.3035>.
- 504 Egidi, E., Delgado-Baquerizo, M., Plett, J.M., Wang, J., Eldridge, D.J., Bardgett, R.D., Maestre, F.T.,
505 Singh, B.K., 2019. A few *Ascomycota* taxa dominate soil fungal communities worldwide. *Nat. Commun.*
506 10, 2369. <https://doi.org/10.1038/s41467-019-10373-z>
- 507 Elsayed, T. R., Jacquiod, S., Nour, E. H., Sørensen, S. J., Smalla, K., 2020. Biocontrol of bacterial wilt
508 disease through complex interaction between tomato plant, antagonists, the indigenous rhizosphere
509 microbiota, and *Ralstonia solanacearum*. *Front. Microbiol.* 10, 2835.
- 510 Fontenelle, A. D. B., Guzzo, S. D., Lucon, C. M. M., Harakava, R., 2011. Growth promotion and
511 induction of resistance in tomato plant against *Xanthomonas euvesicatoria* and *Alternaria solani* by
512 *Trichoderma* spp. *Crop Prot.* 30, 1492-1500.
- 513 Fujinaga, M., Ogiso, H., Tsuchiya, N., Sait, H., 2001. Physiological Specialization of *Fusarium*
514 *oxysporum* f.sp. *lactucae*, a causal organism of Fusarium Root Rot of Crisp Head Lettuce in Japan. *J.*
515 *Gen. Plant Pathol.* 67, 205-206.
- 516 Fujinaga, M., Ogiso, H., Tuchiya, N., Saito, H., Yamanaka, S., Nozue, M., Kojima, M., 2003. Race 3, a
517 new race of *Fusarium oxysporum* f. sp. *lactucae* determined by a differential system with commercial
518 cultivars. *J. Gen. Plant Pathol.* 69, 23–28.

- 519 Garibaldi, A., Gilardi, G., Gullino, M. L., 2004a. Seed transmission of *Fusarium oxysporum* f. sp.
520 *lactucae*. *Phytoparasitica* 32, 61–65
- 521 Garibaldi, A., Gilardi, G., Gullino, M. L., 2004b. Varietal resistance of lettuce to *Fusarium oxysporum*
522 f. sp. *lactucae*. *Crop Prot.* 23, 845–851.
- 523 Garibaldi, A., Gilardi, G., Gullino, M. L., 2002. First report of *Fusarium oxysporum* on lettuce in Europe.
524 *Plant Dis.* 86, 1052–1052.
- 525 Gilardi, G., Franco Ortega, S., Van Rijswijk, P. C. J., Ortu, G., Gullino, M. L., Garibaldi, A., 2017. A
526 new race of *Fusarium oxysporum* f. sp. *lactucae* of lettuce. *Plant Pathol.* 66, 677-688.
- 527 Gilardi, G., Pugliese, M., Gullino, M. L., Garibaldi, A., 2020. Effect of biocontrol agents and potassium
528 phosphite against *Phytophthora* crown rot, caused by *Phytophthora capsici*, on zucchini in a closed
529 soilless system. *Sci. Hortic.* 265, 109207.
- 530 Gilardi, G., Pugliese, M., Gullino, M. L., Garibaldi, A., 2019. Nursery treatments with resistant inducers,
531 soil amendments and biocontrol agents for the management of the *Fusarium* wilt of lettuce under
532 glasshouse and field conditions. *J. Phytopathol.* 167, 98-110.
- 533 Gilardi, G., Pugliese, M., Gullino, M. L., Garibaldi, A., 2016. Effect of different organic amendments on
534 lettuce *Fusarium* wilt and on selected soilborne microorganisms. *Plant Pathol.* 65, 704–712.
535 <https://doi.org/10.1111/ppa.12460>
- 536 Gordon, T. R., 2017. *Fusarium oxysporum* and the *Fusarium* wilt syndrome. *Annu. Rev. Phytopathol.*
537 55, 23-39.
- 538 Gullino, M.L., Gilardi, G., Garibaldi, A., 2019. Ready-to-eat salad crops: a plant pathogen's heaven.
539 *Plant Dis.* 103, 2153-2170.
- 540 Hou, S.P., Ai, C., Zhou, W., Liang, G.Q., He, P., 2018. Structure and assembly cues for rhizospheric
541 nirK- and nirS-type denitrifier communities in long-term fertilized soils. *Soil Biol. Biochem.* 119, 32-

- 542 40. Hubband, J.C., Gerik, J.S., 1993. A new wilt disease of lettuce incited by *Fusarium oxysporum* f. sp.
543 *lactucum* formae specialis nov. Plant Dis. 77, 750-754.
- 544 Hui, C., Jiang, H., Liu, B., Wei, R., Zhang, Y., Zhang, Q., Liang, Y., Zhao, Y., 2020. Chitin degradation
545 and the temporary response of bacterial chitinolytic communities to chitin amendment in soil under
546 different fertilization regimes. Sci. Total Environ. 705, 136003.
- 547 Khati, P., Sharma, A., Chaudhary, P., Singh, A.K., Gangola, S., Kumar, R., 2019. High throughput
548 sequencing approach to access the impact of nanozeolite treatment on species richness and evens of soil
549 metagenome. Biocatal. Agri. Biotech. 20, 101249.
- 550 Kumari, S., Sharma, A., Chaudhary, P., Khati, P., 2020. Management of plant vigor and soil health using
551 two agrisuable nanocompounds and plant growth promotory rhizobacteria in Fenugreek. 3 biotech 10,
552 461.
- 553 Larkin, R.P., Griffin, T.S., 2006. Control of soilborne potato diseases using brassica green manures. Crop
554 Prot. 26, 1067–1077.
- 555 Liu, X., Mao, B., Gu, J., Wu, J., Cui, S., Wang, G., Zhao, J., Zhang, H., Chen, W., 2021. *Blautia*—a new
556 functional genus with potential probiotic properties? Gut Microbes 13, 1-21.
- 557 Liu, D., Yang, Q., Ge, K., Hu, Z., Guozhen, Q., Binghai, D., Liu, K., Ding, Y., 2017. Promotion of iron
558 nutrition and growth on peanut by *Paenibacillus illinoisensis* and *Bacillus* sp. strains in calcareous soil.
559 Brazilian J. of Microbiol. 48, 656–670.
- 560 Martin, M., 2011. Cutadapt removes adaptor sequences from high-throughput sequencing reads
561 EMBnetjournal 17, 10-12. <https://doi.org/10.14806/ej.17.1.200>
- 562 Ma, A., Zhuang, X., Wu, J., Cui, M., Lv, D., Liu, C., Zhuang, G., 2013. *Ascomycota* members dominate
563 fungal communities during straw residue decomposition in arable soil. PloS one 8, e66146.

- 564 Matheron, M., Gullino, M. L., 2012. Fusarium wilt of lettuce and other salad crops. In: Gullino, M. L.,
565 Katan, J., Garibaldi, A. (Eds.), Fusarium wilts of greenhouse vegetable and ornamental crops, APS Press,
566 St. Paul, pp. 175-183.
- 567 Nguyen, N.H., Song, Z., Bates, S.T., Branco, S., Tedersoo, L., Menke, J., Schilling, J.S., Kennedy, P.G.,
568 2016. FUNGuild: an open annotation tool for parsing fungal community data sets by ecological guild.
569 Fung. Ecol. 20, 241–248.
- 570 Nihorimbere, V., Ongena, M., Smargiassi, M., Thonart, P., 2011. Beneficial effect of the rhizosphere
571 microbial community for plant growth and health. Biotechnol. Agron. Soc. 15, 327-337.
- 572 Nilsson, R.H., Larsson, K-H., Taylor, A.F.S., Bengtsson-Palme, J., Jeppesen, T.S., Schigel, D., Kennedy,
573 P., Picard, K., Glöckner, F.O., Tedersoo, L., Saar, I., Kõljalg, U., Abarenkov, K., 2019. The UNITE
574 database for molecular identification of fungi: handling dark taxa and parallel taxonomic classifications.
575 Nucleic Acids Res. 47, D259–64.
- 576 Ongena, M., Jourdan, E., Adam, A., Paquot, M., Brans, A., Joris, B., Arpigny, J.L., Thonart, P., 2007.
577 Surfactin and fengycin lipopeptides of *Bacillus subtilis* as elicitors of induced systemic resistance in
578 plants. Environ. Microbiol. 9, 1084-1090.
- 579 Pal, K.K., Gardener, B.M., 2006. Biological control of plant pathogens. Plant Health Instr. 2, 1117-1142.
- 580 Papatotiriou, F.G., Varypatakis, K.G., Christofi, N., Tjamos, S.E., Paplomatas, E.J., 2013. Olive mill
581 wastes: a source of resistance for plants against *Verticillium dahliae* and a reservoir of biocontrol agents.
582 Biol. Control 67, 51–60. <https://doi.org/10.1016/j.biocontrol.2013.07.008>
- 583 Pieterse, C.M., Zamioudis, C., Berendsen, R.L., Weller, D.M., Van Wees, S.C., Bakker, P.A., 2014.
584 Induced systemic resistance by beneficial microbes. Annu. Rev. Phytopathol. 52, 347-75.
- 585 Pugliese, M., Liu, B., Gullino, M. L., Garibaldi, A., 2011. Microbial enrichment of compost with
586 biological control agents to enhance suppressiveness to four soil-borne diseases in greenhouse. J. Plant
587 Dis. Protect. 118, 45-50.

- 588 Pugliese, M., Gilardi, G., Garibaldi, A., Gullino, M. L., 2015. Organic amendments and soil
589 suppressiveness: results with vegetable and ornamental crops. In: Meghvansi, M., Varma, A. (Eds),
590 Organic Amendments and Soil Suppressiveness in Plant Disease Management, Springer, New York, pp.
591 495–509. https://doi.org/10.1007/978-3-319-23075-7_24
- 592 Quast, C., Pruesse, E., Yilmaz, P., Gerken, J., Schweer, T., Yarza, P., Peplies, J., Glöckner, F.O., 2013.
593 The SILVA ribosomal RNA gene database project: improved data processing and web-based tools.
594 Nucleic Acids Res. 41, D590–6.
- 595 R.C. Team, 2016. R: A language and environment for statistical computing. R Found. Stat. Comput.
- 596 Ramírez-Cariño, H.F., Guadarrama-Mendoza, P.C., Sánchez-López, V., Cuervo-Parra, J.A., Ramírez-
597 Reyes, T., Dunlap, C. A., Valadez-Blanco, R., 2020. Biocontrol of *Alternaria alternata* and *Fusarium*
598 *oxysporum* by *Trichoderma asperelloides* and *Bacillus paralicheniformis* in tomato plants. A. Van
599 Leeuw. J. Microb. 113, 1247–1261. <https://doi.org/10.1007/s10482-020-01433-2>
- 600 Reuveni, R., Raviv, M., Krasnovsky, A., Freiman, L., Medina, S., Bar, A., Orion, D., 2002. Compost
601 induces protection against *Fusarium oxysporum* in sweet basil. Crop Protect. 21, 583–587.
602 [https://doi.org/10.1016/S0261-2194\(01\)00149-1](https://doi.org/10.1016/S0261-2194(01)00149-1)
- 603 Sawant, I.S., Wadkar, P.N., Ghule, S.B., Salunkhe, V.P., Chavan, V., Sawant, S. D., 2020. Induction of
604 systemic resistance in grapevines against powdery mildew by *Trichoderma asperelloides* strains.
605 Australas. Plant Path. 49, 107–117. <https://doi.org/10.1007/s13313-020-00679-8>
- 606 Simmons, T., Caddell, D. F., Deng, S., Coleman-Derr, D., 2018. Exploring the root microbiome:
607 extracting bacterial community data from the soil, rhizosphere, and root endosphere. J. Vis. Exp. 135,
608 57561.
- 609 Sultana, N., Zhao, J., Zheng, Y., Cai, Y., Faheem, M., Peng, X., Wang, W., Jia, Z., 2019. Stable isotope
610 probing of active methane oxidizers in rice field soils from cold regions. Biol. Fertil. Soils 55, 243-250.

- 611 Teuber, M., Geis, A., 2006. The Genus *Lactococcus*. In: Dworkin, M., Falkow, S., Rosenberg, E.,
612 Schleifer, K.H., Stackebrandt, E. (Eds), *The Prokaryotes*. Springer, New York, pp. 205-228.
613 https://doi.org/10.1007/0-387-30744-3_7
- 614 Tilston, E. L., Pitt, D., Groenhof, A. C., 2002. Composted recycled organic matter suppresses soil-borne
615 diseases of field crops. *New Phytol.* 154, 731–740. <https://doi.org/10.1046/j.1469-8137.2002.00411.x>
- 616 Thweatt, J. L., Canniffe, D. P., Bryant, D. A., 2019. Biosynthesis of chlorophylls and
617 bacteriochlorophylls in green bacteria. *Adv. Bot. Res.* 90, 35-89.
- 618 Walters, D. R., Fountaine, J. M., 2009. Practical application of induced resistance to plant diseases: an
619 appraisal of effectiveness under field conditions. *J. Agr. Sci.* 147, 523.
- 620 Vo, N., Tsai, T. C., Maxwell, C., Carbonero, F., 2017. Early exposure to agricultural soil accelerates the
621 maturation of the early-life pig gut microbiota. *Anaerobe* 45, 31-39.
- 622

623 **Table 1.** Coding names, technical formulation, dosage and timing of administration of the three IPM
 624 strategies used in this work for in fields experiments. T0 refers to the day of sowing, the other timing
 625 here presented are referred in days. Dosage column refers to the amount of commercial product used to
 626 treat seedling or soil. Comp_Tricho was given as compost as such, while Bac_Tricho and Tricho were
 627 given in a volume of 1 l per treatment.

Treatment	Formulation	Commercial name	Dosage	Number of applications	Timing	Application
Comp_Tricho	Ant Compost + <i>Trichoderma virens</i> TW2	ANT's CM	400 g/100 seedlings	2	T0	sowing
			1 kg/m ² for Moretta; 0.5 kg/m ² for Carmagnola		Immediately before transplant	field
Bac_Tricho	<i>Bacillus amyloliquefaciens</i> (former subtilis) QST 713	Serenade MAX	8 ml/l	2	T0	sowing
					T10	nursery
	<i>Trichoderma gamsii</i> icc 080 + <i>T. asperellum</i> <i>asperellum</i> icc 012	Remedier	2.5 g/l	2	T5	sowing
					T15	nursery
Potassium phosphite	Alexin	2.5 g/l	2	T0	sowing	
				T15	nursery	
Tricho	<i>Trichoderma polysporum</i> IMI 206039 + <i>Trichoderma atroviride</i> IMI 206040	Binab solution	1.7 g/l	3	T0	nursery
					T7	nursery
					T15	nursery

628

629

630

631

632

633

634

635

636

637

638

639

640

641

642

643

644

645

646

647

648


649


650

651

652 **Fig 1.** (a) Efficacy of the three IPM strategies to reduce *Fusarium oxysporum* f. sp. *lactucae* disease on
 653 lettuce plants expressed as disease severity (%) and (b) productivity expressed as fresh biomass (Kg/m²).
 654 Evaluation made at the final survey. Different letters indicate statistical differences between the four
 655 thesis, as obtained with the ANOVA test and Tukey's post hoc test (p<0.05).


656





657
 658 **Fig 2.** Box and whisker plots showing distribution of diversity indices, number of species, number of
 659 reads and Shannon Index, for bacteria community, for each treatment divided per site and year of
 660 rhizosphere sampling.


661




662

663 **Fig 3.** Box and whisker plots showing distribution of diversity indices, number of species, number of
 664 reads and Shannon Index, for fungal community, for each treatment divided per site and year of
 665 rhizosphere sampling.


666




667
 668 **Fig 4.** Stacked bar chart of relative abundance for (a) bacterial and (b) fungal communities at phylum
 669 level. Data are averaged among three biological replicates for each treatment and are here presented
 670 grouped for site and year.
 671



672
673 **Fig 5.** Bi-dimensional Non-metric MDS of the bacterial and fungal communities in the rhizosphere
674 collected at the end of the trials. Data are averaged for year and site of sampling, following the colour
675 coding of the legend.



676
 677 **Fig 6.** Correlation base network analysis showing potential interactions between bacterial and fungal
 678 families in (a) Moretta and (b) Carmagnola fields. The lines connecting nodes (edges) represent positive
 679 (blue) or negative (red) co-occurrence relationship. The intensity of the colour and the length of the edges
 680 represent the strength of correlation. Square nodes correspond to fungal ASVs and circle nodes
 681 correspond to bacterial ASVs. The connection stands for a strong (Spearman's $\rho > 0.6$ and $\rho < -0.6$) and
 682 significant ($P\text{-value} < 0.05$) correlation. The size of each node is proportional to the ASV relative
 683 abundance, only the top 50 ASVs were kept. The nodes were coloured by phylum level. Data of the years
 684 and the treatments were averaged.



685
 686 **Fig 7.** Stacked bar chart of relative abundance of fungal guilds based on FunGuilds. Data are averaged
 687 among three biological replicates for each treatment and are here presented grouped for site and year.

688 **Appendix A. Supplementary materials**

689

690 **Table S1.** Schedule of the trials conducted in 2019 and in 2020 in both farm and average temperatures
691 registered.

year	Farm	Sowing	Transplantation	Trial end	Field trial duration (days)
2019	Carmagnola	06/5/2019	30/05/2019	10/07/2019	41
2020		26/06/2020	21/07/2020	24/08/2020	34
2019	Moretta	06/5/2019	29/05/2019	01/07/2019	33
2020		07/05/2020	01/06/2020	06/07/2020	36

692

693

694 **Table S2.** Result of Permanova significance test across treatments during the two years of the
695 experiment, in two different fields. Treatments were used as the fixed factor (N° of permutation 999).
696 Here reported main significant variation between the studied treatments for fungal and bacterial
697 communities. Test of significance is based on Bray-Curtis similarity values. Significance level is fixed
698 for p-value below 0.05.

699

	Fungal community	Bacterial community
Permutation N	999	999
Total sum of squares	8.067	3.435
Within-group sum of squares	4.092	1.104
F	2.073	4.504
p	0.004*	0.001*

700 (*Significance level is at 0.05)

701

702


703

704


705 **Table S3.** Result of amplicon sequence analysis in the bacterial and fungal community for each sample
706 collected from rhizosphere at the end of the trials.

Commentato [MP1]: Excel file

707
708




709 **Fig S1.** Stacked bar chart of relative abundance of fungal trophic mode. Data are averaged among
710 biological replicates for each treatment and are here presented grouped for site and year.
711




712
 713 **Fig S2.** Heatmap of Pearson's correlation between the 50 most frequent amplicon sequence variant
 714 (ASV) in the bacterial and fungal community and the disease index and growth (fresh biomass). Data are
 715 expressed as average of every sample per each site.

716

717



719 **Fig S3.** Heatmap showing relative abundance of the 50 most frequent amplicon sequence variant (ASV)
 720 in the bacterial community for each sample collected from rhizosphere at the end of the trials.
 721 Hierarchical clustering of variables is based on an association index. Samples follow the colour coding
 722 of the legend.



724 **Fig S4.** Heatmap showing relative abundance of the 50 most frequent amplicon sequence variant (ASV)
 725 in the fungal community for each sample collected from rhizosphere at the end of the trials. Hierarchical
 726 clustering of variables is based on an association index. Samples follow the colour coding of the legend.
 727