

AperTO - Archivio Istituzionale Open Access dell'Università di Torino

New concepts in anaerobic digestion processes: recent advances and biological aspects

This is the author's manuscript

Original Citation:

Availability:

This version is available <http://hdl.handle.net/2318/1895153> since 2023-03-08T14:27:41Z

Published version:

DOI:10.1007/s00253-018-9039-9

Terms of use:

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)

Title: New concepts in anaerobic digestion processes: recent advances and biological aspects

Authors:

Antonio Castellano-Hinojosa^{1,‡}, Caterina Armato^{2,3,‡}, Clementina Pozo¹, Alejandro González-Martínez⁴, Jesús González-López^{1,*}

Affiliations:

¹ Department of Microbiology and Institute of Water Research, University of Granada, Granada, Spain

² Department of Public Health and Pediatrics, University of Torino, Torino, Italy

³ Centre for Sustainable Future Technologies (CSFT@PoliTo), IstitutoItaliano di Tecnologia, Torino, Italy

⁴ Department of Built Environment, School of engineering, Aalto University, Espoo, Finland

[‡]Antonio Castellano-Hinojosa and Caterina Armato contributed equally to this work.

* Corresponding author:

Department of Microbiology and Institute of Water Research, University of Granada, Granada, Spain

E-mail address: jgl@ugr.es (J. González-López).

Abstract

Waste treatment and the simultaneous production of energy have gained great interest in the world. In the last decades, scientific efforts have focused largely on improving and developing sustainable bioprocess solutions for energy recovery from challenging waste. Anaerobic digestion (AD) has been developed as a low-cost organic waste treatment technology with a simple set-up and relatively limited investment and operating costs. Different technologies such as, one-stage and two-stage AD have been developed. The viability and performance of these technologies have been extensively reported, showing the supremacy of two-stage AD in terms of overall energy recovery from biomass under different substrates, temperatures and pH conditions. However, a comprehensive review of the advantages and disadvantages of these technologies is still lacking. Since microbial ecology is critical to developing successful AD, many studies have shown the structure and dynamics of archaeal and bacterial communities in this type of system. However, the role of Eukarya groups remains largely unknown to date. In this review, we provide a comprehensive review of the role, abundance, dynamics and structure of archaeal, bacterial and eukaryal communities during the AD process. The information provided could help researchers to select the adequate operational parameters to obtain the best performance and biogas production results.

Keywords: anaerobic digestion; one stage vs two stage; microbiome; Archaea, Bacteria and Eukarya communities

Introduction

Energy production from renewable sources and efficient waste treatment are two of the more relevant scientific and social challenges nowadays (De Vrieze et al. 2017). In the last two decades, anaerobic digestion (AD) has been proven to be a valuable method able to solve both of these issues, combining recycling of different waste materials with the production of biogas (Oslaj et al. 2010; Tyagi and Lo 2013). Current systems based on AD aim to convert organic matter into biogas. During this process, hydrolyzing microorganisms hydrolyze organic polymers (i.e. fats and proteins) producing simple molecules (i.e. sugars, amino acids and fatty acids); acidogenic microorganisms consume free monomers generating volatile fatty acids (VFAs) and alcohols; acetogenic microorganisms transform VFA and alcohols into acetic acid, CO₂, and H₂; methanogenic archaea consume acetic acid or hydrogen to generate CH₄ (Gonzalez-Martinez et al. 2016a; Zhang et al. 2016b).

AD is a process that can be applied to almost any organic waste. Many different substrates have been discussed in the literature: agricultural waste, food waste, animal manure, feed waste, energy crops and plant residues, such as brewery wastewater (Pozo et al. 2002; Chen et al. 2008; Meulepas et al. 2010). In addition to the digestion of individual substrates, AD reactors can be loaded with mixtures of different residues. This approach, which is usually termed ‘co-digestion’ or ‘co-fermentation’, offers various technical and commercial advantages. One example is the biostimulating effect coming from the overproduction of nutrients, which can accelerate the degradation of solid waste (Beyene et al. 2018). Moreover, the application of mono or co-digestion is an efficient alternative to obtain a stabilized solid waste that can be applied as soil conditioner (Rolando et al. 2011; Gómez et al. 2006).

The aim of this review is threefold. First, we will discuss relevant features of AD: the structure of the plants (one-stage vs two-stage AD), the operational temperature (mesophilic vs thermophilic) and other technologies in biogas production. A second section will be devoted to describe the role of the microbiome (Archaea, Bacteria and Eukarya communities) involved in AD and its link to operational and performance parameters and biogas production. Finally, we will discuss future implications and prospective biotechnologies in AD.

Digester configurations: advantages and disadvantages

Since the appearance of AD, a wide variety of digester configurations has been tested such as thermophilic/mesophilic digestion, dry/wet digestion, one-phase/two-phase digestion or one-stage/two-stage digestion (Møller et al. 2009; Nizami et al. 2009; Khalid et al. 2011; Mao et al. 2015; Sun et al. 2015; Chen et al. 2016). Among these, the most relevant comparison, as well as the one most debated in the literature, is that based on the number of stages. However, independently of the digester configuration to obtain a high digestion efficiency, anaerobic bioreactors should allow a continuously high and sustainable organic load rate operating with short (Khalid et al. 2011) or long (Bergland et al. 2015) hydraulic retention time (HRT) depending on the substrate.

The simplest possible configuration is the one-stage AD batch reactor, in which the tank is filled with the feedstock and let stand for a period after which it is emptied (Khalid et al. 2011). Although this kind of system has very low operational cost, it exhibits some limitations such as high fluctuations in gas production, biogas losses during emptying the bioreactors and restricted bioreactor heights (Khalid et al. 2011; Zhang et al. 2015; Sunyoto et al. 2016). A more widely used type of one-stage AD bioreactor is commonly defined ‘one-stage continuously fed systems’ (Khalid et al. 2011). In one-stage AD system, hydrolysis, acidogenesis, acetogenesis

and methanogenesis take place in the same tank. This implies that acidogenic and methanogenic microbiota have to cohabit despite the existence of marked differences regarding growth factors and kinetics, nutritional needs and environmental conditions such as pH and temperature (Gonzalez-Martinez et al. 2016b; De Gioannis et al. 2017). In this context, although the ideal pH range for AD has been reported to be between 6.8–7.4, it is known that in one-stage AD bioreactor the operational pH sometimes can affect the digestive progress and products directly. However, two-stage AD process separating the hydrolysis/acidification and acetogenesis/methanogenesis processes, provides optimal conditions for each of the microbiota, since the optimal pH levels for acidogenic (5.5–6.5) and methanogenic (7.0) microorganisms can be controlled to increase the efficiency of the process (Mao et al. 2015). Consequently, in these kinds of systems, the different sub-processes of AD take place in separate sequential reactors. The most common configuration is the two-stage continuously fed system, although three-stage systems have been proposed (Angelidaki et al. 2003). Two-stage AD were originally conceived by Pohland and Ghosh (1971), and soon gained popularity, particularly for laboratory applications (Nizami et al. 2009). Although overall performance supremacy of two-stage AD has been variously reported in the literature, one-stage AD are far from being replaced (Møller et al. 2009). According to Rapport et al. (2012), 90% of the total capacity of the full-scale AD plants installed in Europe at that time was covered by one-stage systems. The main reasons behind this are probably the simpler structural features and lower operating costs. On the other hand, two-stage AD provides higher substrate conversion and better energy recovery, as well as better process stability, resilience and reliability (Salvador et al. 2013; De Gioannis et al. 2017; Shen et al. 2017).

Multiple-stage reactors have been developed to improve process stability and efficiency (Achinas et al. 2017). In this sense, Kim et al. (2011) demonstrated significantly higher digestion efficiency of a four-stage AD system using activated sludge than a single-stage system. Likewise, a novel alternative technique based on a high working pressure (up to 100 bar), permits the production of biogas with more than 95% methane content. This technique integrate in a single process both biogas production and *in situ* increased-pressure purification, generating a clean biogas (99% methane) that can be fed directly into the natural gas networks. However, the effect of the working pressure on microbiome structure is still unknown (Lindeboom et al. 2011). The complexity and high cost of this novel technologies are barriers to commercial use and until date, few multiple-stage AD units operate on a commercial scale.

Thermophilic and mesophilic conditions

A further relevant way to classify AD systems is to consider their operating temperature. Although the biogas process can proceed at different temperatures, mesophilic (30–40°C) and thermophilic (50–60 °C) conditions are commonly used (Møller et al. 2009; Wang et al. 2018). Temperature is, indeed, one of the main environmental factors affecting physical parameters such as viscosity, surface tension and mass transfer properties. Moreover, small changes in the temperature can result in a reduction in process efficiency, so its stability is also important (Angelidaki et al. 2003). Above all, temperature must be considered in relation to microbial growth and reactions (Amani et al. 2010; Gonzalez-Martinez et al. 2017) and changes in the structure and dynamics of prokaryotic and eukaryotic groups (see Section 2). The groups of microbes that have been identified for AD are mesophilic and thermophilic strains. While great diversity exists between mesophilic and thermophilic bacteria, with the latter

showing both higher specific growth and decay rates, methanogen growth is mostly favoured by both mesophilic and thermophilic temperatures (Li et al. 2015; Kundu et al. 2017).

Neither of the two conditions (i.e. mesophilic or thermophilic) is absolutely preferable. Although mesophilic digestion has some disadvantages (i.e. lower metabolic rate, lower rate and efficiency of particulate matter hydrolysis, smaller degree of pathogen deactivation and lower biogas production yields) (Liu et al. 2017), it has important advantages, such as a lower VFA concentration in the final effluents, maintenance of a higher organic loading rate (OLR) (Bayr et al. 2012) and a more stable performance (Guo et al. 2014), compared to thermophilic digestion (Appels et al. 2008; Wang et al. 2017). On the other hand, thermophilic temperatures can produce large quantities of dissolved solids in the digester supernatant and more odours, and have acidification potential and higher energy requirements. For these reasons, two-stage AD offers the opportunity to operate thermophilic hydrolysis/acidogenesis and mesophilic methanogenesis, as a good compromise. Of note, a different approach not requiring an extra heat supply, named '*ambient/seasonal temperature AD*', has also been used for organic waste. However, the changes in temperature induce less stability and lower methane production compared with the mesophilic process (Mao et al. 2015).

Biogas production

Currently, AD is implemented in various ways worldwide. In the Western world there are, to date, about ten thousands of operational AD plants (Yousuf et al. 2016; Vasco-Correa et al. 2018). A comparable amount can be found in Asia, where rural communities use small-scale household digesters for domestic necessities (Surendra et al. 2014). Similar small-scale digesters have also been installed in rural regions of Latin America and Africa during the last few years (REN21, 2016). Laws on the subject of environmental protection and waste treatment, as well as new emerging candidate substrates and innovative technologies, will surely guide the evolution of AD.

Different compositions of mixed substrates have been reported to increase the production of biogas, such as mixing municipal solid waste with industrial sludge (Ağdağ and Sponza 2007) or olive mill wastewater with olive mill solid waste (Fezzani and Cheikh 2010). In addition, co-digestion has been proved to stabilize reactor performance (Lo et al. 2010; Beyene et al. 2018). Interestingly, the use of this approach with substrates rich in carbon has been proposed as a solution to reduce ammonia and other toxic substances (Rajagopal et al. 2013; Fitamo et al. 2017). Moreover, co-digestion is an efficient strategy to degrade those kinds of waste that are difficult to process as a unique substrate. Recently, Park et al. (2016) tested different mixtures in order to optimize the processing of sewage sludge, obtaining optimal results in combination with food waste. As a further solution, Shen et al. (2017) proved that the combination of sewage sludge and pyro-biochar can improve biomethane production, compared with the digestion of sewage sludge alone.

As an example, the Korean government recently solicited the use as an AD substrate of organic waste from ocean dumping or landfill, with the aim to produce renewable energy; this raises the issue of efficiently degrading septage and sewage sludge, and the consequent investigation of different mixtures for co-digestion approaches (Park et al. 2016). Otherwise, good availability of a specific kind of waste can turn it into a candidate substrate. In Colombia, for example, the massive production of coffee generates a large amount of coffee mucilage, a crop residue rich in carbohydrates. This organic matter has been successfully used in co-digestion with pig manure to produce biohydrogen, taking advantage of two types of organic waste readily available in the

same geographical region (Hernández et al. 2014). Finally, technical innovations will help the scale-up of currently experimental systems.

Biohythane is a promising sustainable alternative to hythane. It is more environmentally friendly, requires a shorter fermentation time and offers better energy recovery than traditional biogas. Despite research interest in the production of this gas, numerous challenges have still to be addressed in order to allow large-scale production of biohythane by means of AD (Liu et al. 2018). Similarly, technical improvements are needed for the realization of full-scale three-stage AD plants. Hitherto, an in-lab preliminary study has proved that this approach could considerably improve the production of methane (Zhang et al. 2017). A further promising strategy to increase biogas yield and system performance is the application of selected microbial consortia, often taken from another operating plant. However, more accurate knowledge concerning adaptation of the inoculum is required in order to maximize the potential advantages of this approach (Wojcieszak et al. 2017).

Archaea, Bacteria and Eukarya communities in anaerobic digestion processes

Integration of microbial aspects within the framework of AD is critical to achieve the desired performance and biogas production. The microbiome as an entity does not work as a randomized mix, and scientific efforts focus largely on linking operational and performance parameters with the structure of microbial communities. Here, we highlight engineering of the microbiome, focusing on the most crucial Archaea, Bacteria and Eukarya groups.

Abundance, structure and dynamics of the microbiome in anaerobic digestion processes

Microbial ecologists and engineers have shown increasing interest concerning insight into the microbiome in anaerobic digesters. So far, the most crucial microorganisms have been identified although few authors have linked operational and performance parameters and microbiome response at laboratory or full-scale conditions (Carballa et al. 2011; Werner et al. 2011; Carballa et al. 2015; Gonzalez-Martinez et al. 2016b; De Vrieze et al. 2017; Kundu et al. 2017; Wang et al. 2018). Since a strong syntrophic relationship exists between acetogenic and methanogenic organisms involved in AD, biomonitoring of the system could be an important feature for engineers to obtain a highly efficient microbiome and to predict and prevent system failure (Amani et al. 2010). For example, Kundu et al. (2013) showed that a high degree of microbial diversity could be indicative of stable AD performance. Recently, a methodological approach to link microbial and operational data has also been described (de Los Reyes III et al. 2015).

The development of next-generation sequencing technologies has offered an opportunity to describe the microorganisms present (DNA) or active (RNA) in engineered ecosystems as well as their abundance (Muñoz-Palazon et al. 2018). Nevertheless, a combined DNA–RNA approach would result in a more accurate methodology to link the microbial community's structure and its metabolic ability requirements (Kaeffer et al. 2014; Maus et al. 2016). Identification of the critical representative species by means of these techniques can help to increase the efficiency and stability of AD (Venkiteshwaran et al. 2015; Dang et al. 2017). In this sense, the presence of sulphate-reducing bacteria in AD can decrease methane production because of substrate competition and sulphide inhibition of the methanogenic community (Chen et al. 2008; Sasaki et al. 2011). Thus, biomonitoring tools can help to prevent inefficiencies in AD.

The AD process comprises four interdependent steps in which microorganisms responsible for a specific stage provide the intermediates for the next. Microbial community structure and dynamics are important to sustain functional redundancy and to maintain a well-balanced process (Allison and Martiny 2008; Ziganshin et al. 2013). Archaea, Bacteria and Eukarya communities form the microbiome of the anaerobic digester and change during the stages of the AD process (Matsubayashi et al. 2017).

Archaea play a central role during methanogenic processes of AD, and it has been reported that these microorganisms can be related to different operational parameters (Zhang et al. 2012; Smith et al. 2014; Hao et al. 2016). Synthesis of CH₄ is carried out both by acetoclastic (e.g. *Methanosaeta*, *Methanosarcina* and *Methanothrix*) and hydrogenotrophic methanogens (e.g. *Methanobacterium*, *Methanomicrobium*, *Methanococcus*, *Methanobrevibacter*, *Methanomassili* and *Methanospirillum*) using acetic acid, or by using H₂ and CO₂ or methyl compounds to synthesize CH₄ (Calderón et al. 2013; Gonzalez-Martinez et al. 2016b). The characteristics and properties of the main methanogens involved in an AD as well as their substrates and products have been reported (McHugh et al. 2003; Amani et al. 2010; Goswami et al. 2016; Kundu et al. 2017). In most of the studies in the literature, Archaea diversity decreases with temperature elevation (Kundu et al. 2012; Guo et al. 2014), an effect more remarkable than changes in OLR which abrupt increase (from 1 to 8 g VS L⁻¹ d⁻¹) seemed to have little influence on the microbial community (Gou et al. 2014). Hao et al. (2016) compared the effect of total solid (TS) concentrations on archaeal diversity in sludge-fed digesters. Under high TS conditions (TS > 44 g/L), the relative abundance of *Methanosarcinaceae* and *Methanobacteriaceae* families increased whereas when digesters operated at lower-TS (TS ≤ 44 g/L) only *Methanosaetaceae* family was favoured. Under the use of continuous lab and full-scale reactors and food waste substrate the genus *Methanosarcina* is dominant under thermophilic conditions, with abundance higher than 80%, although *Methanothermobacter* and *Methanoculleus* are also favoured (Cho et al. 2013; Wang et al. 2018), whereas *Methanosaeta* is dominant under mesophilic conditions (accounting for >25% of relative abundance) (Gonzalez-Martinez et al. 2016b). On the other hand, *Methanosaeta* instead of *Methanosarcina* is favoured under low acid concentrations. Since VFA accumulation results in lower values for pH, Guo et al. (2014) showed a decrease in archaeal diversity when VFAs produced in the hydrolytic step are not consumed by methanogens. In fact, acetoclastic methanoarchaea have a positive correlation with VFAs and NH₄⁺ (Lin et al. 2012). Methanogen diversity is also sensitive to a pH value lower than 6.5, particularly during acid and acetate accumulation (Bräuer et al. 2006). In general, lower hydraulic retention time values decrease archaeal diversity by selecting organisms with a high growth rate and poor substrate affinity. In this sense, *Methanosaetaceae* (slower growth rate) predominate when HRT > 5 days, while *Methanosarcinaceae*, *Methanobacteriales* and *Methanomicrobiales* (faster growth rate) become dominant at HRT < 2 days (Padmasiri et al. 2007; Chelliapan et al. 2011). Regueiro et al. (2014) reported that *Methanosaeta* is crucial for reaching stable reactor performance although the archaeal community structure is affected by substrate type. Moreover, taking into account operational performance parameters, Kundu et al. (2017) indicated *Methanosaetaceae* as the best candidate for biomonitoring based on its sensitivity to fluctuations in the AD process.

The presence of bacterial genera such as *Desulfotomaculum*, *Desulfovibrio*, *Syntrophobacter*, *Syntrophomonas*, *Syntrophospora*, *Clostridium*, *Bacteroides*, *Bifidobacterium*, *Butyrivibrio*, *Pseudomonas*, *Bacillus*, *Streptococcus* and *Eubacterium* has been related to acid formation and hydrogen release (Yamada et al. 2006; Gonzalez-Martinez et al. 2016a), and synergistic cooperation with methanogenic archaeal groups in

methanogenesis bioreactors has also been considered (Demirel and Scherer 2008). González-Martínez et al. (2016b) studied archaeal and bacterial community dynamics of a bench-scale two-stage anaerobic digester. An overview of the response of key archaeal and bacterial phylotypes to changes in performance parameters is presented in Fig. 1a and 1b, respectively.

In the acidogenic phase, organic matter is biodegraded to VFAs by bacterial communities. During this phase, *Bacteroidetes*, *Chloroflexi*, *Cloacimonetes*, *Firmicutes* and *Proteobacteria* are the predominant phyla. Moreover, *Microthrix* spp. are usually associated with operational dysfunction while *Firmicutes* species in the digesters are important acetogens utilizing simple and complex carbohydrates (Tracy et al. 2012). *Synergistetes* spp. can utilize amino acids as an energy source to produce VFAs for methanogens (Vartoukian et al. 2007), whereas *Proteobacteria* have been recognized as one of the main consumers of VFAs (Ariesyady et al. 2007). Moreover, *Syntrophomonas* strains are present during this phase and are able to syntrophically degrade straight-chain fatty acids (4–8 carbon atoms) into propionate, acetate and methane in co-culture with methanogens (Zhang et al. 2005).

Changes in operational and performance parameters influence bacterial diversity. Hao et al. (2016) found that under high TS conditions, the relative abundance of *Thermoanaerobacteraceae*, *Syntrophomonadaceae*, *Rhodobacteraceae*, *Comamonadaceae* and *Xanthomonadaceae* families were enriched. In contrast, digesters at lower-TS favoured *Syntrophaceae*, *Syntrophobacteraceae*, *Anaerolineaceae*, *Rikenellaceae* and *WCHB01-69* and *Candidatus Cloacamonas* families. Under thermophilic and mesophilic conditions, Guo et al. (2014) found that *Firmicutes* was the common phylum appearing at both temperatures, accounting for 10–20% of relative abundance. *Thermotogae* (60–80% of relative abundance) and *Bacteroidetes* (5–45% of relative abundance) were the dominant taxa under both conditions, respectively. *Proteobacteria* were present in limited amounts and only in thermophilic AD whereas *Synergistetes* appeared in both reactors. Although the relative abundance of *Chloroflexi*, *Actinobacteria* and *Spirochaetes* was higher than that in thermophilic AD, they were poorly represented, accounting for <3% of relative abundance. Finally, *Gelria*, uncultured *Lachnospiraceae*, *Ruminococcaceae Incertae Sedis*, *Sporanaerobacter*, *Tepidanaerobacter*, *Petrobacter* and *Anaerobaculum* were related to performance variations with OLR elevation.

Adaptation of bacterial communities during the start-up stage of thermophilic and mesophilic AD was explored by Wu et al. (2016) and González-Martínez et al. (2016b), respectively. Under thermophilic conditions, the relative abundance of *Firmicutes* increased gradually; on the contrary, *Proteobacteria* and *Thermotogae* decreased. Under mesophilic conditions, the more abundant microorganisms were related to *Clostridiaceae* (21.49%), *Treponema* (5.10%), *Synergistetes* (4.11%) and *Paenibacillaceae* (3.25%) whereas *Cloacamonas* and *Comamonas* were present at >3% abundance only at the beginning of AD, decreasing after that. Zhang et al. (2016a) analysed the microbial community in the anaerobic co-digestion of food waste and sewage sludge. *Firmicutes*, *Proteobacteria*, *Bacteroidetes* and *Actinobacteria* were found as the predominant phyla in the bacterial community. *Firmicutes* increased significantly on day 5 at acidification phase corresponding to VFAs accumulation. After that, the abundance of *Firmicutes* and *Bacteroidetes* increased much more from day 12 at the active methane production phase. *Proteobacteria* and *Actinobacteria* decreased significantly during the experimental period. The greatest changes in these four dominant phyla all appeared on day 5, which could be an indicator of the acidification phase corresponding to VFA accumulation. Hydrolytic bacteria are known to have a lower sensitivity to changes in environmental factors, such as pH and temperature, than methanogens.

Although the role of eukaryotes in performance, predation on bacteria and excess sludge production has been reported during aerobic treatment processes (Ntougias et al. 2011), it is also important to investigate the diversity, roles and functions of eukaryotes in AD. Few authors have reported on diversity and roles/functions in AD (Luo et al. 2005; Matsubayashi et al. 2017). Under mesophilic AD, *Rotifera* and *Phragmoplastophyta* are the most representative phyla and the majority of eukaryal phylotypes belong to Fungi (42.2%), followed by Animalia (28.8%), Protista (13.3%) and finally Plantae (8.9%). In addition, Luo et al.(2005) described the microeukaryotic community in anaerobic sulphide- and sulphur-rich springs, whereas Matsubayashi et al. (2017) constructed clone libraries by sequencing the 18S rRNA gene in anaerobic sludge digesters (Table 1). The latter study suggested that prokaryotic and eukaryotic community structures do not work independently, and that the functional features of both communities are closely related.

Very limited information on the physiology of anaerobic or facultative anaerobic eukaryotic organisms is available to date. Some of the Fungi found in AD contribute to the degradation of some organic matter in anaerobic environments and they could be implicated in the hydrolysis of organic matter in anaerobic sludge digestion processes. Previous studies have demonstrated that phylotypes in Plantae, Animalia and Fungi can produce CH₄ (Liu et al. 2015; Gorrasi et al. 2014).

Regarding the dynamics of the microbiome during AD, contrasting results have been obtained, showing large changes (>25%) from bench-scale mesophilic anaerobic digesters inoculated with sludge from wastewater treatment plants (Schauer-Gimenez et al. 2010; De Vrieze et al. 2013) or high consistency from reactors with an upflow configuration with anaerobic granular biomass (Werner et al. 2011). Given the presence of a wide variety of microorganisms in the influent of AD, dynamic changes in community diversity are likely the result of proliferation of organisms that are adapted to the selective pressures in each bioreactor. However, a core microbiome dominates the reactors, showing the strong selective pressures present in this type of environment (Town et al. 2014; Gonzalez-Martinez et al. 2016b). Maspolim et al. (2015) compared the microbial community dynamics in single-stage and 2-phase anaerobic AD systems treating municipal sludge and the analysis revealed that microbial adaptation occurred as the sludge formed a different community in each reactor at 30 d HRT but no significant microbial changes occurred at lower HRTs. Engineering of the microbiome by adjusting operational parameters leads to a stable microbial structure (Vanwonterghem et al. 2014; De Vrieze et al. 2016). Accurate monitoring of the microbial community requires that the metabolic potential and mode of interaction between the different microorganisms are distinguished from sudden unwanted changes related to unfavourable operational conditions. While generalist microorganisms are able to occupy a broad range of niches based on their greater phenotypic plasticity (van Tienderen 1997), specialists occupy only a limited number of niches and show high levels of specificity. The former can be considered as keepers of process stability (Matias et al. 2013) whereas the latter may drive evolution towards new traits in the microbial community and could be of direct interest in the search for new potential.

The dynamics of prokaryotic organisms have been described during the start-up stage of AD (Gonzalez-Martinez et al. 2016b) as showing substantial changes under unstable conditions. Thus, a challenge exists to develop a useful biomonitoring tool for environmental engineers. Many studies have indicated that *Methanosaeta* and *Methanosarcina* are competitive genera in the AD process. During the acidification phase, *Methanosaeta*, an acetoclastic methanogen, is the dominant genus but its abundance decreases significantly during the methane production phase. In the latter phase, the acetoclastic methanogen *Methanosarcina* increases

significantly. *Methanosarcina* is more tolerant to inhibitors than *Methanosaeta* (Cho et al. 2013). At the end of AD, *Methanoculleus*, a hydrogenotrophic methanogen, becomes dominant because of the exhaustion of acetic acid. Previous studies have reported that for continuous and fed-batch systems, bacterial community dynamics show larger changes than those for the archaeal community, but there is similar diversity, and VFA-producers show greater relative abundance. Generally considered, the hydrolysing bacterial groups *Bacteroides*, *Cloacamonas*, *Clostridiaceae* and *Rikenellaceae* are dominant at the beginning of AD and finally change to other bacterial groups such as *Clostridiaceae*, *Fervidobacterium*, *Paenibacillus* and *Spirochaetes* (Ghasimi et al. 2015;Gonzalez-Martinez et al. 2016b).

Microbial and Eukaryal groups involved in biogas production

AD for methane production has already been widely adopted (Cavinato et al. 2013; Carrere et al. 2016) using methanogenic microorganisms able to utilize simple organic substrates, such as acetate, CO₂/H₂, methanol and formate (de Bok et al. 2004). A deep insight into the main archaeal and bacterial phylotypes of AD involved in biogas production under different operational conditions can be seen in Hao et al. (2016). There are three main types of methanogen, namely acetoclastic, hydrogenotrophic and methylotrophic. Most archaea produce methane by the hydrogenotrophic route and only those belonging to the order *Methanosarcinales* produce it by the acetoclastic route. *Methanobacterium*, *Methanothermobacter*, and *Methanospirillum* are the most commonly identified hydrogenotrophic methanogens in the AD process. Acetoclastic methanogens belong to two genera: *Methanosaeta* and *Methanosarcina* (Venkiteshwaran et al. 2015; Gonzalez-Martinez et al 2016b). *Methanosaeta* can be considered a key methanogen in the AD process, given its unique morphology and physiology (De Vrieze et al. 2012; 2015), catalysing renewable energy production from organic waste streams.

Bacteria can support methane production during the process of methanogenesis by hydrolysis of organic matter. Positive correlation of *Cytophaga*, *Herbaspirillum*, *Symbiobacterium*, *Comamonas* and *Allochromatium* with biogas production has been found (Gonzalez-Martinez et al. 2016b). The genera *Cytophaga* and *Symbiobacterium* are important organic matter degraders in AD in the hydrolysis and acidogenesis processes, respectively (Panichnumsin et al. 2012; Yi et al. 2014).The importance of *Herbaspirillum* sp. remains widely unclear due to its inability to carry out fermentation (Schmid et al. 2006), but its relationship to biogas production (Gonzalez-Martinez et al. 2016b) and degradation of complex organic matter has been reported (Guo et al. 2015).

The role of Eukarya in the production of methane remains largely unknown although Plantae, Animalia and Fungi eukaryal phylotypes have been reported to direct produce CH₄,even in the presence of oxygen (Liu et al. 2015; Gorrasi et al. 2014). However, the mechanisms involved in this pathway remain largely unclear and it has been proposed that CH₄ originates from organic methyl-type compounds in response to environmental stresses. Although it is estimated that plants could contribute around 3–24% to the global CH₄ budget, an estimate of CH₄ production by animals and fungi is still lacking. Consequently, Eukarya are not considered as a CH₄ source by the Intergovernmental Panel on Climate Change (IPCC), and their role in biogas production could be useful for better quantitation of the global CH₄ budget. The influence of rumen fungi for improvement of biogas production from animal manure on anaerobic digesters have gained attention as a biological pre-treatment option of various polymeric substances. These microorganisms are able to effectively digest lignocellulosic compounds, providing energy due to symbiotic associations with rumen microorganisms (Yıldırım et al. 2017). For instance, Gorrasi et

al. (2014) demonstrated the potential application of chitinolytic fungi to obtain H and Ma et al. (2015) determined that rumen microorganisms have higher hydrolytic and acidogenic activity than other microbial species using lignocellulosic biomass as substrates.

Future implications and prospective biotechnologies

New advances in monitoring AD will require the application of control strategies to redirect the microbiome to reach a stable functionality. Until now, microbial process control actions have usually taken place by altering basic operational parameters, such as pH and temperature. For example, increases in AD efficiency were done using different ways: bioaugmentation, as a suitable alternative to increase VFA removal (Town and Dumonceaux 2016) or hydrolysis (Martin-Ryals et al. 2015); microwave (MW) pre-treatment, as an effective way of enhancing biogas production and solids removal (Coelho et al. 2011). However, to engage direct steering of the microbiome to sustain process stability, this knowledge has to be integrated into advanced monitoring and control strategies. For example, the ratio of syntrophic acetate-oxidizing bacteria or methanogenic archaea to total bacteria has been suggested as a possible microbial community monitoring strategy for AD (De Vrieze et al. 2012). This has to be based on specific genes and/or their transcripts, such as the methyl co-enzyme M reductase (*mcrA*) gene for methanogens (Wilkins et al. 2015) and the formyl tetrahydrofolate synthetase (FTHFS) gene for syntrophic acetate-oxidizing bacteria (Akuzawa et al. 2011; Hori et al. 2011).

The study of biogeochemical cycles in natural ecosystems can drive innovation in bioenergetics applications to support improvements of AD. In this sense, Izzo et al. (2014) explored the potentials offered by the structural and functional microbial biodiversity in hypertrophic lagoons characterised by rapid and huge biomass blooms and decomposition. They selected the microbial communities as inoculum and successfully tested for hydrogen production on different kinds of organic wastes.

To decrease the cost of the treatment is of vital importance in AD. This can be achieved by using raw material with lower water content and running the process with a higher dry matter content. The biogas produced can often be utilized to cover the need for process energy. Thus, the economy of a biogas plant is directly linked to the amount of biogas produced per unit of raw material treated in the plant. In short, advanced and direct monitoring of the microbiome is possible through the application of different microbial techniques. Accurate and quick decision tools have to be developed. The integration of existing physicochemical techniques and microbiome-based monitoring is necessary to increase product recovery and the overall energy efficiency of microbial processes.

Compliance with ethical standards

Conflict of interest The authors declare that they have no conflict of interest.

Ethical approval This article does not contain any studies with human participants or animals performed by any of the authors.

References

Ağdağ ON, Sponza DT (2007) Co-digestion of mixed industrial sludge with municipal solid wastes in anaerobic simulated landfilling bioreactors. J Hazard Mater 140:75–85 . doi: 10.1016/J.JHAZMAT.2006.06.059

395 Achinas S, Achinas V, Euverink GJ(2017) A technological overview of biogas production from biowaste.
 396 Engineering 3:299–307. doi:doi.org/10.1016/J.ENG.2017.03.002

397 Akuzawa M, Hori T, Haruta S, Ueno Y, Ishii M, Igarashi Y (2011) Distinctive responses of metabolically active
 398 microbiota to acidification in a thermophilic anaerobic digester. *Env Microbiol* 61:595–605 . doi:
 399 10.1007/s00248-010-9788-1

400 Allison SD, Martiny JBH (2008) Resistance, resilience, and redundancy in microbial communities. *PNAS*
 401 105:11512–11519 . doi: www.pnas.org/cgi/doi/10.1073/pnas.0801925105

402 Amani T, Nosrati M, Sreekrishnan TR, Sreekrishnan TR (2010) Anaerobic digestion from the viewpoint of
 403 microbiological, chemical, and operational aspects — a review. *Environ Rev* 18:255–278 . doi:
 404 10.1139/A10-011

405 Angelidaki I, Ellegaard L, Ahring BK (2003) Applications of the anaerobic digestion process. In: Scheper T (ed)
 406 Biomethanation II. *Advances in biochemical engineering/biotechnology*. Springer, Berlin, Heidelberg,
 407 Berlin, Germany, pp 1–33

408 Appels L, Baeyens J, Degrève J, Dewil R (2008) Principles and potential of the anaerobic digestion of waste-
 409 activated sludge. *Prog Energy Combust Sci* 34:755–781 . doi: 10.1016/J.PECS.2008.06.002

410 Ariesyady HD, Ito T, Okabe S (2007) Functional bacterial and archaeal community structures of major trophic
 411 groups in a full-scale anaerobic sludge digester. *Water Res* 41:1554–1568 . doi:
 412 10.1016/J.WATRES.2006.12.036

413 Bayr S, Rantanen M, Kaparaju P, Rintala J (2012) Mesophilic and thermophilic anaerobic co-digestion of
 414 rendering plant and slaughterhouse wastes. *Bioresour Technol* 104:28–36 . doi:
 415 10.1016/j.biortech.2011.09.104

416 Bergland WH, Dinamarca C, Toradzadegan C, Nordgard ASR, Bakke I, Bakke R(2015) High rate manure
 417 supernatant digestion. *Water Res* 76:1–9. doi: 10.1016/j.watres.2015.02.051

418 Beyene HD, Werkneh AA, Ambaye TG (2018) Current updates on waste to energy (WtE) technologies: a
 419 review. *Renew Energy Focus* 24:1–11 . doi: 10.1016/j.ref.2017.11.001

420 Bräuer SL, Cadillo-Quiroz H, Yashiro E, Yavitt JB, Zinder SH (2006) Isolation of a novel acidiphilic
 421 methanogen from an acidic peat bog. *Nature* 442: . doi: 10.1038/nature04810

422 Calderón K, González-Martínez A, Gómez-Silván C, Osorio F, Rodelas BN, González-López J (2013) Archaeal
 423 diversity in biofilm technologies applied to treat urban and industrial wastewater: Recent advances and
 424 future prospects. *Int J Mol Sci* 14:18572–18598 . doi: 10.3390/ijms140918572

425 Carballa M, Regueiro L, Lema JM (2015) Microbial management of anaerobic digestion: exploiting the
 426 microbiome-functionality nexus. *Curr Opin Biotechnol* 33:103–111 . doi: 10.1016/J.COPBIO.2015.01.008

427 Carballa M, Smits M, Etchebehere C, Boon N, Verstraete W (2011) Correlations between molecular and
 428 operational parameters in continuous lab-scale anaerobic reactors. *Appl Microbiol Biotechnol* 89:303–314
 429 . doi: 10.1007/s00253-010-2858-y

430 Carrere H, Antonopoulou G, Affes R, Passos F, Battimelli A, Lyberatos G, Ferrer I (2016) Review of feedstock

431 pretreatment strategies for improved anaerobic digestion: From lab-scale research to full-scale application.
432 *Bioresour Technol* 199:386–397 . doi: 10.1016/j.biortech.2015.09.007

433 Cavinato C, Bolzonella D, Pavan P, Fatone F, Cecchi F (2013) Mesophilic and thermophilic anaerobic co-
434 digestion of waste activated sludge and source sorted biowaste in pilot- and full-scale reactors. *Renew*
435 *Energy* 55:260–265 . doi: 10.1016/j.renene.2012.12.044

436 Chelliapan S, Wilby T, Yuzir A, Sallis PJ (2011) Influence of organic loading on the performance and microbial
437 community structure of an anaerobic stage reactor treating pharmaceutical wastewater. *Desalination*
438 271:257–264 . doi: 10.1016/j.desal.2010.12.045

439 Chen Y, Cheng JJ, Creamer KS (2008) Inhibition of anaerobic digestion process: A review. *Bioresour Technol*
440 99:4044–4064 . doi: 10.1016/J.BIORTECH.2007.01.057

441 Chen C, Guo WS, Ngo HH, Lee DJ, Tung KL, Jin PK, Wang J, Wu Y. (2016) Challenges in biogas production
442 from anaerobic membrane bioreactors. *Renew Energ* 98:120–134 . doi: 10.1016/j.renene.2016.03.095

443 Cho S-K, Im W-T, Kim D-H, Kim M-H, Shin H-S, Oh S-E (2013) Dry anaerobic digestion of food waste under
444 mesophilic conditions: Performance and methanogenic community analysis. *Bioresour Technol* 131:210–
445 217 . doi: 10.1016/J.BIORTECH.2012.12.100

446 Coelho NMG, Droste RL, Kennedy KJ (2011) Evaluation of continuous mesophilic, thermophilic and
447 temperature phased anaerobic digestion of microwaved activated sludge. *Water Res* 45:2822–2834 . doi:
448 10.1016/j.watres.2011.02.032

449 Dang Y, Sun D, Woodard TL, Wang L-Y, Nevin KP, Holmes DE (2017) Stimulation of the anaerobic digestion
450 of the dry organic fraction of municipal solid waste (OFMSW) with carbon-based conductive materials.
451 *Bioresour Technol* 238:30–38 . doi: 10.1016/j.biortech.2017.04.021

452 de Bok FAM, Plugge CM, Stams AJM (2004) Interspecies electron transfer in methanogenic propionate
453 degrading consortia. *Water Res* 38:1368–1375 . doi: 10.1016/j.watres.2003.11.028

454 De Gioannis G, Muntoni A, Poletini A, Pomi R, Spiga D (2017) Energy recovery from one- and two-stage
455 anaerobic digestion of food waste. *Waste Manag* 68:595–602 . doi: 10.1016/J.WASMAN.2017.06.013

456 de Los Reyes III FL, Weaver JE, Wang L (2015) A methodological framework for linking bioreactor function to
457 microbial communities and environmental conditions. *Curr Opin Biotechnol* 33:112–118 . doi:
458 10.1016/j.copbio.2015.02.002

459 De Vrieze J, Christiaens MER, Verstraete W (2017) The microbiome as engineering tool: Manufacturing and
460 trading between microorganisms. *N Biotechnol* 39:206–214 . doi: 10.1016/j.nbt.2017.07.001

461 De Vrieze J, Hennebel T, Boon N, Verstraete W (2012) *Methanosarcina*: The rediscovered methanogen for
462 heavy duty biomethanation. *Bioresour Technol* 112:1–9 . doi: 10.1016/j.biortech.2012.02.079

463 De Vrieze J, Raport L, Roume H, Vilchez-Vargas R, Auregui RJ, Pieper DH, Boon N (2016) The full-scale
464 anaerobic digestion microbiome is represented by specific marker populations. *Water Res* 104:101–110 .
465 doi: 10.1016/j.watres.2016.08.008

466 De Vrieze J, Saunders AM, He Y, Fang J, Nielsen PH, Verstraete W, Boon N (2015) Ammonia and temperature

determine potential clustering in the anaerobic digestion microbiome. *Water Res* 75:312–323 . doi: 10.1016/J.WATRES.2015.02.025

De Vrieze J, Verstraete W, Boon N (2013) Repeated pulse feeding induces functional stability in anaerobic digestion. *Microb Biotechnol* 6:414–424 . doi: 10.1111/1751-7915.12025

Demirel B, Scherer P (2008) The roles of acetotrophic and hydrogenotrophic methanogens during anaerobic conversion of biomass to methane: a review. *Rev Environ Sci Bio/Technology* 7:173–190 . doi: 10.1007/s11157-008-9131-1

Fezzani B, Cheikh R Ben (2010) Two-phase anaerobic co-digestion of olive mill wastes in semi-continuous digesters at mesophilic temperature. *Bioresour Technol* 101:1628–1634 . doi: 10.1016/j.biortech.2009.09.067

Fitamo T, Treu L, Boldrin A, Sartori C, Angelidaki I, Scheutz C (2017) Microbial population dynamics in urban organic waste anaerobic co-digestion with mixed sludge during a change in feedstock composition and different hydraulic retention times. *Water Res* 118:261–271 . doi: 10.1016/J.WATRES.2017.04.012

Ghasimi DSM, Tao Y, de Kreuk M, Zandvoort MH, van Lier JB (2015) Microbial population dynamics during long-term sludge adaptation of thermophilic and mesophilic sequencing batch digesters treating sewage fine sieved fraction at varying organic loading rates. *Biotechnol Biofuels* 8:171. doi: 10.1186/s13068-015-0355-3

Gómez X, Cuetos MJ, Cara J, Mora A, Garcíá AI (2006) Anaerobic co-digestion of primary sludge and the fruit and vegetable fraction of the municipal solid wastes Conditions for mixing and evaluation of the organic loading rate. *Renew Energy* 31:2017–2024 . doi: 10.1016/j.renene.2005.09.029

Gonzalez-Martinez A, Calderón K, González-López J (2016a) New concepts of microbial treatment processes for the nitrogen removal: effect of protein and amino acids degradation. *Amino Acids* 48:1123–1130 . doi: 10.1007/s00726-016-2185-4

Gonzalez-Martinez A, Garcia-Ruiz MJ, Rodriguez-Sanchez A, Osorio F, Gonzalez-Lopez J (2016b) Archaeal and bacterial community dynamics and bioprocess performance of a bench-scale two-stage anaerobic digester. *Appl Microbiol Biotechnol* 100:6013–6033 . doi: 10.1007/s00253-016-7393-z

Gonzalez-Martinez A, Muñoz-Palazon B, Rodriguez-Sanchez A, Maza-Márquez P, Mikola A, Gonzalez-Lopez J, Vahala R (2017) Start-up and operation of an aerobic granular sludge system under low working temperature inoculated with cold-adapted activated sludge from Finland. *Bioresour Technol* 239:180–189 . doi: 10.1016/j.biortech.2017.05.037

Gorrasi S, Izzo G, Massini G, Signorini A, Bargini P, Fenice M (2014) From polluting seafood wastes to energy. Production of hydrogen and methane from raw chitin material by a two-phase process. *J Environ Prot Ecol* 75:526–36.

Goswami R, Chattopadhyay P, Shome A, Banerjee SN, Chakraborty AK, Mathew AK, Chaudhury, S (2016) An overview of physico-chemical mechanisms of biogas production by microbial communities: a step towards sustainable waste management. *3 Biotech* 6:72 . doi: 10.1007/s13205-016-0395-9

503 Gou C, Yang Z, Huang J, Wang H, Xu H, Wang L (2014) Effects of temperature and organic loading rate on the
504 performance and microbial community of anaerobic co-digestion of waste activated sludge and food waste.
505 *Chemosphere* 105:146–151 . doi: 10.1016/j.chemosphere.2014.01.018

506 Guo J, Peng Y, Ni B-J, Han X, Fan L, Yuan Z (2015) Dissecting microbial community structure and methane-
507 producing pathways of a full-scale anaerobic reactor digesting activated sludge from wastewater treatment
508 by metagenomic sequencing. *Microb Cell Fact* 14:33–44 . doi: 10.1186/s12934-015-0218-4

509 Guo X, Wang C, Sun F, Zhu W, Wu W (2014) A comparison of microbial characteristics between the
510 thermophilic and mesophilic anaerobic digesters exposed to elevated food waste loadings. *Bioresour*
511 *Technol* 152:420–428 . doi: 10.1016/j.biortech.2013.11.012

512 Hao L, Bize A, Conteau D, Chapleur O, Courtois S, Kroff P, Desmond-LeQuémener E, Bouchez T, Mazeas L
513 (2016) New insights into the key microbial phylotypes of anaerobic sludge digesters under different
514 operational conditions. *Water Res* 102:158–169 . doi: 10.1016/j.watres.2016.06.014

515 Hernández MA, Susa MR, Andres Y (2014) Use of coffee mucilage as a new substrate for hydrogen production
516 in anaerobic co-digestion with swine manure. *Bioresour Technol* 168:112–118 . doi:
517 10.1016/j.biortech.2014.02.101

518 Hori T, Sasaki D, Haruta S, Shigematsu T, Ueno Y, Ishii M, Igarashi Y (2011) Detection of active, potentially
519 acetate-oxidizing syntrophs in an anaerobic digester by flux measurement and formyltetrahydrofolate
520 synthetase (FTHFS) expression profiling. *Microbiology* 157:1980–1989 . doi: 10.1099/mic.0.049189-0

521 Izzo G, Rosa S, Massini G, Patriarca C, Fenice M, Fiocchetti F, Marone A, Varrone C, Signorini A (2014) From
522 hypertrophic lagoons to bioenergy production. *J Env Prot Ecol* 15:537-546.

523 Kaever A, Landesfeind M, Feussner K, Morgenstern B, Feussner I, Meinicke P, Gill AC (2014) Meta-Analysis
524 of Pathway Enrichment: Combining Independent and Dependent Omics Data Sets. *PLoS One* 9:e89297 .
525 doi: 10.1371/journal.pone.0089297

526 Khalid A, Arshad M, Anjum M, Mahmood T, Dawson L (2011) The anaerobic digestion of solid organic waste.
527 *Waste Manag* 31:1737–1744 . doi: 10.1016/J.WASMAN.2011.03.021

528 Kim J, Novak JT, Higgins MJ (2011) Multi-staged anaerobic sludge digestion processes. *J Environ Eng*
529 137:0000372. doi: 10.1061/(ASCE)EE.1943-7870.0000372

530 Kundu K, Bergmann I, Hahnke S, Klocke M, Sharma S, Sreekrishnan TR (2013) Carbon source — A strong
531 determinant of microbial community structure and performance of an anaerobic reactor. *J Biotechnol*
532 168:616–624 . doi: 10.1016/j.jbiotec.2013.08.023

533 Kundu K, Sharma S, Sreekrishnan TR (2017) Influence of process parameters on anaerobic digestion
534 microbiome in bioenergy production: Towards an improved understanding. *Bioenergy Res* 10:288–303 .
535 doi: 10.1007/s12155-016-9789-0

536 Kundu K, Sharma S, Sreekrishnan TR (2012) Effect of operating temperatures on the microbial community
537 profiles in a high cell density hybrid anaerobic bioreactor. *Bioresour Technol* 118:502–511 . doi:
538 10.1016/j.biortech.2012.05.047

539 Lin J, Zuo J, Ji R, Chen X, Liu F, Wang K, Yang Y (2012) Methanogenic community dynamics in anaerobic co-
540 digestion of fruit and vegetable waste and food waste. *J Environ Sci* 24:1288–1294 . doi: 10.1016/S1001-
541 0742(11)60927-3

542 Lindeboom REF, Fermoso FG, Weijma J, Zagt K, van Lier JB (2011) Autogenerative high pressure digestion:
543 Anaerobic digestion and biogas upgrading in a single step reactor system. *Water Sci Technol* 64:647–53.
544 doi: 10.2166/wst.2011.664

545 Liu C, Wang W, Anwar N, Ma Z, Liu G, Zhang R (2017) Effect of Organic Loading Rate on Anaerobic
546 Digestion of Food Waste under Mesophilic and Thermophilic Conditions. *Energy & Fuels* 31:2976–2984 .
547 doi: 10.1021/acs.energyfuels.7b00018

548 Liu J, Chen H, Zhu Q, Shen Y, Wang X, Wang M, Peng C (2015) A novel pathway of direct methane production
549 and emission by eukaryotes including plants, animals and fungi: An overview. *Atmos Environ* 115:26–35 .
550 doi: 10.1016/j.atmosenv.2015.05.019

551 Liu Z, Si B, Li J, He J, Zhang C, Lu Y, Zhang Y, Xing XH (2018) Bioprocess engineering for biohythane
552 production from low-grade waste biomass: technical challenges towards scale up. *Curr Opin Biotechnol*
553 50:25–31 . doi: 10.1016/j.copbio.2017.08.014

554 Lo HM, Kurniawan TA, Sillanpää MET, Pai TY, Chiang CF, Chao KP, Liu MH, Chuang SH, Banks CJ, Wang
555 SC, Lin KC, Lin CY, Liu WF, Cheng PH, Chen CK, Chiu HY, Wu HY (2010) Modeling biogas
556 production from organic fraction of MSW co-digested with MSWI ashes in anaerobic bioreactors.
557 *Bioresour Technol* 101:6329–6335 . doi: 10.1016/j.biortech.2010.03.048

558 Luo Q, Krumholz LR, Najjar FZ, Peacock AD, Roe BA, White DC, Elshahed MS (2005) Diversity of the
559 microeukaryotic community in sulfide-rich zodletone spring (Oklahoma). *Appl Environ Microbiol*
560 71:6175–6184 . doi: 10.1128/AEM.71.10.6175–6184.2005

561 Ma J, Zhao QB, Laurens LL, Jarvis EE, Nagle NJ, Chen S, Frear CS(2015)Mechanism, kinetics and
562 microbiology of inhibition caused by long-chain fatty acids in anaerobic digestion of algal biomass.
563 *Biotechnol Biofuels* 8:141 .doi: 10.1186/s13068-015-0322-z

564 Mao C, Feng Y, Wang X, Ren G (2015) Review on research achievements of biogas from anaerobic digestion.
565 *Renew Sustain Energy Rev* 45:540–555 . doi: 10.1016/J.RSER.2015.02.032

566 Martin-Ryals A, Schideman L, Li P, Wilkinson H, Wagner R (2015) Improving anaerobic digestion of a
567 cellulosic waste via routine bioaugmentation with cellulolytic microorganisms. *Bioresour Technol* 189:62–
568 70 . doi: 10.1016/j.biortech.2015.03.069

569 Maspolim Y, Zhou Y, Guo C, Xiao K, Ng WJ (2015) Comparison of single-stage and two-phase anaerobic
570 sludge digestion systems–Performance and microbial community dynamics. *Chemosphere* 140:54–62.
571 doi:10.1016/j.chemosphere.2014.07.028

572 Matias MG, Combe M, Barbera C, Mouquet N (2013) Ecological strategies shape the insurance potential of
573 biodiversity. *Front Microbiol* 3:3–9 . doi: 10.3389/fmicb.2012.00432

574 Matsubayashi M, Shimada Y, Li YY, Harada H, Kubota K (2017) Phylogenetic diversity and in situ detection of

575 eukaryotes in anaerobic sludge digesters. *PLoS One* 12:1–13 . doi: 10.1371/journal.pone.0172888

576 Maus I, Koeck DE, Cibis KG, Hahnke S, Kim YS, Langer T, Kreubel J, Erhard M, Bremges A, Off S, Stolze Y,
577 Jaenicke S, Goesmann A, Sczyrba A, Scherer P, König H, Schwarz WH, Zverlov V V, Liebl W, Pühler A,
578 Schlüter A, Klocke M (2016) Unraveling the microbiome of a thermophilic biogas plant by metagenome
579 and metatranscriptome analysis complemented by characterization of bacterial and archaeal isolates.
580 *Biotechnol Biofuels* 9:1–28 . doi: 10.1186/s13068-016-0581-3

581 McHugh S, Carton M, Mahony T, O'Flaherty V (2003) Methanogenic population structure in a variety of
582 anaerobic bioreactors. *FEMS Microbiol Lett* 219: 297–304 . doi:10.1016/S0378-1097(03)00055-7

583 Meulepas RJW, Jagersma CG, Khadem AF, Stams AJM, Lens PNL (2010) Effect of methanogenic substrates on
584 anaerobic oxidation of methane and sulfate reduction by an anaerobic methanotrophic enrichment. *Appl*
585 *Microbiol Biotechnol* 87:1499–1506 . doi: 10.1007/s00253-010-2597-0

586 Møller J, Boldrin A, Christensen TH (2009) Anaerobic digestion and digestate use: Accounting of greenhouse
587 gases and global warming contribution. *Waste Manag Res* 27:813–824 . doi: 10.1177/0734242X09344876

588 Muñoz-Palazon B, Rodriguez-Sanchez A, Castellano-Hinojosa A, Gonzalez-Lopez J, van Loosdrecht MCM,
589 Vahala R, Gonzalez-Martinez A (2018) Quantitative and qualitative studies of microorganisms involved in
590 full-scale autotrophic nitrogen removal performance. *AIChE J* 64:457–467 . doi: 10.1002/aic.15925

591 Nizami A-S, Korres NE, Murphy JD (2009) Review of the Integrated Process for the Production of Grass
592 Biomethane. *Environ Sci Technol* 43:8496–8508 . doi: 10.1021/es901533j

593 Ntougias S, Tanasidis S, Melidis P (2011) Microfaunal indicators, Ciliophora phylogeny and protozoan
594 population shifts in an intermittently aerated and fed bioreactor. *J Hazard Mater* 186:1862–1869 . doi:
595 10.1016/j.jhazmat.2010.12.099

596 Oslaj M, Mursec B, Vindis P (2010) Biogas production from maize hybrids. *Biomass and Bioenergy* 34:1538–
597 1545 . doi: 10.1016/j.biombioe.2010.04.016

598 Padmasiri SI, Zhang J, Fitch M, Norddahl B, Morgenroth E, Raskin L (2007) Methanogenic population
599 dynamics and performance of an anaerobic membrane bioreactor (AnMBR) treating swine manure under
600 high shear conditions. *Water Res* 41:134–144 . doi: 10.1016/j.watres.2006.09.021

601 Panichnumsin P, Ahring B, Nopharatana A, Chaiprasert P (2012) Microbial community structure and
602 performance of an anaerobic reactor digesting cassava pulp and pig manure. *Water Sci Technol* 66:1590 .
603 doi: 10.2166/wst.2012.358

604 Park KY, Jang HM, Park M-R, Lee K, Kim D, Kim YM (2016) Combination of different substrates to improve
605 anaerobic digestion of sewage sludge in a wastewater treatment plant. *Int Biodeterior Biodegradation*
606 109:73–77 . doi: 10.1016/J.IBIOD.2016.01.006

607 Pohland FG, Ghosh S (1971) Developments in anaerobic stabilization of organic wastes - The two-phase
608 concept. *Environ Lett* 1:255–266 . doi: 10.1080/00139307109434990

609 Pozo C, Martínez-Toledo M V., Rodelas B, González-López J (2002) Effects of culture conditions on the
610 production of polyhydroxyalkanoates by *Azotobacter chroococcum* H23 in media containing a high

611 concentration of alpechín (wastewater from olive oil mills) as primary carbon source. *J Biotechnol*
612 97:125–131 . doi: 10.1016/S0168-1656(02)00056-1

613 Rajagopal R, Massé DI, Singh G (2013) A critical review on inhibition of anaerobic digestion process by excess
614 ammonia. *Bioresour Technol* 143:632–641 . doi: 10.1016/J.BIORTECH.2013.06.030

615 Rapport JL, Zhang R, Williams RB, Jenkins BM (2012) Anaerobic Digestion technologies for the treatment of
616 Municipal Solid Waste. *Int J Environ Waste Manag* 9:100 . doi: 10.1504/IJEW.2012.044163

617 Regueiro L, Veiga P, Figueroa M, Lema JM, Carballa M (2014) Influence of transitional states on the microbial
618 ecology of anaerobic digesters treating solid wastes. *Appl Microbiol Biotechnol* 98:2015–2027 . doi:
619 10.1007/s00253-013-5378-8

620 REN21 (2016) Renewables 2016 global status report 2016. REN21, Paris

621 Rolando C, Elba V, Carlos R (2011) Anaerobic mono-digestion of turkey manure: Efficient revaluation to obtain
622 methane and soil conditioner. *J Water Resource Prot T*: 3:584–589 . doi: 10.4236/jwarp.2011.38067

623 Sasaki K, Hirano S, Morita M, Sasaki D, Matsumoto N, Ohmura N, Igarashi Y (2011) Bioelectrochemical
624 system accelerates microbial growth and degradation of filter paper. *Appl Microbiol Biotechnol* 89:449–
625 455 . doi: 10.1007/s00253-010-2972-x

626 Schauer-Gimenez AE, Zitomer DH, Maki JS, Struble CA (2010) Bioaugmentation for improved recovery of
627 anaerobic digesters after toxicant exposure. *Water Res* 44:3555–3564 . doi:
628 10.1016/J.WATRES.2010.03.037

629 Schmid M, Baldani JJ, Hartmann A (2006) The Genus *Herbaspirillum*. In: *The Prokaryotes*. Springer New York,
630 New York, NY, pp 141–150

631 Shen Y, Forrester S, Koval J, Urgun-Demirtas M (2017) Yearlong semi-continuous operation of thermophilic
632 two-stage anaerobic digesters amended with biochar for enhanced biomethane production. *J Clean Prod*
633 167:863–874 . doi: 10.1016/J.JCLEPRO.2017.05.135

634 Smith AM, Sharma D, Lappin-Scott H, Burton S, Huber DH (2014) Microbial community structure of a pilot-
635 scale thermophilic anaerobic digester treating poultry litter. *Appl Microbiol Biotechnol* 98:2321–2334 .
636 doi: 10.1007/s00253-013-5144-y

637 Sun Q, Li H, Yan J, Liu L, Yu Z, Yu X (2015) Selection of appropriate biogas upgrading technology-a review of
638 biogas cleaning, upgrading and utilisation. *Renew Sustain Energy Rev* 51:521–532 . doi:
639 10.1016/j.rser.2015.06.029

640 Sunyoto NMS, Zhu M, Zhang Z, Zhang D (2016) Effect of biochar addition on hydrogen and methane
641 production in two-phase anaerobic digestion of aqueous carbohydrates food waste. *Bioresour Technol*
642 219:29–36 . doi: 10.1016/J.BIORTECH.2016.07.089

643 Surendra K, Takara D, Hashimoto AG, Kumar Khanal S (2014) Biogas as a sustainable energy source for
644 developing countries: Opportunities and challenges. *Renew Sustain Energy Rev* 31:846–859 . doi:
645 10.1016/j.rser.2013.12.015

646 Town JR, Dumonceaux TJ (2016) Laboratory-scale bioaugmentation relieves acetate accumulation and

stimulates methane production in stalled anaerobic digesters. *Appl Microbiol Biotechnol* 100:1009–1017 .
doi: 10.1007/s00253-015-7058-3

Town JR, Links MG, Fonstad TA, Dumonceaux TJ (2014) Molecular characterization of anaerobic digester
microbial communities identifies microorganisms that correlate to reactor performance. *Bioresour Technol*
151:249–257 . doi: 10.1016/J.BIORTECH.2013.10.070

Tracy BP, Jones SW, Fast AG, Indurthi DC, Papoutsakis ET (2012) Clostridia: the importance of their
exceptional substrate and metabolite diversity for biofuel and biorefinery applications. *Curr Opin*
Biotechnol 23:364–381 . doi: 10.1016/J.COPBIO.2011.10.008

Tyagi VK, Lo S-L (2013) Sludge: A waste or renewable source for energy and resources recovery? *Renew*
Sustain Energy Rev 25:708–728 . doi: 10.1016/J.RSER.2013.05.029

van Tienderen PH (1997) Generalists, specialists, and the evolution of phenotypic plasticity in sympatric
populations of distinct species. *Evolution (N Y)* 51:1372–1380 . doi: 10.1111/j.1558-5646.1997.tb01460.x

Vanwonterghem I, Jensen PD, Ho DP, Batstone DJ, Tyson GW (2014) Linking microbial community structure,
interactions and function in anaerobic digesters using new molecular techniques. *Curr Opin Biotechnol*
27:55–64 . doi: 10.1016/J.COPBIO.2013.11.004

Vartoukian SR, Palmer RM, Wade WG (2007) The division “Synergistes.” *Anaerobe* 13:99–106 . doi:
10.1016/J.ANAEROBE.2007.05.004

Vasco-Correa J, Khanal S, Manandhar A, Shah A (2018) Anaerobic digestion for bioenergy production: Global
status, environmental and techno-economic implications, and government policies. *Bioresour Technol*
247:1015–1026 . doi: 10.1016/J.BIORTECH.2017.09.004

Venkateshwaran K, Bocher B, Maki J, Zitomer D (2015) Relating Anaerobic Digestion Microbial Community
and Process Function. *Microbiol insights* 8:37–44 . doi: 10.4137/MBI.S33593

Wang P, Wang H, Qiu Y, Ren L, Jiang B (2018) Microbial characteristics in anaerobic digestion process of food
waste for methane production-A review. *Bioresour Technol* 248:29–36 . doi:
10.1016/j.biortech.2017.06.152

Werner JJ, Knights D, Garcia ML, Scalfone NB, Smith S, Yarasheski K, Cummings TA, Beers AR, Knight R,
Angenent LT (2011) Bacterial community structures are unique and resilient in full-scale bioenergy
systems. *Proc Natl Acad Sci* 108:4158–4163 . doi: 10.1073/pnas.1015676108

Wilkins D, Lu X-Y, Shen Z, Chen J, Lee PKH (2015) Pyrosequencing of *mcrA* and archaeal 16S rRNA genes
reveals diversity and substrate preferences of methanogen communities in anaerobic digesters. *Appl*
Environ Microbiol 81:604–613 . doi: 10.1128/AEM.02566-14

Wojcieszak M, Pyzik A, Poszytek K, Krawczyk PS, Sobczak A, Lipinski L, Roubinek O, Palige J, Sklodowska
A, Drewniak L (2017) Adaptation of methanogenic inocula to anaerobic digestion of maize silage. *Front*
Microbiol 8:1–12 . doi: 10.3389/fmicb.2017.01881

Wu B, Wang X, Deng Y-Y, He X-L, Li Z-W, Li Q, Qin H, Chen J-T, He M-X, Zhang M, Hu G-Q, Yin X-B
(2016) Adaption of microbial community during the start-up stage of a thermophilic anaerobic digester

- treating food waste. *Biosci Biotechnol Biochem* 80:2025–2032 . doi: 10.1080/09168451.2016.1191326
- Yamada T, Sekiguchi Y, Hanada S, Imachi H, Ohashi A, Harada H, Kamagata Y (2006) *Anaerolinea thermolimos* sp. nov., *Levilinea saccharolytica* gen. nov., sp. nov. and *Leptolinea tardivitalis* gen. nov., sp. nov., novel filamentous anaerobes, and description of the new classes *Anaerolineae* classis nov. and *Caldilineae* classis nov. in the. *Int J Syst Evol Microbiol* 56:1331–1340 . doi: 10.1099/ijs.0.64169-0
- Yi J, Dong B, Jin J, Dai X (2014) Effect of increasing total solids contents on anaerobic digestion of food waste under mesophilic conditions: performance and microbial characteristics analysis. *PLoS One* 9:e102548 . doi: 10.1371/journal.pone.0102548
- Yıldırım E, Ince O, Aydın S, Ince B (2017) Improvement of biogas potential of anaerobic digesters using rumen fungi. *Renewable Energy* 109:346–353 . doi: 10.1016/j.renene.2017.03.021
- Yousuf A, Khan MR, Pirozzi D, Ab Wahid Z (2016) Financial sustainability of biogas technology: Barriers, opportunities, and solutions. *Energy Sources, Part B Econ Planning, Policy* 11:841–848 . doi: 10.1080/15567249.2016.1148084
- Zhang C, Liu X, Dong X (2005) *Syntrophomonas erecta* sp. nov., a novel anaerobe that syntrophically degrades short-chain fatty acids. *Int J Syst Evol Microbiol* 55:799–803 . doi: 10.1099/ijs.0.63372-0
- Zhang D, Zhu M, Zhou W, Yani S, Zhang Z, Wu J, Zhang D, Zhu M, Zhou W, Yani S, Zhang Z, Wu J (2015) A two-phase anaerobic digestion process for biogas production for combined heat and power generation for remote communities. In: *Handbook of Clean Energy Systems*. John Wiley & Sons, Ltd, Chichester, UK, pp 1–17
- Zhang D, Zhu W, Tang C, Suo Y, Gao L, Yuan X, Wang X, Cui Z (2012) Bioreactor performance and methanogenic population dynamics in a low-temperature (5–18 °C) anaerobic fixed-bed reactor. *Bioresour Technol* 104:136–143 . doi: 10.1016/J.BIORTECH.2011.10.086
- Zhang J, Loh K-C, Lee J, Wang C-H, Dai Y, Tong YW (2017) Three-stage anaerobic co-digestion of food waste and horse manure. *Sci Rep* 7:1269–1278 . doi: 10.1038/s41598-017-01408-w
- Zhang J, Lv C, Tong J, Liu J, Liu J, Yu D, Wang Y, Chen M, Wei Y (2016a) Optimization and microbial community analysis of anaerobic co-digestion of food waste and sewage sludge based on microwave pretreatment. *Bioresour Technol* 200:253–261 . doi: 10.1016/J.BIORTECH.2015.10.037
- Zhang Q, Hu J, Lee D-J (2016b) Biogas from anaerobic digestion processes: Research updates. *Renew Energy* 98:108–119 . doi: 10.1016/J.RENENE.2016.02.029
- Ziganshin AM, Liebetrau J, Pröter J, Kleinstüber S (2013) Microbial community structure and dynamics during anaerobic digestion of various agricultural waste materials. *Appl Microbiol Biotechnol* 97:5161–5174 . doi: 10.1007/s00253-013-4867-0

716 **Table 1** Main eukaryal phylotypes found in anaerobic digesters. Data were taken from Matsubayashi et al. (2017).

<i>Kingdom/ Superphylum</i>	<i>Phylum</i>	
<i>Alveolata</i>	<i>Perkinsozoa</i>	<i>A31</i>
<i>Amoebozoa</i>	<i>Discosea</i>	<i>Order Dactylopodida</i>
	<i>Gastrotricha</i>	<i>Chaetonotus cf.</i>
<i>Animalia</i>	<i>Gastrotricha</i>	<i>Chaetonotus cf.</i>
<i>Archaeplastida</i>	<i>Chlorophyta</i>	<i>ANI-3</i>
	<i>Chlorophyta</i>	<i>Family Chlorellaceae</i>
	<i>Chlorophyta</i>	<i>Prototheca zopfi</i>
	<i>Ciliophora</i>	<i>Acaryophrya sp.</i>
	<i>Ciliophora</i>	<i>Vorticellides aquadulcis</i>
<i>Fungi</i>	<i>Arthropoda</i>	<i>Allonothrus sp.</i>
	<i>Arthropoda</i>	<i>Boletoglyphus sp.</i>
	<i>Arthropoda</i>	<i>Naidacarus arboricola</i>
	<i>Arthropoda</i>	<i>Rhizoglyphus sp.</i>
	<i>Ascomycota</i>	<i>Candida sp.</i>
	<i>Ascomycota</i>	<i>Exophiala equine</i>
	<i>Ascomycota</i>	<i>Family Dipodascaceae</i>
	<i>Ascomycota</i>	<i>Penicillium chrysogenum</i>
	<i>Ascomycota</i>	<i>Phoma sp.</i>
	<i>Ascomycota</i>	<i>Xenobotrytis sp.</i>
	<i>Basidiomycota</i>	<i>Lentinus sp.</i>
	<i>Basidiomycota</i>	<i>Trichosporum cutaneum</i>
	<i>Cryptomycota</i>	<i>LKM11</i>
	<i>Cryptomycota</i>	<i>LKM15</i>
<i>Metazoa</i>	<i>Platyhelminthes</i>	<i>Gieysztoria sp.</i>
	<i>Rotifera</i>	<i>Brachionus calyciflorus</i>
<i>Rhizaria</i>	<i>Cercozoa</i>	<i>Rhogostoma minus</i>
<i>Stramenopiles</i>	<i>Hyphochytriomycetes</i>	<i>Rhizidiomyces apophysatus</i>

717

718 **Figure legends**

719

720 **Fig. 1** Multivariate redundancy analyses relating performance parameters (dried sludge, volatile dried sludge,
721 pH, acid/alkalinity ratio AC/AL, O₂, CO₂ CH₄ and biogas production) with changes in diversity or abundance of
722 the most representative archaeal **(a)** and bacterial **(b)** phylotypes in anaerobic digestion. Data were taken from
723 González-Martínez et al. (2016b)