

AperTO - Archivio Istituzionale Open Access dell'Università di Torino

On the inability of Gaussian process regression to optimally learn compositional functions

This is the author's manuscript

Original Citation:

Availability:

This version is available <http://hdl.handle.net/2318/1901493> since 2023-04-26T20:38:38Z

Publisher:

Curran Associates, Inc.

Terms of use:

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)

On the inability of Gaussian process regression to optimally learn compositional functions

Matteo Giordano*
Department of Statistics
University of Oxford
matteo.giordano@stats.ox.ac.uk

Kolyan Ray*
Department of Mathematics
Imperial College London
kolyan.ray@imperial.ac.uk

Johannes Schmidt-Hieber*
Department of Applied Mathematics
University of Twente
a.j.schmidt-hieber@utwente.nl

Abstract

We rigorously prove that deep Gaussian process priors can outperform Gaussian process priors if the target function has a compositional structure. To this end, we study information-theoretic lower bounds for posterior contraction rates for Gaussian process regression in a continuous regression model. We show that if the true function is a generalized additive function, then the posterior based on any mean-zero Gaussian process can only recover the truth at a rate that is strictly slower than the minimax rate by a factor that is polynomially suboptimal in the sample size n .

1 Introduction

Hierarchical methods are at the heart of modern machine learning and provide state of the art performance for a variety of different problems. While the role of the depth in deep neural networks has been extensively investigated [40, 47, 48, 27, 36], the theory underlying deep Gaussian process priors is still in its infancy. In this work we identify a structured recovery problem for which deep Gaussian process priors are known to achieve near optimal behavior. By proving that for this problem any Gaussian process prior will lead to a sub-optimal posterior shrinkage behavior, we provide a theoretical justification for the use of deep methods for Bayesians.

Gaussian process (GP) priors are arguably the most widely used Bayesian nonparametric priors in machine learning, having found success in multiple settings [29]. Reasons for their popularity include their interpretability, ability to flexibly model target functions, model spatial or temporal correlations, incorporate prior knowledge such as stationarity, periodicity or smoothness, and their provision of uncertainty quantification. This work establishes information-theoretic bounds on the inability of GPs to optimally model generalized additive models, thereby quantifying certain limitations of GP methods.

We will study the performance of GPs in a continuous analogue of nonparametric regression via quantifying the speed of posterior concentration/contraction around the true regression function. This is a frequentist assessment that measures the typical behaviour of the posterior distribution under the true generative model. The performance of Bayesian methods is known to be sensitive to the choice of prior, particularly in high- and infinite-dimensional settings [12, 22], and it is therefore

*All authors contributed equally.

crucial to understand under what conditions and prior calibrations Bayesian inference is reliable. The frequentist analysis of Bayesian methods has become a standard tool for such tasks in both the statistics and machine learning literatures (e.g. [39, 28, 46, 45, 6, 30, 31]), see the monograph [14]. Precise definitions and further discussion are provided below.

If the regression function is a generalized additive model, it has been shown that deep neural networks with properly chosen network architecture achieve the optimal minimax rate of convergence [35]. We show here that for this structural constraint, *any* mean-zero Gaussian process, irrespective of the choice of covariance kernel, will concentrate around the truth at a rate that is strictly slower than the minimax rate by a factor that is polynomially suboptimal in the sample size n . Thus any posterior GP places a significant portion of its posterior probability on functions that are further from the truth in L^2 -distance than the minimax rate of estimation, which has negative implications for both Bayesian estimation and uncertainty quantification.

In contrast, it has recently been shown that deep Gaussian processes [26, 9], which consist of iterations of Gaussian processes and can be viewed as a Bayesian analogue of deep neural networks, are able to attain the minimax rate of contraction for arbitrary compositional classes [11]. As these compositional classes contain generalized additive models as a special case, we have shown that deep GPs provably outperform GPs from a statistical perspective and should thus be preferred.

In related work, it has been established that (not necessarily Bayesian) linear estimators can only attain polynomially suboptimal *linear estimation rates* over certain spatially inhomogeneous Besov classes [10] or for functions with discontinuities [23]. Agapiou and Wang [1] use the results of [10] to show that GPs also cannot concentrate faster than the linear rate for such Besov classes, see also [34] for some related examples. Castillo [4] establishes lower bounds for concentration rates in terms of abstract conditions on the ‘concentration function’ of the GP, which can be verified for certain specific GP choices. The lower bound ideas of [10, 23] have also been extended to various spatially inhomogeneous and discontinuous functions settings to demonstrate the theoretical advantages of deep learning over linear estimators [17–19, 37, 38, 41]. Finally, for generalized additive models, [35] shows that (possibly nonlinear) wavelet methods can only achieve suboptimal rates.

Most of the above works exploit the results from [10, 23] showing that minimax rates for linear estimators are slower than full minimax rates in these settings. Since linear rates are unavailable for generalized additive functions, and seem difficult to prove using existing techniques, we instead use a different and novel proof approach. We exploit the specific structure of the GP posterior mean in regression to construct a direct lower bound for its prediction risk. In an effort to develop the new tools needed to solve such problems, we further provide a second proof in the supplement that involves first reducing the regression setting to a simplified sparse sequence model in which we derive linear rates. While these do not imply linear rates in the full model we consider, they prove sufficient to construct lower bounds for arbitrary GP posterior means.

2 Main results

2.1 Problem setup and posterior contraction rates

In the usual multivariate nonparametric regression model we observe n training samples $(X_1, Y_1), \dots, (X_n, Y_n)$ arising as

$$Y_i = f(X_i) + w_i, \quad w_i \stackrel{iid}{\sim} N(0, 1), \quad (1)$$

where the design points X_i are either independent and uniformly distributed random variables on $[0, 1]^d$ or equally spaced lattice points in $[0, 1]^d$. To simplify technical arguments due to the discretization and provide a clearer exposition, a standard approach in statistical theory is to consider instead the ‘continuous’ analogue of this model [21]. The discrete model (1) is asymptotically equivalent [33] (as $n \rightarrow \infty$) to observing a realisation $Y = Y^n = (Y_x : x \in [0, 1]^d)$ of the multidimensional Gaussian white noise model

$$dY_x = f(x)dx + \frac{1}{\sqrt{n}}dW_x, \quad x \in [0, 1]^d, \quad (2)$$

where $W = (W_x : x \in [0, 1]^d)$ is a d -dimensional Brownian motion. For large n , the two models thus behave identically from a statistical perspective.

Notation: For a domain D and $1 \leq p \leq \infty$, denote by $L^p(D)$ the space of all measurable functions $f : D \rightarrow \mathbb{R}$ that have finite norm $\|f\|_{L^p(D)} := (\int |f(u)|^p du)^{1/p}$ if $p < \infty$, and are essentially bounded on D if $p = \infty$. If there is no ambiguity, we write L^2 for $L^2[0, 1]^d$ and $\langle f, g \rangle_2 = \int_{[0, 1]^d} f(u)g(u)du$ for the corresponding inner product. Let $P_f = P_f^n$ denote the probability distribution of $Y = Y^n$ arising from (2) with corresponding expectation E_f .

We consider the Bayesian setting where we assign to f a (possibly n -dependent) mean-zero Gaussian process prior $\Pi = \Pi_n$ on $L^2[0, 1]^d$ with covariance kernel K , i.e. Π is a (Borel) Gaussian probability measure on $L^2[0, 1]^d$ such that for $f \sim \Pi$, we have

$$E_f(x) = 0, \quad E[f(x)f(y)] = K(x, y), \quad x, y \in [0, 1]^d. \quad (3)$$

The posterior $\Pi_n(\cdot|Y)$ is then computed as usual using Bayes formula and is again a GP by conjugacy, exactly as in the discrete model (1) [29], see (7) for the exact expression. We make the following frequentist assumption:

Assumption 1. *There is a true $f_0 \in L^2[0, 1]^d$ generating the data $Y \sim P_{f_0}$ according to (2).*

We study the behaviour of the posterior distribution $\Pi_n(\cdot|Y)$ under Assumption 1, in which case it can be treated as a random probability distribution whose (frequentist) randomness depends on Y . We next introduce the notion of uniform posterior contraction.

Definition 1. *We say that the posterior contracts about f_0 at rate $\varepsilon_n \rightarrow 0$ uniformly over a sequence of classes $\mathcal{F}_n \subset L^2[0, 1]^d$ if as $n \rightarrow \infty$*

$$\sup_{f_0 \in \mathcal{F}_n} E_{f_0} \Pi_n(f : \|f - f_0\|_{L^2} \geq \varepsilon_n | Y) \rightarrow 0.$$

Posterior contraction is often stated in the equivalent form

$$\Pi_n(f : \|f - f_0\|_{L^2} \geq \varepsilon_n | Y) \rightarrow 0 \quad \text{in } P_{f_0}\text{-probability}$$

as $n \rightarrow \infty$, which says that the posterior puts all but a vanishingly small amount of probability on L^2 -balls of shrinking radius ε_n about the true f_0 generating the data [14]. We are typically interested in the size of the *smallest* such L^2 -ball, i.e. the fastest rate $\varepsilon_n \rightarrow 0$ such that Definition 1 holds. Such results not only quantify the typical distance between a point estimator (e.g. posterior mean or median) and the truth, but also the typical spread of the posterior about the truth. Ideally, most of the posterior probability should concentrate on a ball centered at the true f_0 with radius proportional to the *minimax* estimation rate, see Section C.1 for more details. Indeed, since posterior contraction at rate ε_n automatically yields a point estimator with convergence rate ε_n ([14], Theorem 8.7), the minimax rate provides an information-theoretic lower bound for ε_n . Posterior contraction at a fast rate is a necessary condition for statistically good Bayesian estimation and uncertainty quantification.

We use the E_{f_0} -expectation formulation of contraction to quantify the *uniformity* over $f_0 \in \mathcal{F}_n$ needed to make our lower bounds precise and meaningful. Such uniformity captures that the true f_0 is unknown and rules out unrealistic and trivial situations, such as taking a GP with prior mean equal to the true f_0 and arbitrarily small covariance. Since we consider *worst case* (uniform) rates in Definition 1 and it is well-known that the covariance kernel K of the GP determines the characteristics of the corresponding GP [29], including its posterior contraction rates [43], we without loss of generality restrict to mean-zero GPs.

There is by now an extensive literature on proving *upper bounds* for contraction rates [14], including for GPs [43]. These contraction rates are usually uniform over suitable function classes \mathcal{F}_n , although the results might not be formulated in this way in the literature.

2.2 Lower bounds for posterior concentration in Gaussian processes regression

While the classical approach in nonparametric statistics is to make smoothness assumptions on the regression function f_0 , recent work in the deep learning literature has studied compositional assumptions. A special case are generalized additive models of the form

$$f_0(x_1, \dots, x_d) = h\left(\sum_{i=1}^d g_i(x_i)\right)$$

with g_1, \dots, g_d, h univariate and Lipschitz continuous functions such that all Lipschitz constants are bounded by Λ and $\|g_1\|_{L^\infty[0,1]}, \dots, \|g_d\|_{L^\infty[0,1]}, \|h\|_{L^\infty(\mathbb{R})} \leq \Lambda$. The class of all functions of this form is denoted by $\mathcal{G}(\Lambda)$ and we define $\mathcal{G} = \mathcal{G}(1)$. Generalized additive models are among the most popular structural constraints in function estimation [16].

Equation (13) in [35] and the subsequent discussion shows that the minimax estimation rate $R_n(\mathcal{G}(\Lambda))$ satisfies

$$cn^{-1/3} \leq R_n(\mathcal{G}(\Lambda)) \leq C(\log n)^3 n^{-1/3}$$

for some $C, c > 0$. Up to logarithmic factors, this rate is obtained by carefully calibrated deep GP priors that first assign a hyperprior to different compositional structures and subsequently put GP priors on all functions in the function composition [11].

We are now ready to state the main lower bound results of this paper, which we emphasize applies to any GP, irrespective of the choice of covariance kernel (which may itself depend on n).

Theorem 1. *Let Π_n be any sequence of mean-zero Gaussian process priors on $L^2[0, 1]^d$. For any $0 < \delta < 1/4$ and $n \geq N(d, \delta)$ large enough, the corresponding posterior distributions satisfy*

$$\sup_{f_0 \in \mathcal{G}} E_{f_0} \Pi_n(f : \|f - f_0\|_{L^2} \geq C_d n^{-\frac{2+d}{4+4d}} | Y) \geq 1/4 - \delta,$$

where $C_d = \frac{1}{10 \cdot 2^d} \left(\frac{1}{2(d+2)!} \right)^{\frac{d}{4+4d}}$. In particular, if the posterior distributions contract at rate ε_n uniformly over \mathcal{G} , that is

$$\sup_{f_0 \in \mathcal{G}} E_{f_0} \Pi_n(f : \|f - f_0\|_{L^2} \geq \varepsilon_n | Y) \rightarrow 0$$

as $n \rightarrow \infty$, then $\varepsilon_n > C_d n^{-\frac{2+d}{4+4d}}$.

For dimension $d \geq 3$, the above rate is strictly slower than the minimax rate $n^{-1/3}$ by the polynomial factor $n^{\frac{d-2}{12+12d}}$. As the domain dimension d tends to infinity, our lower bound is approximately $n^{-1/4}$, in which case Theorem 1 says the GP posterior will place at least probability $1/4 - \delta$ on posterior draws $f \sim \Pi_n(\cdot | Y)$ having suboptimal reconstruction error $\|f - f_0\|_{L^2} \gtrsim n^{-1/4}$. In contrast, posteriors based on properly calibrated deep GPs will place almost all their probability on draws with minimax optimal reconstruction error of order $n^{-1/3}$ [11]. Thus GP posteriors provide poor reconstructions of $f_0 \in \mathcal{G}$ compared to deep GPs.

Theorem 1 says that no GP can attain a faster rate than $n^{-\frac{2+d}{4+4d}}$. It does not say that a given GP even attains this suboptimal rate, and specific GPs can perform much worse than this bound. We in fact conjecture that our lower bound can be improved to show that even this rate is not attainable by any GP, see Section 2.3 for more discussion. Nonetheless the main message from Theorem 1 is already clear, namely that GPs underperform in this setting by an already significant factor that is polynomial in the sample size n .

The last result says that generic posterior draws will be poor reconstructions of f_0 with significant posterior probability. The same holds for the posterior mean.

Theorem 2. *Let Π_n be any sequence of mean-zero Gaussian process priors on $L^2[0, 1]^d$. Then the corresponding posterior means $\bar{f}_n = E^{\Pi_n}[f | Y]$ satisfy for $n \geq 2(d+2)!$,*

$$\sup_{f_0 \in \mathcal{G}} E_{f_0} \|\bar{f}_n - f_0\|_{L^2} \geq C'_d n^{-\frac{2+d}{4+4d}},$$

where $C'_d = \frac{1}{2^{d+1}} \left(\frac{1}{2(d+2)!} \right)^{\frac{d}{4+4d}}$.

Theorem 2 states that the posterior mean \bar{f}_n will also be a suboptimal reconstruction. Since the posterior is also Gaussian by conjugacy, and hence centered at \bar{f}_n , this captures the notion that posteriors draws from the ‘center’ of the posterior will also be poor - this is thus generic behaviour and not an artifact of unusual behaviour in the posterior tails.

Our results also have implications for Bayesian uncertainty quantification (UQ) through the use of credible sets, which is one the main motivations for using GPs. Bayesian credible sets are often

used as frequentist confidence sets, a practice which can be rigorously justified in low dimensional settings ([42], Section 10.2), but is much more subtle in high-dimensional settings like GPs [12]. Whether a resulting credible set contains the truth will depend on the size of the posterior standard deviation relative to the distance between the posterior mean and the truth, which Theorem 2 says is greater than the minimax rate with high-probability. In the best case scenario, the former is greater than the latter and thus the credible set will provide honest and reliable UQ. However, in this case the posterior standard deviation must be at least of the order of the contraction rate, and hence the resulting credible sets will be (polynomially) much too large and uninformative. In the worse case, when the posterior standard deviation is smaller, the credible sets will not contain the truth and provide misleading (useless) UQ.

Note that we are considering UQ for the *entire* function f_0 - it may still be possible for the GP posterior to provide efficient UQ for certain low-dimensional functionals of interest, which is itself a current topic of research [5, 32].

The proofs are based on a number of inequalities that are easy to state. Relying on an exponential concentration inequality and Anderson's inequality, we show that Theorem 1 follows from the lower bound for the posterior mean in Theorem 2 (cf. Section 4.1). For Theorem 2 we provide two different proofs. In the first, we lower bound the L^2 -risk of the posterior mean over arbitrary finite subsets $\{f_1, \dots, f_m\} \subset L^2[0, 1]^d$ (cf. Lemma 7), with the lower bound expressed in terms of expansions of the functions f_i in the basis coming from the Karhunen-Loève expansion of the GP. We then construct a suitable collection of orthogonal generalized additive models for which the lower bound is maximized *independently* of the choice of GP basis, as is needed to deal with arbitrary GPs (cf. Section 4.3). Given a *specific* choice of GP, our lower bound techniques can yield more refined results by tailoring the alternatives f_1, \dots, f_m to be especially poorly aligned (difficult to learn) for that specific covariance function, see Section 2.3.

In a second proof of Theorem 2, presented in Section B in the supplement, we reduce the problem of studying orthogonal generalized additive models to the sub-problem of studying the risk of linear estimators in a one-sparse sequence model (cf. Section B.1). For the latter, we exactly compute the linear minimax risk in Lemma 11, which, by exploiting the orthogonality, in turn implies the lower bound in Theorem 2. This second proof sheds additional light on the limitations of GP regression by furnishing new tools to study problems that can be connected to sparse models.

2.3 A sharper lower bound for a specific class of Gaussian processes

While Theorem 1 holds for arbitrary GPs, we now show that our methods can yield sharper lower bounds in the case of specific GPs, implying an even stronger suboptimality in the associated reconstruction error. We illustrate this by considering Gaussian wavelet series priors. Wavelets have been extensively used in Bayesian nonparametrics to construct priors in function spaces, including applications in inverse problems [44, 2], and image and signal processing [3, 20] among the others; see e.g. [25] for an overview.

Let $\{\psi_\gamma, \gamma \in \Gamma\}$, with $\Gamma = \{\gamma = (j, k) : j = -1, 0, 1, \dots, k \in I_j\}$, be a compactly supported wavelet basis of $L^2(\mathbb{R})$ restricted (with boundary correction) to $[0, 1]$, cf. [7] or also [15, Chapter 4]. The collection $\{\psi_\gamma, \gamma = (\gamma_1, \dots, \gamma_d) \in \Gamma^d\}$, with $\Gamma^d = \Gamma \times \dots \times \Gamma$ (d times) and $\psi_\gamma(x_1, \dots, x_d) := \prod_{i=1}^d \psi_{\gamma_i}(x_i)$ provides a tensor product wavelet basis of $L^2[0, 1]^d$.

Consider a mean-zero Gaussian wavelet series prior on $L^2[0, 1]^d$ with Karhunen-Loève expansion

$$f(x) = \sum_{\gamma \in I} \sqrt{\lambda_\gamma} \xi_\gamma \psi_\gamma(x), \quad x = (x_1, \dots, x_d) \in [0, 1]^d, \quad (4)$$

for some arbitrary index set $I \subseteq \Gamma^d$, coefficients $\lambda_\gamma > 0$ and $\xi_\gamma \stackrel{iid}{\sim} N(0, 1)$. Compared to the proof of the general lower bound in Theorem 1, fixing a specific GP basis allows to consider compositional functions that are especially difficult for that GP to approximate. For wavelet bases, such approximation theory was developed in [35], which we employ to obtain the next lower bound.

Theorem 3. *Let Π_n be any sequence of mean-zero Gaussian wavelet series priors on $L^2[0, 1]^d$, defined as in (4) with possibly n -dependent $\lambda_\gamma > 0$. For any $0 < \delta < 1/4$ and $n \geq N(d, \delta)$ large enough, the corresponding posterior distributions satisfy*

$$\sup_{f_0 \in \mathcal{G}} E_{f_0} \Pi_n(f : \|f - f_0\|_{L^2} \geq Cn^{-\frac{1}{2+d}} |Y) \geq 1/4 - \delta,$$

where $C = C(d, \psi) > 0$ only depends on d and the wavelet basis. Furthermore, the corresponding posterior means $\bar{f}_n = E^{\Pi_n}[f|Y]$ satisfy for all n

$$\sup_{f_0 \in \mathcal{G}} E_{f_0} \|\bar{f}_n - f_0\|_{L^2} \geq C' n^{-\frac{1}{2+d}},$$

where $C' = C'(d, \psi) > 0$ only depends on d and the wavelet basis.

Thus, for the class of Gaussian wavelet series priors, Theorem 3 provides a sharper lower bound than Theorem 1, as $n^{-\frac{1}{2+d}} \gg n^{-\frac{2+d}{4+4d}}$ for all d . Furthermore, the former rate deteriorates as d grows, showing that the performance of Gaussian wavelet priors suffers from a form of curse of dimensionality. In particular, when d is large, the fastest rate $n^{-\frac{1}{2+d}}$ achievable by Gaussian wavelet priors is much slower than the minimax rate $n^{-1/3}$ achieved by deep GPs.

3 Discussion

In this work, we showed that no mean-zero Gaussian process can attain the minimax estimation rate over a class of generalized additive models. We proved this by developing new tools for lower bounding posterior contraction rates for GPs. We provided two different proofs, each of which sheds different light on these phenomena.

It remains open whether even sharper lower bounds can be obtained. In this context, an important question is whether all GPs suffer from a curse of dimensionality in their posterior contraction rate, which we showed was incurred for a specific wavelet-based GP. Our lower bound $n^{-(2+d)/(4+4d)}$ for arbitrary GPs in Theorem 1 is never slower than $n^{-1/4}$, irrespective of the domain dimension d . While the obtained lower bound already demonstrates the suboptimality of GPs by a polynomial factor in n , say when compared to deep GPs, our stronger lower bound $n^{-1/(2+d)}$ for Gaussian wavelet series priors becomes arbitrarily slow with increasing d . We thus proved that for this subclass of GPs, a stronger curse of dimensionality is unavoidable.

To avoid the curse of dimensionality, a common two-stage approach is to first map the input features to a learned lower dimensional space and then apply Gaussian process regression to these new features. Our results do not apply to this scenario, in particular when the feature extraction map can be nonlinear, since the transformed underlying function may no longer be compositional in terms of the new learned features.

We expect similar results to hold for non-uniformly distributed input variables and heteroscedastic noise. Let σ be a positive function and ρ a density on $[0, 1]^d$. If ρ, σ are bounded away from zero and infinity, we conjecture that, up to constants, the same lower bounds can be derived working in the more general nonparametric regression model (1) with, conditionally on X_i , independent noise variables $w_i|X_i \sim N(0, \sigma^2(X_i))$ and $X_i \sim^{iid} \rho$.

Ideally, one wants to develop a broader understanding of which structures in the data-generating distribution or regression function (deep) Gaussian processes can or cannot adapt to. In order to understand the advantages of deep learning there is an extensive theoretical literature comparing shallow to deep nets. More refined versions show that for certain approximation rates a minimal network depth is necessary [27]. To gain some insights into the role of depth in deep Gaussian processes, it would be desirable to explore similar limitations if the depth is fixed and a highly structured function needs to be learned.

4 Proofs

We first start with some preliminary facts concerning GPs and the Gaussian white noise model (2). For Π a GP prior on $L^2[0, 1]^d$, we can express its covariance kernel K in (3) in terms of the normalized eigenfunctions $\{\phi_k : k \geq 1\}$ of the compact, self-adjoint, trace-class operator $g \mapsto \int_{[0, 1]^d} K(\cdot, y)g(y)dy$ on $L^2[0, 1]^d$ and its associated eigenvalues $\lambda_k > 0$, which are summable, via the expansion $K(x, y) = \sum_{k=1}^{\infty} \lambda_k \phi_k(x)\phi_k(y)$. In particular, we can realize the Gaussian random element $f \sim \Pi$ through its series expansion

$$f = \sum_{k=1}^{\infty} \sqrt{\lambda_k} \xi_k \phi_k, \quad \xi_k \sim^{iid} N(0, 1), \quad (5)$$

known as the *Karhunen-Loève expansion*. The corresponding reproducing kernel Hilbert space (RKHS) \mathbb{H} of Π is

$$\mathbb{H} = \left\{ h = \sum_{k=1}^{\infty} h_k \phi_k : \|h\|_{\mathbb{H}}^2 := \sum_{k=1}^{\infty} \frac{h_k^2}{\lambda_k} < \infty \right\}.$$

For any orthonormal basis $\{\phi_k : k \geq 1\}$ of $L^2[0, 1]^d$, in particular the Karhunen-Loève basis of the prior (5), the Gaussian white noise model (2) is equivalent to observing the sequence model ($Y_k : k \geq 1$):

$$Y_k = \int_{[0,1]^d} \phi_k(x) dY_x = \langle \phi_k, f \rangle_{L^2} + \frac{1}{\sqrt{n}} \int_{[0,1]^d} \phi_k(x) dW_x =: \theta_k + \frac{w_k}{\sqrt{n}}, \quad (6)$$

for $k = 1, 2, 3, \dots$ and where $w_k \stackrel{\text{iid}}{\sim} N(0, \|\phi_k\|_{L^2}^2) = N(0, 1)$. To be precise, the observations ($Y_k : k \geq 1$) in the last equation and the original white noise model (2) are equivalent in the sense that each can be perfectly recovered from the other, see Section C.2 in the supplement for more details.

Viewing the prior Π through its series expansion (5), Π can be viewed as a prior on the space of coefficients in the basis expansion of $\{\phi_k\}$ leading to the prior distribution $f = (\theta_k)_k \sim \otimes_{k=1}^{\infty} N(0, \lambda_k)$. Denoting by $P_f = P_f^n$ the distribution of the sequence representation (6), we have likelihood

$$P_f = \otimes_{k=1}^{\infty} N(\theta_k, 1/n).$$

Using this representation, one can use posterior conjugacy to show that the posterior takes the form

$$f = \sum_{k=1}^{\infty} \theta_k \phi_k, \quad \theta_k | Y_k \sim \text{ind } N\left(\frac{n\lambda_k}{n\lambda_k + 1} Y_k, \frac{\lambda_k}{n\lambda_k + 1}\right), \quad (7)$$

see Section C.2 in the supplement for full details. In particular, the posterior mean is

$$\bar{f}_n = E^{\Pi_n}[f|Y] = \sum_{k=1}^{\infty} \frac{n\lambda_k}{n\lambda_k + 1} Y_k \phi_k. \quad (8)$$

4.1 Reducing the lower bound for posterior contraction to one for the L^2 -risk of the posterior mean

We will show that a lower bound on the L^2 -risk of the posterior mean implies a corresponding lower bound for the posterior contraction rate, so that it will suffice to consider the former. The proofs of the following lemmas are deferred to Section A.1 in the supplement due to space constraints. We start by showing that the squared estimation error of the posterior mean is of the order of its expectation with high probability.

Lemma 4. *Let $f_0 \in L^2[0, 1]^d$, $\bar{f}_n = E^{\Pi}[f|Y]$ denote the posterior mean based on a mean-zero Gaussian process prior Π on $L^2[0, 1]^d$ and set $\mu_n^2 = E_{f_0} \|\bar{f}_n - f_0\|_{L^2}^2$. Then for $n \geq 1$,*

$$P_{f_0}(\|\bar{f}_n - f_0\|_{L^2}^2 \leq \mu_n^2/4) \leq 4e^{-\frac{n\mu_n^2}{32}}.$$

We next show that a contraction rate for Gaussian process priors implies that the corresponding posterior means converge to the truth at the same rate with frequentist probability tending to one.

Lemma 5. *Let Π_n be a sequence of mean-zero Gaussian process priors on $L^2[0, 1]^d$. Then the corresponding posterior means $\bar{f}_n = E^{\Pi_n}[f|Y]$ satisfy*

$$P_{f_0}(\|\bar{f}_n - f_0\|_{L^2} \geq 2\gamma_n) \leq 2\sqrt{E_{f_0} \Pi_n(f : \|f - f_0\|_{L^2} \geq \gamma_n | Y)}$$

for any sequence γ_n and $n \geq 1$.

Combining the last two lemmas allows us to transfer an L^2 -risk bound for the posterior mean to one for posterior contraction.

Lemma 6. Let $f_0 \in L^2[0, 1]^d$, Π_n be a sequence of mean-zero Gaussian process priors with corresponding posterior means $\bar{f}_n = E^{\Pi_n}[f|Y]$, and set $\mu_n^2 = E_{f_0} \|\bar{f}_n - f_0\|_{L^2}^2$. Then for $n \geq 1$,

$$E_{f_0} \Pi_n(f : \|f - f_0\|_{L^2} \geq \mu_n/4|Y) \geq \frac{1}{4} \left(1 - 4e^{-\frac{n\mu_n^2}{32}}\right)^2.$$

Furthermore, suppose $\mathcal{F}_n \subset L^2$ satisfy $\sup_{f_0 \in \mathcal{F}_n} E_{f_0} \|\bar{f}_n - f_0\|_{L^2}^2 \geq \gamma_n^2 > 0$ for some sequence γ_n for which $n\gamma_n^2 \rightarrow \infty$ as $n \rightarrow \infty$. Then for any $0 < \delta < 1/4$ and n such that $n\gamma_n^2 \geq 32 \log\left(\frac{5}{1-\sqrt{1-4\delta}}\right)$,

$$\sup_{f_0 \in \mathcal{F}_n} E_{f_0} \Pi_n(f : \|f - f_0\|_{L^2} \geq \gamma_n/5|Y) \geq 1/4 - \delta.$$

In particular, the posteriors cannot contract at rate $\gamma_n/5$, uniformly over \mathcal{F}_n .

4.2 A general lower bound for the L^2 -risk of the posterior mean

By Lemma 6, it suffices to prove a lower bound for the L^2 -risk of the posterior mean. We next prove such a general lower bound.

Lemma 7. Let \bar{f}_n be an estimator of the form (8) with corresponding orthonormal basis $\{\phi_k : k \geq 1\}$. For functions $f_1, \dots, f_m \in L^2[0, 1]^d$,

$$\max_{j=1, \dots, m} E_{f_j} \|\bar{f}_n - f_j\|_{L^2}^2 \geq \sum_{k=1}^{\infty} \left(\frac{1}{m} \sum_{j=1}^m \langle f_j, \phi_k \rangle_{L^2}^2 \right) \wedge \frac{1}{n}. \quad (9)$$

Moreover, if f_1, \dots, f_m are orthogonal and $\max_j \|f_j\|_{L^2}^2 \leq m/n$, then

$$\max_{j=1, \dots, m} E_{f_j} \|\bar{f}_n - f_j\|_{L^2}^2 \geq \frac{1}{m} \sum_{j=1}^m \|f_j\|_{L^2}^2.$$

In particular, if $cm/n \leq \|f_j\|_{L^2}^2 \leq m/n$ for all j and some $0 < c \leq 1$, then one can take the lower bound to be cm/n .

Proof. Using the bias-variance decomposition, the expected prediction loss of an estimator of the form (8) is

$$E_f \|\bar{f}_n - f\|_{L^2}^2 = \sum_{k=1}^{\infty} \left(\frac{n\lambda_k}{1+n\lambda_k} - 1 \right)^2 \langle f, \phi_k \rangle_{L^2}^2 + \left(\frac{n\lambda_k}{1+n\lambda_k} \right)^2 \frac{1}{n}.$$

Setting $a_k := n\lambda_k/(1+n\lambda_k)$ and $T_k = \frac{1}{m} \sum_{j=1}^m \langle f_j, \phi_k \rangle_{L^2}^2$, the last equation yields

$$\begin{aligned} \max_{j=1, \dots, m} E_{f_j} \|\bar{f}_n - f_j\|_{L^2}^2 &\geq \frac{1}{m} \sum_{j=1}^m E_{f_j} \|\bar{f}_n - f_j\|_{L^2}^2 = \frac{1}{m} \sum_{j=1}^m \sum_{k=1}^{\infty} (a_k - 1)^2 \langle f_j, \phi_k \rangle_{L^2}^2 + \frac{a_k^2}{n} \\ &= \sum_{k=1}^{\infty} (a_k - 1)^2 T_k + \frac{a_k^2}{n}. \end{aligned}$$

Optimizing over a_k shows that the right-hand side is minimized at $a_k^* = nT_k/(1+nT_k)$. For this choice, we obtain using $1+nT_k \geq 1 \vee (nT_k)$,

$$\max_{j=1, \dots, m} E_{f_j} \|\bar{f}_n - f_j\|_{L^2}^2 \geq \sum_k \frac{T_k}{(1+nT_k)^2} + \frac{T_k^2 n}{(1+nT_k)^2} = \sum_k \frac{T_k}{1+nT_k} \geq \sum_k T_k \wedge \frac{1}{n},$$

proving the first claim.

To see the second claim, observe that $\|f_j\|_{L^2}^2 \leq m/n$ and the orthogonality of the f_j imply

$$T_k = \frac{1}{m} \sum_{j=1}^m \|f_j\|_{L^2}^2 \left\langle \frac{f_j}{\|f_j\|_{L^2}}, \phi_k \right\rangle^2 \leq \frac{1}{n} \sum_{j=1}^m \left\langle \frac{f_j}{\|f_j\|_{L^2}}, \phi_k \right\rangle^2 \leq \frac{\|\phi_k\|_{L^2}^2}{n} = \frac{1}{n}.$$

Therefore $T_k \wedge 1/n = T_k$ and using (9) and that $(\phi_k)_k$ is a basis,

$$\max_{j=1, \dots, m} E_{f_j} \|\bar{f}_n - f_j\|_{L^2}^2 \geq \sum_k \frac{1}{m} \sum_{j=1}^m \langle f_j, \phi_k \rangle_{L^2}^2 = \frac{1}{m} \sum_{j=1}^m \sum_k \langle f_j, \phi_k \rangle_{L^2}^2 = \frac{1}{m} \sum_{j=1}^m \|f_j\|_{L^2}^2. \quad \square$$

4.3 Construction of local functions and proof of Theorems 1 and 2

Lemma 8. *Let $m = k^d$ for some integer $k \geq 1$. Then there exist functions $f_1, \dots, f_m \in \mathcal{G}$ having disjoint support and satisfying $\|f_j\|_{L^2}^2 = \frac{1}{2(d+2)!} m^{-\frac{2+d}{d}}$ for $j = 1, \dots, m$.*

Proof. For $\ell = 1, \dots, k$, consider the midpoints $z_\ell := (\ell - 1/2)/k$ of the intervals $[(\ell - 1)/k, \ell/k]$. Overall, there are $m = k^d$ points in the grid

$$I := \{(z_{\ell_1}, \dots, z_{\ell_d})^\top : \ell_1, \dots, \ell_d \in \{1, \dots, k\}\}.$$

Consider the functions,

$$f_a(x) = \left(\frac{1}{2k} - |x - a|_1 \right)_+, \quad a = (a_1, \dots, a_d) \in I.$$

These functions can be viewed as generalized additive models $f_a(x) = h(\sum_{i=1}^d g_{ai}(x_i))$ with $h(u) = (1/(2k) - u)_+$ and $g_{ai}(x_i) = |x_i - a_i|$. All these functions have Lipschitz constant bounded by 1. Moreover, $\max_{a \in I, i=1, \dots, d} \|g_{ai}\|_{L^\infty[0,1]} \leq 1$ and $\|h\|_{L^\infty(\mathbb{R})} = 1/(2k) \leq 1$. Thus, $f_a \in \mathcal{G}$.

Since $|x - a|_1 \geq |x - a|_\infty$, the support of the function f_a is contained in the hypercube $\times_{i=1}^d [a_i - 1/(2k), a_i + 1/(2k)]$. Since a_i is of the form $(\ell - 1/2)/k$ with ℓ an integer, the functions f_a must have disjoint support and are therefore orthogonal.

It thus remains to compute the L^2 norm of these functions, which by translation are the same as that of $f_0(x) = (1/(2k) - |x|_1)_+$ on \mathbb{R}^d . Note that the support of f_0 can be decomposed as $\text{supp}(f_0) = \cup_{\epsilon \in \{-1,1\}^d} R_\epsilon$, where

$$R_\epsilon = \left\{ x \in \mathbb{R}^d : \text{sign}(x_i) = \epsilon_i, \sum_{j=1}^d |x_j| \leq \frac{1}{2k} \right\}.$$

Since the L^2 -norm of f_0 on each of these 2^d quadrants is equal, we have

$$\begin{aligned} \|f_0\|_{L^2}^2 &= 2^d \int_{R_1} f_0(x)^2 dx \\ &= 2^d \int_{x_1=0}^{1/(2k)} \int_{x_2=0}^{1/(2k)-x_1} \dots \int_{x_d=0}^{1/(2k)-x_1-\dots-x_{d-1}} \left(\frac{1}{2k} - x_1 - \dots - x_d \right)^2 dx_1 dx_2 \dots dx_d. \end{aligned}$$

Using repeatedly that $\int_0^{M-r} (M-r-y)^t dy = \frac{1}{t+1} (M-r)^{t+1}$ for $0 \leq r \leq M$ and $t > 0$, we obtain

$$\|f_0\|_{L^2}^2 = 2^d \frac{2}{(d+2)!} \left(\frac{1}{2k} \right)^{d+2} = \frac{1}{2(d+2)!} k^{-d-2}.$$

□

Proof of Theorems 1 and 2. Let $r_d = \frac{1}{2(d+2)!}$, $k = \lceil (r_d n)^{\frac{1}{2d+2}} \rceil$ and $m = k^d$. By Lemma 8, there exist orthogonal functions $f_1, \dots, f_m \in \mathcal{G}$ with $\|f_j\|_{L^2}^2 = r_d m^{-\frac{2+d}{d}}$. For such a choice of k and m , $cm/n \leq \|f_j\|_{L^2}^2 \leq m/n$ holds if and only if

$$c^{\frac{1}{2d+2}} \leq \frac{(r_d n)^{\frac{1}{2d+2}}}{\lceil (r_d n)^{\frac{1}{2d+2}} \rceil}$$

after rearranging the inequalities. Now the function $\varphi(x) = x/\lceil x \rceil$ satisfies $\varphi(x) \geq \frac{k}{k+1}$ for all $x \geq k \in \mathbb{N}$, so that the right side is lower bounded by $1/2$ for $(r_d n)^{\frac{1}{2d+2}} \geq 1$, or equivalently $n \geq 1/r_d$. For $n \geq 1/r_d$, it therefore holds that $cm/n \leq \|f_j\|_{L^2}^2 \leq m/n$ with $c = (1/2)^{2d+2}$. Applying Lemma 7 thus gives the risk bound

$$\sup_{f_0 \in \mathcal{G}} E_{f_0} \| \bar{f}_n - f_0 \|_{L^2}^2 \geq \max_{j=1, \dots, m} E_{f_j} \| \bar{f}_n - f_j \|_{L^2}^2 \geq \frac{cm}{n} \geq \frac{1}{2^{2d+2}} r_d^{\frac{d}{2d+2}} n^{-\frac{2+d}{2d+2}},$$

which proves Theorem 2.

Set

$$N(d, \delta) = 2(d+2)! 2^{\frac{(2d+2)^2}{d}} \left[32 \log \left(\frac{5}{1 - \sqrt{1 - 4\delta}} \right) \right]^{\frac{d+2}{d}},$$

and note that $N(d, \delta) \geq 1/r_d$. Theorem 1 then follows from the second last display and applying Lemma 6 with $\mathcal{F}_n = \{f_1, \dots, f_m\} \subset \mathcal{G}$ and $\gamma_n^2 = 2^{-(2d+2)} r_d^{\frac{d}{2d+2}} n^{-\frac{2+d}{2+2d}}$. \square

Acknowledgments. We are grateful to three anonymous referees for helpful comments that improved the manuscript. The research of MG has been supported by the European Research Council under ERC grant agreement No.834275 (GTBB). The research of JSH has been supported by the NWO/STAR grant 613.009.034b and the NWO Vidi grant VI.Vidi.192.021.

Supplementary material to “On the inability of Gaussian process regression to optimally learn compositional functions”

In this supplement we provide additional background material and all the technical results and proofs not included in the main article. For the convenience of the reader, we repeat the statements of results that we prove here. We continue the equation numbering scheme from the main document to the supplement.

A Remaining proofs: Lemmas 4-6, and Theorem 3

A.1 Proof of Lemmas 4-6

Lemma 4. Let $f_0 \in L^2[0, 1]^d$, $\bar{f}_n = E^\Pi[f|Y]$ denote the posterior mean based on a mean-zero Gaussian process prior Π on $L^2[0, 1]^d$ and set $\mu_n^2 = E_{f_0} \|\bar{f}_n - f_0\|_{L^2}^2$. Then for $n \geq 1$,

$$P_{f_0} (\|\bar{f}_n - f_0\|_{L^2}^2 \leq \mu_n^2/4) \leq 4e^{-\frac{n\mu_n^2}{32}}.$$

Proof. Using (8), under P_{f_0} the posterior mean equals

$$\bar{f}_n = \sum_k \frac{n\lambda_k}{n\lambda_k + 1} Y_k \phi_k = \sum_k \frac{n\lambda_k}{n\lambda_k + 1} f_{0k} \phi_k + \sum_k \frac{n\lambda_k}{n\lambda_k + 1} \frac{w_k}{\sqrt{n}} \phi_k.$$

Since $f_0 = \sum_k f_{0k} \phi_k$, we deduce

$$\begin{aligned} \|\bar{f}_n - f_0\|_{L^2}^2 &= \sum_k \left(\frac{\sqrt{n}\lambda_k}{n\lambda_k + 1} w_k - \frac{1}{n\lambda_k + 1} f_{0k} \right)^2 \\ &= \sum_k \frac{1}{(n\lambda_k + 1)^2} [n\lambda_k^2(w_k^2 - 1) + n\lambda_k^2 - 2\sqrt{n}\lambda_k w_k f_{0k} + f_{0k}^2] \\ &= I + II + III + IV. \end{aligned}$$

Taking E_{f_0} -expectation yields

$$E_{f_0} \|\bar{f}_n - f_0\|_{L^2}^2 = \sum_k \frac{n\lambda_k^2}{(n\lambda_k + 1)^2} + \frac{f_{0k}^2}{(n\lambda_k + 1)^2} = II + IV.$$

We will now show that $I + III$ are of smaller order with high P_{f_0} -probability, so that $\|\bar{f}_n - f_0\|_{L^2}^2$ is close to its expectation, and hence the size of its expectation drives its behaviour.

I: Using Lemma 1 of [24] with $a_k = \frac{n\lambda_k^2}{(n\lambda_k + 1)^2}$, we get

$$P(|I| \geq 2\|a\|_{\ell_2} \sqrt{x} + 2\|a\|_{\ell_\infty} x) \leq 2e^{-x}$$

for any $x > 0$ (we write P instead of P_{f_0} here to emphasize the above probability does not depend on f_0). Further set $\alpha_k = \frac{\lambda_k}{\lambda_k + 1/n}$, so that $n\alpha_k = \alpha_k^2$. Then

$$n^2 \|a\|_{\ell_2}^2 = \|\alpha\|_{\ell_4}^4 \leq \sup_k |\alpha_k|^2 \sum_k \alpha_k^2 = \sup_k \left| \frac{\lambda_k^2}{(\lambda_k + 1/n)^2} \right| \|\alpha\|_{\ell_2}^2 \leq \|\alpha\|_{\ell_2}^2.$$

Using that $\|\alpha\|_{\ell_2}^2 = nII$, this implies $\|a\|_{\ell_2} \leq \frac{1}{n} \|\alpha\|_{\ell_2} = \sqrt{II/n}$. For the ℓ_∞ term,

$$\|a\|_{\ell_\infty} = \frac{1}{n} \sup_k \left| \frac{n^2 \lambda_k^2}{(n\lambda_k + 1)^2} \right| \leq \frac{1}{n}.$$

In conclusion, we have the exponential inequality

$$P \left(|I| \geq 2\sqrt{\frac{IIx}{n}} + \frac{2x}{n} \right) \leq 2e^{-x}.$$

III: We have $III \sim N(0, \sum_k \frac{4n\lambda_k^2 f_{0k}^2}{(n\lambda_k + 1)^4})$, whose variance is bounded by $\frac{4}{n} \sup_k \frac{n^2 \lambda_k^2}{(n\lambda_k + 1)^2} \sum_k \frac{f_{0k}^2}{(n\lambda_k + 1)^2} \leq \frac{4}{n} IV$. Thus using the standard Gaussian tail bound, for any $x \geq 0$,

$$P_{f_0} \left(|III| \geq 2\sqrt{IV/nx} \right) \leq 2e^{-x^2/2},$$

uniformly over $f_0 \in L^2$.

The triangle inequality gives $|\|\bar{f}_n - f_0\|_{L^2}^2 - E_{f_0} \|\bar{f}_n - f_0\|_{L^2}^2| = |I + III| \leq |I| + |III|$. Since for any real numbers a, b , $P_{f_0}(|I| + |III| \geq a + b) \leq P_{f_0}(\{|I| \geq a\} \cup \{|III| \geq b\}) \leq P_{f_0}(|I| \geq a) + P_{f_0}(|III| \geq b)$, combining the bounds for I and III , we get

$$\sup_{f_0 \in L^2} P_{f_0} \left(|\|\bar{f}_n - f_0\|_{L^2}^2 - E_{f_0} \|\bar{f}_n - f_0\|_{L^2}^2| \geq 2\sqrt{\frac{IIx}{n}} + \frac{2x}{n} + \sqrt{\frac{8IVx}{n}} \right) \leq 4e^{-x}$$

for all $x > 0$, a non-asymptotic inequality. Using $ab \leq (a^2 + b^2)/2$ gives $\sqrt{8IVx/n} \leq IV/2 + 4x/n$ and $2\sqrt{IIx/n} \leq II/2 + 2x/n$. Since $E_{f_0} \|\bar{f}_n - f_0\|_{L^2}^2 = II + IV$, it holds that for any $x > 0$,

$$\sup_{f_0 \in L^2} P_{f_0} \left(|\|\bar{f}_n - f_0\|_{L^2}^2 - E_{f_0} \|\bar{f}_n - f_0\|_{L^2}^2| \geq \frac{1}{2} E_{f_0} \|\bar{f}_n - f_0\|_{L^2}^2 + \frac{8x}{n} \right) \leq 4e^{-x}.$$

Set $x = n\mu_n^2/32$ so that $8x/n = \mu_n^2/4$. Then the last inequality implies that for all $f_0 \in L^2$,

$$P_{f_0} \left(\|\bar{f}_n - f_0\|_{L^2}^2 \leq \frac{1}{2} E_{f_0} \|\bar{f}_n - f_0\|_{L^2}^2 - \frac{\mu_n^2}{4} \right) \leq 4e^{-\frac{n\mu_n^2}{32}}.$$

Substituting in $\mu_n^2 = E_{f_0} \|\bar{f}_n - f_0\|_{L^2}^2$ gives the result. \square

Lemma 5. Let Π_n be a sequence of mean-zero Gaussian process priors on $L^2[0, 1]^d$. Then the corresponding posterior means $\bar{f}_n = E^{\Pi_n}[f|Y]$ satisfy

$$P_{f_0} (\|\bar{f}_n - f_0\|_{L^2} \geq 2\gamma_n) \leq 2\sqrt{E_{f_0} \Pi_n(f : \|f - f_0\|_{L^2} \geq \gamma_n | Y)}$$

for any sequence γ_n and $n \geq 1$.

Proof. Write $v_n = E_{f_0} \Pi_n(f : \|f - f_0\|_{L^2} \geq \gamma_n | Y)$. We may assume $v_n < 1/4$, otherwise the right side is greater than one and there is nothing to prove. Using Markov's inequality, the events $A_{n, f_0} = \{\Pi_n(f : \|f - f_0\|_{L^2} \geq \gamma_n | Y) \leq \sqrt{v_n}\}$ then satisfy $P_{f_0}(A_{n, f_0}^c) \leq \sqrt{v_n}$.

Recall that the posterior distributions $\Pi_n(f|Y)$ are also Gaussian by conjugacy, see (7). Since $\{f \in L^2 : \|f\|_{L^2} \leq \gamma_n\}$ is convex and symmetric, Anderson's inequality (e.g. [15], Theorem 2.4.5) implies that

$$1 - v_n = E_{f_0} \Pi_n(f : \|f - f_0\|_{L^2} \leq \gamma_n | Y) \leq E_{f_0} \Pi_n(f : \|f - \bar{f}_n\|_{L^2} \leq \gamma_n | Y).$$

Using the same argument as above, we thus have $P_{f_0}(B_n^c) \leq \sqrt{v_n}$ for $B_n = \{\Pi_n(f : \|f - \bar{f}_n\|_{L^2} \geq \gamma_n | Y) \leq \sqrt{v_n}\}$, so that $P_{f_0}(A_{n,f_0} \cap B_n) \geq 1 - 2\sqrt{v_n}$. Now on the event $A_{n,f_0} \cap B_n$, we have

$$\begin{aligned} \Pi_n(f : \|f - f_0\|_{L^2} \leq \gamma_n, \|f - \bar{f}_n\|_{L^2} \leq \gamma_n | Y) &\geq 1 - \Pi_n(f : \|f - f_0\|_{L^2} \geq \gamma_n | Y) \\ &\quad - \Pi_n(f : \|f - \bar{f}_n\|_{L^2} \geq \gamma_n | Y) \\ &\geq 1 - 2\sqrt{v_n}. \end{aligned}$$

The right-hand side is strictly positive for $v_n < 1/4$ and hence so is the left posterior probability. Thus there must exist $f \in L^2$ in the left set, in which case

$$\|\bar{f}_n - f_0\|_{L^2} \leq \|\bar{f}_n - f\|_{L^2} + \|f - f_0\|_{L^2} \leq 2\gamma_n.$$

We have thus shown that $P_{f_0}(\|\bar{f}_n - f_0\|_{L^2} \leq 2\gamma_n) \geq 1 - 2\sqrt{v_n}$ as required. \square

Lemma 6. *Let $f_0 \in L^2[0, 1]^d$, Π_n be a sequence of mean-zero Gaussian process priors with corresponding posterior means $\bar{f}_n = E^{\Pi_n}[f|Y]$, and set $\mu_n^2 = E_{f_0}\|\bar{f}_n - f_0\|_{L^2}^2$. Then for $n \geq 1$,*

$$E_{f_0}\Pi_n(f : \|f - f_0\|_{L^2} \geq \mu_n/4 | Y) \geq \frac{1}{4} \left(1 - 4e^{-\frac{n\mu_n^2}{32}}\right)_+.$$

Furthermore, suppose $\mathcal{F}_n \subset L^2$ satisfy $\sup_{f_0 \in \mathcal{F}_n} E_{f_0}\|\bar{f}_n - f_0\|_{L^2}^2 \geq \gamma_n^2 > 0$ for some sequence γ_n for which $n\gamma_n^2 \rightarrow \infty$ as $n \rightarrow \infty$. Then for any $0 < \delta < 1/4$ and n such that $n\gamma_n^2 \geq 32 \log\left(\frac{5}{1-\sqrt{1-4\delta}}\right)$,

$$\sup_{f_0 \in \mathcal{F}_n} E_{f_0}\Pi_n(f : \|f - f_0\|_{L^2} \geq \gamma_n/5 | Y) \geq 1/4 - \delta.$$

In particular, the posteriors cannot contract at rate $\gamma_n/5$, uniformly over \mathcal{F}_n .

Proof. Using Lemma 5 and then Lemma 4, the square-root of the left-side of the first display is lower bounded by

$$\frac{1}{2} P_{f_0}(\|\bar{f}_n - f_0\|_{L^2} \geq \mu_n/2) \geq \frac{1}{2} \left(1 - 4e^{-\frac{n\mu_n^2}{32}}\right).$$

Squaring everything then gives the first result.

By hypothesis, for any $\eta > 0$, there exists $f_{0,n} \in \mathcal{F}_n$ such that $\mu_n^2 = E_{f_{0,n}}\|\bar{f}_n - f_{0,n}\|_{L^2}^2 \geq \gamma_n^2 - \eta$. Let $\eta = \eta_n$ be small enough that $e^{-\frac{\eta_n}{32}} \leq 5/4$ and $\sqrt{\gamma_n^2 - \eta_n}/4 \geq \gamma_n/5$. By the first part of the lemma,

$$E_{f_{0,n}}\Pi_n(f : \|f - f_{0,n}\|_{L^2} \geq \sqrt{\gamma_n^2 - \eta_n}/4 | Y) \geq \frac{1}{4} \left(1 - 5e^{-\frac{n\gamma_n^2}{32}}\right)_+.$$

After rearranging, the right-side is at least $1/4 - \delta$ for $n\gamma_n^2 \geq 32 \log\left(\frac{5}{1-\sqrt{1-4\delta}}\right)$. The second part then follows from upper bounding the left-side of the last display by $E_{f_{0,n}}\Pi_n(f : \|f - f_{0,n}\|_{L^2} \geq \gamma_n/5 | Y)$. \square

A.2 Proof of Theorem 3

Similar to the proof of Theorems 1 and 2, we derive a lower bound for the L^2 -risk of the posterior mean over a suitable subset of the class \mathcal{G} .

For a wavelet prior Π_n defined as in (4) (with possibly n -dependent λ_γ) and \bar{f}_n the corresponding posterior mean, the conclusion of Lemma 7 reads, for $m = 1$,

$$E_f \|\bar{f}_n - f\|_{L^2}^2 \geq \sum_{\gamma \in \Gamma^d} \langle f, \psi_\gamma \rangle_{L^2}^2 \wedge \frac{1}{n}, \quad \text{any } f \in L^2[0, 1]^d. \quad (10)$$

By applying Lemma 2 in [35] (with $\alpha = K = 1$), for a constant $c(d, \psi) > 0$ only depending on d and the wavelet basis, letting $j_n \in \mathbb{N}$ satisfy

$$\frac{1}{n} \leq c(d, \psi) 2^{-j_n(2+d)} \leq \frac{2^{2+d}}{n}$$

there exists for each n a Lipschitz function $h_{j_n} : [0, d] \rightarrow [0, 1]$ with Lipschitz constant bounded by 1 such that the generalized additive function $f_{j_n}(x_1, \dots, x_d) := h_{j_n}(x_1 + \dots + x_d)$ satisfies for all $p_1, \dots, p_d \in \{0, 1, \dots, 2^{j_n - q - \nu} - 1\}$,

$$\langle f_{j_n}, \psi_{(j_n, 2^{q+\nu} p_1), \dots, (j_n, 2^{q+\nu} p_d)} \rangle_{L^2}^2 = c(d, \psi)^2 2^{-j_n(2+d)} \geq \frac{1}{n}.$$

Above, $q > 0$ is such that the mother wavelet ψ is supported within $[0, 2^q]$ and $\nu = \lceil \log_2 d \rceil + 1$. Noting that with our choice of j_n we have $2^{j_n} \geq \frac{1}{2}(c(d, \psi)^2 n)^{\frac{1}{2+d}}$, it follows from (10) that

$$\begin{aligned} E_{f_{j_n}} \|\bar{f}_n - f_{j_n}\|_{L^2}^2 &\geq \sum_{p_1, \dots, p_d \in \{0, 1, \dots, 2^{j_n - q - \nu} - 1\}} \frac{1}{n} \\ &= 2^{-(q+d+\nu)d} \frac{1}{n} 2^{j_n d} \geq 2^{-(q+\nu+1)d} (c(d, \psi))^{\frac{2d}{2+d}} n^{-\frac{2}{2+d}}. \end{aligned}$$

Since $f_{j_n} \in \mathcal{G}$, this proves the second claim of Theorem 3. Another application of Lemma 6 with $\mathcal{F}_n = \{f_{j_n}\} \subset \mathcal{G}$, $\gamma_n^2 = 2^{-(q+\nu+1)d} (c(d, \psi))^{\frac{2d}{2+d}} n^{-\frac{2}{2+d}}$ then proves the first claim taking $N(d, \delta) \geq 32 \log \left(\frac{5}{1 - \sqrt{1 - 4\delta}} \right) / \gamma_n^2$ and $C = C(d, \psi) = 2^{-(q+\nu+1)d/2} (c(d, \psi))^{\frac{d}{2+d}} / 5$.

B An alternative proof of Theorems 1 and 2

As part of the proofs of Theorems 1 and 2 in Section 4, we directly lower bounded the L^2 -risk of the posterior mean in Lemma 7. We now provide an alternative strategy to prove a lower bound via first reducing the regression setting to a one-sparse sequence model in which we can explicitly evaluate the minimax risk for linear estimators.

B.1 Reduction to a one-sparse sequence model

Consider the Gaussian white noise model (2) with finite parameter space $\mathcal{F}_n = \{f_1, \dots, f_m\}$, $m = m_n$, satisfying

$$\langle f_i, f_j \rangle_{L^2} = \delta_{ij} c_n^2, \quad c_n > 0 \quad (11)$$

[we will ultimately take $f_1, \dots, f_m \in \mathcal{G}$ as in Lemma 8, but this approach may be useful for other function classes]. Denote by $f \in \mathcal{F}_n$ the regression function driving (2) and define

$$y_i := \frac{1}{c_n^2} \int_{[0,1]^d} f_i(x) dY_x = \frac{1}{c_n^2} \langle f_i, f \rangle_{L^2} + \frac{1}{\sqrt{n} c_n^2} \int_{[0,1]^d} f_i(x) dW_x, \quad (12)$$

$i = 1, \dots, m$. Further write

$$\theta_i := \frac{1}{c_n^2} \langle f_i, f \rangle_{L^2}, \quad w_i := \frac{1}{c_n} \int_{[0,1]^d} f_i(x) dW_x \sim^{iid} N(0, 1), \quad \sigma_n := \frac{1}{c_n \sqrt{n}}. \quad (13)$$

Since $f \in \mathcal{F}_n$ in our statistical model satisfy (11), $\theta = (\theta_1, \dots, \theta_m)^T$ is a *one-sparse* vector. The present statistical model thus yields an observation from the finite Gaussian sequence model

$$y_i = \theta_i + \sigma_n w_i, \quad i = 1, \dots, m, \quad (14)$$

with $\theta \in \Theta_m = \{e_1, \dots, e_m\}$ for e_i the i^{th} basis vector, that is $e_{ij} = \delta_{ij}$.

We now relate estimation in the full Gaussian white noise model with $f \in \mathcal{F}_n$ with estimation in (14). Since $\theta_i = \langle f_i, f \rangle_{L^2} / c_n^2$, f_i / c_n are L^2 -normalized and orthogonal, and $f \in \mathcal{F}_n$ trivially lies in the linear span of $\{f_1, \dots, f_m\}$, we can express the function f in terms of $\theta = (\theta_1, \dots, \theta_m)^T$ via

$$f = \sum_{i=1}^m \frac{\langle f_i, f \rangle_{L^2}}{c_n} \frac{f_i}{c_n} = \sum_{i=1}^m \theta_i f_i.$$

Any estimator $\hat{\theta} = (\hat{\theta}_1, \dots, \hat{\theta}_m)^T$ for θ thus yields a series estimator $\hat{f}_{\hat{\theta}} = \sum_{i=1}^m \hat{\theta}_i f_i$ for $f \in \mathcal{F}_n$, and

$$\|\hat{f}_{\hat{\theta}} - f\|_{L^2}^2 = \sum_{i=1}^m (\hat{\theta}_i - \theta_i)^2 \|f_i\|_{L^2}^2 = c_n^2 \sum_{i=1}^m (\hat{\theta}_i - \theta_i)^2 = c_n^2 |\hat{\theta} - \theta|^2. \quad (15)$$

We next show that in the full Gaussian white noise model with restricted parameter space \mathcal{F}_n , the risk of the posterior mean is larger than that of an estimator of the form $\hat{f}_{\hat{\theta}}$ with $\hat{\theta} = Ay$ a linear function of the observations $y = (y_1, \dots, y_m)^T$ in (12).

Lemma 9. Let Π_n be a sequence of mean-zero Gaussian process priors on $L^2[0, 1]^d$ with corresponding posterior means $f_n = E^{\Pi_n}[f|Y]$. Then there exists a matrix sequence $A_n \in \mathbb{R}^{m \times m}$ such that for every n and every $f \in \mathcal{F}_n = \{f_1, \dots, f_m\}$ satisfying (11),

$$E_f \|\bar{f}_n - f\|_{L^2} \geq E_f \|\hat{f}_{A_n} - f\|_{L^2},$$

where $\hat{f}_{A_n} = \sum_{i=1}^m (A_n y)_i f_i$ and y_1, \dots, y_m are defined in (12).

Proof. Recall from (8) that the posterior mean takes the form $\bar{f}_n = \sum_{k=1}^{\infty} a_k Y_k \phi_k$, where $a_k = \frac{n\lambda_k}{1+n\lambda_k}$ with $(\lambda_k), (\phi_k)$ the eigenvalues/vectors arising in the Karhunen-Loève expansion (5), and $Y_k = \langle Y, \phi_k \rangle_{L^2}$ defined in (6). We can decompose each ϕ_k as

$$\phi_k = \psi_k + g_k, \quad \psi_k = \sum_{i=1}^m \psi_{k,i} f_i \in \text{span}\{f_1, \dots, f_m\}, \quad g_k \perp \{f_1, \dots, f_m\}$$

using the orthogonality relation (11), giving

$$\bar{f}_n = \sum_{k=1}^{\infty} a_k \langle Y, \psi_k + g_k \rangle_{L^2} \psi_k + \sum_{k=1}^{\infty} a_k \langle Y, \psi_k + g_k \rangle_{L^2} g_k =: \bar{f}_1 + \bar{f}_2.$$

Define the matrix $A_n \in \mathbb{R}^{m \times m}$ by

$$(A_n)_{ij} = c_n^2 \sum_{k=1}^{\infty} a_k \psi_{k,i} \psi_{k,j}.$$

Using that $\langle Y, f_j \rangle_{L^2} = c_n^2 y_j$ from (12), we can rewrite

$$\begin{aligned} \bar{f}_1 &= \sum_{k=1}^{\infty} a_k \left(\sum_{j=1}^m \psi_{k,j} \langle Y, f_j \rangle_{L^2} + \langle Y, g_k \rangle_{L^2} \right) \sum_{i=1}^m \psi_{k,i} f_i \\ &= \sum_{i=1}^m \left(\sum_{j=1}^m \left[\sum_{k=1}^{\infty} a_k \psi_{k,i} \psi_{k,j} \right] c_n^2 y_j + \sum_{k=1}^{\infty} a_k \psi_{k,i} \langle Y, g_k \rangle_{L^2} \right) f_i \\ &= \sum_{i=1}^m \left((A_n y)_i + \sum_{k=1}^{\infty} a_k \psi_{k,i} \langle Y, g_k \rangle_{L^2} \right) f_i. \end{aligned}$$

Since $\bar{f}_2 \perp f$ for all $f \in \mathcal{F}_n$, $\|f - \bar{f}_n\|_{L^2}^2 = \|f - \bar{f}_1\|_{L^2}^2 + \|\bar{f}_2\|_{L^2}^2 \geq \|f - \bar{f}_1\|_{L^2}^2$. Writing $f = \sum_{i=1}^m \theta_i f_i$, where $\theta_i = \langle f, f_i \rangle_{L^2} / c_n^2$, and recalling $\langle f_i, f_j \rangle_{L^2} = \delta_{ij} c_n^2$,

$$\begin{aligned} \|f - \bar{f}_1\|_{L^2}^2 &= \left\| \sum_{i=1}^m \left(\theta_i - (A_n y)_i - \sum_{k=1}^{\infty} a_k \psi_{k,i} \langle Y, g_k \rangle_{L^2} \right) f_i \right\|_{L^2}^2 \\ &= c_n^2 \sum_{i=1}^m \left(\theta_i - (A_n y)_i - \sum_{k=1}^{\infty} a_k \psi_{k,i} \langle Y, g_k \rangle_{L^2} \right)^2. \end{aligned}$$

Using the independence between $y_i = \langle Y, f_i \rangle_{L^2} / c_n^2$ and $\langle Y, g_k \rangle_{L^2}$ for all $i = 1, \dots, m$ and $k \geq 1$, and that $E_f \langle Y, g_k \rangle_{L^2} = 0$, we obtain that for all $f \in \mathcal{F}_n$,

$$\begin{aligned} E_f \|f - \bar{f}_n\|_{L^2}^2 &\geq c_n^2 E_f \sum_{i=1}^m \left\{ (\theta_i - (A_n y)_i)^2 + \left(\sum_{k=1}^{\infty} a_k \psi_{k,i} \langle Y, g_k \rangle_{L^2} \right)^2 \right\} \\ &\geq c_n^2 \sum_{i=1}^m E_f [\theta_i - (A_n y)_i]^2 = E_f \|f - \hat{f}_{A_n}\|_{L^2}^2. \end{aligned}$$

□

Using Lemma 9 and (15), we thus obtain

$$\inf_{f_n} \max_{f \in \mathcal{F}_n} E_f \|\bar{f}_n - f\|_{L^2}^2 \geq c_n^2 \inf_{A \in \mathbb{R}^{m \times m}} \max_{\theta \in \Theta_m} |A y - \theta|^2, \quad (16)$$

where the infimum on the left side is over all posterior means based on mean-zero Gaussian processes. It thus suffices to lower bound the right-side, which is the minimax risk for *linear* estimators in the one-sparse sequence model (14).

B.2 Minimax risk for linear estimators in the one-sparse Gaussian sequence model

We now study the minimax risk for linear estimators in the one-sparse model (14) for arbitrary noise level $\sigma_n > 0$. Recall that $\theta = (\theta_1, \dots, \theta_m)^T$ has exactly one non-zero coordinate, which is equal to one, so that the parameter space is $\Theta_m = \{e_1, \dots, e_m\}$ for e_i the i^{th} basis vector, i.e. $e_{ij} = \delta_{ij}$.

A linear estimator of θ in model (14) takes the form

$$\hat{\theta}_A = Ay,$$

for some matrix $A \in \mathbb{R}^{m \times m}$ and $y = (y_1, \dots, y_m)^T$. We now show that in this one-sparse model, such estimators are dominated by *diagonal homogeneous* linear estimators in terms of their maximal mean-squared error $E_\theta |\hat{\theta}_A - \theta|^2$, where $|\cdot|$ denotes the usual Euclidean norm on \mathbb{R}^m and E_θ the expectation in model (14) with true parameter θ . Hence the minimax risk for linear estimators is attained by a diagonal homogeneous linear estimator.

Lemma 10. *Let $A = (a_{ij}) \in \mathbb{R}^{m \times m}$ be any matrix and let $\bar{a}^2 = \frac{1}{m} \sum_{j=1}^m a_{jj}^2$. Then*

$$\max_{\theta \in \Theta_m} E_\theta |\hat{\theta}_A - \theta|^2 \geq \max_{\theta \in \Theta_m} E_\theta |\hat{\theta}_{\bar{a}I_m} - \theta|^2.$$

In particular,

$$\inf_{A \in \mathbb{R}^{m \times m}} \max_{\theta \in \Theta_m} E_\theta |\hat{\theta}_A - \theta|^2 = \inf_{a \in \mathbb{R}} \max_{\theta \in \Theta_m} E_\theta |\hat{\theta}_{aI_m} - \theta|^2.$$

Proof. Using the bias-variance decomposition,

$$\begin{aligned} E_\theta |\hat{\theta}_A - \theta|^2 &= |E_\theta \hat{\theta}_A - \theta|^2 + \text{tr}[\text{Cov}_\theta(\hat{\theta}_A)] \\ &= |(A - I_m)\theta|^2 + \text{tr}[A \text{Cov}_\theta(y) A^T] \\ &= |(A - I_m)\theta|^2 + \sigma_n^2 \text{tr}[A A^T] \\ &= \sum_{i=1}^m \left(\sum_{j=1}^m (a_{ij} - \delta_{ij}) \theta_j \right)^2 + \sigma_n^2 \sum_{i,j=1}^m a_{ij}^2, \end{aligned}$$

since $\text{Cov}_\theta(y) = \text{Cov}_\theta(w) = \sigma_n^2 I_m$. Since $\theta \in \Theta_m = \{e_1, \dots, e_m\}$, let $j^* \in \{1, \dots, m\}$ be the index such that $\theta = e_{j^*}$. Then

$$E_\theta |\hat{\theta}_A - \theta|^2 = \sum_{i=1}^m (a_{ij^*} - \delta_{ij^*})^2 + \sigma_n^2 \sum_{i,j=1}^m a_{ij}^2.$$

However, applying this last expression also with $\tilde{A} = \text{diag}(A)$, the diagonal matrix with entries $\tilde{a}_{ij} = a_{ij} \delta_{ij}$, gives

$$E_\theta |\hat{\theta}_{\tilde{A}} - \theta|^2 = (a_{j^*j^*} - 1)^2 + \sigma_n^2 \sum_{i=1}^m a_{ii}^2 \leq E_\theta |\hat{\theta}_A - \theta|^2,$$

a bound which holds for all $\theta \in \Theta_m$. Thus we need only consider the estimator $\hat{\theta}_{\tilde{A}}$ with diagonal matrix \tilde{A} . Using the last display,

$$\max_{\theta \in \Theta_m} E_\theta |\hat{\theta}_{\tilde{A}} - \theta|^2 = \max_{j=1, \dots, m} (a_{jj} - 1)^2 + \sigma_n^2 m \bar{a}^2.$$

Since the matrix $\bar{a}I_m$ is also diagonal, this further yields

$$\max_{\theta \in \Theta_m} E_\theta |\hat{\theta}_{\bar{a}I_m} - \theta|^2 = (\bar{a} - 1)^2 + \sigma_n^2 m \bar{a}^2,$$

so that it is enough to show $(\bar{a} - 1)^2 \leq \max_j (a_{jj} - 1)^2$. Since the function $\varphi(x) = (\sqrt{x} - 1)^2$ is convex on $(0, \infty)$, Jensen's inequality implies

$$(\bar{a} - 1)^2 = \left(\sqrt{\frac{1}{m} \sum_{i=1}^m a_{ii}} - 1 \right)^2 \leq \frac{1}{m} \sum_{i=1}^m (|a_{ii}| - 1)^2 \leq \max_{j=1, \dots, m} (a_{jj} - 1)^2$$

as desired. \square

The last lemma immediately gives the minimax risk for linear estimators in this model.

Lemma 11 (Linear minimax risk in the one-sparse model). *Consider model (14). For $\hat{\theta}_A = Ay$, we have*

$$\inf_{A \in \mathbb{R}^{m \times m}} \max_{\theta \in \Theta_m} E_\theta |\hat{\theta}_A - \theta|^2 = \frac{m\sigma_n^2}{1 + m\sigma_n^2}.$$

Proof. Lemma 10 implies that a linear estimator of θ with minimal maximal risk over Θ_m necessarily has the form $\hat{\theta}_{aI_m} = ay$ for some $a \in \mathbb{R}$. For such an estimator and any $\theta \in \Theta_m$, the bias-variance decomposition gives

$$E_\theta |\hat{\theta}_{aI_m} - \theta|^2 = (a - 1)^2 + m\sigma_n^2 a^2,$$

which can be explicitly minimized at $a^* = \frac{1}{1 + m\sigma_n^2}$ with corresponding minimal risk

$$E_\theta |\hat{\theta}_{a^*I_m} - \theta|^2 = \frac{(m\sigma_n^2)^2}{(1 + m\sigma_n^2)^2} + \frac{m\sigma_n^2}{(1 + m\sigma_n^2)^2} = \frac{m\sigma_n^2}{1 + m\sigma_n^2}.$$

□

B.3 Proof of Theorems 1 and 2

We do not keep explicit track of constants in this version of the proof, noting simply they will finally depend only on the dimension d .

Proof of Theorems 1 and 2. Let $r_d = \frac{1}{2(d+2)}$, $k = \lceil (r_d n)^{\frac{1}{2d+2}} \rceil$ and $m = k^d$. By Lemma 8, there exist orthogonal function $f_1, \dots, f_m \in \mathcal{G}$ with $\|f_j\|_{L^2}^2 = r_d m^{-\frac{2+d}{d}}$. Set $\mathcal{F}_n = \{f_1, \dots, f_n\}$ so that (11) is satisfied with $c_n^2 = r_d m^{-\frac{2+d}{d}}$. Using (16) and Lemma 11,

$$\sup_{f_0 \in \mathcal{G}} E_f \|\bar{f}_n - f\|_{L^2}^2 \geq \max_{f \in \mathcal{F}_n} E_f \|\bar{f}_n - f\|_{L^2}^2 \geq c_n^2 \frac{m\sigma_n^2}{1 + m\sigma_n^2} = \frac{m}{n} \frac{1}{1 + m/(nc_n^2)}$$

since $\sigma_n^2 = \frac{1}{nc_n^2}$ by (13). Since $m = m_n$ satisfies $\frac{m}{c_n^2 n} \simeq n^{-1+1/d} \lesssim 1$, we deduce

$$\max_{f \in \mathcal{F}_n} E_f \|\bar{f}_n - f\|_{L^2}^2 \gtrsim \frac{m}{n} \simeq n^{-\frac{2+d}{2+2d}},$$

which proves Theorem 2. Theorem 1 then follows by applying Lemma 6 with $\mathcal{F}_n = \{f_1, \dots, f_m\} \subset \mathcal{G}$ and $\gamma_n^2 \simeq n^{-\frac{2+d}{2+2d}}$. □

C Background material

C.1 Minimax estimation

Describing the large sample behavior of the smallest achievable worst case risk over all estimation procedures, minimax (estimation) rates are a standard tool to establish statistical optimality of a method. For a statistical model $(P_\theta^n : \theta \in \Theta)$ with sample size n and a loss function ℓ , the minimax risk is $R_n = \inf_{\hat{\theta}_n} \sup_{\theta \in \Theta} E_\theta[\ell(\hat{\theta}_n, \theta)]$, where the infimum is taken over all estimators $\hat{\theta}_n$. The minimax rate is any sequence $(r_n)_n$ such that $r_n \asymp R_n$. Any estimator $\tilde{\theta}_n$ with $R_n \asymp \sup_{\theta \in \Theta} E_\theta[\ell(\tilde{\theta}_n, \theta)]$ is called minimax rate optimal. By definition, the risk of minimax rate optimal estimators is at most a constant factor larger than the minimax risk R_n .

If the posterior contraction rate is ε_n , then under very weak assumptions one can find an estimator with worst case risk of the order ε_n , see Theorem 2.5 [13]. This in turn implies that the posterior cannot contract faster than the minimax rate.

C.2 Sequence representation and posterior mean

We provide here some explanation behind the sequence representation (6) of the Gaussian white noise model (2) and the derivation of the posterior distribution (7). Recall that we can realize a random element $f \sim \Pi$ distributed according to a GP prior on $L^2[0, 1]^d$ via its series expansion (5), namely

$$f = \sum_{k=1}^{\infty} \sqrt{\lambda_k} \xi_k \phi_k, \quad \xi_k \sim^{iid} N(0, 1),$$

known as the *Karhunen-Loève expansion*.

The Brownian motion W in (2) can equivalently be viewed via the action of integration on test functions $g \in L^2[0, 1]^d$ through $W_g = \int_{[0,1]^d} g(x) dW_x$, leading to the mean-zero Gaussian process $W = (W_g : g \in L^2[0, 1]^d)$ indexed by $L^2[0, 1]^d$ with covariance $E(W_g W_h) = \langle g, h \rangle_{L^2}$. In this form, the Gaussian white noise model (2) can be interpreted as observing the Gaussian process $Y = (Y_g : g \in L^2[0, 1]^d)$ with

$$Y_g = \langle f, g \rangle_{L^2} + n^{-1/2} W_g, \quad g \in L^2[0, 1]^d.$$

It is then statistically equivalent to observe the subprocess $(Y_k = Y_{\phi_k} : k \geq 1)$ for any orthonormal basis $\{\phi_k : k \geq 1\}$ of $L^2[0, 1]^d$, in particular the basis corresponding to the Karhunen-Loève expansion of the prior. The white noise model (2) is thus equivalent to observing

$$Y_k = \int_{[0,1]^d} \phi_k(x) dY_x = \langle \phi_k, f \rangle_{L^2} + \frac{1}{\sqrt{n}} \int_{[0,1]^d} \phi_k(x) dW_x =: \theta_k + \frac{w_k}{\sqrt{n}},$$

for $k = 1, 2, 3, \dots$ and where $w_k \sim^{iid} N(0, \|\phi_k\|_{L^2}^2) = N(0, 1)$. The observations $(Y_k : k \geq 1)$ in the last equation and the original white noise model (2) are equivalent in the sense that each can be perfectly recovered from the other as we now explain. One can clearly obtain (Y_k) as in the last display from the whole trajectory $(Y_x : x \in [0, 1]^d)$ by simply computing the integrals $(\int \phi_k(x) dY_x : k \geq 1)$. Conversely, suppose one observes $(Y_k : k \geq 1)$. For any $g \in L^2[0, 1]^d$, one can recover Y_g in the second last display using the basis expansion $g = \sum_k \langle g, \phi_k \rangle_{L^2} \phi_k$ via

$$Y_g = \int g(x) dY_x = \int \sum_{k=1}^{\infty} \langle g, \phi_k \rangle_{L^2} \phi_k(x) dY_x = \sum_{k=1}^{\infty} \langle g, \phi_k \rangle_{L^2} Y_k.$$

Since this holds for arbitrary $g \in L^2[0, 1]^d$, one can reconstruct the process $(Y_g : g \in L^2[0, 1]^d)$ and thus the whole trajectory $(Y_x : x \in [0, 1]^d)$ as in (2). This shows these two representations are equivalent, and thus we may consider either as our ‘data’.

Viewing the prior Π through its series expansion (5), Π can be viewed as a prior on the space of coefficients in the basis expansion of $\{\phi_k\}$ leading to the prior distribution $f = (\theta_k)_k \sim \otimes_{k=1}^{\infty} N(0, \lambda_k)$. Denoting by $P_f = P_f^n$ the distribution of the sequence representation (6), we have

$$P_f = \otimes_{k=1}^{\infty} N(\theta_k, 1/n).$$

Using Kakutani’s product martingale theorem ([8], Theorem 2.7), the measures $(P_f : (\theta_k)_k \in \ell_2)$ are absolutely continuous with respect to $\otimes_{k=1}^{\infty} N(0, 1/n)$ with density

$$e^{\ell_n(f)} = \frac{dP_f}{dP_0} = \exp \left(\sqrt{n} \sum_{k=1}^{\infty} \theta_k Y_k - \frac{n}{2} \sum_{k=1}^{\infty} \theta_k^2 \right) = \exp \left(\sqrt{n} \langle f, Y \rangle_{L^2} - \frac{n}{2} \|f\|_{L^2}^2 \right),$$

so that $e^{\ell_n(f)}$ and $\ell_n(f)$ are the likelihood and log-likelihood of the model, respectively. Note that the likelihood is invariant to the choice of basis $\{\phi_k\}$, but the particular choice can (and will) provide a convenient representation.

Since the likelihood factorizes in terms of the coefficients $(\theta_k)_k$, a prior that makes the $(\theta_k)_k$ independent as in (5) will yield similar independence in the posterior. The posterior distribution is therefore conjugate and takes the form

$$f = \sum_{k=1}^{\infty} \theta_k \phi_k, \quad \theta_k | Y_k \sim^{ind} N \left(\frac{n \lambda_k}{n \lambda_k + 1} Y_k, \frac{\lambda_k}{n \lambda_k + 1} \right),$$

where the exact form follows from standard one-dimensional conjugate computations for the normal likelihood $Y_k | \theta_k = N(\theta_k, 1/n)$ with normal prior $\theta_k \sim N(0, \lambda_k)$. This gives the form of the posterior (7) and its mean (8).

References

- [1] AGAPIOU, S., AND WANG, S. Laplace priors and spatial inhomogeneity in Bayesian inverse problems, 2021.
- [2] AYKROYD, R. G., AND ALJOHANI, H. A Bayesian approach to wavelet-based modelling of discontinuous functions applied to inverse problems. *Comm. Statist. Simulation Comput.* 49, 1 (2020), 207–225.
- [3] BIOUCAS-DIAS, J. M. Bayesian wavelet-based image deconvolution: a GEM algorithm exploiting a class of heavy-tailed priors. *IEEE Trans. Image Process.* 15, 4 (2006), 937–951.
- [4] CASTILLO, I. Lower bounds for posterior rates with Gaussian process priors. *Electron. J. Stat.* 2 (2008), 1281–1299.
- [5] CASTILLO, I., AND ROUSSEAU, J. A Bernstein–von Mises theorem for smooth functionals in semiparametric models. *Ann. Statist.* 43, 6 (2015), 2353–2383.
- [6] CHÉRIEF-ABDELLATIF, B.-E. Convergence rates of variational inference in sparse deep learning. In *Proceedings of the 37th International Conference on Machine Learning* (13–18 Jul 2020), vol. 119 of *Proceedings of Machine Learning Research*, PMLR, pp. 1831–1842.
- [7] COHEN, A., DAUBECHIES, I., AND VIAL, P. Wavelets on the interval and fast wavelet transforms. *Appl. Comput. Harmon. Anal.* 1, 1 (1993), 54–81.
- [8] DA PRATO, G. *An introduction to infinite-dimensional analysis*. Universitext. Springer-Verlag, Berlin, 2006. Revised and extended from the 2001 original by Da Prato.
- [9] DAMIANOU, A., AND LAWRENCE, N. D. Deep Gaussian processes. In *Proceedings of the Sixteenth International Conference on Artificial Intelligence and Statistics* (29 Apr–01 May 2013), vol. 31 of *Proceedings of Machine Learning Research*, PMLR, pp. 207–215.
- [10] DONOHO, D. L., AND JOHNSTONE, I. M. Minimax estimation via wavelet shrinkage. *Ann. Statist.* 26, 3 (1998), 879–921.
- [11] FINOCCHIO, G., AND SCHMIDT-HIEBER, J. Posterior contraction for deep Gaussian process priors, 2021.
- [12] FREEDMAN, D. On the Bernstein-von Mises theorem with infinite-dimensional parameters. *Ann. Statist.* 27, 4 (1999), 1119–1140.
- [13] GHOSAL, S., GHOSH, J. K., AND VAN DER VAART, A. W. Convergence rates of posterior distributions. *Ann. Statist.* 28, 2 (2000), 500–531.
- [14] GHOSAL, S., AND VAN DER VAART, A. W. *Fundamentals of Nonparametric Bayesian Inference*. Cambridge University Press, New York, 2017.
- [15] GINÉ, E., AND NICKL, R. *Mathematical foundations of infinite-dimensional statistical models*. Cambridge Series in Statistical and Probabilistic Mathematics, [40]. Cambridge University Press, New York, 2016.
- [16] HASTIE, T. J., AND TIBSHIRANI, R. J. *Generalized additive models*, vol. 43 of *Monographs on Statistics and Applied Probability*. Chapman and Hall, Ltd., London, 1990.
- [17] HAYAKAWA, S., AND SUZUKI, T. On the minimax optimality and superiority of deep neural network learning over sparse parameter spaces. *Neural Networks* 123 (2020), 343–361.
- [18] IMAIZUMI, M., AND FUKUMIZU, K. Deep neural networks learn non-smooth functions effectively. In *The 22nd International Conference on Artificial Intelligence and Statistics, AISTATS 2019* (2019), vol. 89 of *Proceedings of Machine Learning Research*, PMLR, pp. 869–878.
- [19] IMAIZUMI, M., AND FUKUMIZU, K. Advantage of deep neural networks for estimating functions with singularity on hypersurfaces, 2020.

- [20] JACOBSEN, R. D., AND MØLLER, J. Frequentist and Bayesian inference for Gaussian-log-Gaussian wavelet trees and statistical signal processing applications. *Stat* 6 (2017), 248–256.
- [21] JOHNSTONE, I. Gaussian estimation: Sequence and wavelet models. available from https://imjohnstone.su.domains/GE_09_16_19.pdf, 2019.
- [22] JOHNSTONE, I. M. High dimensional Bernstein–von Mises: simple examples. In *Borrowing strength: theory powering applications—a Festschrift for Lawrence D. Brown*, vol. 6 of *Inst. Math. Stat. (IMS) Collect.* Inst. Math. Statist., Beachwood, OH, 2010, pp. 87–98.
- [23] KOROSTELĚV, A. P., AND TSYBAKOV, A. B. *Minimax theory of image reconstruction*, vol. 82 of *Lecture Notes in Statistics*. Springer-Verlag, New York, 1993.
- [24] LAURENT, B., AND MASSART, P. Adaptive estimation of a quadratic functional by model selection. *Ann. Statist.* 28, 5 (2000), 1302–1338.
- [25] MÜLLER, P., AND VIDAKOVIC, B., Eds. *Bayesian inference in wavelet-based models*, vol. 141 of *Lecture Notes in Statistics*. Springer-Verlag, New York, 1999.
- [26] NEAL, R. M. *Bayesian Learning for Neural Networks*. Springer-Verlag, Berlin, Heidelberg, 1996.
- [27] PETERSEN, P., AND VOIGTLAENDER, F. Optimal approximation of piecewise smooth functions using deep ReLU neural networks. *Neural Networks* 108 (2018), 296–330.
- [28] POLSON, N. G., AND ROČKOVÁ, V. Posterior concentration for sparse deep learning. In *Advances in Neural Information Processing Systems* (2018), pp. 930–941.
- [29] RASMUSSEN, C. E., AND WILLIAMS, C. K. I. *Gaussian processes for machine learning*. Adaptive Computation and Machine Learning. MIT Press, Cambridge, MA, 2006.
- [30] RAY, K., SZABO, B., AND CLARA, G. Spike and slab variational Bayes for high dimensional logistic regression. In *Advances in Neural Information Processing Systems* (2020), vol. 33, Curran Associates, Inc., pp. 14423–14434.
- [31] RAY, K., AND SZABÓ, B. Variational Bayes for high-dimensional linear regression with sparse priors. *Journal of the American Statistical Association* 0, 0 (2021), 1–12.
- [32] RAY, K., AND VAN DER VAART, A. Semiparametric Bayesian causal inference. *Ann. Statist.* 48, 5 (2020), 2999–3020.
- [33] REISS, M. Asymptotic equivalence for nonparametric regression with multivariate and random design. *Ann. Statist.* 36, 4 (2008), 1957–1982.
- [34] ROCKOVA, V., AND ROUSSEAU, J. Ideal Bayesian spatial adaptation, 2021.
- [35] SCHMIDT-HIEBER, J. Nonparametric regression using deep neural networks with ReLU activation function. *Ann. Statist.* 48, 4 (2020), 1875–1897.
- [36] SCHMIDT-HIEBER, J. The Kolmogorov–Arnold representation theorem revisited. *Neural Networks* 137 (2021), 119–126.
- [37] SUZUKI, T. Adaptivity of deep ReLU network for learning in Besov and mixed smooth Besov spaces: optimal rate and curse of dimensionality. In *7th International Conference on Learning Representations, ICLR 2019, New Orleans, LA, USA, May 6-9, 2019* (2019), OpenReview.net.
- [38] SUZUKI, T., AND NITANDA, A. Deep learning is adaptive to intrinsic dimensionality of model smoothness in anisotropic Besov space. In *Advances in Neural Information Processing Systems* (2021), vol. 34, Curran Associates, Inc., pp. 3609–3621.
- [39] TANG, J., MENG, Z., NGUYEN, X., MEI, Q., AND ZHANG, M. Understanding the limiting factors of topic modeling via posterior contraction analysis. In *Proceedings of the 31st International Conference on Machine Learning* (Beijing, China, 22–24 Jun 2014), vol. 32 of *Proceedings of Machine Learning Research*, PMLR, pp. 190–198.

- [40] TELGARSKY, M. Benefits of depth in neural networks. In *29th Annual Conference on Learning Theory* (Columbia University, New York, New York, USA, 23–26 Jun 2016), vol. 49 of *Proceedings of Machine Learning Research*, PMLR, pp. 1517–1539.
- [41] TSUJI, K., AND SUZUKI, T. Estimation error analysis of deep learning on the regression problem on the variable exponent Besov space. *Electronic Journal of Statistics* 15, 1 (2021), 1869 – 1908.
- [42] VAN DER VAART, A. W. *Asymptotic statistics*, vol. 3 of *Cambridge Series in Statistical and Probabilistic Mathematics*. Cambridge University Press, Cambridge, 1998.
- [43] VAN DER VAART, A. W., AND VAN ZANTEN, J. H. Rates of contraction of posterior distributions based on Gaussian process priors. *Ann. Statist.* 36, 3 (2008), 1435–1463.
- [44] VÄNSKÄ, S., LASSAS, M., AND SILTANEN, S. Statistical X-ray tomography using empirical Besov priors. *Int. J. Tomogr. Stat.* 11, S09 (2009), 3–32.
- [45] WANG, Y., AND BLEI, D. Variational Bayes under model misspecification. In *Advances in Neural Information Processing Systems* 32. 2019, pp. 13357–13367.
- [46] WANG, Y., AND BLEI, D. M. Frequentist consistency of variational Bayes. *J. Amer. Statist. Assoc.* 114, 527 (2019), 1147–1161.
- [47] YAROTSKY, D. Error bounds for approximations with deep ReLU networks. *Neural Networks* 94 (2017), 103–114.
- [48] YAROTSKY, D. Optimal approximation of continuous functions by very deep ReLU networks. In *Proceedings of the 31st Conference On Learning Theory* (06–09 Jul 2018), vol. 75 of *Proceedings of Machine Learning Research*, PMLR, pp. 639–649.