

AperTO - Archivio Istituzionale Open Access dell'Università di Torino

**NK(1) receptor activation leads to enhancement of inhibitory neurotransmission in spinal substantia gelatinosa neurons of mouse**

**This is the author's manuscript**

*Original Citation:*

*Availability:*

This version is available <http://hdl.handle.net/2318/1908> since

*Terms of use:*

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)



## UNIVERSITÀ DEGLI STUDI DI TORINO

This Accepted Author Manuscript (AAM) is copyrighted and published by Elsevier. It is posted here by agreement between Elsevier and the University of Turin. Changes resulting from the publishing process - such as editing, corrections, structural formatting, and other quality control mechanisms - may not be reflected in this version of the text. The definitive version of the text was subsequently published in

***NK1 receptor activation leads to enhancement of inhibitory neurotransmission in spinal substantia gelatinosa neurons of mouse, Pain, 2 (1-2), 2004, <http://dx.doi.org/10.1016/j-pain.2004.07.018>***

You may download, copy and otherwise use the AAM for non-commercial purposes provided that your license is limited by the following restrictions:

- (1) You may use this AAM for non-commercial purposes only under the terms of the CC-BY-NC-ND license.
- (2) The integrity of the work and identification of the author, copyright owner, and publisher must be preserved in any copy.
- (3) You must attribute this AAM in the following format: Creative Commons BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/deed.en>), ,  
***<http://dx.doi.org/10.1016/j-pain.2004.07.018>***

# NK<sub>1</sub> receptor activation leads to enhancement of inhibitory neurotransmission in spinal substantia gelatinosa neurons of mouse

Angela Maria Vergnano<sup>a</sup>, Chiara Salio<sup>a</sup>, Adalberto Merighi<sup>a,b,\*</sup>

<sup>a</sup>Department of Veterinary Morphophysiology, Via Leonardo da Vinci 44, 10095 Grugliasco, Italy

<sup>b</sup>Rita Levi-Montalcini Center for Brain Repair, Via Leonardo da Vinci 44, 10095 Grugliasco, Italy

## Abstract

Substance P (SP) is a well-established pain messenger in the spinal cord, although its role in substantia gelatinosa (lamina II) still remains elusive. We carried out patch-clamp recordings on lamina II neurons from transverse mouse spinal cord slices (P8–12), using the selective NK<sub>1</sub> receptor agonist [Sar<sup>9</sup>,Met(O<sub>2</sub>)<sup>11</sup>]-SP (SM-SP, 3–5 μM) in the presence of NBQX. Activation of NK<sub>1</sub> receptors was confirmed after pre- incubation with selective NK<sub>1</sub> antagonist L732,138 (4 μM) that consistently blocked the effects of SM-SP (nine neurons). After SM-SP challenge and spontaneous inhibitory post-synaptic current (sIPSC) analysis, 50% of recorded neurons (15 out of 30) were found to display a transient increase in frequency; in five neurons this was also associated with increase of peak amplitude. Five out of eight neurons displayed pure GABA<sub>A</sub> receptor-mediated sIPSCs, whereas the remaining ones showed mixed GABAergic/glycinergic events. After miniature IPSC analysis, a significant increase in frequency was observed in three out of 14 SM-SP responsive neurons. At least four different morphological types were apparent among NK<sub>1</sub>-responsive neurons after filling with Lucifer Yellow/biocytin: fusiform with dorso-ventral dendritic arbors (i); round-to-oval with dendritic arborization mainly directed to lamina I (ii) or III (iii), and round-to-oval with dendrites sparsely distributed all around the cell body (iv). Thus, there was no correlation between morphology and electrophysiological properties of responsive neurons. Our observations provide new insights on the processing of sensory neurotransmission in spinal cord, and indicate that activation of NK<sub>1</sub> receptors is involved in the maintenance of the inhibitory tone of substantia gelatinosa interneurons.

Keywords: NK<sub>1</sub> receptors; substantia gelatinosa neurons; spontaneous inhibitory post-synaptic current; substance P

\* Corresponding author. Address: Department of Veterinary Morphophysiology, Via Leonardo da Vinci 44, 10095 Grugliasco Italy. Cell: +39-011-6709-118; fax: C39-011-6709-138. E-mail address: adalberto.merighi@unito.it (A. Merighi).

## 1. Introduction

Several lines of evidence have implicated substance P (SP) as a major neurotransmitter of pain (see Afrah et al., 2001; Basbaum, 1999; Mantyh, 2002; Saria, 1999; Urban and Gebhart, 1999; Woolf et al., 1998). SP is synthesized in nociceptors of the dorsal root ganglia which send their axons (C and Ad fibers) to dorsal horn projection neurons in laminae I and IV–V and to nocispecific interneurons in laminae II–III (see Millan, 1999), and is released in vivo upon activation of C and Ad fibers (Afrah et al., 2001; Duggan and Furnidge, 1994; Hua et al., 1986; Teoh et al., 1996). Administration of morphine or opioid peptides specifically blocks this release (Jessell and Iversen, 1977; Takano et al., 1993; Yaksh et al.,

1980), while in hyperalgesic conditions release of SP is increased (Coderre and Yashpal, 1994; Garry and Hargreaves, 1992; Meller and Gebhart, 1994; Oku et al., 1987a,b). SP evokes slow excitatory post-synaptic potentials (EPSPs) in second-order sensory neurons in the dorsal horn (De Koninck and Henry, 1991; De Koninck et al., 1992; Otsuka and Yoshioka, 1993), and mediates central sensitization (Xu et al., 1992), leading to hyperalgesia and allodynia (McMahon et al., 1993; Woolf and Costigan, 1999; Yaksh et al., 1999). These altered conditions of pain sensitivity are attenuated by blocking of the SP preferred receptor NK<sub>1</sub> (Honor et al., 1999; Mantyh et al., 1997; Nichols et al., 1999).

Despite the wealth of data, mechanisms of SP action remain elusive in the substantia gelatinosa (SG—lamina II), which is traditionally considered to play a pivotal role in modulating nociceptive transmission (Ribeiro-Da-Silva, 2003; Willis and Coggeshall, 1991; Yoshimura, 1996). Different studies have concluded that SP was responsible for the generation of slow EPSPs elicited by C and Ad fiber stimulation in the dorsal horn (De Koninck and Henry, 1991; Otsuka and Yoshioka, 1993; Yoshimura and Jessell, 1989, 1990; Yoshimura and Nishi, 1993; Yoshimura et al., 1993). However, administration of SP or synthetic agonists of the SP NK<sub>1</sub> receptor evoked an excitatory response in a very limited subset of SG neurons in other studies (Bleazard et al., 1994; Yoshimura et al., 1993), and it was thus proposed that the peptide does not play a role in lamina II. This lack of effect has been considered resulting from the low density of NK<sub>1</sub> receptor-immunoreactive neurons in the SG (Ribeiro-Da-Silva et al., 2001) but this explanation is weakened by the observation that SP activates more than a single intracellular pathway: it displays maximal affinity for its preferred NK<sub>1</sub> receptor (Ikeda et al., 2003), but also binds other NK receptors (Severini et al., 2002), and modulates N-methyl-D-aspartate (NMDA) neurotransmission (Budai and Larson, 1996; Chizh et al., 1995; Cumberbatch et al., 1995; Rusin et al., 1993).

To further analyze the effects of SP in SG, we made electrophysiological recordings from mouse spinal cord slices. We found that an NK<sub>1</sub> receptor-specific agonist enhances inhibitory neurotransmission in lamina II neurons.

## 2. Methods

### 2.1. Animals

Studies were performed on 35 neonatal mice (8–12 days old). All experimental procedures were approved by the Committee of Bioethics and Animal Welfare of the University of Torino. After deep sodium pentobarbital anesthesia (30 mg/kg), animals were decapitated, a laminectomy was performed and the spinal cord quickly removed and placed in an ice-cold artificial cerebrospinal fluid (ACSF) containing (in mM): 125 NaCl, 2.5 KCl, 25 NaHCO<sub>3</sub>, 1 NaHPO<sub>4</sub>, 25 glucose, 1 MgCl<sub>2</sub>, and 2 CaCl<sub>2</sub>, saturated with 95% O<sub>2</sub>:5% CO<sub>2</sub>. Transverse vibratome slices 350 μm thick were kept in ACSF at 35 °C for 30 min. Slices were subsequently maintained at room temperature until used for the experiments.

### 2.2. Electrophysiological recordings

All experiments were performed at room temperature by placing a slice in a recording chamber constantly perfused at 2 ml/min with oxygenated ACSF. Neurons were visually identified using a fixed-stage microscope (Axioskop 1, Zeiss, Germany), equipped with infrared gradient contrast optics (Luigs and Neumann, Germany) and a 63x insulated water immersion objective (Achromplan, Zeiss). Patch-clamp whole cell recordings were obtained with an Axopatch 200B amplifier (Axon Instruments, USA). Patch pipettes were pulled from borosilicate glass tubing (WPI, USA) and had a resistance of 4–7 MU when filled with the intracellular solution. To study spontaneous excitatory post-synaptic currents (sEPSCs), pipettes were filled with an intracellular solution containing (in mM): 145 Kgluconate, 5 EGTA, 2 MgCl<sub>2</sub>, 10 HEPES, 2 ATPNa, and 0.2 GTPNa, pH 7.2 (with KOH).

To study spontaneous inhibitory post-synaptic currents (sIPSCs), the intracellular solution contained (in

mM): 145 KCl, 5 EGTA, 2 MgCl<sub>2</sub>, 10 Hepes, 2 ATPNa, and 0.2 GTPNa, pH 7.2 (with KOH). Under these conditions sIPSCs were detected as inward currents. In some cases, 0.5% biocytin (Sigma, USA) and 0.1% Lucifer Yellow (LY, Sigma, USA) were added.

Lamina II neurons were patched under visual control, and only cells with membrane potential more negative than K55 mV were considered for the recordings. Currents were sampled at 10 kHz, filtered at 2 kHz, and analyzed with MiniAnalysis (Synaptosoft, USA) and pClamp 8 (Axon Instruments, USA) software. On an average, 30–80 IPSCs were analyzed from each cell in order to obtain the mean kinetic and amplitude parameters. From the average of these events, we measured the rise time, calculated from 10 to 90% peak amplitude, and the values of decay time constants (expressed as t<sub>1</sub> and t<sub>2</sub>) by fitting the 10–90% region of the decay phase with a biexponential function. All electrophysiological values were expressed as mean GSEM with n, indicating the number of cells.

### 2.3. Drugs

All drugs were bath applied. [Sar<sup>9</sup>,Met(O<sub>2</sub>)<sup>11</sup>]-substance P (SM-SP), an analog of SP that shows greater selectivity for NK<sub>1</sub> receptors and greater potency and duration of action than SP *in vivo* (Regoli et al., 1988; Sakurada et al., 1994; Tousignant et al., 1989) was used to study the effects of NK<sub>1</sub> activation in slices. SM-SP was delivered continuously during the recording in all experiments. N-Acetyl-L-tryptophan-3,5-bistrifluoromethyl benzyl ester (L732,138), a potent selective antagonist of the NK<sub>1</sub> receptor (Cascieri et al., 1994) was also used in certain experiments to block the effects of SM-SP and thus assess the specificity of NK<sub>1</sub> receptor activation. 2,3-Dioxo-6-nitro-1,2,3,4-tetrahydrobenzo[f]quinoxaline-7-sulfonamide disodium salt (NBQX), SM-SP and L732,138 were from Tocris (UK) and tetrodotoxin (TTX), bicuculline methiodide, and strychnine methiodide from Sigma (USA).

### 2.4. Histological procedures

After recording, slices were fixed in 0.1 M phosphate buffer (pH 7.4) containing 4% paraformaldehyde and 2.5% glutaraldehyde overnight at 4 °C, and subsequently rinsed several times in phosphate-buffered saline (PBS, pH 7.4). Slices were then mounted in a fluorescence-free medium or processed further for the visualization of intracellular biocytin by the ABC method (Vector, USA). Briefly, slices were incubated in PBS containing 10% methanol and 3% hydrogen peroxide for 30 min to suppress endogenous peroxidase activity, thoroughly rinsed in PBS, and then incubated in ABC for 2.5 h at room temperature. After extensive washing in PBS, the biocytin-bound peroxidase was revealed using nickel ammonium sulphate-intensified 3,3'-diaminobenzidine (DAB) for 60 min at room temperature (0.025% DAB, 1% nickel ammonium sulfate, 0.04% H<sub>2</sub>O<sub>2</sub> in PBS).

Labeled neurons were observed with a light microscope (Axiophot 1, Zeiss) under bright field or fluorescence illumination and photographed with a high-resolution digital camera (Coolpix 995, Nikon, Japan). Digital images were further processed using Adobe Photoshop 5 (USA).

The immunocytochemical visualization of NK<sub>1</sub> receptor distribution within the dorsal horn was performed on free-floating neonatal mouse spinal cord sections incubated in PBS containing 10% methanol and 3% hydrogen peroxide for 30 min to suppress endogenous peroxidase activity, and then blocked in 0.02 M PBS containing 6% normal goat serum (NGS) for 30 min at room temperature, before treatment with an affinity-purified polyclonal rabbit anti-NK<sub>1</sub> receptor antibody (Sigma, USA; Mantyh et al., 1995), diluted 1:30,000 in PBS containing 1% NGS. After washing in PBS, sections were incubated in 1:250 biotinylated anti-rabbit IgG (Vector, USA) and then in 1:100 avidin-biotinylated-peroxidase complex (Vector, USA). The peroxidase reaction was developed using 0.025% (w/v) DAB (Sigma, USA) and 0.04% (v/v) hydrogen peroxide. Sections were then transferred onto glass slides and mounted in PBS-glycerol.

### 3. Results

#### 3.1. Electrophysiology

Seventy one neurons in lamina II were voltage-clamped at a holding potential ( $V_h$ ) of K63 mV and challenged with SM-SP (3–5  $\mu$ M). Since both sEPSCs and sIPSCs were evident after SM-SP superfusion in the absence of specific receptor antagonists, we employed different intracellular solutions to better discriminate between the two types of current.

Using the low  $Cl^-$  intracellular solution, bath-application of SM-SP induced a transient increase in both amplitude and frequency of sEPSCs (from 1.5 to 5 Hz) in two out of 18 cells recorded (Fig. 1).

sIPSCs were recorded using the high  $Cl^-$  intracellular solution in the presence of 10  $\mu$ M NBQX to block AMPA/kainate neurotransmission (illustrative traces are shown in Fig. 2A–C).




Fig. 1. Representative traces of sEPSCs recorded from a SG neuron under effects of SM-SP. Whole cell recordings of sEPSCs from a SG neuron in control (A), and under bath-applied SM-SP (B). Upon SM-SP perfusion this cell displayed an increase in both frequency and amplitude of sEPSCs. The holding potential ( $V_h$ ) was K63 mV.

In 15 out of 30 neurons recorded (Fig. 2F), activation of NK<sub>1</sub> receptors elicited a strong increase in sIPSC frequency (from  $0.6 \pm 0.2$  to  $1.6 \pm 0.3$  Hz,  $P < 0.05$ ). However, this was not observed over the entire SM-SP application: increase in the sIPSC frequency lasted only for about 60–120 s, as shown in Fig. 2A. This transitory effect was likely due to internalization of NK<sub>1</sub>. Studies *in vivo* (Mantyh et al., 1995) have indeed shown that NK<sub>1</sub> receptors are internalized after noxious stimulation, and that, as internalization proceeds, there are parallel changes in pain-related behaviors (Mantyh, 2002).

Rise and decay time constants of sIPSCs were not affected by application of SM-SP. The mean rise time for the control was  $1.1 \pm 0.3$  and  $1.2 \pm 0.4$  ms in the presence of SM-SP ( $n = 15$ ,  $P > 0.05$ ). The mean decay values were:  $t_1 = 6.6 \pm 1.1$  ms,  $t_2 = 40 \pm 5$  ms for the control and  $t_1 = 5.9 \pm 1.2$  ms,  $t_2 = 42 \pm 5$  ms in the presence of SM-SP ( $n = 15$ ,  $P > 0.05$ ). Average events calculated from one neuron recorded under control conditions and in the presence of SM-SP are shown in D and E, respectively.

In five neurons, the presence of SM-SP also led to an increase in sIPSC peak amplitude. In these cells, the mean amplitude values were  $37.4 \pm 1.4$  pA in controls and  $52.8 \pm 1.7$  pA in the presence of SM-SP ( $P < 0.05$ , Fig. 2G). This change was unlikely to arise via a post-synaptic effect, as SM-SP still increased the frequency in the presence of TTX, without affecting the amplitude (see below).

To document that sIPSCs were indeed generated by specific NK<sub>1</sub> receptor signaling, slices were superfused with 4  $\mu$ M of the specific NK<sub>1</sub> receptor antagonist L732,138 for at least 8 min before SM-SP challenge (Kombian, 2003a,b; Yang et al., 2000). Nine cells were successfully recorded and their outputs statistically analyzed (Fig. 3). In these cells, the frequency of sIPSCs passed from  $0.26 \pm 0.04$  Hz upon L732,138 superfusion to  $0.29 \pm 0.06$  Hz following SM-SP (5  $\mu$ M;  $P > 0.05$ ), and amplitude from  $41.08 \pm 6.77$  to  $41.31 \pm 5.48$  pA ( $P > 0.05$ ). This lack of statistical significance demonstrates that the effects of SM-SP on inhibitory neurotransmission are specifically due to NK<sub>1</sub> receptor activation; it

also rules out concerns about




Fig. 2. Effects of SM-SP on sIPSCs recorded from SG neurons. (A–C) Whole cell recordings of sIPSCs. (A) In the presence of 10  $\mu$ M NBQX, SM-SP (5  $\mu$ M) evoked a transient increase in frequency and amplitude of sIPSCs which returned to control values after about 100 s still in the presence of the agonist. The arrowhead indicates the start of SM-SP application.  $V_h$  ZK63 mV. (B and C) Effect of SM-SP on the frequency and amplitude of sIPSCs recorded from another SG neuron (B, control; C, SM-SP). (D and E) Averages of sIPSCs detected in control (D) and in presence of SM-SP (E); same neuron as in B and C. Note that superfusion with SM-SP elicited an increase in both amplitude and frequency without affecting sIPSC kinetics. (F) Pooled data from 15 lamina II neurons reveal a significant increase in sIPSC frequency upon SM-SP superfusion. The mean frequency passed from  $0.6 \pm 0.2$  Hz in control to  $1.6 \pm 0.3$  Hz in the presence of the agonist ( $n=15$ ,  $P<0.05$ ). (G) In five neurons, the increase in sIPSC frequency was accompanied by a significant increase in amplitude (from  $37.4 \pm 1.4$  pA in control to  $52.8 \pm 1.7$  pA under SM-SP) ( $n=5$ ,  $P<0.05$ ).

selectivity of the agonist at the relatively high concentration employed here in comparison with most electrophysiological studies (1–4  $\mu$ M) on rat brain/spinal cord slices or isolated neurons (King et al., 1997; Kombian, 2003a, b; Li and Zhao, 1998). Nonetheless it should be noted that up to 20  $\mu$ M, SM-SP has been used to study substance P-mediated excitation and expression of the transcription factor FOS in rat dorsal horn neurons in vitro without any noticeable loss of agonist specificity (Badie-Mahdavi et al., 2001). The possible existence of interspecies differences in NK receptor agonist affinities should also be considered (Nsa Allogho et al., 1997).

In order to further study the sIPSCs, 10  $\mu$ M bicuculline and/or 1  $\mu$ M strychnine was added to the bath solution at the end of each experiment performed on eight SM-SP responsive neurons (Fig. 4). In five of these cells, sIPSCs were completely blocked by bicuculline, revealing that they were mediated by

GABA<sub>A</sub> receptor activation only (Fig. 4B). The remaining cells (3/8) displayed mixed GABAergic and glycinergic events, although the latter were very rare (Fig. 4D). This was not surprising since: (i) co-existence of GABA/glycine has been widely reported in the superficial dorsal horn (Keller et al., 2001; Maxwell et al., 1995; Todd et al., 1996); and (ii) functionally active




Fig. 3. SM-SP effects on sIPSCs are blocked by the selective NK1 antagonist L732,138. (A) sIPSCs recorded from a lamina II neuron in the presence of 4  $\mu$ M L732,138 (left) and 4  $\mu$ M L732,138 + 5  $\mu$ M SM-SP (right).  $V_h$  ZK63 mV. When slices were pre-treated with the selective NK1 receptor antagonist, SM-SP no longer elicited a significant increase in sIPSC frequency and amplitude. (B) Pooled data from nine lamina II neurons. The frequency of sIPSCs was  $0.26 \pm 0.04$  Hz upon L732,138 superfusion and  $0.29 \pm 0.06$  Hz during SM-SP administration ( $P > 0.05$ ), and mean amplitude passed from  $41.08 \pm 6.77$  to  $41.31 \pm 5.48$  pA ( $P > 0.05$ ).

GABA/glycine co-synapses exist in the dorsal horn, at least in certain stages of development (Jonas et al., 1998; Keller et al., 2001).

To better characterize the nature of NK1 activation in lamina II neurons, we recorded miniature IPSCs (mIPSCs) in the presence of 1  $\mu$ M TTX. The mean mIPSC frequency was  $0.2 \pm 0.1$  Hz ( $n=14$ ), and a clear effect of SM-SP was only observed in three out of 14 neurons recorded (21.4%). In these cells, a significant increase ( $150 \pm 20\%$ ) in mIPSC frequency was detected, compared to the control (Fig. 5). Moreover one of these three neurons responded to SM-SP application by a slow inward current that peaked at 30 pA and was associated with an increase in amplitude (from  $18 \pm 7$  to  $34 \pm 12$  pA).

### 3.2. Morphological characterization of recorded SG neurons

Fourteen of the recorded neurons, were successfully injected with LY or biocytin to reveal their morphology. All labeled neurons were located in the SG (Fig. 6A–C), and exhibited different dendritic arrangements and orientations. On this basis, we observed several different morphological types: (i) fusiform neurons with two funnel-shaped dorso-ventrally oriented dendritic arbors; (ii) rounded neurons with a funnel-shaped dendritic arbor originating from a main dendritic trunk primarily directed to lamina I;






Fig. 4. sIPSCs of SM-SP-responsive neurons are mainly mediated by GABA<sub>A</sub> receptor. (A and B) Recordings from a lamina II neuron in the presence of 10  $\mu$ M NBQX and SM-SP challenge. Block of GABA<sub>A</sub> receptors with 10  $\mu$ M bicuculline (B) totally suppressed activity,  $V_h = -63$  mV. (C–E) Recordings from another lamina II neuron in the presence of 10  $\mu$ M NBQX and SM-SP challenge. The block of GABA<sub>A</sub> receptors with 10  $\mu$ M bicuculline (D) did not completely suppress inhibitory activity.  $V_h = -63$  mV. Note a single glycinergic event of small amplitude (arrow) in about 30 s. The frequency of glycinergic events in this cell was 0.01 Hz. (E) Co-application of bicuculline and strychnine completely suppressed sIPSCs.

(iii) rounded neurons with a funnel-shaped dendritic arbor originating from a main dendritic trunk primarily directed to lamina III; and (iv) oval-to-rounded neurons with an overall spherical distribution of the dendritic arborization that originated from several independent dendrites. The fusiform neurons (three in all) were located in lamina II<sub>O</sub> (Fig. 6A). Only one of them was SM-SP-responsive. All had a long dorso-ventrally oriented axis with primary beaded dendrites arising from the two opposite poles of the cell body (Fig. 6A and B). Dendritic branches were mainly restricted to lamina II, and displayed sparse spines. Of the seven rounded neurons, four were located in lamina II<sub>O</sub>, and three in II<sub>I</sub> (Fig. 6C). Three rounded neurons




Fig. 5. mIPSCs recorded from SG neurons. (A) mIPSCs recorded from a lamina II neuron in the presence of 1  $\mu$ M TTX. (B) Activation of NK<sub>1</sub> receptors induced an increase in mIPSC frequency;  $V_{h} = -63$  mV. (C) Cumulative amplitude plot of mIPSCs recorded from a lamina II neuron in control (gray line) and in the presence of 5  $\mu$ M SM-SP (black line). Note that the mean peak amplitude was not significantly affected by perfusion of the NK<sub>1</sub> agonist.

were SM-SP-responsive. Irrespective of their sublaminar location these cells had two or more primary beaded dendrites projecting to lamina I (Fig. 6D) or lamina III (not shown). The dendrites had no spines, and had a rather thin constant diameter along their whole length (Fig. 6D). The axon, when visible, projected to lamina III.

Of the four oval-to-rounded neurons, three were located in lamina II<sub>O</sub>, and one in lamina II<sub>j</sub>. Two oval-to-rounded neurons were SM-SP-responsive. Their spherical dendritic tree was confined within the SG, and the axon, when filled by the tracer, was seen to project ventrally for a short distance.

### 3.3. Distribution of NK<sub>1</sub> receptor immunoreactivity in the dorsal horn

Labeling of mouse spinal cord sections with the anti-NK<sub>1</sub> receptor antibody (Fig. 6E and F) resulted in intense staining within the superficial dorsal horn, particularly in lamina I, where a dense meshwork of neuronal processes, mainly oriented on a transversal plane, was detected at all levels of the cord. The superficial dorsal horn also contained numerous immunoreactive fusiform cell bodies with bipolar dendritic arborization.

Staining in lamina II was far less prominent, revealing a looser network of processes, mainly oriented on a longitudinal plane and more densely packed in lamina II<sub>O</sub>. Some NK<sub>1</sub>-immunoreactive cell bodies were also detected in this lamina; they were oval-to-fusiform, and gave rise to positive dorso-ventrally oriented dendrites.

In deeper laminae, immunostaining was limited to scattered processes and isolated neurons of larger size. These cells usually displayed an irregularly polygonal perykarion, with few long dendrites traveling across to more superficial or ventral laminae of the dorsal horn.

## 4. Discussion

### 4.1. Effects of SM-SP administration

Whereas anatomical, functional, and recombinant DNA studies have shown that SP is a fundamental messenger of certain types of pain conveyed to supraspinal centers by lamina I and III–V projection neurons (Basbaum, 1999; De Felipe et al., 1998; Honor et al., 1999; Ikeda et al., 2003; Mantyh, 2002; Mantyh et al., 1997; McLeod et al., 1999; Millan, 1999; Nichols et al., 1999; Todd, 2002; Todd et al., 2002; Woolf et al., 1998), the role of SP in lamina II has remained unclear. In this study, about 50% of recorded neurons in mouse SG responded to NK1 receptor stimulation with a SP synthetic analog by showing a transient increase in inhibitory spontaneous activity. Such an effect was mainly due to GABA<sub>A</sub> receptor activation. Nonetheless, in neurons displaying mixed GABAergic/glycinergic activity, activation of NK1 receptors was likely to enhance the release of glycine also, but because of the very low frequency of isolated glycinergic events we did not pursue this issue.

Although an inhibitory role of SP might be surprising in light of its well-known role as a positive pain modulator, numerous reports in an array of different experimental contexts and species support our present findings. When the release of amino acids evoked by SP in neonatal rats was examined in the isolated spinal cord (Sakuma et al., 1991) or in a hemisectioned spinal cord preparation (Maehara et al., 1995), it was demonstrated that 10  $\mu$ M SP evoked a significant increase in the basal release of GABA and glycine. Moreover in cats, SP is released from primary nociceptive afferent terminals, and activates second-order GABAergic interneurons in the dorsal horn (Wei and Zhao, 1996). Similarly, SP facilitated the strychnine-sensitive glycine response in neurons acutely dissociated from the rat sacral dorsal commissural nucleus, leading to conclude that SP may suppress nociception in the spinal cord (Wang et al., 1999). Finally, in CNS areas other than the spinal cord, SP was also shown to stimulate GABA-mediated inhibitory synaptic transmission in vitro (Kombian, 2003a, b; Kouznetsova and Nistri, 1998; Maubach et al., 2001; Ogier and Raggenbass, 2003; Stacey et al., 2002). Taken together, these studies demonstrate that release of GABA/ glycine by SP is relevant in vivo, and our observations show that SG is a further CNS site in which such a release is also relevant.




Fig. 6. Morphology of SG neurons recorded, and immunocytochemical distribution of the NK1 receptor. (A–D) Biocytin-injected neurons in lamina II of transverse spinal cord slices. A SM-SP-responsive neuron in lamina II<sub>o</sub> (A) is shown at higher magnification in (B). This neuron, fusiform in shape, had a long axis extended dorso-ventrally with primary dendrites arising from the two poles of the cell body. Arrows point to beaded dendrites. An SM-SP-responsive neuron in lamina II<sub>j</sub> (C) is shown at higher magnification in (D). This neuron, round in shape, had a dendritic arborization (arrows) mainly directed to lamina I. An axon process, emerging from the soma, projected to lamina III (arrowheads). (E and F) Distribution of NK1 receptor immunoreactivity. A dense meshwork of immunoreactive neuronal processes is spread throughout in lamina I. An NK1-positive fusiform neuron in

lamina II displays an immunoreactive dendrite directed to deeper laminae (E, arrow). A fusiform lamina III neuron, with a long dendrite projecting to lamina II, is intensely labeled for NK<sub>1</sub> (F, arrowhead). Note that lamina I neurons (F, arrows) and processes are also NK<sub>1</sub>-immunoreactive. Scale bars: A, C=200 μm; B, D=50 μm; E, F=40 μm.

#### 4.2. Morphological characterization of SM-SP responsive neurons in SG

Numerous studies have examined the morphology of neurons in the SG of the spinal cord (Beal and Bice, 1994; Beal et al., 1988; Bennett et al., 1980; Bicknell and Beal, 1984; Gobel, 1975, 1978; Gobel et al., 1980; Todd and Lewis, 1986). Several different types and differences have been reported among species (see Ribeiro-Da-Silva, 2003). The two most common neuronal types described by Cajal (1952), i.e. the central cells, which are widely distributed throughout the lamina, and the marginal cells, which are concentrated at the border with lamina I, have been referred to, respectively, as islet and stalked cells in cats (Gobel, 1975) and rats (Todd and Lewis, 1986). Of the SM-SP-responsive neurons in this study, three could be classified as stalked cells, two as islet cells, and one did not belong to any of these categories. This lack of structure-to-function correlation may be due to the fact that maturation of lamina II neuron dendrites occurs after birth, with extensive rearrangement until the adult pattern is attained (Bicknell and Beal, 1984; Falls and Gobel, 1979). However, in keeping with our findings, numerous other studies have failed to show a correlation between morphology and electrophysiological properties of SG neurons (see Ribeiro-Da-Silva, 2003).

#### 4.3. Circuitry involved in the response of SG neurons to SM-SP

SP, neurokinin A (NKA) and neurokinin B (NKB) belong to a family of peptides, the tachykinins, all implicated in nociception, and they, respectively, act upon preferred receptors, namely NK<sub>1</sub>, NK<sub>2</sub> and NK<sub>3</sub>, and cross-talk between different tachykinins at different receptors is known to occur (Regoli et al., 1987, 1994; Saria, 1999; Severini et al., 2002). Existence of NK<sub>1</sub> receptor subtypes has been proposed: two types of NK<sub>1</sub> receptors were originally described with preference for either septide or Sar9/-SP-sulphone (Fox et al., 1996). More recently, a septide-sensitive and a 'classical' conformation of NK<sub>1</sub> receptor have been proposed to explain different tachykinin affinities and signal transduction pathways (Maggi and Schwartz, 1997). Since SM-SP selectively targets the NK<sub>1</sub> receptor in its classical conformation, a proportion of SP-sensitive SG neurons might have been missed in this study. Nonetheless this does not invalidate, but rather reinforces, the functional significance of our findings *in vivo* where it seems possible that SP's effect on inhibitory neurotransmission is greater than in our experimental conditions.

As to the circuitry involved in the response of SG neurons to SM-SP, one can hypothesize that: (i) responsive cells receive a direct synaptic input from GABAergic (and/or glycinergic) neurons expressing NK<sub>1</sub> receptors at their membrane; and/or (ii) a polysynaptic chain is activated consisting (in its simplest configuration) of a first synapse between a pre-synaptic excitatory NK<sub>1</sub>-positive element and a post-synaptic GABAergic neuron, that, in turn acts at a second synapse upon the SG-responsive cell.

One is also led to ask what is (are) the neuronal domain(s) in which pre-synaptic NK<sub>1</sub> receptors are located. Our results show that they are mainly (but not exclusively) located at cell soma and/or dendrites, since blockade of Na<sup>+</sup>-dependent action potentials with TTX was largely effective in blocking the effects of SM-SP (11 out of 14 neurons). This observation is in agreement with localization studies, which failed to show NK<sub>1</sub> receptor immunoreactivity in axons (McLeod et al., 1998; Naim et al., 1997; Ribeiro-Da-Silva et al., 2001, and this study). However, the increase in the frequency of mIPSCs in a few neurons indicates that pre-synaptic NK<sub>1</sub> receptors could also be expressed in the axonal domain.

Co-expression of NK<sub>1</sub> and GABA has been reported in brain areas other than the spinal cord (Echevarria et al., 1997). However, in rat, most spinal neurons which possessed NK<sub>1</sub> receptor-immunoreactivity were not GABA- or glycine-immunoreactive (Littlewood et al., 1995). Nonetheless,

these authors described a few GABA (and glycine)-immunoreactive neurons in lamina III and, to a lesser extent, lamina IV (see Fig. 4 in Littlewood et al., 1995). NK<sub>1</sub>/GABA-positive neurons of the deep dorsal horn have dorsally oriented dendrites branching in laminae I–II, where they receive an input from tachykinin-releasing primary afferent fibers (PAFs – Brown et al., 1995; Littlewood et al., 1995; Liu et al., 1994; Naim et al., 1997). It is therefore conceivable that mouse laminae III–IV neurons expressing NK<sub>1</sub> receptors at dendrites are capable of dendritic release of GABA (and/or glycine) upon SM-SP challenge. Anatomically, dendritic spines filled with GABA-immunoreactive vesicles have been observed to contact other vesicle-containing dendrites (Carlton and Hayes, 1991; Powell and Todd, 1992). Functionally, dendritic release of GABA has not been described in the spinal cord, but is relevant in other CNS areas such as the olfactory bulb (Halabisky and Strowbridge, 2003; Isaacson and Vitten, 2003). Putative dendro-dendritic synapses between NK<sub>1</sub> expressing GABAergic (and glycinergic) neurons and SM-SP-responsive post-synaptic targets may be an important means to regulate the activity of the SG neurons expressing GABA<sub>A</sub> (and glycine<sub>A</sub>) receptors. In addition, considering that PAFs release a mixture of SP and NKA, and that selectivity of the NK receptor subtypes is not absolute for naive peptides, one cannot exclude that enhancement of inhibitory neurotransmission in SG could also be due in vivo to NKA release (Li and Zhuo, 2001; Trafton et al., 2001). It is also possible that in the intact spinal cord SP activates other NK receptor subtypes besides NK<sub>1</sub>, considering that both NK<sub>2</sub> and NK<sub>3</sub> receptors are expressed in the dorsal horn, albeit at lower levels than NK<sub>1</sub> (Ding et al., 2002; Fleetwoodwalker et al., 1990; Fox et al., 1996; Ribeiro-Da-Silva et al., 2001).

#### 4.4. Functional implications

Inhibitory interneurons in SG regulate the flow of nociceptive information to supraspinal centers. They probably do this by directly targeting neurons of the wide dynamic range and nocispecific projection types, and/or primary afferent terminals. The overall organization involved allows for both pre- and post-synaptic inhibitory modulation (Basbaum, 1999; Millan, 1999). Inhibitory interneurons are themselves targeted by C and Ad fibers (Bernardi et al., 1995; Todd and Spike, 1993; Todd et al., 1994; Yoshimura and Nishi, 1995). This suggests a direct C and Ad counter-regulatory inhibitory feed-back control upon their parallel excitation of neurons of the wide dynamic range and nocispecific projection types. Our results indicate that SP is involved in such a feed-back circuit: thus, pain elicited upon direct activation of projection neurons by C and Ad fibers may be limited by simultaneous recruitment of inhibitory interneurons, at least when the stimulus is of short duration and sub-maximal intensity.

A further element which adds to the complexity of spinal modulation of pain processing by SP is the notion that the peptide can also be released from sources other than PAFs, such as intersegmental neurons and descending fibers (Ribeiro-da-Silva and Hökfelt, 2000). Independently from this, loss of physiological inhibitory tone in the dorsal horn following injuries and/or experimental manipulations has serious effects, altering pain perception and leading to conditions such as hyperalgesia and allodynia (Basbaum, 1999).

#### Acknowledgements

This work was supported by local grants from the University of Torino and Italian MIUR (Fondi Cofin and FIRB 2001). We are greatly indebted to Drs G. Carmignoto for help with patch-clamp experiments, R. Bardoni for useful discussion and suggestions and R.J. Milne for reading the manuscript.

#### References

Afrah AW, Stiller CO, Olgart L, Brodin E, Gustafsson H. Involvement of spinal N-methyl-D-aspartate receptors in capsaicin-induced in vivo release of substance P in the rat dorsal horn. *Neurosci Lett*

2001;316: 83–6.

- Badie-Mahdavi H, Worsley MA, Ackley MA, Asghar AU, Slack JR, King AE. A role for protein kinase intracellular messengers in substance P- and nociceptor afferent-mediated excitation and expression of the transcription factor Fos in rat dorsal horn neurons in vitro. *Eur J Neurosci* 2001;14:426–34.
- Basbaum AI. Spinal mechanisms of acute and persistent pain. *Reg Anesth Pain Med* 1999;24:59–67.
- Beal JA, Bice TN. Neurogenesis of spinothalamic and spinocerebellar tract neurons in the lumbar spinal cord of the rat. *Dev Brain Res* 1994;78: 49–56.
- Beal JA, Russell CT, Knight DS. Morphological and developmental characterization of local-circuit neurons in lamina III of the rat spinal cord. *Neurosci Lett* 1988;86:1–5.
- Bennett GJ, Abdelmoumene M, Hayashi H, Dubner R. Physiology and morphology of substantia gelatinosa neurons intracellularly stained with horseradish peroxidase. *J Comp Neurol* 1980;194:809–27.
- Bernardi PS, Valtschanoff JG, Weinberg RJ, Schmidt HHHW, Rustioni A. Synaptic interactions between primary afferent terminals and GABA and nitric oxide-synthesizing neurons in superficial laminae of the rat spinal cord. *J Neurosci* 1995;15:1363–71.
- Bicknell HR, Beal JA. Axonal and dendritic development of substantia gelatinosa neurons in the lumbosacral spinal cord of the rat. *J Comp Neurol* 1984;226:508–22.
- Bleazard L, Hill RG, Morris R. The correlation between the distribution of the NK1 receptor and the actions of tachykinin agonists in the dorsal horn of the rat indicates that substance P does not have a functional role on substantia gelatinosa (lamina II) neurons. *J Neurosci* 1994;14: 7655–64.
- Brown JL, Liu H, Maggio JE, Vigna SR, Mantyh PW, Basbaum AI. Morphological characterization of substance P receptor-immunoreactive neurons in the rat spinal cord and trigeminal nucleus caudalis. *J Comp Neurol* 1995;356:327–44.
- Budai D, Larson AA. Role of substance P in the modulation of C-fiber-evoked responses of spinal dorsal horn neurons. *Brain Res* 1996;710: 197–203.
- Cajal SRY. In: *Histologie du système nerveux de l'homme et des vertébrés* (French translation by Azouley), vol. 1. Madrid: L.C.S.I.C.; 1952.
- Carlton SM, Hayes ES. GABAergic vesicle-containing dendrites and spines: a critical element in processing sensory input in the monkey dorsal horn. *Neurosci Lett* 1991;121:40–2.
- Cascieri MA, McLeod AM, Underwood D, Shiao L-L, Ber E, Sadowski S, Yu H, Merchant KJ, Swain CJ, Strader CD, Fong TM. Characterization of the interaction of N-acyl-L-tryptophan benzyl ester neurokinin antagonists with the human neurokinin-1 receptor. *J Biol Chem* 1994;269: 6587–91.
- Chizh BA, Cumberbatch MJ, Birch PJ, Headley PM. Endogenous modulation of excitatory amino acid responsiveness by tachykinin NK1 and NK2 receptors in the rat spinal cord. *Br J Pharmacol* 1995;115: 1013–9.
- Coderre TJ, Yashpal K. Intracellular messengers contributing to persistent nociception and hyperalgesia induced by L-glutamate and substance P in the rat formalin pain model. *Eur J Neurosci* 1994;6:1328–34.
- Cumberbatch MJ, Chizh BA, Headley PM. Modulation of excitatory amino acid responses by tachykinins and selective tachykinin receptor agonists in the rat spinal cord. *Br J Pharmacol* 1995;115:1005–12.
- De Felipe C, Herrero JF, O'Brien JA, Palmer JA, Doyle CA, Smith AJ, Laird JMA, Ben-Ari Y, Cervero F, Hunt SP. Altered nociception, analgesia and aggression in mice lacking the receptor for substance P. *Nature* 1998;392:394–7.
- De Koninck Y, Henry JL. Substance P-mediated slow excitatory postsynaptic potential elicited in dorsal horn neurons in vivo by noxious stimulation. *Proc Natl Acad Sci USA* 1991;88:11344–8.
- De Koninck Y, Ribeiro-Da-Silva A, Henry JL, Cuello AC. Spinal neurons exhibiting a specific nociceptive response receive abundant substance P-containing synaptic contacts. *Neurobiology*

- 1992;89:5073–7.
- Ding YQ, Lu CR, Wang H, Su CJ, Chen LW, Zhang YQ, Ju G. Two major distinct subpopulations of neurokinin-3 receptor-expressing neurons in the superficial dorsal horn of the rat spinal cord. *Eur J Neurosci* 2002; 16:551–6.
- Duggan AW, Furnidge LJ. Probing the brain and spinal cord with neuropeptides in pathways related to pain and other functions. *Front Neuroendocrinol* 1994;15:275–300.
- Echevarria D, Matute C, Albus K. Neurons in the rat occipital cortex co-expressing the substance P-receptor and GABA: a comparison between in vivo and organotypic cultures. *Eur J Neurosci* 1997;9:1530–5.
- Falls W, Gobel S. Golgi and EM studies of the formation of dendritic and axonal arbors: the interneurons of the substantia gelatinosa of Rolando in newborn kittens. *J Comp Neurol* 1979;172:511–28.
- Fleetwoodwalker SM, Mitchell R, Hope PJ, Elyassir N, Molony V, Bladon CM. The involvement of neurokinin receptor subtypes in somatosensory processing in the superficial dorsal horn of the cat. *Brain Res* 1990;519:169–82.
- Fox AJ, Naeem S, Patel IA, Walpole C, Urban L. Receptors mediating tachykinin-evoked depolarisations of neurons in the neonatal rat spinal cord. *Acta Biol Hung* 1996;47:129–44.
- Garry MG, Hargreaves KM. Enhanced release of immunoreactive CGRP and substance P from spinal dorsal horn slices occurs during carrageenan inflammation. *Brain Res* 1992;582:139–42.
- Gobel S. Golgi studies of the substantia gelatinosa neurons in the spinal trigeminal nucleus. *J Comp Neurol* 1975;162:397–416.
- Gobel S. Golgi studies of the neurons of layer II of the dorsal horn of the medulla (trigeminal nucleus caudalis). *J Comp Neurol* 1978;180: 375–93.
- Gobel S, Falls WM, Bennett GJ, Abdelmoumene M, Hayashi H, Humprey E. An EM analysis of the synaptic connections of horseradish peroxidase filled stalked cells and islet cells in the substantia gelatinosa of adult cat spinal cord. *J Comp Neurol* 1980;194:781–807.
- Halabisky B, Strowbridge BW.  $\gamma$ -frequency excitatory input to granule cells facilitates dendrodendritic inhibition in the rat olfactory bulb. *J Neurophysiol* 2003;90(2):644–54.
- Honor P, Menning PM, Rogers SD, Nichols ML, Basbaum AI, Besson JM, Mantyh PW. Spinal substance P receptor expression and internalization in acute, short-term, and long-term inflammatory pain states. *J Neurosci* 1999;19:7670–8.
- Hua XY, Saria A, Gamse R, Theodorsson-Norheim E, Brodin E, Lundberg JM. Capsaicin induced release of multiple tachykinins (substance P, neurokinin A and eledoisin-like material) from guinea-pig spinal cord and ureter. *Neuroscience* 1986;19:313–9.
- Ikeda H, Heinke B, Ruscheweyh R, Sandkuhler J. Synaptic plasticity in spinal lamina I projection neurons that mediate hyperalgesia. *Science* 2003;299:1237–40.
- Isaacson JS, Vitten H. GABA(B) receptors inhibit dendrodendritic transmission in the rat olfactory bulb. *J Neurosci* 2003;23:2032–9.
- Jessell TM, Iversen LL. Opiate analgesics inhibit substance P release from rat trigeminal nucleus. *Nature* 1977;268:549–51.
- Jonas P, Bischofberger J, Sandkuhler J. Corelease of two fast neurotransmitters at a central synapse. *Science* 1998;281:419–24.
- Keller AF, Coull JA, Chery N, Poisbeau P, De Koninck Y. Region-specific developmental specialization of GABA–glycine cosynapses in laminae I–II of the rat spinal dorsal horn. *J Neurosci* 2001;21:7871–80.
- King AE, Slack JR, Lopez-Garcia JA, Ackley MA. Tachykinin actions on deep dorsal horn neurons in vitro: an electrophysiological and morphological study in the immature rat. *Eur J Neurosci* 1997;9: 1037–46.
- Kombian SB, Ananthlakshmi KV, Parvathy SS, Matowe WC. Substance P depresses excitatory synaptic

- transmission in the nucleus accumbens through dopaminergic and purinergic mechanisms. *J Neurophysiol* 2003a;89:728–37.
- Kombian SB, Ananthalakshmi KV, Parvathy SS, Matowe WC. Dopamine and adenosine mediate substance P-induced depression of evoked IPSCs in the rat nucleus accumbens in vitro. *Eur J Neurosci* 2003b;18:303–11.
- Kouznetsova M, Nistri A. Modulation by substance P of synaptic transmission in the mouse hippocampal slice. *Eur J Neurosci* 1998;10:3076–84.
- Li HS, Zhao ZQ. Small sensory neurons in the rat dorsal root ganglia express functional NK-1 tachykinin receptor. *Eur J Neurosci* 1998;10:1292–9.
- Li P, Zhuo M. Substance P and neurokinin A mediate sensory synaptic transmission in young rat dorsal horn neurons. *Brain Res Bull* 2001;55:521–31.
- Littlewood NK, Todd AJ, Spike RC, Watt C, Shehab SA. The types of neuron in spinal dorsal horn which possess neurokinin-1 receptors. *Neuroscience* 1995;66:597–608.
- Liu H, Brown JL, Jasmin L, Maggio JE, Vigna SR, Mantyh PW, Basbaum AI. Synaptic relationship between substance P and the substance P receptor: light and electron microscopic characterization of the mismatch between neuropeptides and their receptors. *Proc Natl Acad Sci USA* 1994;91:1009–13.
- Maehara T, Suzuki H, Yoshioka K, Otsuka M. Characteristics of substance P-evoked release of amino acids from neonatal rat spinal cord. *Neuroscience* 1995;68:577–84.
- Maggi CA, Schwartz TW. The dual nature of the tachykinin NK1 receptor. *Trends Pharmacol Sci* 1997;18:351–5.
- Mantyh PW. Neurobiology of substance P and the NK1 receptor. *J Clin Psychiatry* 2002;63:6–10.
- Mantyh PW, Allen CJ, Ghilardi JR, Rogers SD, Mantyh CR, Liu H, Basbaum AI, Vigna SR, Maggio JE. Rapid endocytosis of a G protein-coupled receptor: substance P-evoked internalization of its receptor in the rat striatum in vivo. *Proc Natl Acad Sci USA* 1995;92:2622–6.
- Mantyh PW, Rogers SD, Honoré P, Allen BJ, Ghilardi JR, Li J, Daughters RS, Lappi DL, Wiley RG, Simone DA. Inhibition of hyperalgesia by ablation of lamina I spinal neurons expressing substance P receptor. *Science* 1997;278:275–9.
- Maubach KA, Martin K, Smith DW, Hewson L, Frankshun RA, Harrison T, Seabrook GR. Substance P stimulates inhibitory synaptic transmission in the guinea pig basolateral amygdala in vitro. *Neuropharmacology* 2001;40:806–17.
- Maxwell DJ, Todd AJ, Kerr R. Colocalization of glycine and GABA in synapses on spinomedullary neurons. *Brain Res* 1995;690:127–32.
- McLeod AL, Krause JE, Cuello AC, Ribeiro-Da-Silva A. Preferential synaptic relationships between substance P-immunoreactive boutons and neurokinin 1 receptor sites in the rat spinal cord. *Proc Natl Acad Sci USA* 1998;95:15775–80.
- McLeod AL, Ritchie J, Cuello AC, Julien JP, Ribeiro-Da-Silva A, Henry JL. Transgenic mice over-expressing substance P exhibit allodynia and hyperalgesia which are reversed by substance P and N-methyl-D-aspartate receptor antagonists. *Neuroscience* 1999;89:891–9.
- McMahon SB, Lewin GR, Wall PD. Central hyperexcitability triggered by noxious inputs. *Curr Opin Neurobiol* 1993;3:602–10.
- Meller ST, Gebhart GF. Spinal mediators of hyperalgesia. *Drugs* 1994;47:10–20.
- Millan MJ. The induction of pain: an integrative review. *Prog Neurobiol* 1999;57:1–164.
- Naim M, Spike RC, Watt C, Shehab SA, Todd AJ. Cells in laminae III and IV of the rat spinal cord that possess the neurokinin-1 receptor and have dorsally directed dendrites receive a major synaptic input from tachykinin-containing primary afferents. *J Neurosci* 1997;17:5536–48. Nichols ML, Allen BJ, Rogers SD, Ghilardi JR, Honoré P, Luger NM, Finke MP, Li J, Lappi DA, Simone DA, Mantyh PW. Transmission of chronic nociception by spinal neurons expressing the substance P receptor. *Science* 1999;286:1558–61.



- Nsa Allogho S, Nguyen-Le XK, Gobell F, Pheng LH, Regoli D. Neurokinin receptors (NK<sub>1</sub>, NK<sub>2</sub>) in the mouse: a pharmacological study. *Can J Physiol Pharmacol* 1997;75:552–7.
- Ogier R, Raggenbass M. Action of tachykinins in the rat hippocampus: modulation of inhibitory synaptic transmission. *Eur J Neurosci* 2003; 17:2639–47.
- Oku R, Satoh M, Fuji N, Otaka A, Yajima H, Takagi H. Calcitonin gene-related peptide promotes mechanical nociception by potentiating release of substance P from spinal dorsal horn in rat. *Brain Res* 1987a;403:350–4.
- Oku R, Satoh M, Takagi H. Release of substance P from the spinal dorsal horn is enhanced in polyarthritic rats. *Neurosci Lett* 1987b;74:315–9.
- Otsuka M, Yoshioka K. Neurotransmitter functions of mammalian tachykinins. *Physiol Rev* 1993;73:229–308.
- Powell JJ, Todd AJ. Light and electron microscope study of GABA-immunoreactive neurones in lamina III of rat spinal cord. *J Comp Neurol* 1992;315:125–36.
- Regoli D, Drapeau G, Dion S, D'Orleans-Juste P. Pharmacological receptors for substance P and neurokinins. *Life Sci* 1987;40:109–17.
- Regoli D, Drapeau G, Dion S, Couture R. New selective agonists for neurokinin receptors: pharmacological tools for receptor characterization. *Trends Pharmacol Sci* 1988;9:290–5.
- Regoli D, Boudon A, Fauchère J-L. Receptors and antagonists for substance P and related peptides. *Pharmacol Rev* 1994;46:551–99.
- Ribeiro-Da-Silva A. Substantia gelatinosa of spinal cord. In: Paxinos G, editor. *The rat nervous system*. Sydney: Academic Press; 2003.
- Ribeiro-da-Silva A, Hökfelt T. Neuroanatomical localisation of substance P in the CNS and sensory neurons. *Neuropeptides* 2000;34:256–71.
- Ribeiro-Da-Silva A, McLeod AL, Krause JE. Neurokinin receptors in the CNS. In: Björklund A, Hökfelt T, editors. *Handbook of chemical neuroanatomy*. Amsterdam: Elsevier; 2001. p. 195–240.
- Rusin KI, Bleakman D, Chard PS, Randic M, Miller RJ. Tachykinin potentiate N-methyl-D-aspartate responses in acutely isolated neurons from the dorsal horn. *J Neurochem* 1993;60:952–60.
- Sakuma M, Yoshioka K, Suzuki H, Yanagisawa M, Onishi Y, Kobayashi N, Otsuka M. Substance P-evoked release of GABA from isolated spinal cord of the newborn rat. *Neuroscience* 1991;45:323–30.
- Sakurada T, Yogo H, Tan-No K, Sakurada S, Kisara K. Behavioral activation of neurokinin-1 agonists in relation to enzymatic degradation in the spinal cord. *J Pharm Sci* 1994;83:2–4.
- Saria A. The tachykinin NK<sub>1</sub> receptor in the brain: pharmacology and putative functions. *Eur J Pharmacol* 1999;375:51–60.
- Severini C, Improta G, Falconieri-Erspamer G, Salvadori S, Erspamer V. The tachykinin peptide family. *Pharmacol Rev* 2002;54:285–322.
- Stacey AE, Woodhall GL, Jones RSG. Activation of neurokinin-1 receptors promotes GABA release at synapses in the rat entorhinal cortex. *Neuroscience* 2002;115:575–86.
- Takano M, Takano Y, Yaksh TL. Release of calcitonin gene-related peptide (CGRP), substance P (SP), and vasoactive intestinal polypeptide (VIP) from rat spinal cord: modulation by  $\alpha_2$  agonists. *Peptides* 1993;14: 371–8.
- Teoh H, Malcangio M, Fowler LF, Bowery NG. Evidence for release of glutamic acid, aspartic acid and substance P but not gamma-aminobutyric acid from primary afferent fibres in rat spinal cord. *Eur J Pharmacol* 1996;302:27–36.
- Todd AJ. Anatomy of primary afferents and projection neurones in the rat spinal dorsal horn with particular emphasis on substance P and the neurokinin 1 receptor. *Exp Physiol* 2002;87:245–9.
- Todd AJ, Lewis SG. The morphology of Golgi-stained neurons in lamina II of the rat spinal cord. *J Anat* 1986;149:113–9.
- Todd AJ, Spike RC. The localization of classical transmitters and neuropeptides within neurons in laminae I–III of the mammalian spinal dorsal horn. *Prog Neurobiol* 1993;41:609–38.

- Todd AJ, Spike RC, Brodbelt AR, Price RF, Shehab SAS. Some inhibitory neurons in the spinal cord develop c-fos-immunoreactivity after noxious stimulation. *Neuroscience* 1994;63:805–16.
- Todd AJ, Watt C, Spike RC, Sieghart W. Colocalization of GABA, glycine, and their receptors at synapses in the rat spinal cord. *J Neurosci* 1996; 16:974–82.
- Todd AJ, Puskar Z, Spike RC, Hughes C, Watt C, Forrest L. Projection neurons in lamina I of rat spinal cord with the neurokinin 1 receptor are selectively innervated by substance P-containing afferents and respond to noxious stimulation. *J Neurosci* 2002;22:4103–13.
- Tousignant C, Guillemette G, Drapeau G, Regoli D. Comparison of binding assay and biological activity on a NK-1 system with new selective agonists. *Neuropeptides* 1989;14:275–83.
- Trafton JA, Abbadie C, Basbaum AI. Differential contribution of substance P and neurokinin A to spinal cord neurokinin-1 receptor signaling in the rat. *J Neurosci* 2001;21:3656–64.
- Urban MO, Gebhart GF. Central mechanisms in pain. *Med Clin North Am* 1999;83:585–96.
- Wang DS, Xu TL, Li JS. Modulation of glycine-activated chloride currents by substance P in rat sacral dorsal commissural neurons. *Sheng Li Xue Bao* 1999;51:361–70.
- Wei F, Zhao ZQ. Blockade of capsaicin-induced reduction of GABA- immunoreactivity by spantide in cat spinal superficial dorsal horn. *Neuroscience* 1996;71:277–83.
- Willis Jr WD, Coggeshall RE. In: *Sensory mechanisms of the spinal cord*. New York: Plenum Press; 1991.
- Woolf CJ, Costigan M. Transcriptional and posttranslational plasticity and the generation of inflammatory pain. *Proc Natl Acad Sci USA* 1999;96: 7723–30.
- Woolf CJ, Mannion RJ, Neuman S. Null mutations lacking substance: elucidating pain mechanisms by genetic pharmacology. *Neuron* 1998; 20:1063–6.
- Xu XJ, Dalsgaard CJ, Wiesenfeld-Hallin Z. Spinal substance P and N- methyl-D-aspartate receptors are coactivated in the induction of central sensitization of the nociceptive flexor reflex. *Neuroscience* 1992;51: 641–8.
- Yaksh TL, Jessell TM, Gamse R, Mudge AW, Leeman SE. Intrathecal morphine inhibits substance P release from spinal cord in vivo. *Nature* 1980;286:155–7.
- Yaksh TL, Hua XY, Kalcheva I, Nozaki-Taguchi N, Marsala M. The spinal biology in humans and animals of pain states generated by persistent small afferent input. *Proc Natl Acad Sci USA* 1999;96:7680–6.
- Yang K, Kumamoto E, Furue H, Li YQ, Yoshimura M. Capsaicin induces a slow inward current which is not mediated by substance P in substantia gelatinosa neurons of the rat spinal cord. *Neuropharmacology* 2000;39: 2185–94.
- Yoshimura M. Slow synaptic transmission in the spinal dorsal horn. *Prog Brain Res* 1996;113:443–62.
- Yoshimura M, Jessell TM. Membrane properties of rat substantia gelatinosa neurons in vitro. *J Neurophysiol* 1989;62:109–18.
- Yoshimura M, Jessell TM. Amino acid-mediated EPSPs at primary afferent synapses with substantia gelatinosa neurones in the rat spinal cord. *J Physiol* 1990;430:315–35.
- Yoshimura M, Nishi S. Blind patch-clamp recordings from substantia gelatinosa neurons in adult rat spinal cord slices: pharmacological properties of synaptic currents. *Neuroscience* 1993;53:519–26.
- Yoshimura M, Nishi S. Primary afferent-evoked glycine- and GABA- mediated IPSPs in substantia gelatinosa neurones in the rat spinal cord in vitro. *J Physiol (London)* 1995;482:29–38.
- Yoshimura M, Shimizu T, Yajiri Y, Inokuchi H, Nishi S. Primary afferent- evoked slow EPSPs and responses to substance P of dorsal horn neurons in the adult rat spinal cord slices. *Regul Pept* 1993;46:407–9.