

Grapevine Crown Gall Suppression Using Biological Control and Genetic Engineering: A Review of Recent Research

Attila Filo,¹ Paolo Sabbatini,^{1*} George W. Sundin,² Thomas J. Zabadal,¹
Gene R. Safir,² and Peter S. Cousins³

Abstract: Crown gall is a devastating grapevine disease often encountered in vineyards prone to winter cold injury. *Agrobacterium vitis*, the predominant causal agent of this disease, moves from the roots via xylem sap flow to freeze injury sites where genetic transformations then occur. Crown gall disrupts the grapevine trunk vascular system, which prevents nutrient flow and leads to plant decline and death. Viticultural practices designed to fight this disease are only partially effective, thereby requiring alternatives. Genetic engineering and biological control could be more desirable approaches for disease prevention. Biological control typically involves antagonistic organisms, which are applied to grapevine roots to reduce the concentration of pathogenic *Agrobacterium* strains. Genetic engineering may prevent infection and tumor formation by modifying grapevines and antagonistic organisms. In the grapevine, this may be achieved by enhancing molecular mechanisms for producing bacterium-specific antimicrobial peptides or preventing transferred deoxyribonucleic acid export, integration, and oncogene expression. Alteration of antagonistic organisms enhances the production of bacteriocins effective against agrobacteria. This article reviews the potential use of biological control options and genetic engineering tools for grapevine crown gall suppression and makes recommendations for further use and research.

Key words: *Agrobacterium vitis*, crown gall, control, biological technology, molecular biology

Crown Gall on Grape and *Agrobacterium* Nomenclature

The unchecked growth of tumors on grape, commonly known as crown gall (CG), is one of the most economically destructive diseases in temperate zone viticulture. The chief disease agent is the bacterium *Agrobacterium vitis*, although other *Agrobacterium* species may also cause CG on grapevine (Szegedi et al. 2005, Kawaguchi and Inoue 2009). Crown gall results in yield reductions, grapevine decline, and vine death (Süle and Burr 1998). It occurs worldwide and varies by viticultural regions, with the most severe impact in temperate climates. The world map of the Köppen-Geiger on climate classification (Kottek et al. 2006) indicates that grapevine CG disease occurs in climate regions coded with Dfb (Tmin \leq -3°C). Severe CG incidences are common in the Great Lakes Region, the Canadian province of Ontario, and

Germany, Hungary, Slovakia, and the Czech Republic. The economic impact of CG varies significantly depending upon region. For example, in the state of Pennsylvania, losses were estimated over a six-year period at US\$46,500 per 0.4 hectare vineyard (Steward and Wenner 2004).

Numerous taxonomic schemes have been proposed for the nomenclature of the *Agrobacterium* species (Sawada et al. 1993, Willems and Collins 1993, de Lajudie et al. 1998). Nomenclature in this review refers to the system proposed by Kerr and Panagopoulos (1977) and Ophel and Kerr (1990). Kerr and Panagopoulos (1977) demonstrated that grapevine *Agrobacterium* isolates form a distinct group and differ from strains isolated from other plants. They classified *Agrobacterium* spp. into three groups: biotype-1, biotype-2, and biotype-3. Ophel and Kerr (1990) demonstrated, through biochemical tests and differential levels of deoxyribonucleic acid (DNA) binding, phenotypic differences between the grapevine *Agrobacterium* isolates and the previously described *A. tumefaciens*, *A. radiobacter*, *A. rhizogenes*, and *A. rubi* spp. They proposed that grapevine biotype-3 isolates form a new separate *Agrobacterium* sp. named *A. vitis*. Based on comparative 16S rDNA sequencing, renaming of agrobacteria to *Rhizobium radiobacter*, *R. rhizogenes*, *R. rubi*, *R. undicola*, and *R. vitis* was proposed (Young et al. 2001, Willems 2006). Currently, the most commonly used nomenclature refers to biotype-1, biotype-2, and biotype-3 strains of *Agrobacterium* as *A. tumefaciens*, *A. rhizogenes*, and *A. vitis*, respectively. Tumorigenic *Agrobacterium* spp. are also classified according to the type of their CG tumor-inducing or root-inducing (Ti/Ri) plasmid, which determines the type of opine(s) synthesized and metabolized by each strain (Paulus et al. 1989, Otten et al. 1996). Classification of *Agrobacterium* strains by opine production is problematic because of unknown or nonexistent opine types

¹Department of Horticulture, Michigan State University, East Lansing, MI 48824; ²Department of Plant Pathology, Michigan State University, East Lansing, MI 48824; and ³USDA-ARS, Grape Genetics Research Unit, 630 W. North Street, Geneva, NY 14456.

*Corresponding author (email: sabbatin@msu.edu; fax: 517 353-0890)

Acknowledgments: This work was partially supported by the O'Keefe Research Fund in the Department of Horticulture, Michigan State University.

The authors thank T.J. Burr and J.E. Creasap (Cornell University) for valuable information on viticulture research on crown gall and L.W. Stobbs (Agriculture and Agri-Food Canada) for the photographs of grapevine crown gall. This manuscript was in partial fulfillment of requirements for the master of science degree for A. Filo in the Department of Horticulture, Michigan State University. Manuscript submitted Feb 2012, revised Jul 2012, accepted Sept 2012. Publication costs of this article defrayed in part by page fees.

Copyright © 2013 by the American Society for Enology and Viticulture. All rights reserved.

doi: 10.5344/ajev.2012.12038

of several Ti/Ri plasmid-induced tumors. Moreover, a single plasmid does not always induce only one specific opine type. In fact, inductions of several opine types in various combinations are most often associated with a single Ti/Ri plasmid (Paulus et al. 1989, Otten et al. 1996). *Agrobacterium* grapevine isolates predominantly metabolize nopaline, vitopine, or octopine/cucumopine and rarely other opine types (Otten et al. 1996). Eleven opine types (octopine, nopaline, succinamopine, agropine, agropine/mannopine, mannopine, chrysopine/succinamopine, chrysopine/nopaline, cucumopine/mikimopine, octopine/cucumopine, and vitopine) are currently being classified (Petit et al. 1983, Szegedi et al. 1988, Dessaux et al. 1998, Pionnat et al. 1999, Moriguchi et al. 2001).

Agrobacterium Ecology and Etiology

Etiology. Agrobacteria commonly inhabit soils and roots of susceptible host plants (Figure 1). *Agrobacterium vitis* survives in grapevine plant material, including roots, trunks, cordons, canes, cuttings, and debris, and in vineyard soil. Agrobacteria are generally harmless to plants. However, they can cause a disease if they possess a large (~200 to 800 kilobase pairs in size; Gelvin 2003) Ti or Ri plasmid. Crown gall is triggered by the integration of transferred-DNA (T-DNA) into the plant nuclear genome and the expression T-DNA genes encoding enzymes of plant hormone biosynthesis. Only the T-DNA, that is only a segment of the Ti plasmid, is transferred into the plant nuclear genome. It is this Ti plasmid that contains the T-DNA and the genes required for its export, integration, and oncogene expression (van Larebeke et al. 1974, De Cleene and De Ley 1976, Portier et al. 2006).

The infection process. Crown gall infection is a three-step process. In step one the pathogen enters the apoplastic

space of the plant. *Agrobacterium vitis* is particularly concentrated in the rhizosphere of grapevines and most commonly infects through roots and underground wound sites (Süle et al. 1995, Burr et al. 1998). Step two centers on the colonization of the xylem by the bacteria. *Agrobacterium vitis* systemically colonize grapevines and disseminate to shoots via xylem sap flow (Szegedi and Bottka 2002, Szegedi and Dula 2006). The pathogen survives and persists in apparently healthy cuttings for extended periods (Lehoczky 1968, 1971, 1978, 1989, Burr and Katz 1984, Burr et al. 1987, 1988). Step three involves the evasion of plant reactions and suppression of plant defense mechanisms. A well-known example of such mechanism of evasion is the degradation of hydrogen peroxide by the bacterium's catalase enzyme encoded by the *kata* gene (Xu and Pan 2000).

T-DNA export to plant cell. Wound healing processes, such as callus formation, development of graft union vascular connection, and tissue repair and remodeling in response to grafting and freeze injury, as well as physical damages caused by vineyard equipment, involve cell division and render grapevine cells competent to *Agrobacterium* transformation. Consequently, these are the primary sites for CG development (Creasap et al. 2005, Otten et al. 2008). The tumorous growth typically occurs around the graft union (Figure 2), on the lower trunk (Figure 3), and at the base and disbudded points of cuttings, as those are prevalent sites of wound healing and highly susceptible to T-DNA export. Many proteins encoded by virulence (*vir*) genes play essential roles in the T-DNA transfer process. Nearly all proteins encoded by the *vir* region of the Ti plasmid are required for T-DNA transfer to occur (Burr and Otten 1999, Gelvin 2003, Matthysse 2006, Otten et al. 2008). VirA and VirG form a




Figure 1 Disease cycle of grapevine crown gall (from Burr et al. 1998; reprinted by permission; © 1998 American Phytopathological Society).

two-component regulatory system. VirA works as sensor, while VirG works as transcriptional regulator. VirA senses the presence of specific plant phenolic compounds induced at wound healing, while VirG increases the level of *vir* gene transcription. VirD4 and VirB proteins are necessary for the transfer of the T-DNA and other Vir proteins. VirD2 serves as a pilot protein that guides the T-DNA through the transfer process, while VirE2 is required for efficient tumor formation. A detailed account of the T-DNA transfer process and the role of individual Vir proteins are provided in a recent review (Lacroix et al. 2008).




Figure 2 Graft unions of Zinfandel grapevines exhibited crown gall shortly after vineyard establishment in the Canadian province of Ontario. Both vines will inevitably die. Crown gall developed at (A) the upper trunk, including the entire scion, and (B) wound sites around the graft union and the base of the shoots (red arrows).



Figure 3 Own-rooted Cayuga White grapevine exhibits crown gall (red arrows). Photo taken at the Horticulture Teaching and Research Center of Michigan State University (East Lansing). The use of multiple trunks helps to reduce the impact of the disease on the entire vine.

T-DNA integration into the plant nuclear genome. Induction of CG formation is unique because it involves the interkingdom transfer of T-DNA from the bacterial pathogen into the plant nuclear genome (Zupan and Zambryski 1995). The bacterial T-DNA is integrated into the plant chromosome where the genetic transformation of the plant cell then begins. Plant cell transformation leads to elevated levels of indole-3-acetic acid (auxin or IAA) and cytokinin production (Morris 1986). Expression of two oncogenes (*iaaM* and *iaaH*) was required for auxin synthesis and tumor induction in grapevines (Thomashow et al. 1986, Huss et al. 1990). Auxin is synthesized in a two-step biochemical pathway. Step one includes the expression of the *iaaM* gene, which induces tryptophan mono-oxygenase production. Tryptophan mono-oxygenase then converts tryptophan to indoleacetamide. Step two involves indoleacetamide hydrolase production by the *iaaH* gene. Indoleacetamide hydrolase then converts indoleacetamide to auxin (Thomashow et al. 1986, Huss et al. 1990). Interestingly, most *A. vitis* strains only encode auxin synthesis in their T-DNA (Fournier et al. 1994). The expression of T-DNA genes within the plant cell and the elevated level of hormone production break the regulation of the cell cycle because the plant cell cannot regulate the expression of the T-DNA genes. These trigger the transformation and abnormal proliferation of plant cells resulting in CG tumor or hairy root formation (Petersen et al. 1989, Gaudin et al. 1994, Costacurta and Vanderleyden 1995).

Management of Grapevine Crown Gall

Traditional viticultural practices. There are no current completely effective methods of controlling CG disease in nurseries and vineyards. The stage is often set for CG during in-nursery callusing and rooting or after the first winter in the vineyard, generating problems during vineyard establishment (Figure 4). Prevention of freeze injury (selecting



Figure 4 Severe crown gall (CG) incidence and consequently grapevine mortality in a 3-year-old vineyard planted with Riesling (clone 239) vines grafted to Couderc 3309 or Millardet et de Grasset 101-14 rootstocks. Photo taken at Chateau Grand Traverse vineyard (Old Mission Peninsula, Michigan) on 11 July 2010. In April 2010, 1776 of 3077 vines (57.7%) exhibited CG and 525 of those 1776 (29.6%) did not exhibit any growth during the summer.

vineyard sites that avoid low temperatures), graft union protection (through soil burial/mulching techniques), frequent trunk renewal (Figure 3), and removal of infected vine parts are major viticulture strategies to suppress CG; however, they are only partially effective (Zabada et al. 2007). Prevention of infection is a critical disease management strategy. This involves producing pathogen-free planting material by *in vitro* apical meristem or shoot-tip tissue culture and inhibiting bacterial infection (soil fumigation and soil solarization) (Burr et al. 1988, Pu and Goodman 1993a, Burr and Otten 1999, Szegedi and Süle 2005, Otten et al. 2008). Other strategies to reduce CG incidence in both vineyard and nursery are heat and chemical treatments (such as hot water submersion and oxyquinoline sulfate treatments) of dormant cuttings prior to grafting (Szegedi 1995, Burr et al. 1996, 1998, Burr and Otten 1999, Otten et al. 2008). Breeding for grapevine CG resistance represents another approach. However, resistant varieties that produce high-quality fruit have not yet been developed (Burr et al. 2003). Cultural practices to reduce CG incidence also include the use of own-rooted vines, cold-hardy cultivars (such as Frontenac and La Crescent), multiple-trunk training systems (Figure 3), irrigation management to avoid freeze injury (Matthysse 2006), and fumigation and solarization to reduce *Agrobacterium* and nematode population density in soil (Süle et al. 1995, Pinkerton et al. 2000). Nematode (*Meloidogyne hapla*) feeding sites on grapevine roots may serve as an entry point for *A. vitis* (Süle et al. 1995). Despite the multiple strategies, none of the currently available cultural practices provide acceptable levels of CG control in temperate viticultural regions.

Novel approaches. The increasing global demand for disease-resistant and stress-tolerant grapevines has prompted significant interest in research into genetic engineering-assisted grapevine breeding. Genetic engineering, also called genetic modification, uses recombinant nucleic acid techniques involving the formation of new combinations of genetic material. This material is produced by inserting externally produced nucleic acid molecules into a virus, bacterial plasmid, or other vector system and then incorporating that vector into a host organism in which they do not naturally occur but in which they are capable of continued propagation (Sands and Galizzi 2006). Organisms derived through genetic engineering are referred to as genetically engineered (GE), genetically modified (GM), and/or transgenic (TG) organisms. For sake of clarity, this review adheres to the most commonly used acronym and refers to such organisms as GM. This approach to genetic engineering-assisted grapevine breeding uses the natural ability of numerous *A. tumefaciens* and *A. rhizogenes* spp. to transfer DNA interkingdom. The refinement of this technique has impacted plant science, led to rapid progress in plant molecular biology, and has now become the essential embodiment of today's crop biotechnology.

This same characteristic of *Agrobacterium* is used in efforts to suppress CG disease in viticulture. Research that addresses the development of CG-resistant grapevines through genetic engineering, predominantly for less cold-hardy varieties (*Vitis vinifera* L.), emerged in the 21st century. Grapevine

CG resistance has been hypothesized to be related to a single gene (Szegedi et al. 1984, Szegedi and Kozma 1984). Research in this era targeted the introduction and expression of foreign genes, the efficacious regeneration of transformed grapevines, and the efficiency of the transformation process (Nakano et al. 1994, Scorza et al. 1996, Mozsár et al. 1998, Torregrosa et al. 2002, Oláh et al. 2003). Deoxyribonucleic acid fingerprint comparisons conducted on ancestral forms of *A. vitis* aimed to provide information on the evolution of the pathogen (Argun et al. 2002). After overcoming the technological hurdles of grapevine transformation, more recent studies addressed specific problems, such as stress tolerance and disease resistance in grapevines (Burr and Otten 1999, Otten et al. 2008).

Efforts to engineer CG resistance into grapevines have focused on three main approaches: blocking infection by expressing antimicrobial peptides in GM plants inhibitory to *A. vitis*; blocking T-DNA export and/or integration; and blocking T-DNA oncogene expression following its export and integration. Genetic engineering may create GM grapevines with resistance to CG and GM biological control agents (BCAs) with higher CG control efficacy.

A biological control approach for CG disease caused by *A. tumefaciens* depends on the use of a nonpathogenic biological control strain *A. radiobacter* K84. This strain produces the bacteriocin agrocin 84 which is toxic to certain strains of *A. tumefaciens* and *A. rhizogenes* (Kerr and Htay 1974, Kim et al. 2006). The mode of action of agrocin 84 is based on the agrocinopine (an analog of agrocin 84) biosynthesis by the plant and its catabolism by pathogenic *A. vitis* strains (Kim et al. 2006). Application of strain K84 results in agrocin 84 production by the K84 strain and agrocinopine biosynthesis by the plant. The Ti plasmid genes of agrocin 84-sensitive tumorigenic strains encode for agrocinopine uptake and catabolism. This agrocinopine-utilization system enables those strains to take up agrocin 84. Once taken up, agrocin 84 is toxic to the bacterial cell (Kim et al. 2006). Unfortunately, strain K84 will not control grapevine CG caused by *A. vitis* (Staphorst et al. 1985, Chen and Xiang 1986, Webster et al. 1986). However, the effectiveness of this BCA in controlling *A. tumefaciens*-mediated CG suggests that biological control is a viable tool if an effective strain is found.

Biological control strategies to prevent infection. One of the most promising grape CG control technologies is biological control, and it has been defined in many ways. This review adheres to the definition of plant disease biological control as the control of a plant disease with a natural biological process or the product of a natural biological process, including biological chemicals extracted from and delivered by living organisms and constitutive and elicited host resistances (Wilson 1997). This definition allows clear dissociation from other means of plant disease control, such as physical, cultural, synthetic chemical, or genetic engineering-assisted control. Research efforts to discover or develop living organisms, or products created by those organisms, for grapevine CG suppression were initiated in 1972 (Kerr 1972).

The remarkable success of *A. radiobacter* strain K84 represents one of the most exhaustively researched and

commercially widely used antagonistic microorganisms (Cooksey and Moore 1982, du Plessis et al. 1985, Farrand et al. 1985, Donner et al. 1993, Vicedo et al. 1993, Peñalver et al. 2000). Effective CG control by the *A. radiobacter* strain was demonstrated on several plants, including *Prunus*, *Rubus*, *Malus*, *Salix*, *Libocedrus*, *Chrysanthemum*, *Crataegus*, *Carya*, *Rosa*, *Pyrus*, and *Humulus* spp. (Moore and Warren 1979). Unfortunately, strain K84 failed to control grapevine CG caused by *A. vitis* (Staphorst et al. 1985, Chen and Xiang 1986, Webster et al. 1986). Consequently, many alternative bacterial strains have now been studied for their ability to inhibit grapevine CG infection and formation with mixed results. We have included a summary of the most important strains (Table 1) and their relevant characteristics (Table 2).

***A. radiobacter* strains MI15 and HLB-2.** One of the BCAs proved to be effective against *A. vitis* strains was isolated from grapevines (Xuemei et al. 1993). The agrocin-producing *A. radiobacter* strain MI15 (isolated from CG on Muscat Hamburg) inhibited tumor formation caused by *A. vitis* strains in vitro and on inoculated grapevine shoots. Colonization of wound sites on grapevine stems by *A. radiobacter* strain MI15 was also reported (Xuemei et al. 1993). In another experiment, the *A. radiobacter* strain HLB-2 inhibited CG formation on inoculated young grapevine shoots (Chen and Xiang 1986). In vitro growth of *A. vitis* strains harboring the nopaline or octopine plasmid was inhibited by co-inoculations (Chen and Xiang 1986).

Further research on *A. radiobacter* strain HLB-2 demonstrated suppression of grapevine CG caused by *A. vitis* strains Ag57, Ag63R, and G-1 (Pu and Goodman 1993b). Seven additional strains of *Agrobacterium* spp. also showed sensitivity to *A. radiobacter* strain HLB-2. Greatest suppression was achieved when *A. radiobacter* strain HLB-2 was applied at 10,000 times higher concentration than the tumorigenic *A. vitis* strain Ag63R. In this case, 87% of inoculated grapevine stems did not exhibit tumor symptoms. When the same *Agrobacterium* strains were co-inoculated with equal or lower concentrations of *A. radiobacter* strain HLB-2, 35 and 0% of co-inoculated grapevine stems remained symptomless, respectively (Pu and Goodman 1993b).

Dipping of grapevine roots with *A. radiobacter* strain HLB-2 just prior to planting into *Agrobacterium*-infested soil prevented CG formation (Pu and Goodman 1993a). Thirty-four months after planting, 100% of the Catawba, Seyval blanc, and Vidal blanc and 78% of the Chancellor root-treated grapevines had no CG symptoms. Twenty percent of the untreated Catawba and Vidal blanc and 100% of the untreated Chancellor had CGs (Pu and Goodman 1993a).

Strain HLB-2 reduced internal populations of *A. vitis*, thus promoting grapevine vitality and increasing graft take (Bazzi et al. 1999). Internal populations of *A. vitis* strain CG49 were reduced from $\sim 1.0 \times 10^9$ to 7.35×10^6 colony-forming units per mL (cfu/mL) following HLB-2 treatment. Less than half the HLB-2 inoculated grapevines were dead or unmarketable

Table 1 List of selected bacterial spp. strains examined for biological control against grapevine crown gall and their relevant characteristics.

Bacterial spp.	Strain	Relevant characteristics	References
<i>Agrobacterium radiobacter</i>	HLB-2	Inhibits CG formation on inoculated grapevine shoots. Root treatments prevent CG formation.	Pu and Goodman 1993a, 1993b, Bazzi et al. 1999
<i>A. radiobacter</i>	K84	Reported ineffective against <i>A. vitis</i> strains. One contradictory report indicated inhibition of a few <i>A. vitis</i> strains in vitro.	Kerr and Htay 1974, Moore and Warren 1979, Donner et al. 1993, Vicedo et al. 1993
<i>A. radiobacter</i>	M115	Inhibits in vitro growth and tumor formation caused by <i>Agrobacterium</i> biotype-3 strains. Effective colonizer of wound sites on grape stems.	Xuemei et al. 1993
<i>A. tumefaciens</i>	J73	Agrocin-producing strain with broad spectrum activity against <i>A. vitis</i> and <i>A. tumefaciens</i> .	Thomson 1986, Webster et al. 1986
<i>A. vitis</i>	E26	In vitro and in vivo activity against <i>A. vitis</i> strains.	Liang et al. 2001, Li et al. 2005, Wang et al. 2008, Wei et al. 2009
<i>A. vitis</i>	F2	Nontumorigenic on grapevines. Effective against grape CG caused by agrocin and non-agrocin producing <i>A. vitis</i> .	Staphorst et al. 1985
<i>A. vitis</i>	F2/5	Nontumorigenic on grapevines. Biological control is specific to grapevine CG. Inhibits growth of all <i>A. vitis</i> strains tested. Causes necrosis of treated grapevine tissues.	Burr and Reid 1994, Burr et al. 1997, 1999, Bazzi et al. 1999, Szegedi et al. 1999, Zheng et al. 2003, Creasap et al. 2005, Hao et al. 2005
<i>A. vitis</i>	VAR03-1	Highly effective inhibitor of <i>A. vitis</i> pathogenic strains in vitro and in vivo. Root colonizer.	Kawaguchi et al. 2005, 2007, 2008
<i>Pseudomonas aureofaciens</i>	B-4117	Inhibits CG development if grape cuttings are submerged in bacterial solution.	Khmel et al. 1998
<i>P. fluorescens</i>	CR330D	Reduces CG incidence and inhibits tumor formation.	Khmel et al. 1998
<i>P. fluorescens</i>	1100-6	Reduces CG incidence and internal populations of <i>A. vitis</i> .	Eastwell et al. 2006
<i>P. corrugata</i>	JC583	Varying levels of tumor formation inhibition in inoculated grapevine.	Bell et al. 1995
<i>Rahnella aquatilis</i>	HX2	Inhibits CG formation. Effective against a number of <i>A. vitis</i> strains. Produces an antimicrobial compound.	Bell et al. 1995, Chen et al. 2007, 2009

Table 2 In vitro growth, crown gall (CG) formation inhibition, colonization, persistence, and other relevant characteristics of selected bacterial spp. strains effective in biological control against grapevine crown gall.

Bacterial spp. strains	In vitro growth and CG formation inhibition	Colonization, persistence, and other characteristics
<i>A. radiobacter</i> HLB-2	CG did not develop on root-treated vines. 75% of root-treated Chancellor did not exhibit CG (Pu and Goodman 1993a). 87% of treated vines did not exhibit CG (Pu and Goodman 1993b).	Reduced internal <i>A. vitis</i> concentration from $\sim 10^9$ to 7.35×10^5 cfu/mL (Bazzi et al. 1999).
<i>A. tumefaciens</i> J73	Inhibited in vitro growth of 9 of 16 <i>Agrobacterium</i> biotype-1, 11 of 14 biotype-2, and 22 of 27 <i>A. vitis</i> strains (Webster et al. 1986). Inhibited in vitro growth of all nopaline, octopine, and agropine type Ti plasmid strains (Webster and Thomson 1988).	Development of Ti plasmid transfer deficient strain J73 accomplished (Webster and Thomson 1988).
<i>A. vitis</i> E26	Inhibited in vitro growth of 20 biotype-3 and one biotype-1 <i>Agrobacterium</i> strains. Inhibited in vivo CG formation of 18 octopine, one nopaline, and one arginine plasmid harboring <i>Agrobacterium</i> biotype-3 strains (Liang et al. 1990a, 1990b). Inhibited in vivo CG formation of all <i>A. vitis</i> strains (Liang et al. 2001). Inhibited in vitro growth of 11 of 12 <i>A. vitis</i> strains (Wang et al. 2003).	Isolated from wound sites 1 month postinoculation (Liang et al. 1990a). $\sim 10^4$ cfu per g of dry soil and fresh roots measured at 5 months postplanting. Blocked attachment of tumorigenic <i>A. vitis</i> strain K308 to grape stem and seedling root cells (Li et al. 2005). The <i>avsl</i> _{E26} mutant strain was as effective in controlling grape CG as the wild-type E26 (Wang et al. 2008).
<i>A. vitis</i> F2	Inhibited in vitro growth of 17 of 25 non-agrocin producing biotype-3 strains (Staphorst et al. 1985). Inhibited in vivo CG formation of 12 <i>A. vitis</i> strains (Burr et al. 1999).	
<i>A. vitis</i> F2/5	Inhibited in vitro growth of 21 of 25 <i>A. vitis</i> strains. Two of 19 <i>A. tumefaciens</i> strains showed sensitivity to F2/5. Reduced <i>A. vitis</i> and <i>A. tumefaciens</i> tumor sizes. Number of sites at which CG developed was not reduced only when <i>A. vitis</i> strain CG78 was used in inoculations (Burr and Reid 1994).	Caused necrosis of treated grape tissues, which inhibited callus and graft union formation and reduced graft take (Bazzi et al. 1999, Creasap et al. 2005). Biological control by F2/5 not associated with agrocin production and competition for attachment sites or with its tartrate or octopine utilization plasmids. Inhibition or prevention of T-DNA transfer is likely the main factor of its CG control mechanism (Burr et al. 1997, Szegedi et al. 1999). Necrosis induction may be regulated by <i>aviR</i> gene (Zheng et al. 2003). Regulatory role determination of <i>clp</i> genes in strain F2/5 in progress (Burr, personal communication, 2010).
<i>A. vitis</i> CG1077 F2/5	CG1077 inhibited pathogenic <i>A. vitis</i> more than strain F2/5 (Burr et al. 1997). Maximum reduction in internal <i>A. vitis</i> concentration, from $\sim 10^9$ to 1.24×10^6 cfu/mL by CG1077 (Bazzi et al. 1999).	
<i>A. vitis</i> F2/5 CG1076 CG1077 CG1078 CG1079	Reduced the percentage of sites at which CG developed by <i>A. vitis</i> strain K306 or CG49 up to 92 and 100%. Reduced gall areas ranged from 3 to 21 mm ² . Average tumor areas and sites at which galls developed for inoculations only with strain K306 or CG49 were 140 and 40 mm ² and 95 and 80%, respectively (Burr et al. 1997).	
<i>A. vitis</i> F2/5 (pT2TFXK)	Exhibited increased efficacy against all tested <i>A. vitis</i> strains compared to wild-type F2/5 (Herlache and Triplett 2002).	Produces trifolitoxin peptide antibiotic (Herlache and Triplett 2002).
<i>A. vitis</i> VAR03-1	Inhibited in vitro growth of all tumorigenic <i>A. vitis</i> strains (Kawaguchi et al. 2005). Reduced the number of sites at which CG developed up to 94.3% (Kawaguchi et al. 2005, 2007). Inhibited in vitro growth of all tested pathogenic <i>Agrobacterium</i> strains. Inhibition zones ranged up to 24.5 mm. Root treatment reduced CG incidence up to 26.7% in vivo and 46.7% in the greenhouse (Kawaguchi et al. 2008).	Eighteen months after root submersion treatment, detectable VAR03-1 ranged from $\sim 10^{6.10}$ to $10^{6.28}$ cfu/g fresh roots (Kawaguchi et al. 2008).
<i>P. aureofaciens</i> B-4117 and <i>P. fluorescens</i> CR330D	In vitro growth inhibition zones ranged up to 14 and 20 mm for <i>A. vitis</i> strain Sz1 and Tm4. Reduced in vivo CG incidence up to 5-fold. Reduced in vivo CG incidence up to 3.5-fold 2 months postplanting submerged and sprayed vines in the vineyard (Khmel et al. 1998).	120 days postinoculation, population density declined from $\sim 10^9$ to 10^4 cfu/g dry root. The $\sim 10^4$ cfu/g dry root population density remained constant for up to 1 year. 24 days postinoculation, the mean colonization level for sterile and nonsterile soil by the antagonists was $\sim 10^4$ cfu/g soil (Khmel et al. 1998).
<i>R. aquatilis</i> HX2	In vitro inhibition zones of <i>A. vitis</i> strains ranged up to 42.7 mm. Eight months after root submersion treatment, in vivo CG incidences were reduced with 62.5, 63.8, and 61.3% (Chen et al. 2007). Prevented CG formation by <i>A. vitis</i> strain K308 for 40 days (Chen et al. 2009).	

compared to the untreated and controls (Bazzi et al. 1999). Although strain HLB-2 reduced internal *A. vitis*, the magnitude of the reduction cannot ensure prevention of CG incidence in the vineyard.

Results of these experiments (Chen and Xiang 1986, Pu and Goodman 1993a, 1993b, Bazzi et al. 1999) cannot be compared because different treatment methods were used, such as grapevine root versus stem, cutting, or shoot inoculations. Each treatment mode aims to assess the efficacy of select BCAs against pathogenic *Agrobacterium* strains. However, deployment of BCAs can be accomplished through different plant tissue systems and by various methods, suggesting that efficacy (including colonization and translocation) of BCAs may vary significantly by plant tissue systems and methods for deployment.

***A. tumefaciens* strain J73.** One of the first demonstrations of the biological control potential of grapevine CG, using *Agrobacterium* strains other than *A. radiobacter* and *A. vitis*, involved *A. tumefaciens* strain J73 (Thomson 1986, Webster et al. 1986). This strain exhibited a broad spectrum of activity against *A. tumefaciens* and *A. vitis* strains *in vitro*. J73 was classified as an agrocin-producing biotype-2 strain with a nopaline-type Ti plasmid. A previous experiment (Staphorst et al. 1985) verified sensitivity of four biotype-3 strains to *A. radiobacter* strain K84. However, *A. radiobacter* strain K84 was effective only against strains harboring nopaline-type Ti plasmids. J73 inhibited all nopaline-, octopine-, and agropine-type Ti plasmid strains (Thomson 1986, Webster et al. 1986).

Strain J73 was pathogenic on a number of plants other than grapevines, which restricted its *in vivo* use as a BCA for grapevine CG. To prevent potential CG elicitation, it had to be cured of its nopaline-type Ti plasmid to disable the interkingdom transfer of strain J73 T-DNA. This was accomplished by the introduction of selectable plasmids carrying the origins of replication of either the nopaline Ti plasmid, pTiC58, or the octopine Ti plasmid, pTi15955 (Webster and Thomson 1988).

***A. vitis* strain E26.** A nonpathogenic, agrocin-producing, *Agrobacterium* biotype-3 strain was isolated in 1989 (Liang et al. 1990a). It was originally designated as *A. radiobacter* strain E26 and is now identified as *A. vitis* strain E26. It was effective *in vitro* and *in vivo* against 20 biotype-3 and one biotype-1 *Agrobacterium* isolates (Liang et al. 1990a, 1990b). Although inhibition levels were not specified, strain E26 exhibited both *in vitro* and *in vivo* inhibition of C58 (biotype-1), K27 (biotype-2), pt12 (biotype-1), A6 (biotype-1), and Bo542 (biotype-2) strains. *In vivo* inhibition was obtained on 18 octopine, 1 nopaline, and 1 arginin plasmid harboring *Agrobacterium* biotype-3 strains when grapevine stems were co-inoculated with equal concentrations of strain E26. Isolation of strain E26 from wounded sites one month after inoculation verified its ability to colonize grapevine tissues. It was also effective when tested against the Ti-plasmid cured avirulent *A. tumefaciens* strain C58. Liang et al. (1990a, 1990b) concluded that resistance to strain E26 is chromosomally encoded.

When grapevines were inoculated with tumorigenic *A. vitis* strains, CG formation was inhibited in all 12 cases by

strain E26 (Liang et al. 2001). Eleven of 12 *A. vitis* strains showed *in vitro* sensitivity to strain E26. An agrocin compound designated as Ar26 was isolated and purified from E26. It was effective against *A. vitis* strain MI3-2 (Liang et al. 2001, Wang et al. 2003).

E26 was able to achieve high-efficacy rhizosphere and rhizoplane colonization (Li et al. 2005). It was also able to block attachment of the tumorigenic *A. vitis* K308 to grape stem and seedling root cells. The average 10^4 cfu E26 population per g of dry soil and fresh roots at 5 months postplanting (Li et al. 2005) also suggests that E26 is suitable for grapevine CG suppression.

Numerous molecular biology studies were done on E26. The role of a LuxR-LuxI type quorum-sensing system involved in the induction of the hypersensitive response (HR) in tobacco by strain E26 was shown (Wang et al. 2008). AvsI_{E26} and AvsR_{E26} components of the wild-type strain E26 are believed to be responsible for long- and short-chain acyl-homoserine lactones signal induction required for HR in tobacco. The avsI_{E26} mutant was unable to elicit the HR. This proved that the AvsI quorum-sensing system was necessary for induction of the HR. Biological control efficacy of the wild-type E26 and the avsI_{E26} mutant strains was not significantly different, as demonstrated in pot trials. Developed galls were very small and their weights, 40 days following inoculation, for grapevines co-inoculated with the pathogenic *A. vitis* strain K308 and strain E26 or avsI_{E26} mutant strain, were greatly reduced as compared to when only K308 was used as inoculum-negative control (Wang et al. 2008).

A genetic study was undertaken on the *vir* regions of strain E26 for risk determination of its commercial use as a BCA in field applications (Wei et al. 2009). None of the five (*virA*, *virG*, *iaaH*, *iaaM*, and *ipt*) genes required for tumorigenicity were detected by PCR and Southern blot analyses. Lack of *virA* and *virG* pathogenic determinants in strain E26 suggests its inability for CG symptom elicitation in both host and non-host plants (Wei et al. 2009).

***A. vitis* strain F2.** Another *A. vitis* strain, first isolated in South Africa and designated as strain F2, redesignated as *A. vitis* strain F2/5 by Burr and Reid (1994), was proven to be nontumorigenic on grapevines and effective against grapevine CG (Staphorst et al. 1985, Burr et al. 1999). Using *in vitro* testing, 17 of 25 serologically different non-agrocin-producing biotype-3 strains showed sensitivity to strain F2. Crown gall formation by 12 biotype-3 strains was prevented in the greenhouse when co-inoculation of 11 different non-agrocin-producing and four agrocin-producing tumorigenic *A. vitis* strains was made with strain F2 (Staphorst et al. 1985, Burr et al. 1999).

Burr and Reid (1994) tested *A. vitis* strain F2/5 against 25 *A. vitis* and 19 *A. tumefaciens* strains *in vitro* and 10 *A. vitis* and three *A. tumefaciens* on inoculated grapevine trunks. *In vitro*, 21 *A. vitis* and two *A. tumefaciens* (biotype-1) strains exhibited sensitivity to strain F2/5. Greatest efficacy of CG suppression was observed on grapevine trunks when suspensions containing equal concentrations of strain F2/5 and the pathogen were co-inoculated. Reduction of gall size for seven

A. vitis and for two *A. tumefaciens* strains was observed. Only in cases where *A. vitis* strain CG78 was used, no reduction was exhibited in the number of inoculated sites in which galls developed (Burr and Reid 1994).

The protective action of antagonist *A. vitis* strains F2/5, CG1077 (the agrocin-minus mutant of F2/5), CG523, and *A. radiobacter* strain HLB-2 was tested (Bazzi et al. 1999). Those antagonists were used against the pathogenic *A. vitis* strain CG49. Influences on grapevine vitality, graft take, and wood and root productions were assessed. Eight months posttreatment, callus tissues were collected close to the graft union and subjected to quantitative isolation of *A. vitis* strain CG4. Maximum decrease in the internal concentration of this strain was achieved in grapevines treated with the antagonist F2/5-mutant strain CG1077. Excessive lignification and severe tissue necrosis at both the graft union and vascular parenchyma, throughout the vines treated with the wild-type strain F2/5, were reported. The strain F2/5 treatment significantly reduced graft take (from 97.5 to 67.5%) while there was only a slight decrease (from 67.1 to 63.2%) for the control. Grapevines treated with the BCAs (other than F2/5) also exhibited fewer necrotic symptoms on woody parenchyma tissues. Bazzi et al. (1999) concluded that establishing vineyards with grapevines containing a high concentration of *A. vitis* may be difficult.

Burr et al. (1997) questioned whether the biological control ability of strain F2/5 is based on agrocin production. Other hypotheses, including competition for attachment sites on grapevine cells and possible prevention of T-DNA transfer by strain F2/5, were also addressed. Chardonnay vines were inoculated with a 1:1 ratio of mixed suspensions containing the *A. vitis* strain CG49 and strain F2/5 or its agrocin-minus mutants CG1076, CG1077, CG1078, or CG1079. The number of cells of strain CG49 that attached to grapevine shoot tissues was significantly reduced (from $\sim 10^5$ to 10^4 cfu/mL) following co-inoculation with either F2/5 or CG1077 strains. Results confirmed that grapevine CG control by strain F2/5 is not associated with agrocin production or with competition for attachment sites (Burr et al. 1997). Moreover, it is not associated with its tartrate or with octopine utilization plasmids (Szegedi et al. 1999).

Strain F2/5 was GM for trifolixotoxin (TFX) peptide antibiotic production (Herlache and Triplett 2002). The strain (*A. vitis* strain F2/5(pT2TFXK)) exhibited an enhanced level of CG disease suppression. It contains a plasmid that is encoded for TFX production. Strain F2/5(pT2TFXK) is more effective against all tested *A. vitis* tumorigenic strains than the wild-type F2/5, as well as the TFX nonproducing negative control, F2/5(pT2TX3K). One month after co-inoculation of *Nicotiana glauca*, with wild-type F2/5 and *A. vitis* strains CG49, CG78, or CG435, gall formation occurred in all plants. However, when co-inoculation was performed in the same manner, but with the GM TFX-producing F2/5(pT2TFXK), gall formation was reduced. In those cases, 10 of 14 (71.4%), 1 of 8 (12.5%), and 0 of 14 (0.0%) of the inoculants exhibited tumor symptoms, respectively (Herlache and Triplett 2002). The competition of TFX compounds with *Agrobacterium* and

Rhizobium spp. is likely an important factor of its biological control mechanism (Robledo et al. 1998, Scupham and Triplett 2006). More work is needed to specify TFX mode of action.

Potential negative effects of strain F2/5 on grapevine tissues were previously reported (Bazzi et al. 1999) that include wound healing and graft-take reductions, partial callus formation, excessive lignification, and tissue necrosis in cuttings. Those negative effects were confirmed (Creasap et al. 2005). Necrotic symptoms were observed on cells generated following cambium activation at wounded sites. This necrosis was suggested as a cause for abnormal wound healing and failure of graft union. The necrosis is regulated by expression of a *luxR* homolog, *aviR* gene (Zheng et al. 2003, Hao et al. 2005). The exact mode of action by which strain F2/5 controls grape CG remains unknown.

***A. vitis* strain VAR03-1.** The nonpathogenic *A. vitis* strain VAR03-1 was the most effective strain tested against pathogenic *A. vitis* isolates in several experiments. In the first series of experiments (Kawaguchi et al. 2005), all the 11 tumorigenic *A. vitis* strains showed in vitro sensitivity to strain VAR03-1. One year after in vivo co-inoculation of Neo Muscat seedling stems, no withered symptoms were observed. In this case, strain VAR03-1 was co-inoculated with the tumorigenic *A. vitis* strain G-Ag-27. In a separate experiment, significant reduction was achieved (as compared to when only G-Ag-27 was used as inoculum-positive control) in the number of sites at which galls developed. Ten grapevine seedling stems were used for each replication and seven inoculations were made on each stem. Co-inoculations were made with suspensions containing a 1:1 ratio of the pathogenic G-Ag-27 and VAR03-1. Tumor formation reductions (TFRs), measured 12 weeks after inoculation, were from 66 of 70 to 0 of 70 (94.3% TFR), from 70 of 70 to 23 of 70 (67.1% TFR), and from 68 of 70 to 10 of 70 (82.9% TFR) (Kawaguchi et al. 2005).

In the second series of experiments, grape CG biological control ability of strain VAR03-1 was assessed against *A. vitis* strains G-Ag-27 and seven other *A. vitis* strains isolated in Japan (Kawaguchi et al. 2007). Inoculation of Neo Muscat cuttings was made with suspensions containing $\sim 10^8$ cfu/mL of pathogenic inoculant-positive control. Co-inoculation was made with a 1:1 ratio of the pathogenic strain and the BCA VAR03-1. Ten grapevine seedling stems were used for each replication and six inoculations were made on each stem. Twelve weeks after inoculations, CG incidence were reduced from 93.3% (highest TFR) to 33.3% (lowest TFR) when the BCA and strain A5-6 or strain A5-1 were used in co-inoculations. Statistically significant reduction in CG incidence by *A. vitis* strain G-Ag-27 was also reported after grapevine root treatments (Kawaguchi et al. 2007). In the third series of experiments, high efficacy of strain VAR03-1 to control grape CG caused by *A. vitis* strains MAFF 211674, MAFF 211676, At-90-23, and G-Ag-27 was reported (Kawaguchi et al. 2008).

Antagonists outside the genus *Agrobacterium*. Although the majority of BCAs tested against grape CG belong to the genus *Agrobacterium*, significant efforts have been made to identify potential biological control organisms outside this genus. In one study, 24 of 851 bacterial isolates exhibited

activity against *A. vitis* strains (Bell et al. 1995). Biolog identification tests revealed that the few effective isolates were strains of *Enterobacter agglomerans*, *Rahnella aquatilis*, and *Pseudomonas* spp. The most rapid decline of population of *A. vitis* strain AA25 in soil was achieved when soil co-inoculations were made using a 10:1 ratio of AA25 and *Pseudomonas corrugata* strain JC583. In vitro and in planta, three *P. corrugata* strains exhibited various levels of biological control ability as measured by growth and tumor formation inhibition of tumorigenic *A. vitis* strains, respectively (Bell et al. 1995).

***Pseudomonas* spp. strains.** *Pseudomonas aureofaciens* strain B-4117 and *P. fluorescens* strain CR330D were tested for biological control activity against *A. vitis* strains Tm4 and Sz1 (Khmel et al. 1998). The pattern of decline over time in numbers of antagonistic bacteria on grapevine roots was similar for all strains used. Rapid decline was observed 120 days after inoculation and then remained constant for up to one year. When cuttings were treated with antagonists, a decrease in CG incidence and lowered pathogenicity, as shown by reduction in tumor sizes, were observed as compared to water-treated control vines (Khmel et al. 1998).

***Rahnella aquatilis* strain HX2.** Although inhibitory effects of *R. aquatilis* bacteria toward various isolates of *A. vitis* were already reported (Bell et al. 1995), only recently have extensive studies on this potential BCA been initiated. *Rahnella aquatilis* strain HX2 isolated in China from vineyard soil was shown to be effective against a number of *A. vitis* isolates (Chen et al. 2007). Experiments confirmed the remarkable efficacy of HX2 for short-term CG suppression (Chen et al. 2007). Further biological control assays and characterizations of strain HX2 have also been made (Chen et al. 2009). Analysis of the antimicrobial compound produced by this strain revealed that the compound is likely to be thermostable, contains one or more kind of sugars, is most active against *Agrobacterium* spp., and inhibits *Clavibacter*, *Pectobacterium*, *Pseudomonas*, and *Xanthomonas* spp. It was determined that its mode of action does not include bacterial cell lysis or cause cytoplasmic material leakage at minimal inhibitory concentration. Analyses conducted on the impacts of HX2 antimicrobial compound(s) suggested that inhibition of RNA and protein synthesis, required for T-DNA transfer and tumor formation processes, is likely to be the main factor involved in its mode of action (Chen et al. 2009).

Genetic engineering strategies to prevent infection. Several experiments targeted the prevention of grape CG infection by expressing antimicrobial peptides in GM plants inhibitory to *A. vitis*. Genetically modified Chardonnay grapevines, which possessed either the natural (*Mag-2*) or the synthetic (*MSI99*) maganin short peptide antimicrobial compound-producing genes, were assayed. Their ability to confer resistance to grape CG (*A. vitis*) and powdery mildew (*Erysiphe necator*) was tested (Vidal et al. 2006). The development of strong resistance was not achieved; nonetheless, regenerated lines expressing the *Mag-2* or *MSI99* genes exhibited significantly less CG incidence than non-GM controls. Enhancement of transgene expression and the spectrum of resistance were suggested to achieve grape CG and powdery

mildew resistance with GM grapevines carrying genes encoded for maganin antimicrobial compound production (Vidal et al. 2006).

Crown gall suppression via GM grapevines carrying the *Mag-2* or *MSI99* genes was demonstrated (Kikkert et al. 2009). Decreased pathogenicity by reduction in gall sizes was observed on greenhouse-grown maganin-expressing Chardonnay inoculated with *A. vitis* strain TM4 or CG450. In vitro inhibition of those *A. vitis* strains by *Mag-2* and *MSI99* compounds was also reported (Kikkert et al. 2009).

These results suggest the high CG resistance potential of grapevines expressing *Mag-2* or *MSI99* genes. Increased resistance to CG may be achieved by expressing the *Mag-2* or *MSI99* peptides in GM grapevines containing truncated *vir* genes, which may result in antimicrobial activity and inhibition of T-DNA export and integration simultaneously. This multiple inhibition of pathogenic strains may prevent CG formation by simultaneous blocking of the infection and genetic transformation. Therefore, further research with the *Mag-2* and *MSI99* peptides in GM grapevines is warranted.

Strategies to prevent T-DNA export to plant cell and integration into the plant nuclear genome. High-efficacy CG-resistant GM grapevine generations were confirmed by transferring and expressing the truncated form of the Ti-plasmid *virE2* gene in grapevine somatic embryos via *Agrobacterium*-mediated transformation (Krastanova et al. 2010). According to several previous *VirE2*-involved T-DNA export and integration blocking studies, the increased resistance is believed to be associated with competition of the mutated *VirE2* proteins with functional *VirE2* proteins. Within the plant cell the competition appears to block the integration of the T-DNA into the plant cell nucleus (Burr and Otten 1999, Gelvin 2003, Matthyse 2006, Otten et al. 2008). The truncated *virE2* genes from *A. tumefaciens* strain C58 and A6 and *A. vitis* strain CG450 were expressed in Teleki 5C, Richter 110, and Couderc 3309 grapevine rootstocks. The truncated and transferred *virE2* genes were present in 314 of 322 (97.5%) GM lines, confirmed by double antibody sandwich enzyme-linked immunosorbent assay (DAS-ELISA), and in 285 of 295 (96.6%) GM lines, confirmed by polymerase chain reaction (PCR). In vitro susceptibility to *Agrobacterium* was evaluated 14 and 21 days postinoculation by visual observation of tumor formation on inoculated shoot internodes. Genetically modified plants with no galls or with galls less than 2 mm diameter on half or less of the total number of explants were scored as resistant. In vitro, shoot internodes of Richter 110 transformed with strain C58-*virE2* gene were inoculated with one of three tumorigenic strains: *A. tumefaciens* C58 (nopaline), A6 (octopine), or *A. vitis* CG450 (vitopine). This resulted in tumor formations being significantly reduced when compared to the non-GM control Richter 110. The highest TFR was achieved when tumorigenic *A. tumefaciens* strain C58 was used in the inoculations of strain C58-*virE2* gene-transformed Richter 110. All of the non-GM control Richter 110 exhibited visible tumors. In vitro, shoot internodes of Teleki 5C, transformed with C58, A6, or CG450-*virE2* gene, were inoculated with homologous or heterologous tumorigenic *A. tumefaciens*

strain C58, A6, or *A. vitis* strain CG450. As in the case before, tumor formations were reduced significantly when compared to the non-GM control Teleki 5C.

The highest TFRs (90.0%), as compared to the non-GM control Teleki 5C (5.0 and 0.0%), were achieved when tumorigenic *A. tumefaciens* strain C58 or *A. vitis* strain CG450 was used in the inoculations of strain C58-*virE2* or CG450-*virE2* gene-transformed Teleki 5C, respectively. In cases of the non-GM control Teleki 5C inoculations, CG developed on 95.0, 94.7, and 100.0% strain C58, A6, or CG450 inoculated shoot internodes, respectively (Krastanova et al. 2010).

This research confirmed the feasibility of GM rootstock grapevine production using truncated *virE2* genes of tumorigenic *A. tumefaciens* strain C58, A6, and *A. vitis* CG450. Due to the highest TFR performances exhibited by C58-*virE2* gene-transformed vines, the truncated C58-*virE2* can be considered one of the most effective genetic resources available for CG-resistant GM grapevine generation. Evaluation of long-term CG resistance of the generated GM vines in vineyard settings and further investigation into the efficacy of truncated *virE2* genes of other previously untested *Agrobacterium* strains are suggested.

The development of CG-resistant Richter 110 rootstock grapevines was achieved by blocking T-DNA integration into the host genome (Holden et al. 2003). Genetically modified grapevines containing the truncated *virE2* genes of *A. tumefaciens* strain C58 or A6 or of *A. vitis* strain CG450 were generated and assessed for CG susceptibility. Seven lines of shoot explants, transformed with the truncated C58-*virE2* gene and inoculated with strain C58, expressed resistance. Two of these seven also showed resistance to the other two *Agrobacterium* strains (Holden et al. 2003).

These results suggest that transformation of grapevines with strain-specific truncated *Agrobacterium virE2* genes can result in grapevines expressing resistance to different *Agrobacterium* biotypes and strains. As reported by Krastanova et al. (2010), this experiment confirmed the importance of truncated *virE2* gene of *A. tumefaciens* strain C58 in CG-resistant grapevine generation.

A total of 928 putative GM grapevine combinations were generated (Xue et al. 1999) and tested for resistance to grape CG, grapevine fanleaf virus (GFLV), and grapevine leafroll-associated closterovirus-3 (GLRaV-3). Transformed vines contained a sense-oriented translatable gene of GFLV, an antisense coat protein gene of truncated HSP90-related (heat-shock protein 90) gene of GLRaV-3, and a *virE2* of B gene (a truncated *virE2* that lacks 215 carboxyl-terminal amino acids; Citovsky et al. 1994) from *A. tumefaciens* strain C58. Transformation efficiency of the five grape rootstocks with seven gene constructs varied. Rates of transformation ranged from 45 to 100%. Transgenes were present in each case as determined by PCR or Southern Blot analysis. Disease and virus resistance evaluations of the GM vines are in progress (Xue et al. 1999).

Strategies to prevent expression of T-DNA oncogenes. Transferred-DNA oncogene expression prevention (often referred to as “gene silencing”), using RNAi technology, in-

volves within-plant cell expression of messenger ribonucleic acid (mRNA) complementary to the mRNA of a gene whose expression inhibition is targeted. Expression inhibition of a gene is accomplished through 21 to 25 nucleotides-long RNA species (RNAs) that are derived to become RNA-induced gene silencing (Agrawal et al. 2003). The use of RNAs leads to the degradation of targeted RNA to nucleotides resulting in T-DNA oncogene expression prevention. Efficacy of gene silencing for high-level CG resistance was confirmed by several research experiments (Ebinuma et al. 1997, Escobar et al. 2001, Kovács et al. 2003, Lee et al. 2003).

Feasibility of silencing the expression of an *A. vitis* oncogene, thus preventing tumor formation, has been investigated (Kovács et al. 2003). The study demonstrated oncogene silencing in GM *Petunia hybrida* containing double-transformed 35S-*iaaM* gene. One line of those transformed plants exhibited reduced *iaaM* RNA levels; however, it failed to attenuate tumorigenesis when challenged with *A. vitis* isolates. A possible explanation for this phenomenon is that the 35S-*iaaM* transgene may have produced a chimeric RNA molecule that was unable to induce silencing of the *A. vitis iaaM* gene (Kovács et al. 2003, Kovács, personal communication, 2009). It was demonstrated that the *iaaM* silencing construct must contain the translation start site of the gene’s sense strand in order for silencing to be effective (Lee et al. 2003).

Conclusion

Whether biological control using natural bacterial strains, biological control by genetically modified antagonistic strains, or genetic engineering of grapevines will lead to control of grape CG disease is still under investigation. Despite current limitations, biological control appears to have the greatest probability for success in grapevine CG control. Moreover, it shall likely prove to be the most palatable strategy for consumers, which would expectedly drive the preference of the industry accordingly. The genetic diversity of *Agrobacterium* grapevine isolates is considered the major limitation for successful biological control. For example, diversity in the *A. vitis* population could affect susceptibility to bacteriocins or antibiotics produced by antagonistic strains. In numerous research experiments, the effectiveness of grapevine CG BCAs was restricted to certain opine-metabolizing tumorigenic *A. vitis* strains, which suggests that the sensitivity of various opine-metabolizing strains to biological control strains varies. Assessments of opine-metabolizing strain sensitivities to BCAs may foster understanding of action mechanisms of biological control strains. Deoxyribonucleic acid fingerprint comparison may also foster understanding of the genetic background of the diversity of *Agrobacterium* spp., which is essential for developing more effective BCAs. These may lead to the development of a BCA effective against each opine-metabolizing *A. vitis* strain type. In addition to efficacy against pathogenic strains, the limitation of biological control strains includes their persistence following deployment, including persistence in various soil types in vivo and in different rooting media. Endophytic colonization by BCAs and systemic movement throughout the plant are some other major limitations to their

effective deployment. Nevertheless, the commercial success of *A. radiobacter* strain K84 and derivatives in controlling CG caused by *A. tumefaciens* confirms that this approach could also be used in grape CG management. Future research, of both previously tested and potential new isolates of BCAs, is necessary to: determine specific mode(s) of action, complete molecular and genetic understanding of biological control mechanisms, discover and develop new BCAs, field-test all effective biological control strains, and develop combinations of various BCAs capable of controlling grapevine CG disease regardless of the diversity of *A. vitis* strains.

Genetic mechanisms that trigger malignant cell growth and underlie CG formation are still not completely understood. Determination of currently unknown grapevine and *Agrobacterium* genes involved in the grape CG infection and genetic transformation processes and their exact roles would enhance understanding of the mechanisms of this disease. Furthermore, technological approaches in GM grapevine production, including the production of GM CG-resistant grapevines, must ensure that GM grapevines will have the same positive horticultural and quality characteristics as the non-GM plants.

Continued interest of the grape industry in CG-resistant grapevines is another factor critically necessary to achieve CG-resistant grapevine production. Virus-resistant GM grapevine production was accomplished with significant support from the grape and wine industry after unacceptable economic losses attributed to grapevine decline due to viral infections. Produced vines have been tested under experimental field conditions in different regions of the world; however, they are not available commercially. Therefore, to achieve CG-resistant grapevine production, close collaboration between scientific institutions and grape growers is pivotal. Improvements in the following areas may further foster the development of CG-resistant GM grapevine production: development of grape variety-specific regeneration methods to increase regeneration efficiency; reduction of the 5- to 10-year time requirement to generate testable GM grapevines; development of more rapid, accurate, and sensitive indexing systems to increase the reliability of presence determination of inserted genes and for the selective detection of tumorigenic and nontumorigenic *Agrobacterium* strains within grapevines; and development of a standardized evaluation system for CG resistance. In addition, successful commercial production of GM CG-resistant grapevines cannot be achieved without public acceptance.

Although biological control and genetic engineering were discussed separately, successful suppression of grapevine CG is likely to be achieved by incorporating and applying strategies from both those areas. This was mainly indicated in this review by describing GM BCAs and their applications. The importance of further collaboration among researchers from those fields and grape industry representatives would seem essential to further progress.

Literature Cited

Agrawal, N., P.V.N. Dasaradhi, A. Mohammed, P. Malhotra, R.K. Bhatnagar, and S.K. Mukherjee. 2003. RNA interference: Biology, mechanism, and applications. *Microbiol. Mol. Biol. Rev.* 67:657-685.

- Argun, N., M.T. Momol, S. Maden, E.A. Momol, C.L. Reid, H. Çelek, and T.J. Burr. 2002. Characterization of *Agrobacterium vitis* strains isolated from Turkish grape cultivars in the Central Anatolia region. *Plant Dis.* 86:162-166.
- Bazzi, C., M. Alexandrova, E. Stefani, F. Anaclerio, and T.J. Burr. 1999. Biological control of *Agrobacterium vitis* using non-tumorigenic agrobacteria. *Vitis* 38:31-35.
- Bell, C.R., G.A. Dickie, and J.W.Y.F. Chan. 1995. Variable response of bacteria isolated from grapevine xylem to control grape crown gall disease in planta. *Am. J. Enol. Vitic.* 46:499-508.
- Burr, T.J., and B.H. Katz. 1984. Grapevine cuttings as potential sites of survival and means of dissemination of *Agrobacterium tumefaciens*. *Plant Dis.* 68:976-978.
- Burr, T.J., and L. Otten. 1999. Crown gall of grape: Biology and disease management. *Ann. Rev. Phytopathol.* 37:53-80.
- Burr, T.J., and C.L. Reid. 1994. Biological control of grape crown gall with non-tumorigenic *Agrobacterium vitis* strain F2/5. *Am. J. Enol. Vitic.* 45:213-219.
- Burr, T.J., C. Bazzi, S. Süle, and L. Otten. 1998. Crown gall of grape: Biology of *Agrobacterium vitis* and the development of disease control strategies. *Plant Dis.* 82:1288-1297.
- Burr, T.J., A.L. Bishop, B.H. Katz, L.M. Blanchard, and C. Bazzi. 1987. A root-specific decay of grapevine caused by *Agrobacterium tumefaciens* and *A. radiobacter* biovar 3. *Phytopathology* 77:1424-1427.
- Burr, T.J., B.H. Katz, A.L. Bishop, C.A. Meyers, and V.L. Mittak. 1988. Effect of shoot age and tip culture propagation of grapes on systemic infestations by *Agrobacterium tumefaciens* biovar 3. *Am. J. Enol. Vitic.* 39:67-70.
- Burr, T. J., L. Kovács, S. Süle, and E. Szegedi. 2003. Breeding for crown gall resistance: Traditional and molecular approaches. *Acta Hort.* 603:441-447.
- Burr, T.J., C.L. Reid, C.E. Adams, and E.A. Momol. 1999. Characterization of *Agrobacterium vitis* strains isolated from feral *Vitis riparia*. *Plant Dis.* 83:102-107.
- Burr, T.J., C.L. Reid, D.F. Splittstoesser, and M. Yoshimura. 1996. Effect of heat treatments on grape bud mortality and survival of *Agrobacterium vitis* in vitro and in dormant grape cuttings. *Am. J. Enol. Vitic.* 47:119-123.
- Burr, T.J., C.L. Reid, E. Tagliati, C. Bazzi, and S. Süle. 1997. Biological control of grape crown gall by strain F2/5 is not associated with agrocin production or competition for attachment sites on grape cells. *Am. Phytopathol. Soc.* 87:706-711.
- Chen, F., Y.B. Guo, J.H. Wang, J.Y. Li, and H.M. Wang. 2007. Biological control of grape crown gall by *Rahnella aquatilis* HX2. *Plant Dis.* 91:957-963.
- Chen, F., J.Y. Li, Y.B. Guo, J.H. Wang, and H.M. Wang. 2009. Biological control of grapevine crown gall: Purification and partial characterization of an antibacterial substance produced by *Rahnella aquatilis* strain HX2. *Eur. J. Plant Pathol.* 124:427-437.
- Chen, X.Y., and W.N. Xiang. 1986. A strain of *Agrobacterium radiobacter* inhibits growth and gall formation by biotype III strains of *A. tumefaciens* from grapevine. *Acta Microbiol. Sin.* 26:193-199.
- Citovsky, V., D. Warnick, and P.C. Zambryski. 1994. Nuclear import of *Agrobacterium* VirD2 and VirE2 proteins in maize and tobacco. *Proc. Natl. Acad. Sci. USA* 91:3210-3214.
- Cooksey, D.A., and L.W. Moore. 1982. Biological control of crown gall with an agrocin mutant of *Agrobacterium radiobacter*. *Am. Phytopathol. Soc.* 72:919-921.
- Costacurta, A., and J. Vanderleyden. 1995. Synthesis of phytohormones by plant associated bacteria. *Crit. Rev. Microbiol.* 21:1-18.
- Creasap, J.E., C.L. Reid, M.C. Goffinet, R. Aloni, C. Ullrich, and T.J. Burr. 2005. Effect of wound position, auxin and *Agrobacterium*

- vitis* strain F2/5 on wound healing and crown gall in grapevine. *Phytopathology* 95:362-367.
- De Cleene, M., and J. De Ley. 1976. The host range of crown gall. *Bot. Rev.* 42:389-466.
- Dessaux, Y., A. Petit, S.K. Farrand, and P.J. Murphy. 1998. Opines and opine-like molecules involved in plant-*Rhizobiaceae* interactions. In *The Rhizobiaceae: Molecular Biology of Model Plant-Associated Bacteria*. H.P. Spaink et al. (eds.), pp. 173-197. Kluwer Academic, Dordrecht, Netherlands.
- Donner, S.C., D.A. Jones, N.C. McClure, G.M. Rosewarne, M.E. Tate, A. Kerr, N.N. Fajardo, and B.G. Clare. 1993. Agrocin 434, a new plasmid encoded agrocin from the biocontrol *Agrobacterium* strains K84 and K1026, which inhibits biovar 2 agrobacteria. *Physiol. Mol. Plant Pathol.* 42:185-194.
- Eastwell, K.C., P.L. Sholberg, and R.J. Sayler. 2006. Characterizing potential bacterial biocontrol agents for suppression of *Rhizobium vitis*, casual agent of crown gall disease in grapevines. *Crop Prot.* 25:1191-1200.
- Ebinuma, H., E. Matsunaga, K. Yamada, and M. Yamakado. 1997. Transformation of hybrid aspen for resistance to crown gall disease. In *Micropropagation, Genetic Engineering, and Molecular Biology of Populus*. N.B. Klopfenstein et al. (eds.), pp. 161-164. USDA Forest Service, Rocky Mountain Forest and Range Experiment Station, Fort Collins, CO.
- Escobar, M.A., E.L. Civerolo, K.R. Summerfelt, and A.M. Dandekar. 2001. RNAi-mediated oncogene silencing confers resistance to crown gall tumorigenesis. *Proc. Natl. Acad. Sci. USA* 98:13437-13442.
- Farrand, S.K., J.E. Slota, J.S. Shim, and A. Kerr. 1985. Tn5 insertions in the agrocin 84 plasmid: The conjugal nature of pAgK84 and the locations of determinants for transfer and agrocin 84 production. *Plasmid* 13:106-117.
- Fournier, P., P. de Ruffray, and L. Otten. 1994. Natural instability of *Agrobacterium vitis* Ti plasmid due to unusual duplication of a 2.3-kb DNA fragment. *Mol. Plant Microbe In.* 7:164-172.
- Gaudin, V., T. Vrain, and L. Jouanin. 1994. Bacterial genes modifying hormonal balances in plants. *Plant Physiol. Biochem.* 32:11-29.
- Gelvin, S.B. 2003. *Agrobacterium*-mediated plant transformation: The biology behind the “gene-jockeying” tool. *Microbiol. Mol. Biol. Rev.* 67:16-37.
- Hao, G., H. Zhang, D. Zheng, and T. J. Burr. 2005. *luxR* homolog *avhR* in *Agrobacterium vitis* affects the development of a grape-specific necrosis and a tobacco hypersensitive response. *J. Bacteriol.* 187:185-192.
- Herlache, T.C., and E.W. Triplett. 2002. Expression of a crown gall biological control phenotype in an avirulent strain of *Agrobacterium vitis* by addition of the trifolixotoxin production and resistance genes. *BMC Biotechnol.* 2:2.
- Holden, M., S. Krasatanova, B. Xue, S. Pang, M. Sekiya, E.A. Momol, D. Gonsalves, and T.J. Burr. 2003. Genetic engineering of grape for resistance to crown gall. *Acta Hort.* 603:481-484.
- Huss, B., B. Tinland, F. Paulus, B. Walter, and L. Otten. 1990. Functional analysis of a complex oncogene arrangement in biotype III *Agrobacterium tumefaciens* strains. *Plant Mol. Biol.* 14:173-186.
- Kawaguchi, A., and K. Inoue. 2009. Grapevine crown gall caused by *Rhizobium radiobacter* (Ti) in Japan. *J. Gen. Plant Pathol.* 75:205-212.
- Kawaguchi, A., K. Inoue, and Y. Ichinose. 2008. Biological control of crown gall of grapevine, rose, and tomato by nonpathogenic *Agrobacterium vitis* strain VAR03-1. *Am. Phytopathol. Soc.* 98:1218-1225.
- Kawaguchi, A., K. Inoue, and H. Nasu. 2005. Inhibition of crown gall formation by *Agrobacterium radiobacter* biovar 3 strains isolated from grapevine. *J. Gen. Plant Pathol.* 71:422-430.
- Kawaguchi, A., K. Inoue, and H. Nasu. 2007. Biological control of grapevine crown gall by nonpathogenic *Agrobacterium vitis* strain VAR03-1. *J. Gen. Plant Pathol.* 73:133-138.
- Kerr, A. 1972. Biological control of crown gall: Seed inoculation. *J. Appl. Microbiol.* 35:493-497.
- Kerr, A., and K. Htay. 1974. Biological control of crown gall through bacteriocin production. *Physiol. Plant Pathol.* 4:37-44.
- Kerr, A., and C.G. Panagopoulos. 1977. Biotypes of *Agrobacterium radiobacter* var. *tumefaciens* and their biological control. *Phytopathol. Z.* 90:172-179.
- Khmel, I.A., T.A. Sorokina, N.B. Lemanova, V.A. Lipasova, O.Z. Metlitski, T.V. Burdeinaya, and L.S. Chernin. 1998. Biological control of crown gall in grapevine and raspberry by two *Pseudomonas* spp. with a wide spectrum of antagonistic activity. *Biocontrol Sci. Tech.* 8:45-57.
- Kikkert, J.R., J.R. Vidal, P.G. Wallace, S. Garcia-Zitter, W.F. Wilcox, D.M. Gadoury, R.C. Seem, T.J. Burr, C.L. Rosenfield, S. Samuelian, and B.I. Reisch. 2009. Disease resistance analyses of transgenic grapevines that contain endochitinase or antimicrobial peptide genes. *Acta Hort.* 827:379-384.
- Kim, J.G., B.K. Park, S.U. Kim, D. Choi, B.H. Nahm, J.S. Moon, J.S. Reader, S.K. Farrand, and I. Hwang. 2006. Bases of biocontrol: Sequence predicts synthesis and mode of action of agrocin 84, the Trojan horse antibiotic that controls crown gall. *Proc. Natl. Acad. Sci. USA* 103:8846-8851.
- Kottek, M., J. Grieser, C. Beck, B. Rudolf, and F. Rubel. 2006. World map of the Köppen-Geiger climate classification updated. *Meteorol. Z.* 15:259-263.
- Kovács, L., S. Fekete, H. Hou, and E. Szegedi. 2003. Silencing the expression of an *Agrobacterium vitis* oncogene in a model plant. *Acta Hort.* 603:519-526.
- Krastanova, S.V., V. Balaji, M.R. Holden, M. Sekiya, B. Xue, E.A. Momol, and T.J. Burr. 2010. Resistance to crown gall disease in transgenic grapevine rootstocks containing truncated *virE2* of *Agrobacterium*. *Transgenic Res.* 19:949-958.
- Lacroix, B., M. Elbaum, V. Citovsky, and T. Tzfira. 2008. Intracellular transport of *Agrobacterium* T-DNA. In *Agrobacterium: From Biology to Biotechnology*. T. Tzfira and V. Citovsky (eds.), pp. 365-394. Springer, New York.
- de Lajudie, P., E. Laurent-Fulele, A. Willems, U. Torck, R. Coopman, M.D. Collins, K. Kersters, D. Dreyfus, and M. Gillis. 1998. Description of *Allorhizobium undicola* gen. nov., sp. nov. for nitrogen-fixing bacteria efficiently nodulating *Neptunia natans* in Senegal. *Int. J. Syst. Bacteriol.* 48:1277-1290.
- van Larebeke, N., G. Engler, M. Holsters, S. Van den Elsacker, I. Zaenen, R.A. Schilperoort, and J. Schell. 1974. Large plasmid in *Agrobacterium tumefaciens* essential for crown gall-inducing ability. *Nature* 252:169-170.
- Lee, H., J.L. Humann, J.S. Pitrak, J.T. Cuperus, T.D. Parks, C.A. Whistler, M.C. Mok, and L.W. Ream. 2003. Translation start sequences affect the efficiency of silencing of *Agrobacterium tumefaciens* T-DNA oncogenes. *Plant Physiol.* 133:966-977.
- Lehoczy, J. 1968. Spread of *Agrobacterium tumefaciens* in the vessels of the grapevine after natural infection. *Phytopathol. Z.* 63:239-246.
- Lehoczy, J. 1971. Further evidences concerning the systemic spreading of *Agrobacterium tumefaciens* in the vascular system of grapevines. *Vitis* 10:215-221.
- Lehoczy, J. 1978. Root-system of the grapevine as a reservoir of *Agrobacterium tumefaciens* cells. In *Proceedings of the Fourth International Conference on Plant Pathogenic Bacteria*. Station de Pathologie Végétale et Phytobactériologie (ed.), pp. 239-243. Institut National de la Recherche Agronomique, Angers, Beaucozéz, France.
- Lehoczy, J. 1989. Inoculation experiment on trellised grapevines with *Agrobacterium tumefaciens* for study of the process of crown gall disease. *Acta Phytopathol. Entomol. Hung.* 24:283-291.

- Li, J.Y., F. Chen, J.H. Wang, and H.M. Wang. 2005. Colonization of crown gall biocontrol agent – *Agrobacterium vitis* strain E26 on grapevine. *Acta Phytopathol. Sin.* 35:78-83.
- Liang, Y.J., Y.B. Di, J.Y. Zhao, and D.Q. Ma. 1990a. Strain E26 of *Agrobacterium radiobacter* efficiently inhibits crown gall formation on grapevines. *Chin. Sci. Bull.* 35:1055-1056.
- Liang, Y.J., J.Y. Zhao, and D.Q. Ma. 1990b. A biotype 3 strain of *Agrobacterium radiobacter* inhibits crown gall formation on grapevine. *Acta Microbiol. Sin.* 30:161-171.
- Liang, Z., H. Wang, and J. Wang. 2001. Preliminary study on effectiveness and the stability of E26 on controlling crown gall disease. *J. Chin. Agric. Univ.* 6:91-95.
- Matthysse, A.G. 2006. The genus *Agrobacterium*. *Prokaryotes* 5:91-114.
- Moore, L.W., and G. Warren. 1979. *Agrobacterium radiobacter* strain 84 and biological control of crown gall. *Ann. Rev. Phytopathol.* 17:163-179.
- Moriguchi, K., Y. Maeda, M. Satou, N.S. Hardayani, M. Kataoka, N. Tanaka, and K. Yoshida. 2001. The complete nucleotide sequence of a plant root-inducing (Ri) plasmid indicates its chimeric structure and evolutionary relationship between tumor-inducing (Ti) and symbiotic (Sym) plasmids in *Rhizobiaceae*. *J. Mol. Biol.* 307:771-784.
- Mozsár, J., O. Viczián, and S. Süle. 1998. *Agrobacterium*-mediated genetic transformation of an interspecific grapevine. *Vitis* 37:127-130.
- Morris, R.O. 1986. Genes specifying auxin and cytokinin biosynthesis in phytopathogens. *Ann. Rev. Plant. Physiol.* 37:509-538.
- Nakano, M., Y. Hoshino, and M. Mii. 1994. Regeneration of transgenic plants of grapevine (*Vitis vinifera* L.) via *Agrobacterium rhizogenes*-mediated transformation of embryogenic calli. *J. Exp. Bot.* 45:649-656.
- Oláh, R., E. Szegedi, S. Ruthner, and J. Korbuly. 2003. Optimization of conditions for regeneration and genetic transformation of rootstock- and scion grape varieties. *Acta Hort.* 603:491-497.
- Ophel, K., and A. Kerr. 1990. *Agrobacterium vitis* sp. nov. for strains of *Agrobacterium* biovar 3 from grapevines. *Int. J. System. Bacteriol.* 40:236-241.
- Otten, L., P. de Ruffray, E.A. Momol, M.T. Momol, and T.J. Burr. 1996. Phylogenetic relationships between *Agrobacterium vitis* isolates and their Ti plasmids. *Mol. Plant Microbe In.* 9:782-786.
- Otten, L., T.J. Burr, and E. Szegedi. 2008. *Agrobacterium*: A disease-causing bacterium. In *Agrobacterium: From Biology to Biotechnology*. T. Tzfira and V. Citovsky (eds.), pp. 1-46. Springer, New York.
- Paulus, F., B. Huss, G. Bonnard, M. Ridé, E. Szegedi, J. Tempé, A. Petit, and L. Otten. 1989. Molecular systematics of biotype III Ti plasmids of *Agrobacterium tumefaciens*. *Mol. Plant Microbe In.* 2:64-74.
- Peñalver, R., B. Vicedo, and M.M. López. 2000. Use of the genetically engineered *Agrobacterium* strain K1026 for biological control of crown gall. *Eur. J. Plant Pathol.* 106:801-810.
- Petersen, S.G., B.M. Stummann, P. Olesen, and K.W. Henningsen. 1989. Structure and function of root-inducing (Ri) plasmids and their relation to tumor-inducing (Ti) plasmids. *Physiol. Plant.* 77:427-435.
- Petit, A., C. David, G.A. Dahl, J.G. Ellis, P. Guyon, F. Casse-Delbart, and J. Tempé. 1983. Further extension of the opine concept: Plasmids in *Agrobacterium rhizogenes* cooperate for opine degradation. *Mol. Gen. Genet.* 190:204-214.
- Pinkerton, J.N., K.L. Ivors, M.L. Miller, and L.W. Moore. 2000. Effect of soil solarization and cover crops on populations of selected soil borne plant pathogens in western Oregon. *Plant Dis.* 84:952-960.
- Pionnat, S., H. Keller, D. Hericher, A. Bettachini, Y. Dessaux, X. Nesme, and C. Poncet. 1999. Ti plasmids from *Agrobacterium* characterize rootstock clones that initiated a spread of crown gall disease in Mediterranean countries. *Appl. Environ. Microbiol.* 65:4197-4206.
- Du Plessis, H.J., M.J. Hattingh, and H.J.J. van Vuuren. 1985. Biological control of crown gall in South Africa by *Agrobacterium radiobacter* strain K84. *Plant Dis.* 69:302-305.
- Portier, P., M. Fischer-Le Saux, C. Mougel, C. Lerondelle, D. Chapuliot, J. Thioulouse, and X. Nesme. 2006. Identification of genomic species in *Agrobacterium* biovar 1 by AFLP genomic markers. *Appl. Environ. Microbiol.* 72:7123-7131.
- Pu, X.A., and R.N. Goodman. 1993a. Effects of fumigation and biological control on infection of indexed crown gall free grape plants. *Am. J. Enol. Vitic.* 44:241-248.
- Pu, X.A., and R.N. Goodman. 1993b. Tumor formation by *Agrobacterium tumefaciens* is suppressed by *Agrobacterium radiobacter* HLB-2 on grape plants. *Am. J. Enol. Vitic.* 44:249-254.
- Robledo, E.A., J. Borneman, and E.W. Triplett. 1998. Effects of bacterial antibiotic production on rhizosphere microbial communities from a culture-independent perspective. *Appl. Environ. Microbiol.* 64:5020-5022.
- Sands, P., and P. Galizzi. 2006. Documents in European Community Environmental Law. Cambridge University Press, Cambridge, UK.
- Sawada, H., H. Ieki, H. Oyaizu, and S. Matsumoto. 1993. Proposal for rejection of *Agrobacterium tumefaciens* and revised descriptions for the genus *Agrobacterium* and for *Agrobacterium radiobacter* and *Agrobacterium rhizogenes*. *Int. J. Syst. Bacteriol.* 43:694-702.
- Scorza, R., J.M. Cordts, D.J. Gray, D. Gonsalves, R.L. Emershad, and D.W. Ramming. 1996. Producing transgenic 'Thompson Seedless' grape (*Vitis vinifera* L.) plants. *J. Am. Soc. Hort. Sci.* 121:616-619.
- Scupham, A.J., and E.W. Triplett. 2006. Determination of the amino acid residues required for the activity of the anti-rhizobial peptide antibiotic trifolixotoxin. *J. Appl. Microbiol.* 100:500-507.
- Staphorst, J.L., F.G.H. van Zyl, B.W. Strijdom, and Z.E. Groenewold. 1985. Agrocin-producing pathogenic and nonpathogenic biotype-3 pathogens. *Current Microb.* 12:45-52.
- Stewart, E.L., and N.G. Wenner. 2004. Grapevine decline in Pennsylvania and New York. *Wine East.* 32:12-21, 51-53.
- Süle, S., and T.J. Burr. 1998. The effect of resistance of rootstocks to crown gall (*Agrobacterium* spp.) on the susceptibility of scions in grape vine cultivars. *Plant Pathol.* 47:84-88.
- Süle, S., J. Lehoczky, G. Jenser, P. Nagy, and T.J. Burr. 1995. Infection of grapevine roots by *Agrobacterium vitis* and *Meloidogyne hapla*. *J. Phytopathol.* 143:169-171.
- Szegedi, E. 1995. A review of the use of thermotherapy in viticulture to eliminate pathogens and pests from propagating material. *Pestic. Sci.* 45:285-286.
- Szegedi, E., and S. Bottka. 2002. Detection of *Agrobacterium vitis* by polymerase chain reaction in grapevine bleeding sap after isolation on a semiselective medium. *Vitis* 41:37-42.
- Szegedi, E., and B.-né Dula. 2006. Az agrobaktériumos fertőzés kimutatása szőlőoltványokból. *Növényvédelem* 2:61-66.
- Szegedi, E., and P. Kozma. 1984. Studies on the inheritance of resistance to crown gall disease of grapevine. *Vitis* 23:121-126.
- Szegedi, E., and S. Süle. 2005. Agrobaktériumos fertőzéstől mentes szőlő szaporítóanyag előállítás. *Növényvédelem* 10:467-476.
- Szegedi, E., S. Bottka, J. Mikulás, L. Otten, and S. Süle. 2005. Characterization of *Agrobacterium tumefaciens* strains isolated from grapevine. *Vitis* 44:49-54.
- Szegedi, E., M. Czakó, L. Otten, and C. Koncz. 1988. Opines in crown gall tumors induced by biotype 3 isolates of *Agrobacterium tumefaciens*. *Physiol. Mol. Plant Pathol.* 32:237-247.
- Szegedi, E., J. Korbuly, and I. Koleda. 1984. Crown gall resistance in East-Asian *Vitis* species and in their *V. vinifera* hybrids. *Vitis* 23:21-26.
- Szegedi, E., S. Süle, and T.J. Burr. 1999. *Agrobacterium vitis* strain F2/5 contains tartrate and octopine utilization plasmids which do not encode functions for tumor inhibition on grapevine. *J. Phytopathol.* 147:665-669.

- Thomashow, M.F., S. Hugly, W.G. Buchholz, and L.S. Thomashow. 1986. Molecular basis for the auxin-independent phenotype of crown gall tumor tissues. *Science* 231:616-618.
- Thomson, J.A. 1986. The potential for biological control of crown gall disease on grapevines. *Trends Biotechnol.* 4:219-224.
- Torregrosa, L., P. Iocco, and M.R. Thomas. 2002. Influence of *Agrobacterium* strain, culture medium, and cultivar on the transformation efficiency of *Vitis vinifera* L. *Am. J. Enol. Vitic.* 53:183-190.
- Vicedo, B., R. Peñalver, M.J. Asins, and M.M. López. 1993. Biological control of *Agrobacterium tumefaciens*, colonization, and pAgK84 transfer with *Agrobacterium radiobacter* K84 and the Tra⁺ mutant strain K1026. *Appl. Environ. Microbiol.* 59:309-315.
- Vidal, J.R., J.R. Kikkert, M.A. Malnoy, P.G. Wallace, J. Barnard, and B.I. Reisch. 2006. Evaluation of transgenic 'Chardonnay' (*Vitis vinifera*) containing magainin genes for resistance to crown gall and powdery mildew. *Transgenic Res.* 15:69-82.
- Wang, H.M., H.X. Wang, T.B. Ng, and J.Y. Li. 2003. Purification and characterization of an antibacterial compound produced by *Agrobacterium vitis* strain E26 with activity against *A. tumefaciens*. *Plant Pathol.* 52:134-139.
- Wang, Y.H., L.Q. Zhang, J.Y. Li, J.H. Wang, and H.M. Wang. 2008. The quorum-sensing system AvsR-AvsI regulates both long-chain and short-chain acyl-homoserine lactones in *Agrobacterium vitis* E26. *Anton. Leeuwenhoek* 93:267-273.
- Webster, J., and J. Thomson. 1988. Genetic analysis of an *Agrobacterium tumefaciens* strain producing an agrocin active against biotype 3 pathogens. *Mol. Gen. Genet.* 214:142-147.
- Webster, J., M.D. Santos, and J.A. Thomson. 1986. Agrocin-producing *Agrobacterium tumefaciens* strain active against grapevine isolates. *Appl. Environ. Microbiol.* 52:217-219.
- Wei, Q., J.Y. Li, J.H. Wang, and H.M. Wang. 2009. Strain E26 of *Agrobacterium vitis*, a biological control agent of grapevine crown gall, does not contain *virA* and *virG* pathogenic determinants. *J. Phytopathol.* 157:657-665.
- Willems, A. 2006. The taxonomy of rhizobia: an overview. *Plant Soil* 287:3-14.
- Willems, A., and M.D. Collins. 1993. Phylogenetic analysis of rhizobia and agrobacteria based on 16S rRNA gene sequences. *Int. J. Syst. Bacteriol.* 43:305-313.
- Wilson, C.L. 1997. Biological control and plant diseases—A new paradigm. *J. Ind. Microbiol. Biotechnol.* 19:158-159.
- Xue, B., K.S. Ling, C.L. Reid, S. Krastanova, M. Sekiya, E.A. Momol, S. Süle, J. Mozsár, D. Gonsalves, and T.J. Burr. 1999. Transformation of five grape rootstocks with plant virus genes and a *virE2* gene from *Agrobacterium tumefaciens*. *In vitro Cell. Dev. Biol. Plant.* 35:226-231.
- Xu, X., and S.Q. Pan. 2000. An *Agrobacterium* catalase is a virulence factor involved in tumorigenesis. *Mol. Microbiol.* 35:407-414.
- Xuemei, X., Y. Jifeng, C. Peimin, and G. Junmei. 1993. On a strain MI 15 of *Agrobacterium radiobacter* for the biological control of grapevine crown gall. *Acta Phytopathol. Sin.* 23:137-141.
- Young, J.M., L.D. Kuykendall, E. Martínez-Romero, A. Kerr, and H. Sawada. 2001. A revision of *Rhizobium* Frank 1889, with an emended description of the genus, and the inclusion of all species of *Agrobacterium* Conn 1942 and *Allorhizobium undicola* de Lajudie et al. 1998 as new combinations: *Rhizobium radiobacter*, *R. rhizogenes*, *R. rubi*, *R. undicola* and *R. vitis*. *Int. J. Syst. Evol. Microbiol.* 51:89-103.
- Zabadal, T.J., I.E. Dami, M.C. Goffinet, T.E. Martinson, and M.L. Chien. 2007. Winter injury to grapevines and methods of protection. Extension Bulletin E2930. Michigan State University, East Lansing.
- Zheng, D., H. Zhang, S. Carle, G. Hao, M.R. Holden, and T.J. Burr. 2003. A *luxR* homolog, *aviR*, in *Agrobacterium vitis* is associated with induction of necrosis on grape and a hypersensitive response on tobacco. *Mol. Plant Microbe In.* 16:650-658.
- Zupan, J.R., and P.C. Zambryski. 1995. Transfer of T-DNA from *Agrobacterium* to the plant cell. *Plant Physiol.* 107:1041-1047.