





















Co-Occurrence of Cytogenetic Abnormalities and High-Risk Disease in Newly Diagnosed and Relapsed/Refractory Multiple Myeloma

Martin F. Kaiser, MD^{1,2} ; Pieter Sonneveld, MD³ ; David A. Cairns, PhD⁴ ; Marc S. Raab, MD⁵ ; Jesús San-Miguel Izquierdo, MD⁶ ; Rick Zhang, PhD⁷; Jorge Acosta, MD⁸; Alessandra Larocca, PhD⁹ ; Rakesh Popat, MD¹⁰ ; Cong Li, PhD¹¹; Marc-A. Baertsch, MD⁵ ; Sarah R. Brown, PhD² ; JuanJose Lahuerta Palacios, MD¹² ; Anita K. Gandhi, PhD¹³; Sandrine Macé, PhD¹⁴; Pellegrino Musto, MD¹⁵ ; Kwee Yong, PhD¹⁰ ; Elias K. Mai, MD⁵ ; Franck Dubin, PharmD¹⁴; Joan Blade, MD¹⁶; Andrea Capra, MScEng¹⁷; Gordon Cook, DSc^{4,18}; Uta Bertsch, MD⁵ ; María-Victoria Mateos, PhD¹⁹ ; Mario Boccadoro, MD²⁰ ; Graham H. Jackson, MD²¹ ; Norma C. Gutiérrez, MD²² ; Francesca Gay, PhD²³ ; and Niels Weinhold, PhD⁵ 

DOI <https://doi.org/10.1200/JCO-24-01253>

ABSTRACT

PURPOSE Survival for patients with multiple myeloma (MM) has improved but outcomes remain heterogeneous. Consistent diagnostic identification of high-risk disease is desirable to address unmet patient need. The aim was to investigate the consistency of association of co-occurrence of high-risk cytogenetic abnormalities (HRCAs) with prognosis in patients with newly diagnosed MM (NDMM) and relapsed/refractory MM (RRMM), and across a range of treatment modalities.

METHODS A systematic review of randomized controlled trials of MM that reported testing for HRCA between January 1, 2000, and December 9, 2021, was performed. Groups were contacted and asked to locally perform a novel, federated analysis of their data for single hit (one HRCA) and double hit (\geq two HRCAs), using a centrally provided algorithm. Analysis results were centrally collated and meta-analyzed to assess the hazard ratio (HR) for progression-free survival (PFS) and overall survival (OS) for one/ \geq two HRCAs across patient subgroups using random-effects models.

RESULTS Twenty-four trials including 13,926 patients were included. The median age of participants was 66.5 years (IQR, 59–72) and 56.5% were male (IQR, 52–60). The HR for PFS was 2.28 (95% CI, 2.05 to 2.54) for patients with \geq two HRCAs and 1.51 (95% CI, 1.38 to 1.65) for patients with one HRCA. The HR for OS was 2.94 (95% CI, 2.49 to 3.47) and 1.69 (95% CI, 1.52 to 1.88) for the two subgroups, respectively. In studies initiated since 2015, the effect abides (\geq two HRCA PFS, HR, 2.39 [95% CI, 1.96 to 2.91]; OS, 3.10 [95% CI, 2.10 to 4.60]) both for NDMM and RRMM. Heterogeneity related to transplant eligibility and relapsed/refractory status was as expected.

CONCLUSION The association of \geq two HRCAs with the poorest outcome in NDMM and RRMM, and across treatment modalities, as demonstrated here for the first time to our knowledge, allows for more focused development of novel approaches to these patients with high unmet need.

ACCOMPANYING CONTENT

 [Data Supplement](#)

Accepted January 14, 2025

Published February 18, 2025

J Clin Oncol 00:1-13

© 2025 by American Society of Clinical Oncology



[View Online Article](#)

Licensed under the Creative Commons Attribution 4.0 License

INTRODUCTION

Survival of patients with multiple myeloma (MM) has markedly improved over the past 20 years. However, outcomes remain highly heterogeneous with approximately 20% of patients experiencing a rapid relapse within the first 2–3 years after initiation of treatment, also termed high-risk MM (HRMM). The increasing availability of highly active

regimens, but also constraints to their access in public health care systems, makes consistent identification of patients with high-risk disease at diagnosis and relapse highly desirable: it holds the prospect of focusing on unmet patient needs within and outside of clinical trials.

Molecular risk reporting by the International Myeloma Working Group consensus¹ has so far been limited to the high-

CONTEXT

Key Objective

Consistent diagnostic identification of patients with high-risk multiple myeloma (HRMM) is important for standard-of-care treatment allocation and clinical research.

Knowledge Generated

In this federated analysis of clinical trials, \geq two high-risk chromosomal aberrations (HRCAs) were consistently associated with adverse outcome across newly diagnosed, transplant-eligible and transplant-ineligible, and relapsed MM. Strikingly, the prognostic impact of \geq two HRCAs was maintained in trials initiated in the past decade with modern combination therapies, highlighting the ongoing unmet need of HRMM.

Relevance (S. Lentzsch)

This study establishes \geq two HRCAs as a consistent diagnostic marker for HRMM, identifying patients with the poorest outcomes who require intensified treatment strategies. By providing a standardized framework for reporting HRMM in clinical trials, it enables better comparability of therapeutic outcomes and guides the development of targeted interventions for this high-risk group.*

*Relevance section written by JCO Associate Editor Suzanne Lentzsch, MD, PhD.

risk cytogenetic abnormalities (HRCAs) t(4;14), t(14;16), and del(17p), but recent research suggests genetic profiling including gain(1q) as prognostically relevant, subsequently termed extended genetic profiling. However, association of individual risk markers with outcome is relatively weak and has been inconsistent, suggesting limited certainty in identifying HRMM in trials and clinical care.² Recent evidence in newly diagnosed MM (NDMM) transplant-eligible (TE) patients suggests that co-occurrence of \geq two HRCAs (double hit, or multihit in some contexts) may more consistently identify patients with HRMM, while patients with one HRCA (single hit) may show less adverse and more variable outcome. There is currently a paucity of evidence regarding the prognostic impact of double hit across transplant-ineligible (TNE) and relapsed/refractory MM (RRMM) patients and a wider range of available treatment regimens.

Here, we addressed this uncertainty on the prognostic impact of double hit through identifying randomized, controlled phase II and phase III interventional clinical trials with genetic information, followed by a centrally coordinated federated analysis including in total 13,926 patients with TE and TNE NDMM, and RRMM and subsequent summary-level meta-analysis. In an academic-led initiative, we contacted study groups and industry collaborators who had reported extended genetic profiles from randomized trials to identify studies with at least a complete set of HRCA results including t(4;14), t(14;16), del(17p), and gain(1q) available in a substantial proportion of enrolled patients to ensure representativeness of results. In a federated analysis approach, participating collaborators were requested to perform analyses using uniform, prespecified methods for progression-free survival (PFS) and overall

survival (OS) for patient groups defined by the presence of double hit or single hit MM for each trial individually. Federated analysis of individual trials was chosen not only to provide best transparency on potential heterogeneity between data sets, but also to incentivize active participating and closer engagement of all parties with their existing data sets and potential value for systematic genetic stratification of patients.

METHODS

Selection Criteria and Search Strategy

For this centrally coordinated and quality controlled federated analysis and meta-analysis, any phase II or III randomized trials reported since 2000 that included PFS and/or OS outcomes and tested for co-occurrence of HRCA were included, for NDMM and RRMM settings. Studies that did not test for HRCA were excluded. The review was not registered.

Studies were identified by searching the bibliographical database PubMed. D.A.C. designed the PubMed search, which sought to identify clinical trials of treatment for MM that reported at least one HRCA to screen for studies that reported multiple HRCAs (Data Supplement, online only). The searches were run on December 9, 2021, and only English language studies were included. A publication date limit of January 1, 2000, to present was applied to all search results. The results of the database searches were downloaded to EndNote (version 20.4.1).

The initial screening (title and abstract) was conducted by one reviewer (D.A.C.). The studies that qualified from first

screen underwent a second full-text screen from two reviewers independently (D.A.C. and M.F.K.). Any questions regarding study eligibility were raised to the other reviewer and a consensus reached where possible with the default action being to exclude the study. At this stage, it was identified that data were rarely available in the required format. For studies where the analysis of PFS and OS by no hit (zero HRCA), single hit (one HRCA), and double hit (\geq two HRCAs) was not available in the published literature, we contacted the investigators or sponsors. A list of those contacted is given in the Data Supplement (Table S2). The data were recorded in Microsoft Excel (Microsoft Corporation, Redmond, WA) in a worksheet provided by the study statistician (Data Supplement).

Data Analysis

In a federated analysis approach, participating collaborators provided summaries and performed analyses following uniform, prespecified methods. Academic and industry collaborators provided results generated with these uniform methods. Study design, data specification, analysis and interpretation, and writing of the report were the responsibility of M.F.K., P.S., D.A.C., M.R., F.G., and N.W.

The following descriptive data were requested: trial name, trial registration number, trial patient population (NDMM TE, NDMM TNE, or RRMM), regimens used, patient characteristics in the overall trial population, and in the subset with complete genetic information, including number of patients, median and range of age (years), percentage male sex, and number in International Staging System (ISS) groups³ (I, II, III, or not available).

The following genetic data were extracted: the complete information to define no hit, single hit, or double hit including the number of patients with t(4;14), gain(1q) or amp(1q), and del(17p). If t(14;16)/MAF translocation testing was performed for all patients with complete genetic information, then this was also included in determining no/single/double hit status. Technical information related to the percentage cutoff for del(17p) positivity, 1q abnormality: how many copies counted as abnormal, and the detection method used was also requested.

Collaborators were requested to locally perform univariate Cox proportional hazard analyses for PFS and OS for patient groups defined by presence of double hit or single hit for each trial individually in a modified intention-to-treat population including those patients with a complete set of genetic results. The following outcome data were extracted for each subgroup defined by single hit or double hit: the estimated hazard ratio (HR) and corresponding 95% CI from Cox proportional hazards models comparing each group with no hit. The number of patients in each trial analysis can be understood by summing the columns no HRCA, one HRCA, and two HRCAs in Table 1. For example, in the model

comparing double hit with no hit, you sum no HRCA and two HRCAs. A Wald *P* value from these models for the linear predictor was also extracted. If possible, relevant estimates of median PFS and OS, and estimates at 24 months and 36 months, with corresponding 95% CI estimated using the Kaplan-Meier method were requested. Kaplan-Meier plots for visualization of results were also requested but were described as optional.

We collated results centrally and performed a meta-analysis using the HR of progression or death for PFS, or death for OS. As the trial populations treated were expected to be heterogeneous, we decided a priori to use a random-effects model using the Der Simonian and Laird method.⁴ Results were presented with forest plots. All analyses were performed using R (version 4.2.1).⁵

We also planned a priori to perform subgroup analysis, considering patients with TE NDMM, TNE NDMM, and RRMM in separate meta-analysis. We undertook sensitivity analysis focusing on studies commencing recruitment since 2015. The funnel plot and the Egger test were used to assess publication and availability bias.⁶ Heterogeneity was assessed with the *I*² statistic⁷ ($[Q - df]/Q \times 100$, where *Q* is the chi-square test for statistical heterogeneity, and *df* its associated degrees of freedom) and 50% or more was defined as substantial heterogeneity on the basis of the Cochrane Handbook for Systematic Reviews of Interventions.⁸

Role of the Funding Source

The funders of the study had no role in study design, data collection, data analysis, data interpretation, or writing of the report. Industry partners were invited collaborators, provided no funding, and had no role in study design or data interpretation.

RESULTS

Search and Data Extraction

The search identified 582 reports for screening (Fig 1). After screening, 33 reports were included⁹⁻⁴¹ that reported 20 randomized studies identified for further consideration (Data Supplement, Table S1). After contacting investigators and sponsors, nine of these studies were excluded (two sponsors declined to participate totaling eight studies, and one study did not have complete genetic information). However, contact with investigators and sponsors identified 15 further studies that were eligible (Data Supplement, Table S3). Hence, 24 studies comprising 13,926 patients were included.⁴²⁻⁶⁶ Two studies were included that contained pathways for TE and TNE NDMM: Medical Research Council-IX and National Cancer Research Institute (NCRI) XI. The study characteristics are provided in Table 1 and brief descriptions of the regimens investigated are provided in the Data Supplement (Table S4).

TABLE 1. Characteristics of the Studies Included in the Meta-Analysis

Study Name	Registration	Years Recruiting	Trial Population	Median Previous Lines	N _{ITT}	n _{HRCA}	Age, Years, Median (range)	Male (%)	ISS 1 (%)	ISS 2 (%)	ISS 3 (%)	No HRCA	One HRCA	Two HRCAs
MRC-IX	ISRCTN68454111	2003-2007	NDMM (TE)		1,111	511	59 (35-78)	64	23	37	33	266	172	73
HOVON-65/GMMG-HD4	ISRCTN64455289	2005-2008	NDMM (TE)		827	335	57 (25-65)	58	37	32	25	187	106	42
GEM2005MENOS65	ClinicalTrials.gov identifier: NCT00461747	2006-2009	NDMM (TE)		386	218	57 (32-95)	58	38	42	20	123	71	24
GMMG-MM5	ISRCTN05622749	2010-2012	NDMM (TE)		502	524	59 (32-70)	60	38	34	28	259	197	68
NCRI-XI	ClinicalTrials.gov identifier: NCT01554852	2010-2016	NDMM (TE)		2568	1,064	61 (28-75)	62	28	43	23	576	354	134
GEM2012	ClinicalTrials.gov identifier: NCT01916252	2013-2015	NDMM (TE)		458	359	58 (31-65)	52	37	28	29	169	144	46
FORTE	ClinicalTrials.gov identifier: NCT02203643	2015-2017	NDMM (TE)		474	403	57 (32-66)	55	48	35	18	171	144	88
GMMG-HD6	ClinicalTrials.gov identifier: NCT02495922	2015-2017	NDMM (TE)		559	459	59 (25-65)	60	42	37	21	228	153	78
Cardamon	ClinicalTrials.gov identifier: NCT02315716	2015-2019	NDMM (TE)		281	233	59 (34-74)	59	46	37	18	137	73	23
MRC-IX	ISRCTN68454111	2003-2007	NDMM (TNE)		849	358	73 (61-89)	57	11	36	43	181	120	57
GIMEMA-03-05	ClinicalTrials.gov identifier: NCT01063179	2006-2008	NDMM (TNE)		511	130	72 (57-87)	54	23	35	24	53	55	22
GEM2005MAS65	ClinicalTrials.gov identifier: NCT00443235	2006-2008	NDMM (TNE)		260	155	72 (65-84)	47	29	40	42	80	61	14
NCRI-XI	ClinicalTrials.gov identifier: NCT01554852	2010-2016	NDMM (TNE)		1,852	750	74 (56-89)	58	15	41	37	409	273	68
GEM2010MAS65	ClinicalTrials.gov identifier: NCT01237249	2011-2013	NDMM (TNE)		233	150	73 (65-88)	51	28	47	24	70	64	16
RV-MM-PI-0752	ClinicalTrials.gov identifier: NCT02215980	2014-2016	NDMM (TNE)		199	134	76 (67-80)	51	31	44	24	64	49	21
EMN10	ClinicalTrials.gov identifier: NCT02586038	2015-2018	NDMM (TNE)		175	139	74 (53-88)	48	27	46	27	64	56	19
GMMG-ReLApsE	ISRCTN16345835	2010-2016	RRMM	1	277	182	61 (29-74)	59	59	25	10	64	88	30
TOURMALINE-MM1	ClinicalTrials.gov identifier: NCT01564537	2012-2014	RRMM	1	722	529	66 (30-91)	56	63	24	13	236	219	74
MUKFive	ISRCTN17354232	2013-2016	RRMM	1	300	171	68 (40-82)	64	48	37	15	85	71	15
OPTIMISMM	ClinicalTrials.gov identifier: NCT01734928	2013-2017	RRMM	1+	559	350	65 (25-85)	55	51	31	6	166	130	54
MUKSeven	ISRCTN24593488	2016-2018	RRMM	3	102	71	72 (44-85)	65	32	31	37	29	33	9
MUKEight	ISRCTN58227268	2016-2018	RRMM	4	112	48	71 (50-80)	48	21	44	33	20	15	13
ICARIA	ClinicalTrials.gov identifier: NCT02990338	2017-2018	RRMM	2+	307	194	67 (36-86)	52	25	32	28	64	99	31
IKEMA	ClinicalTrials.gov identifier: NCT03275285	2017-2019	RRMM	2+	302	257	64 (33-90)	56	53	31	15	108	105	44

Abbreviations: EMN, European Myeloma Network; GEM, Grupo Español de Mieloma; GIMEMA, Gruppo Italiano Malattie Ematologiche Maligne dell'Adulto; GMMG, German Multicenter Myeloma Group; HOVON, Dutch-Belgian Hemato-Oncology Cooperative Group; HRCA, high-risk cytogenetic abnormality; ISS, International Staging System as assessed at trial entry; MM, multiple myeloma; MRC, Medical Research Council; MUK, Myeloma UK; NCRI, National Cancer Research Institute; NDMM, newly diagnosed MM; N_{HRCA}, number with complete data for HRCA; N_{ITT}, number in the intention-to-treat population; RRMM, relapsed/refractory MM; TE, transplant-eligible; TNE, transplant-ineligible.




FIG 1. Study selection. ^aTwo studies (MRC Myeloma IX and NCRI Myeloma XI were platform trials including pathways for transplant-eligible and transplant-ineligible NDMM). IMiD, immunomodulatory drug; MRC, Medical Research Council; NCRI, National Cancer Research Institute; NDMM, newly diagnosed multiple myeloma; PI, proteasome inhibitor.

Study Characteristics

In total, results from the federated analysis on HRCA co-occurrence were successfully provided on 7,724 patients (out of 13,926, 55.5%) enrolled in 24 trials and results as per prespecified patient subgroups were included in the meta-analysis: 4,106 patients from nine trials conducted in TE NDMM, 1,816 from seven trials conducted in TNE NDMM, and 1,802 from eight trials conducted in RRMM. The median age of participants was 66.5 years (IQR, 59–72) and 56.5% were male (IQR, 52–60). The percentage of patients with ISS-I, ISS-II, and ISS-III were 34.5% (IQR, 27–47), 37% (IQR, 34–41), and 24% (IQR, 18–30), respectively. Frequencies of double hit (median, 13.8%; IQR, 12.2–16.1) and single hit (median, 37.4%; IQR, 33.5–41.4) were comparable across trials.

Meta-Analysis

Meta-analysis of all studies showed highly consistent separation of risk groups by co-occurrence of HRCAs: PFS HR for double hit was 2.28 (95% CI, 2.05 to 2.54; $P < 10^{-49}$) and for single hit was 1.51 (95% CI, 1.38 to 1.65; $P < 10^{-18}$) compared with those without HRCA (Fig 2). The I^2 for PFS for single hit was 46.4% and for double hit was 33.8%. OS HRs were 2.94 (95% CI, 2.49 to 3.47; $P < 10^{-36}$) and 1.69 (95% CI,

1.52 to 1.86; $P < 10^{-21}$), respectively, compared with those without HRCA (Fig 3). The I^2 for OS was 35.4% for single hit and 56.5% for double hit, which is substantial by accepted definitions.⁷ However, 95% CIs did not overlap between double hit and single hit groups, providing evidence for consistent prognostic discrimination by number of HRCAs.

Considering clinical subgroups separately, double hit MM was consistently associated with the most adverse outcomes across patients with TE NDMM, TNE NDMM, and RRMM. The prognostic effect size was largest in patients with TE NDMM (PFS, 2.53 [95% CI, 2.26 to 2.84]; OS HR, 4.17 [95% CI, 3.34 to 5.22]), followed by patients with TNE NDMM (PFS, 1.97 [95% CI, 1.55 to 2.50]; OS HR, 2.31 [95% CI, 1.92 to 2.78]) and patients with RRMM (PFS, 2.05 [95% CI, 1.66 to 2.54]; OS HR, 2.21 [95% CI, 1.83 to 2.67]). In TE NDMM, the I^2 for PFS for single hit was 57.7% and double hit was 0.0%, and for OS for single hit was 36.1% and double hit was 47.3%. In TNE NDMM, the I^2 for PFS for single hit was 14.1% and double hit was 44.8%, and for OS for single hit was 0.0% and double hit was 0.0%. In RRMM, the I^2 for PFS for single hit was 44.8% and double hit was 39.5%, and for OS for single hit was 0.0% and double hit was 0.0%.

Considering only those studies commencing recruitment since 2015, including 2,312 patients (Data Supplement, Table




FIG 2. Random-effects meta-analysis of all studies considering PFS, by patient group and overall. EMN, European Myeloma Network; GEM, Grupo Español de Mieloma; GIMEMA, Gruppo Italiano Malattie Ematologiche Maligne dell'Adulto; GMMG, German Multicenter Myeloma Group; HOVON, Dutch-Belgian Hemato-Oncology Cooperative Group; HR, hazard ratio; MRC, Medical Research Council; MM, multiple myeloma; MUK, Myeloma UK; NCRI, National Cancer Research Institute; NDMM, newly diagnosed MM; PFS, progression-free survival; RE, random-effects; RRMM, relapsed/refractory MM; TE, transplant-eligible; TNE, transplant-ineligible.




FIG 3. Random-effects meta-analysis of all studies considering OS, by patient group and overall. EMN, European Myeloma Network; GEM, Grupo Español de Mieloma; GIMEMA, Gruppo Italiano Malattie Ematologiche Maligne dell'Adulto; GMMG, German Multicenter Myeloma Group; HOVON, Dutch-Belgian Hemato-Oncology Cooperative Group; HR, hazard ratio; MM, multiple myeloma; MRC, Medical Research Council; MUK, Myeloma UK; NCRI, National Cancer Research Institute; NDMM, newly diagnosed MM; OS, overall survival; RE, random-effects; RRMM, relapsed/refractory MM; TE, transplant-eligible; TNE, transplant-ineligible.

Downloaded from ascopubs.org by Universit degli Studi di Torino on February 21, 2025 from 130.192.098.153
Copyright © 2025 American Society of Clinical Oncology. All rights reserved.

S4), double hit MM was consistently associated with the most adverse outcomes overall (PFS, 2.39 [95% CI, 1.96 to 2.91]; OS HR, 3.10 [95% CI, 2.10 to 4.60]) and across patients with NDMM (PFS, 2.62 [95% CI, 2.05 to 3.35]; OS HR, 4.81 [95% CI, 2.85 to 8.13]) and RRMM (PFS, 2.34 [95% CI, 1.66 to 3.30]; OS HR, 2.45 [95% CI, 1.61 to 3.73]).

Considering each study separately showed consistent separation of Kaplan–Meier estimated survivor functions by co-occurrence of HRCAs for PFS and OS in double hit and single hit for TE NDMM (Figs 4A and 5A; Data Supplement, Fig S1), TNE NDMM (Figs 4B and 5B; Data Supplement, Fig S2), and RRMM (Figs 4C and 5C; Data Supplement, Fig S3). Of note, the separation was consistently observed in studies considering combination of proteasome inhibitors and immunomodulatory agents such as FORTE for NDMM and OPTIMISM in RRMM, as well as anti-CD38 monoclonal antibody combination therapy such as ICARIA.

Risk of Bias

In terms of publication and availability bias, the funnel plots for PFS (Data Supplement, Fig S4) did not indicate bias, which was confirmed by Egger's test for single hit ($z = 1.5718$, $P = .1160$) and double hit ($z = -0.0240$, $P = .9809$). Findings were similar for TE NDMM (single hit: $z = 1.0828$, $P = .2789$; double hit: $z = 1.8265$, $P = .0678$), TNE NDMM (single hit: $z = 0.5595$, $P = .5758$; double hit: $z = -0.4292$, $P = .6678$), and RRMM (single hit: $z = 0.6195$, $P = .5356$; double hit: $z = 1.4207$, $P = .1554$). There was no evidence of asymmetry in the funnel plot that might indicate small-study effect, but NCRI-XI was an example of a large study that provided a smaller effect size than might be expected, particularly in the TE pathway patients. The funnel plots for OS (Data Supplement, Fig S5) did not indicate bias, which was confirmed by Egger's test for single hit ($z = 1.5988$, $P = .1099$) and double hit ($z = -0.3667$, $P = .7139$). Findings were similar for TE NDMM (single hit: $z = 1.4946$, $P = .1350$), TNE NDMM (single hit: $z = 0.1835$, $P = .8544$; double hit: $z = -1.0793$, $P = .2805$), and RRMM (single hit: $z = 0.7654$, $P = .4440$; double hit: $z = 0.5380$, $P = .5906$). However, for TE NDMM double hit, there was evidence of asymmetry supported by Egger's test (double hit: $z = 1.9754$, $P = .0482$). This might indicate smaller-study effects, particularly notable in GEM2012 and FORTE.

DISCUSSION

Over the past 20 years, evidence has accumulated that certain cytogenetic abnormalities, including del(17p), translocation t(4;14), translocation t(14;16), and gain(1q), are associated with poor prognosis in patients with MM. However, despite the considerable data available from trials and real-world experience, the association between outcome and co-occurrence of cytogenetic abnormalities in different patient groups remains unclear. Our analysis, which included 24 randomized controlled trials comprising 13,926 patients, addresses this question. We show that co-occurrence of two

or more cytogenetic abnormalities confers significant poor prognosis in patients with NDMM that were TE or TNE, and in patients with RRMM.

Prognostic association was consistent across trials investigating immunomodulatory, proteasome inhibitor, or anti-CD38 monoclonal antibody therapies and combinations. Effect sizes were consistent in trials commencing since 2015, but among the highest for the most contemporaneous trials such as FORTE (OS HR, 4.29 [95% CI, 2.11 to 8.73]) for TE NDMM, and OPTIMISM (OS HR, 2.71 [95% CI, 1.76 to 4.17]) and ICARIA (OS HR, 2.50 [95% CI, 1.49 to 4.19]) for RRMM. This may suggest a proportionately higher benefit from current regimens for patients with standard risk and highlights the ongoing unmet need for patients with more aggressive disease, despite the marked improvement of therapies. We observed mildly higher heterogeneity in RRMM trials, which inherently include clinically diverse patient populations; the relevance of co-occurrence of cytogenetic lesions should accordingly be contextualized in these trials with other important prognostic factors, including, but not limited to, exposure and refractoriness to previous treatments. Relatively lower effect sizes in TNE NDMM would be consistent with the previously described higher frequency and impact of frailty and comorbidity on outcome in these patients,³⁵ although our study was not designed to investigate these further.

Strengths of our study include the diversity of trials, treatments, and patient populations included in the meta-analysis. We intentionally did not contrast individual treatment arms of the randomized trials, which could incentivize scientifically inappropriate cross-trial comparisons, but designed the study to generate evidence on the usefulness of extended genetic profiling and the co-occurrence of HRCAs. Participating groups rated the novel, federated analysis approach highly, which allowed all teams to gain local expertise and novel insight into their own data, while still contributing to a collaborative aim. A limitation of our study is the absence of trials that include frontline use of anti-CD38 in TE NDMM in the context of triplet and quadruplet combinations. It would also be of interest to include studies that have compared transplant and nontransplant strategies in more depth. However, it is likely that double hit retains its negative prognostic association with PFS and OS, while the impact on single hit is not clear, as demonstrated recently in studies ineligible (nonrandomized)⁶⁷ or unavailable to this meta-analysis (sponsor declined participation).⁶⁸

Our analysis did not set out to incorporate nongenetic prognostic factors such as beta-2-microglobulin, or next-generation sequencing genomic data, nor did it aim to use such information to build a more complex classifier, such as those recently proposed.⁶⁹⁻⁷¹ The analysis did not aim to exclusively and definitively define high risk on co-occurrence of markers only; single genetic lesions can have marked prognostic impact, including those resulting in




FIG 4. Kaplan-Meier estimates of survivor function for PFS separated by co-occurrence of cytogenetic abnormality for representative trials in (A) NDMM TE—HOVON-65/GMMG-HD4, (B) NDMM TNE—NCRI Myeloma XI, and (C) RRMM—OPTIMISMM. GMMG, German Multicenter Myeloma Group; HOVON, Dutch-Belgian Hemato-Oncology Cooperative Group; MM, multiple myeloma; NCRI, National Cancer Research Institute; NDMM, newly diagnosed MM; PFS, progression-free survival; RRMM, relapsed/refractory MM; TE, transplant-eligible; TNE, transplant-ineligible.

homozygous inactivation of tumor suppressor genes such as TP53.^{72,73} However, these events are relatively rare, and they have not yet been consistently tested and reported across international NDMM and RRMM trials. We did not consider

the type of HRCA, but rather any two or more HRCA. It is possible that additional HRCAs such as del(1p) should be considered as well, and that some combinations of specific HRCAs are worse than others. Furthermore, it would be




FIG 5. Kaplan-Meier estimates of survivor function for OS separated by co-occurrence of cytogenetic abnormality for representative trials in (A) TE—HOVON-65/GMMG-HD4, (B) TNE—NCRI Myeloma XI, and (C) RRMM—OPTIMISMM. GMMG, German Multicenter Myeloma Group; HOVON, Dutch-Belgian Hemato-Oncology Cooperative Group; NCRI, National Cancer Research Institute; OS, overall survival; TE, transplant-eligible; TNE, transplant-ineligible; RRMM, relapsed/refractory multiple myeloma.

interesting to understand the impact of lesions acquired in the NDMM to RRMM transition as we explored in a longitudinal study and found a clear added prognostic impact of lesions acquired or detectable for the first time at relapse.³⁷ Studies investigating this in more detail are eagerly awaited.

The analysis did aim to evaluate the enhanced utility of combining a set of established, accessible genetic markers across current treatment modalities and indications. We believe that our results strongly support wider patient access to these tests, both within and outside of clinical trials. These results accordingly also form a basis and framework for

future detailed and delineated exploration of single hit disease.

The heterogeneity observed in this study reflects the diversity of trial participants and is therefore a strength of our findings. The I^2 statistic is a measure of the impact of heterogeneity on the summary effect estimate. In each of the subgroup meta-analyses undertaken in this report, I^2 is greater than zero and, in the TE NDMM double hit analysis, it can be classified as substantial. However, when considering the clinical subgroups separately, I^2 for PFS and OS for each of single hit and double hit was <50%. This indicates that the observed heterogeneity is due to the diversity of patients in

each of these subgroups and the estimates of prognostic association should be considered alongside the estimates of heterogeneity. As a corollary to this, the estimates in each of the patient subgroups are not affected as strongly by heterogeneity, and hence can be considered as such.

In conclusion, co-occurrence of cytogenetic abnormalities is associated with poor prognosis in patients with MM. This association is consistent in key patient subgroups and across widely accessible treatment modalities, across TE NDMM and RRMM, supporting improved access to testing and reporting of risk marker co-occurrence across patient groups.

AFFILIATIONS

¹The Royal Marsden Hospital NHS Foundation Trust, London, United Kingdom

²Myeloma Molecular Therapy Group, The Institute of Cancer Research, London, United Kingdom

³Department of Hematology, Erasmus University Medical Center, Rotterdam, the Netherlands

⁴Leeds Cancer Research UK Clinical Trials Unit, University of Leeds, Leeds, United Kingdom

⁵Internal Medicine V, Hematology, Oncology and Rheumatology, Heidelberg University Hospital, Heidelberg, Germany

⁶Clinica Universidad de Navarra, Pamplona, Spain

⁷Sanofi, Cambridge, MA

⁸Clinical Development Haematology, Celgene International Sàrl, a Bristol Myers Squibb Company, Boudry, Switzerland

⁹Division of Hematology, Department of Molecular Biotechnology and Health Sciences, University of Torino, Torino, Italy

¹⁰Department of Haematology, University College London, London, United Kingdom

¹¹Takeda Development Center Americas, Inc, Deerfield, IL

¹²Instituto de Investigación Hospital Universitario 12 de Octubre, Madrid, Spain

¹³Hematology Translational Medicine, Bristol Myers Squibb, Summit, NJ

¹⁴Sanofi, Vitry-sur-Seine, France

¹⁵Department of Precision and Regenerative Medicine and Ionian Area, "Aldo Moro" University School of Medicine, Unit of Hematology and Stem Cell Transplantation, AOUC Policlinico, Bari, Italy

¹⁶Hematology Department, Hospital Clinic, Institut d'Investigacions Biomèdiques August Pi I Sunyer (IDIBAPS), Barcelona, Spain

¹⁷European Myeloma Network Trial Office, Torino, Italy

¹⁸Leeds Cancer Centre, St James's University Hospital, Leeds, United Kingdom

¹⁹Haematology Department, University of Salamanca, Salamanca, Spain

²⁰Department of Molecular Biotechnology and Health Sciences, University of Torino, Torino, Italy

²¹Department of Haematology, University of Newcastle, Newcastle upon Tyne, United Kingdom

²²Department of Hematology, University Hospital of Salamanca, IBSAL, Cancer Research Center-IBMCC (USAL-CSIC), CIBERONC, Salamanca, Spain

²³Division of Hematology 1, AOU Città della Salute e della Scienza, Department of Molecular Biotechnology and Health Sciences, University of Torino, Torino, Italy

CORRESPONDING AUTHOR

Martin F. Kaiser, MD; Twitter: @MyMKaiser; e-mail: martin.kaiser@icr.ac.uk.

DISCLAIMER

The views expressed are those of the author(s) and not necessarily those of the NHS, the NIHR, or the Department of Health and Social Care.

EQUAL CONTRIBUTION

N.C.G., F.G., and N.W. contributed equally.

PRIOR PRESENTATION

Presented in part at the American Society of Hematology 64th Annual Meeting and Exposition, New Orleans, LO, December 10-13, 2022.

SUPPORT

Supported in part by a Jacquelin Forbes-Nixon Fellowship (to M.F.K.) via Myeloma UK and by Core Clinical Trials Unit Infrastructure from Cancer Research UK (to D.A.C.; C7852-A25447). M.F.K. was supported in part by the National Institute of Health and Care Research (NIHR) Biomedical Research Center (BRC) at the Royal Marsden Hospital and the Institute of Cancer Research. D.A.C. and G.C. are supported in part by the NIHR Leeds BRC (NIHR203331). R.P. is supported by the NIHR University College London Hospitals BRC.

AUTHORS' DISCLOSURES OF POTENTIAL CONFLICTS OF INTEREST

Disclosures provided by the authors are available with this article at DOI <https://doi.org/10.1200/JCO-24-01253>.

DATA SHARING STATEMENT

All data included in this manuscript are available in tables, figures, and the Data Supplement. Code used to undertake this analysis is available from the authors upon request.

AUTHOR CONTRIBUTIONS

Conception and design: Martin F. Kaiser, David A. Cairns, Jesús San-Miguel Izquierdo, JuanJose Lahuerta Palacios, Pellegrino Musto, Elias

K. Mai, María-Victoria Mateos, Graham H. Jackson, Francesca Gay, Niels Weinhold

Administrative support: Martin F. Kaiser

Provision of study materials or patients: Martin F. Kaiser, Marc S. Raab, Jorge Acosta, Alessandra Larocca, Rakesh Popat, JuanJose Lahuerta Palacios, Elias K. Mai, Joan Blade, María-Victoria Mateos, Graham H. Jackson, Francesca Gay

Collection and assembly of data: Martin F. Kaiser, Pieter Sonneveld, David A. Cairns, Marc S. Raab, Jesús San-Miguel Izquierdo, Jorge Acosta, Rakesh Popat, Cong Li, Marc-A. Baertsch, Sarah R. Brown, Anita K. Gandhi, Sandrine Macé, Kwee Yong, Elias K. Mai, Franck Dubin, Joan Blade, Andrea Capra, Uta Bertsch, María-Victoria Mateos, Graham H. Jackson, Norma C. Gutiérrez, Niels Weinhold

Data analysis and interpretation: Martin F. Kaiser, Pieter Sonneveld, David A. Cairns, Jesús San-Miguel Izquierdo, Rick Zhang, Jorge Acosta,

Alessandra Larocca, Cong Li, Anita K. Gandhi, Sandrine Macé, Elias K. Mai, Franck Dubin, Gordon Cook, María-Victoria Mateos, Mario Boccadoro, Graham H. Jackson, Norma C. Gutiérrez, Francesca Gay, Niels Weinhold

Manuscript writing: All authors

Final approval of manuscript: All authors

Accountable for all aspects of the work: All authors

ACKNOWLEDGMENT

The authors thank all participating study groups, investigators and centers and the participating patients and their families.

REFERENCES

- Sonneveld P, Avet-Loiseau H, Lonial S, et al: Treatment of multiple myeloma with high-risk cytogenetics: A consensus of the International Myeloma Working Group. *Blood* 127:2955-2962, 2016
- Weinhold N, Heuck CJ, Rosenthal A, et al: Clinical value of molecular subtyping multiple myeloma using gene expression profiling. *Leukemia* 30:423-430, 2016
- Greipp PR, Miguel JS, Durie BGM, et al: International Staging System for multiple myeloma. *J Clin Oncol* 23:3412-3420, 2005
- DerSimonian R, Laird N: Meta-analysis in clinical trials. *Control Clin Trials* 7:177-188, 1986
- R Core Team: R: A Language and Environment for Statistical Computing. Vienna, Austria, R Foundation for Statistical Computing, 2022
- Egger M, Smith GD, Schneider M, et al: Bias in meta-analysis detected by a simple, graphical test. *BMJ* 315:629-634, 1997
- Higgins JP, Thompson SG, Deeks JJ, et al: Measuring inconsistency in meta-analyses. *BMJ* 327:557-560, 2003
- Higgins J, Thomas J, Chandler J, et al (eds): *Cochrane Handbook for Systematic Reviews of Interventions*. Cochrane, 2023. <https://training.cochrane.org/handbook>
- Avet-Loiseau H, Attal M, Moreau P, et al: Genetic abnormalities and survival in multiple myeloma: the experience of the Intergroupe Francophone du Myélome. *Blood* 109:3489-3495, 2007
- Moreau P, Attal M, Garban F, et al: Heterogeneity of t(4;14) in multiple myeloma. Long-term follow-up of 100 cases treated with tandem transplantation in IFM99 trials. *Leukemia* 21:2020-2024, 2007
- Harousseau JL, Avet-Loiseau H, Attal M, et al: Achievement of at least very good partial response is a simple and robust prognostic factor in patients with multiple myeloma treated with high-dose therapy: Long-term analysis of the IFM 99-02 and 99-04 trials. *J Clin Oncol* 27:5720-5726, 2009
- Neben K, Jauch A, Bertsch U, et al: Combining information regarding chromosomal aberrations t(4;14) and del(17p13) with the International Staging System classification allows stratification of myeloma patients undergoing autologous stem cell transplantation. *Haematologica* 95:1150-1157, 2010
- Boyd KD, Ross FM, Chiecchio L, et al: Gender disparities in the tumor genetics and clinical outcome of multiple myeloma. *Cancer Epidemiol Biomarkers Prev* 20:1703-1707, 2011
- Boyd KD, Ross FM, Tapper WJ, et al: The clinical impact and molecular biology of del(17p) in multiple myeloma treated with conventional or thalidomide-based therapy. *Genes Chromosomes Cancer* 50:765-774, 2011
- Mateos MV, Gutiérrez NC, Martín-Ramos ML, et al: Outcome according to cytogenetic abnormalities and DNA ploidy in myeloma patients receiving short induction with weekly bortezomib followed by maintenance. *Blood* 118:4547-4553, 2011
- Avet-Loiseau H, Attal M, Campion L, et al: Long-term analysis of the IFM 99 trials for myeloma: cytogenetic abnormalities [t(4;14), del(17p), 1q gains] play a major role in defining long-term survival. *J Clin Oncol* 30:1949-1952, 2012
- Boyd KD, Ross FM, Chiecchio L, et al: A novel prognostic model in myeloma based on co-segregating adverse FISH lesions and the ISS: Analysis of patients treated in the MRC Myeloma IX trial. *Leukemia* 26:349-355, 2012
- Ho PJ, Brown RD, Spencer A, et al: Thalidomide consolidation improves progression-free survival in myeloma with normal but not up-regulated expression of fibroblast growth factor receptor 3: Analysis from the Australasian Leukaemia and Lymphoma Group MM6 clinical trial. *Leuk Lymphoma* 53:1728-1734, 2012
- Nemec P, Zemanova Z, Kuglik P, et al: Complex karyotype and translocation t(4;14) define patients with high-risk newly diagnosed multiple myeloma: Results of CMG2002 trial. *Leuk Lymphoma* 53:920-927, 2012
- Walker BA, Wardell CP, Melchor L, et al: Intraclonal heterogeneity and distinct molecular mechanisms characterize the development of t(4;14) and t(11;14) myeloma. *Blood* 120:1077-1086, 2012
- Moreau P, Cavo M, Sonneveld P, et al: Combination of international scoring system 3, high lactate dehydrogenase, and t(4;14) and/or del(17p) identifies patients with multiple myeloma (MM) treated with front-line autologous stem-cell transplantation at high risk of early MM progression-related death. *J Clin Oncol* 32:2173-2180, 2014
- Kuiper R, van Duin M, van Vliet MH, et al: Prediction of high- and low-risk multiple myeloma based on gene expression and the International Staging System. *Blood* 126:1996-2004, 2015
- Leleu X, Karlin L, Macro M, et al: Pomalidomide plus low-dose dexamethasone in multiple myeloma with deletion 17p and/or translocation (4;14): IFM 2010-02 trial results. *Blood* 125:1411-1417, 2015
- Pawlyn C, Melchor L, Murison A, et al: Coexistent hyperdiploidy does not abrogate poor prognosis in myeloma with adverse cytogenetics and may precede IGH translocations. *Blood* 125:831-840, 2015
- Avet-Loiseau H, Fonseca R, Siegel D, et al: Carfilzomib significantly improves the progression-free survival of high-risk patients in multiple myeloma. *Blood* 128:1174-1180, 2016
- Wu SP, Pfeiffer RM, Ahn IE, et al: Impact of genes highly correlated with MMSET myeloma on the survival of non-MMSET myeloma patients. *Clin Cancer Res* 22:4039-4044, 2016
- Avet-Loiseau H, Bahlis NJ, Chng WJ, et al: Ixazomib significantly prolongs progression-free survival in high-risk relapsed/refractory myeloma patients. *Blood* 130:2610-2618, 2017
- GroΔ JP, Nattenmüller J, Hemmer S, et al: Body fat composition as predictive factor for treatment response in patients with newly diagnosed multiple myeloma—Subgroup analysis of the prospective GMMG MM5 trial. *Oncotarget* 8:58460-58471, 2017
- Shah V, Sherborne AL, Walker BA, et al: Prediction of outcome in newly diagnosed myeloma: A meta-analysis of the molecular profiles of 1905 trial patients. *Leukemia* 32:102-110, 2018
- Cook G, Royle KL, O'Connor S, et al: The impact of cytogenetics on duration of response and overall survival in patients with relapsed multiple myeloma (long-term follow-up results from BSBMT/UKMF Myeloma X Relapse [Intensive]): A randomised, open-label, phase 3 trial. *Br J Haematol* 185:450-467, 2019
- Perrot A, Lauwers-Cances V, Tournay E, et al: Development and validation of a cytogenetic prognostic index predicting survival in multiple myeloma. *J Clin Oncol* 37:1657-1665, 2019
- Dash AB, Zhang J, Shen L, et al: Clinical benefit of ixazomib plus lenalidomide-dexamethasone in myeloma patients with non-canonical NF-κB pathway activation. *Eur J Haematol* 105:274-285, 2020
- Kaufman JL, Dimopoulos MA, White D, et al: Daratumumab, lenalidomide, and dexamethasone in relapsed/refractory myeloma: A cytogenetic subgroup analysis of POLLUX. *Blood Cancer J* 10:111, 2020
- Larocca A, Mina R, Offidani M, et al: First-line therapy with either bortezomib-melphalan-prednisone or lenalidomide-dexamethasone followed by lenalidomide for transplant-ineligible multiple myeloma patients: A pooled analysis of two randomized trials. *Haematologica* 105:1074-1080, 2020
- Pawlyn C, Cairns D, Kaiser M, et al: The relative importance of factors predicting outcome for myeloma patients at different ages: Results from 3894 patients in the Myeloma XI trial. *Leukemia* 34:604-612, 2020
- Weisel K, Spencer A, Lentzsch S, et al: Daratumumab, bortezomib, and dexamethasone in relapsed or refractory multiple myeloma: Subgroup analysis of CASTOR based on cytogenetic risk. *J Hematol Oncol* 13:115, 2020
- Croft J, Ellis S, Sherborne AL, et al: Copy number evolution and its relationship with patient outcome—an analysis of 178 matched presentation-relapse tumor pairs from the Myeloma XI trial. *Leukemia* 35:2043-2053, 2021

38. Harrison SJ, Perrot A, Alegre A, et al: Subgroup analysis of ICARIA-MM study in relapsed/refractory multiple myeloma patients with high-risk cytogenetics. *Br J Haematol* 194:120-131, 2021
39. Richard S, Chari A, Delimpasi S, et al: Selinexor, bortezomib, and dexamethasone versus bortezomib and dexamethasone in previously treated multiple myeloma: Outcomes by cytogenetic risk. *Am J Hematol* 96:1120-1130, 2021
40. Usmani SZ, Hoering A, Ailawadhi S, et al: Bortezomib, lenalidomide, and dexamethasone with or without elotuzumab in patients with untreated, high-risk multiple myeloma (SWOG-1211): Primary analysis of a randomised, phase 2 trial. *Lancet Haematol* 8:e45-e54, 2021
41. Zaccaria GM, Bertamini L, Petrucci MT, et al: Development and validation of a simplified score to predict early relapse in newly diagnosed multiple myeloma in a pooled dataset of 2,190 patients. *Clin Cancer Res* 27:3695-3703, 2021
42. Mateos MV, Oriol A, Martinez-Lopez J, et al: Bortezomib, melphalan, and prednisone versus bortezomib, thalidomide, and prednisone as induction therapy followed by maintenance treatment with bortezomib and thalidomide versus bortezomib and prednisone in elderly patients with untreated multiple myeloma: A randomised trial. *Lancet Oncol* 11:934-941, 2010
43. Morgan GJ, Davies FE, Gregory WM, et al: Cyclophosphamide, thalidomide, and dexamethasone (CTD) as initial therapy for patients with multiple myeloma unsuitable for autologous transplantation. *Blood* 118:1231-1238, 2011
44. Morgan GJ, Davies FE, Gregory WM, et al: Cyclophosphamide, thalidomide, and dexamethasone as induction therapy for newly diagnosed multiple myeloma patients destined for autologous stem-cell transplantation: MRC Myeloma IX randomized trial results. *Haematologica* 97:442-450, 2012
45. Rosinol L, Oriol A, Teruel AI, et al: Superiority of bortezomib, thalidomide, and dexamethasone (VTD) as induction pretransplantation therapy in multiple myeloma: A randomized phase 3 PETHEMA/GEM study. *Blood* 120:1589-1596, 2012
46. Morabito F, Brinthen S, Larocca A, et al: Bortezomib, melphalan, prednisone (VMP) versus melphalan, prednisone, thalidomide (MPT) in elderly newly diagnosed multiple myeloma patients: A retrospective case-matched study. *Am J Hematol* 89:355-362, 2014
47. Palumbo A, Brinthen S, Larocca A, et al: Bortezomib-melphalan-prednisone-thalidomide followed by maintenance with bortezomib-thalidomide compared with bortezomib-melphalan-prednisone for initial treatment of multiple myeloma: Updated follow-up and improved survival. *J Clin Oncol* 32:634-640, 2014
48. Moreau P, Masszi T, Grzasko N, et al: Oral ixazomib, lenalidomide, and dexamethasone for multiple myeloma. *N Engl J Med* 374:1621-1634, 2016
49. Lahuerta JJ, Paiva B, Vidriales MB, et al: Depth of response in multiple myeloma: A pooled analysis of three PETHEMA/GEM clinical trials. *J Clin Oncol* 35:2900-2910, 2017
50. Goldschmidt H, Lokhorst HM, Mai EK, et al: Bortezomib before and after high-dose therapy in myeloma: Long-term results from the phase III HOVON-65/GMMG-HD4 trial. *Leukemia* 32:383-390, 2018
51. Attal M, Richardson PG, Rajkumar SV, et al: Isatuximab plus pomalidomide and low-dose dexamethasone versus pomalidomide and low-dose dexamethasone in patients with relapsed and refractory multiple myeloma (ICARIA-MM): A randomised, multicentre, open-label, phase 3 study. *Lancet* 394:2096-2107, 2019
52. Richardson PG, Oriol A, Beksac M, et al: Pomalidomide, bortezomib, and dexamethasone for patients with relapsed or refractory multiple myeloma previously treated with lenalidomide (OPTIMISMM): A randomised, open-label, phase 3 trial. *Lancet Oncol* 20:781-794, 2019
53. Rosinol L, Oriol A, Rios R, et al: Bortezomib, lenalidomide, and dexamethasone as induction therapy prior to autologous transplant in multiple myeloma. *Blood* 134:1337-1345, 2019
54. Goldschmidt H, Mai EK, Durig J, et al: Response-adapted lenalidomide maintenance in newly diagnosed myeloma: Results from the phase III GMMG-MM5 trial. *Leukemia* 34:1853-1865, 2020
55. Gay F, Musto P, Rota-Scalabrini D, et al: Carfilzomib with cyclophosphamide and dexamethasone or lenalidomide and dexamethasone plus autologous transplantation or carfilzomib plus lenalidomide and dexamethasone, followed by maintenance with carfilzomib plus lenalidomide or lenalidomide alone for patients with newly diagnosed multiple myeloma (FORTE): A randomised, open-label, phase 2 trial. *Lancet Oncol* 22:1705-1720, 2021
56. Goldschmidt H, Baertsch MA, Schlenzka J, et al: Salvage autologous transplant and lenalidomide maintenance vs. lenalidomide/dexamethasone for relapsed multiple myeloma: The randomized GMMG phase III trial ReLApsE. *Leukemia* 35:1134-1144, 2021
57. Jackson GH, Davies FE, Pawlyn C, et al: Lenalidomide before and after autologous stem cell transplantation for transplant-eligible patients of all ages in the randomized, phase III, Myeloma XI trial. *Haematologica* 106:1957-1967, 2021
58. Jackson GH, Pawlyn C, Cairns DA, et al: Carfilzomib, lenalidomide, dexamethasone, and cyclophosphamide (KRdc) as induction therapy for transplant-eligible, newly diagnosed multiple myeloma patients (Myeloma XI+): Interim analysis of an open-label randomised controlled trial. *PLoS Med* 18:e1003454, 2021
59. Jackson GH, Pawlyn C, Cairns DA, et al: Optimising the value of immunomodulatory drugs during induction and maintenance in transplant ineligible patients with newly diagnosed multiple myeloma: Results from Myeloma XI, a multicentre, open-label, randomised, phase III trial. *Br J Haematol* 192:853-868, 2021
60. Larocca A, Bonello F, Gaidano G, et al: Dose/schedule-adjusted Rd-R vs continuous Rd for elderly, intermediate-fit patients with newly diagnosed multiple myeloma. *Blood* 137:3027-3036, 2021
61. Mina R, Falcone AP, Brinthen S, et al: Ixazomib-based induction regimens plus ixazomib maintenance in transplant-ineligible, newly diagnosed multiple myeloma: The phase II, multi-arm, randomized UNITO-EMN10 trial. *Blood Cancer J* 11:197, 2021
62. Moreau P, Dimopoulos MA, Mikhael J, et al: Isatuximab, carfilzomib, and dexamethasone in relapsed multiple myeloma (IKEMA): A multicentre, open-label, randomised phase 3 trial. *Lancet* 397:2361-2371, 2021
63. Yong K, Hinsley S, Auner HW, et al: Carfilzomib or bortezomib in combination with cyclophosphamide and dexamethasone followed by carfilzomib maintenance for patients with multiple myeloma after one prior therapy: Results from a multicenter, phase II, randomized, controlled trial (MUKfive). *Haematologica* 106:2694-2706, 2021
64. Auner HW, Brown SR, Walker K, et al: Ixazomib with cyclophosphamide and dexamethasone in relapsed or refractory myeloma: MUKeight phase II randomised controlled trial results. *Blood Cancer J* 12:52, 2022
65. Yong K, Wilson W, de Tute RM, et al: Upfront autologous haematopoietic stem-cell transplantation versus carfilzomib-cyclophosphamide-dexamethasone consolidation with carfilzomib maintenance in patients with newly diagnosed multiple myeloma in England and Wales (CARDAMON): A randomised, phase 2, non-inferiority trial. *Lancet Haematol* 10:e93-e106, 2023
66. Mai EK, Goldschmidt H, Miah K, et al: Elotuzumab, lenalidomide, bortezomib, dexamethasone, and autologous haematopoietic stem-cell transplantation for newly diagnosed multiple myeloma (GMMG-HD6): Results from a randomised, phase 3 trial. *Lancet Haematol* 11:e101-e113, 2024
67. Costa LJ, Chhabra S, Medvedova E, et al: Minimal residual disease response-adapted therapy in newly diagnosed multiple myeloma (MASTER): Final report of the multicentre, single-arm, phase 2 trial. *Lancet Haematol* 10:e890-e901, 2023
68. Callander NS, Silbermann R, Kaufman JL, et al: Daratumumab-based quadruplet therapy for transplant-eligible newly diagnosed multiple myeloma with high cytogenetic risk. *Blood Cancer J* 14:69, 2024
69. D'Agostino M, Cairns DA, Lahuerta JJ, et al: Second Revision of the International Staging System (R2-ISS) for overall survival in multiple myeloma: A European Myeloma Network (EMN) report within the HARMONY project. *J Clin Oncol* 40:3406-3418, 2022
70. Maura F, Rajanna AR, Ziccheddu B, et al: Genomic classification and individualized prognosis in multiple myeloma. *J Clin Oncol* 42:1229-1240, 2024
71. Maura F, Boyle EM, Coffey D, et al: Genomic and immune signatures predict clinical outcome in newly diagnosed multiple myeloma treated with immunotherapy regimens. *Nat Cancer* 4:1660-1674, 2023
72. Shah V, Johnson DC, Sherborne AL, et al: Subclonal TP53 copy number is associated with prognosis in multiple myeloma. *Blood* 132:2465-2469, 2018
73. Corre J, Perrot A, Caillot D, et al: del(17p) without TP53 mutation confers a poor prognosis in intensively treated newly diagnosed patients with multiple myeloma. *Blood* 137:1192-1195, 2021

AUTHORS' DISCLOSURES OF POTENTIAL CONFLICTS OF INTEREST

Co-Occurrence of Cytogenetic Abnormalities and High-Risk Disease in Newly Diagnosed and Relapsed/Refractory Multiple Myeloma

The following represents disclosure information provided by authors of this manuscript. All relationships are considered compensated unless otherwise noted. Relationships are self-held unless noted. I = Immediate Family Member, Inst = My Institution. Relationships may not relate to the subject matter of this manuscript. For more information about ASCO's conflict of interest policy, please refer to www.asco.org/rwc or ascopubs.org/jco/authors/author-center.

Open Payments is a public database containing information reported by companies about payments made to US-licensed physicians ([Open Payments](#)).

Martin F. Kaiser

Honoraria: Janssen Oncology, Sanofi

Consulting or Advisory Role: Janssen Oncology, Bristol Myers Squibb, AbbVie, GlaxoSmithKline, Pfizer, Regeneron, Poolbeg, Sanofi, Roche

Research Funding: Celgene (Inst)

Pieter Sonneveld

Leadership: European Myeloma Network

Consulting or Advisory Role: Celgene, Janssen, Amgen, Karyopharm Therapeutics, CARsgen Therapeutics

Research Funding: Janssen (Inst), Amgen (Inst), Skyline Diagnostics (Inst), BMS (Inst)

David A. Cairns

Research Funding: Celgene (Inst), Merck Sharp & Dohme (Inst), Amgen (Inst), Takeda (Inst), Sanofi (Inst)

Travel, Accommodations, Expenses: Celgene

Marc S. Raab

Honoraria: AbbVie, Bristol Myers Squibb/Celgene, Janssen, Sanofi

Consulting or Advisory Role: Bristol Myers Squibb/Celgene (Inst), Amgen (Inst), GlaxoSmithKline (Inst), Janssen (Inst), Sanofi (Inst), Pfizer (Inst)

Research Funding: Bristol Myers Squibb/Celgene (Inst), Janssen (Inst), Sanofi (Inst)

Travel, Accommodations, Expenses: AbbVie, Bristol Myers Squibb/Celgene, Amgen, Janssen, Sanofi, Pfizer

Jesús San-Miguel Izquierdo

Consulting or Advisory Role: Amgen (Inst), Celgene (Inst), Takeda (Inst), Bristol Myers Squibb (Inst), MSD (Inst), Novartis (Inst), Sanofi (Inst), Janssen (Inst), Roche (Inst), AbbVie (Inst), GlaxoSmithKline (Inst), Karyopharm Therapeutics (Inst), Secura Bio (Inst), Regeneron (Inst), Haemalogix (Inst), Pfizer (Inst), Kite, a Gilead company (Inst), Dr Reddy's (Inst)

Rick Zhang

Stock and Other Ownership Interests: Sanofi

Jorge Acosta

Employment: Bristol Myers Squibb/Celgene

Stock and Other Ownership Interests: Bristol Myers Squibb/Celgene

Alessandra Larocca

Honoraria: Janssen-Cilag, Bristol Myers Squibb, Amgen, Takeda, Oncopeptides, GlaxoSmithKline, Sanofi, Karyopharm Therapeutics, Menarini

Consulting or Advisory Role: Janssen-Cilag, Bristol Myers Squibb, Amgen, Takeda, Oncopeptides, GlaxoSmithKline, Sanofi, Karyopharm Therapeutics, Menarini

Rakesh Popat

Honoraria: Bristol Myers Squibb/Celgene

Consulting or Advisory Role: GlaxoSmithKline, Roche, Janssen, Galapagos NV, Sanofi, Regeneron

Speakers' Bureau: GlaxoSmithKline, Johnson & Johnson/Janssen, Pfizer, Bristol Myers Squibb/Celgene, AbbVie

Research Funding: GlaxoSmithKline (Inst), Pfizer (Inst)

Travel, Accommodations, Expenses: Janssen, GlaxoSmithKline

Cong Li

Employment: Takeda

Stock and Other Ownership Interests: Takeda

Travel, Accommodations, Expenses: Takeda

Marc-A. Baertsch

Stock and Other Ownership Interests: Novartis

Research Funding: Sanofi

Travel, Accommodations, Expenses: Johnson & Johnson/Janssen

Sarah R. Brown

Research Funding: AstraZeneca (Inst), Janssen (Inst), Bristol Myers Squibb/Celgene (Inst), Adlai Nortye (Inst), GlaxoSmithKline (Inst)

JuanJose Lahuerta Palacios

Travel, Accommodations, Expenses: GlaxoSmithKline

Anita K. Gandhi

Employment: Bristol Myers Squibb/Celgene

Stock and Other Ownership Interests: Bristol Myers Squibb/Celgene

Patents, Royalties, Other Intellectual Property: biomarker for DLBCL

Sandrine Macé

Employment: Sanofi

Pellegrino Musto

Honoraria: Celgene/Bristol Myers Squibb, Janssen, Amgen, Sanofi, Takeda, GlaxoSmithKline

Kwee Yong

Honoraria: Sanofi, Pfizer

Consulting or Advisory Role: Janssen-Cilag

Speakers' Bureau: Sanofi, Johnson and Johnson

Research Funding: Autolus, Sanofi, Bristol Myers Squibb/Celgene

Elias K. Mai

Honoraria: Janssen, Takeda, Bristol Myers Squibb/Celgene, Sanofi, GlaxoSmithKline, Stemline Therapeutics, Oncopeptides
Consulting or Advisory Role: Janssen, Bristol Myers Squibb/Celgene, Takeda, Sanofi, GlaxoSmithKline, Stemline Therapeutics, Oncopeptides
Research Funding: Janssen, Bristol Myers Squibb/Celgene, Takeda, Sanofi, GlaxoSmithKline
Travel, Accommodations, Expenses: Janssen, Bristol Myers Squibb/Celgene, Takeda, GlaxoSmithKline, Sanofi, Stemline Therapeutics

Franck Dubin

Employment: Sanofi
Stock and Other Ownership Interests: Sanofi

Joan Blade

Honoraria: Janssen, Celgene, Amgen, Takeda, Oncopeptides

Gordon Cook

Honoraria: Takeda, Janssen-Cilag, Bristol Myers Squibb, Amgen
Consulting or Advisory Role: Janssen, Bristol Myers Squibb, Amgen, Takeda, Karyopharm Therapeutics, Oncopeptides
Speakers' Bureau: Takeda, Janssen-Cilag, Takeda
Research Funding: Takeda (Inst), Celgene (Inst), J&J Innovative Medicines (Inst)
Travel, Accommodations, Expenses: Takeda

María-Victoria Mateos

This author is a member of the *Journal of Clinical Oncology* Editorial Board. Journal policy recused the author from having any role in the peer review of this manuscript.
Honoraria: Janssen-Cilag, Celgene, Amgen, GlaxoSmithKline, AbbVie/Genentech, Sanofi, Pfizer
Consulting or Advisory Role: Takeda, Janssen-Cilag, Celgene, Amgen, AbbVie, GlaxoSmithKline, Pfizer, Regeneron, Roche/Genentech, Stemline Therapeutics, Kite, a Gilead company

Mario Boccadoro

Honoraria: Sanofi, Celgene, Amgen, Janssen, Novartis, Bristol Myers Squibb, AbbVie
Consulting or Advisory Role: Janssen, GlaxoSmithKline
Research Funding: Sanofi (Inst), Celgene (Inst), Amgen (Inst), Janssen (Inst), Novartis (Inst), Bristol Myers Squibb (Inst), Mundipharma (Inst)

Graham H. Jackson

Honoraria: J and J, Sanofi, BMS, Takeda, Amgen, Pfizer, Dr Reddy's Laboratories
Consulting or Advisory Role: Oncopeptides, Janssen Oncology, Sanofi, Celgene, Takeda, Amgen

Norma C. Gutiérrez

Honoraria: Amgen, Sanofi
Travel, Accommodations, Expenses: AbbVie, Kite, a Gilead company

Francesca Gay

Honoraria: Amgen, Celgene, Janssen, Takeda, Bristol Myers Squibb, AbbVie, GlaxoSmithKline, Sanofi
Consulting or Advisory Role: Amgen, Celgene, Janssen, Takeda, Bristol Myers Squibb, AbbVie, GlaxoSmithKline, Roche, Oncopeptides, Pfizer, Sanofi, AstraZeneca, Kite, a Gilead company

Niels Weinhold

Honoraria: GlaxoSmithKline
Consulting or Advisory Role: Sanofi
Research Funding: Celgene/Bristol Myers Squibb

No other potential conflicts of interest were reported.

Downloaded from ascopubs.org by Universit degli Studi di Torino on February 21, 2025 from 130.192.098.153
 Copyright © 2025 American Society of Clinical Oncology. All rights reserved.