AperTO - Archivio Istituzionale Open Access dell'Università di Torino

Interacting effects of physical environment and anthropogenic disturbances on the structure of European larch (*Larix decidua* Mill.) forests

This is the author's manuscript

Original Citation:

Availability:

This version is available <http://hdl.handle.net/2318/56356> since 2017-05-25T12:12:45Z

Published version:

DOI:10.1016/j.foreco.2008.12.031

Terms of use:

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)

This is the author's final version of the contribution published as:

[Garbarino M., Weisberg P.J., Motta R. 2009. Interacting effects of physical environment and anthropogenic disturbances on the structure of European larch (*Larix decidua* Mill.) forests. *Forest Ecology and Management* 2009. ISSN: 0378-1127, doi: 10.1016/j.foreco.2008.12.031]

The publisher's version is available at:

[<http://www.sciencedirect.com/science/article/pii/S0378112709000711>]

When citing, please refer to the published version.

Link to this full text:

[<http://hdl.handle.net/2318/56356>]

This full text was downloaded from iris-AperTO: <https://iris.unito.it/>

Interacting effects of physical environment and anthropogenic disturbances on the structure of European larch (*Larix decidua* Mill.) forests

Matteo Garbarino* (1), Peter J. Weisberg (2), Renzo Motta (1)

(1) Department of Agronomy, Forest and Land Management, University of Turin, Via L. da Vinci 44, I-10095 Grugliasco, TO, Italy;

(2) Department of Natural Resources and Environmental Science, University of Nevada, Reno 1000 Valley Road, Mail Stop 186, Reno, Nevada 89512, USA

* Corresponding author. Tel: +39 11 6705535; Fax: +39 11 6705556; Email: matteo.garbarino@unito.it

Abstract

Most forested landscapes have been strongly influenced by humans, and hence prediction of response to future perturbations or climate change requires understanding of the interaction among human influences and the abiotic environment. Environmental and anthropogenic influences on forest structure in *Larix decidua* Mill. stands were investigated in two watersheds of the central Italian Alps (Valtellina, SO). We related three data sets (forest structure, anthropogenic influences, and topography) using ordination methods. Path models of correlative and causal relationships between these sets of variables were developed and used to differentiate the two watersheds with respect to levels of historical human influence. The two study areas (Musella and Ventina) were characterized by strong climatic and topographic gradients as well as a long history of human settlement, although historical intensity of agricultural activities was much greater for Musella. We

hypothesized a weaker influence of abiotic variables on forest structure where the intensity of the primary human disturbance factors, forest thinning and cattle grazing, had been strongest. Stand structure types varied from sparse, larch-dominated forests at high elevations, to denser stands at lower elevations dominated by spruce. Correlations of environmental variables with dominant trends in stand structure were low with the exception of elevation. Anthropogenic variables were unimportant at Ventina, whereas the interactive effects of both anthropogenic and abiotic variables were needed to explain stand structure in Musella. The best-fit model indicated a negative effect of elevation and anthropogenic variables on overall tree size. Stands with greater tree density and canopy height layer diversity were located further from roads. Both watersheds were characterized by a strong dominance of larch stands, but their structure and spatial pattern differed greatly. Sparse larch forests were exclusively associated with moraines and upper elevations at Ventina, but were also common near low-elevation farms at Musella. Historical human influences were difficult to measure and may play a greater role in determining forest structure than was suggested. Our study emphasizes the importance of landscape context for interpreting the relative strengths of anthropogenic and abiotic influences on stand development pathways.

Key words

Stand structure, *Larix decidua*, Path analysis, anthropogenic disturbances, central Italian Alps.

1. Introduction

An important goal of ecology is to clarify causal relationships that determine plant species distribution and forest structure over broad spatial scales. A traditional approach to this topic, stemming from efforts of early biogeographers to map the world's biomes (von Humboldt, 1808;

Hawkins et al., 2003), is to consider the abiotic template (climate and topography) as the principal constraint acting on vegetation dynamics and plant successional processes (Curtis and McIntosh, 1951; Bray and Curtis, 1957; Stephenson, 1990; Urban et al., 2000). However, for most regions of the world anthropogenic influences play a critical role (Leduc et al., 1992). The resulting cultural landscapes are semi-natural systems developed as a result of the tight interaction between human traditions and the natural environment (Antrop, 1997; Naveh, 1982, 1995).

Such cultural landscapes have only recently become a focus of ecological study, despite their prevalence and importance. In Europe and North America, a disproportionate number of studies is conducted in areas that have only been minimally influenced by humans (Spies et al., 1990; Peterken, 1992, 1996; Bradshaw et al., 1994; Bergeron and Harvey, 1997), notwithstanding the difficulty in generalizing from such pristine study areas to the more prevalent condition (i.e. human-dominated landscapes). In the European Alps, human-induced disturbances have been profound and of long duration (c. 5000 years) (Stern, 1983; Carcaillet, 1998; Motta and Nola, 2001; Motta et al., 2002). In North America, there is a growing understanding that many if not most contemporary landscapes can only be understood in the context of historical human land use and disturbance, in some cases predating Euro-American settlement (White and Mladenoff, 1994; Hunter, 1996; Swetnam et al., 1999; Parshall et al., 2003). Native American influences were important for many forested landscapes, particularly with regard to vegetation management using fire in the western United States (Barrett and Arno, 1982; Baker, 2002; Whitlock and Knox, 2002; Hessburg and Agee, 2003). Although the time span of Euro-American impact has been relatively short (c. 200-400 years), many studies concerning cultural influences on ecological systems have since been conducted, particularly in the northeastern United States (Marsh, 1864; Foster, 1992; Turner and Meyer, 1993; Motzkin et al., 1999; Turner et al., 2001; Hall et al., 2002; Foster et al., 2003).

A decline in traditional agricultural practices, due to depopulation and marginalization of mountainous areas, is documented in many European countries (Baldock et al., 1996; MacDonald et al., 2000; Dullinger et al., 2003; Bätzing, 2003; Gellrich and Zimmermann, 2007). Marginalization is "a process driven by a combination of social, economic, political and environmental factors by which in certain areas farming ceases to be viable under an existing land use and socio-economic structure and no other agricultural options are available, so the process ends at land abandonment" (Baldock et al., 1996). This process has been particularly relevant in the southern side of the European Alps (Bätzing et al., 1996; Lehringer et al., 2003; Lingua et al., 2007). Since land abandonment, encroachment of tree and shrub species into former agricultural pastures has severely modified those landscapes that had been intensively managed. At high altitudes, in the subalpine belt of the European Alps, secondary succession following cessation of grazing is causing widespread changes for extensive areas of sparsely wooded "larch meadow" vegetation types (Piussi, 2000) that were traditionally grazed in the past (Didier, 2001; Maurer et al., 2006; Bolli et al., 2007).

It is well accepted that the abiotic template and natural and anthropogenic disturbances act together to constrain biological processes. Both human-dominated and biophysical systems exhibit complex nonlinear dynamics (Ellis and Swift, 1988; Sprugel, 1991; Wu and Loucks, 1995; Carpenter et al., 1999). The key factors that create vegetation mosaic structures in cultural landscapes are therefore a challenge to disentangle (Leduc et al., 1992) and generalizations have been elusive.

In this research we investigated the relative importance of natural and anthropogenic factors for shaping forest structure of two watersheds of the central Italian Alps, which represented different historical land use intensities. We conducted a landscape level analysis of the distribution of stand structural types, classified using hierarchical cluster analysis, in order to evaluate the relative

influences of anthropogenic disturbances and biophysical factors. Path analysis was used to elucidate direct and indirect effects of anthropogenic and abiotic influences on forest structure. We expected a weaker influence of abiotic variables on forest structure where the intensity of the primary human disturbance factors, forest thinning and cattle grazing, had been strongest.

2. Methods

2.1. Study area

The research area consists of two watersheds of the upper Val Malenco, an inner valley of Valtellina (Central Alps, Lombardy, Italy). The Musella study area occupies 1150 ha in the eastern Malenco valley (45° 27' N; 28° 41' E) and elevation ranges from 1650 m a.s.l. and 3050 m a.s.l. The Ventina study area occupies 1124 ha in the western Malenco valley (45° 26' N; 28° 33' E) with an elevation range from 1650 m a.s.l. to 3570 m a.s.l. (Figure 1).

Musella has 470 ha of forested area, Ventina 170 ha. Moraines and glaciers cover a majority of the Ventina watershed, which is steeper than Musella. The bedrock is silicate and serpentine is the predominant rock. Both study areas are inner valleys of the “endalpic district” (Del Favero, 2002) characterized by a continental climate. Annual precipitation from 1921 to 1990 has varied from 668 mm to 1551 mm, averaging 974.9 mm (Lanzada, 1000 m a.s.l.). In both catchments European larch (*Larix decidua* Mill.) is the dominant tree species with Norway spruce (*Picea abies* (L.) H.Karst), Swiss stone pine (*Pinus cembra* L.) and mountain pine (*Pinus mugo* subsp. *uncinata*) as co-dominant species throughout the subalpine zone. Two subalpine shrub species are locally abundant: dwarf mountain pine (*Pinus mugo* subsp. *mugo* Turra) and green alder (*Alnus viridis* (Chaix) D.C.) ones.

#Figure 1 approximately here#

Grazing activity existed in the area even before 1447 (Bergomi, 2006). Cattle grazing was commonly limited by stockyards within the alpine pastures, while goats were permitted to roam freely as long as they did not damage pastures. Grazing data from the beginning of twentieth century (Società Agraria di Lombardia, 1901) revealed that grazing pressure was much higher at Musella than Ventina: 4.1 and 1.1 LUha⁻¹ respectively (Livestock Units; i.e. 600 kg body weight according to BL/BUWAL, 1994). During the 1978-80 period, grazing declined in both areas (to 0.6 LUha⁻¹ at Musella and 0.5 LUha⁻¹ at Ventina). More recently, grazing activity has ended at Ventina (Della Marianna et al., 2004). Previously grazed open stands that have developed without grazing pressure for decades coexist with newly established forests on the glacial moraines. However, forests at Musella are still partially grazed with varying intensities. A substantial increase of livestock units (to 1.1 LUha⁻¹) in 2006 was likely a consequence of a recent restoration policy applied to mountain pastures throughout the Lombardy region. For this reason we expected a weaker influence of abiotic variables on forest structure at Musella, where anthropogenic disturbances are still intense.

2.2. Sampling design and data collection

High resolution (50-cm) color aerial orthoimages (Sondrio province, 2003) were photo-interpreted, and a stratified random sampling was applied to locate sample plots. An object-oriented image analysis, based on homogeneous patches and not on single pixels, was implemented in eCognition (v. 4.0) software to perform a segmentation of the aerial orthoimages (Benz et al., 2004; Weisberg et al., 2007). The “image objects” resulting from segmentation were then manually photo-interpreted. The segmentation process facilitated separation of homogeneous patches of forest from non-forest land uses. This binary classification was developed over a “patch” scale (scale parameter = 10; 0.8 ha mapping resolution) and a centroid from each forested patch was derived in a GIS environment.

Approximately seventy circular plots (40 for Ventina and 28 for Musella) were established by using the patch centroids as center points for sample plots. Plots of 12-m radius were used for the tree (DBH ≥ 5 cm) layer survey, and subplots with a radius of 6 m were established within each plot for the understory and sapling (DBH < 5 cm and height > 10 cm) layers. The following parameters were measured for all trees: diameter at 1.30 m, total height, crown length and crown radius projection to the ground in four directions. For saplings, only density, composition and height were collected. Cover and abundance of understory shrub species were recorded for each subplot. All trees were mapped and the three larches with the greatest diameter were cored upslope at a height of 50 cm in order to estimate stand age.

2.3. Stand descriptors

A combination of structural diversity indices and classical stand structure measures was used to classify different stand types (Table 1). Diversity and structural diversity measures included species richness, relative dominance of larch, diameter standard deviation, and vertical evenness (Neumann and Starlinger, 2000). An index of aggregation (Clark and Evans, 1954; Neumann and Starlinger, 2000) was used to measure the horizontal structure within each plot. Vertical structure was assessed with the Vertical Evenness Index that applies the Shannon evenness formula to the proportions of crown projection area of four height layers (Neumann and Starlinger, 2000).

#Table 1 approximately here#

2.4. Data analysis

Three data sets were used in this study: forest structure data collected in the field, anthropogenic variables derived from thematic maps, and topographic variables derived from a 10-m resolution digital elevation model (DEM). Proximities to man-made features (buildings and roads) were calculated in ArcGIS using Euclidean distances and considered proxy variables for human

pressure (Roath and Krueger, 1982, Nakamura et al., 2000, Kawamura et al., 2005). Buildings were classified as “malghe” (a local name for shepherd’s huts) and “other buildings.” A greater weight was assigned to malghe because they represent an important source of grazing disturbance. The roads category comprised a network of roads and trails data derived from maps or photo-interpreted. Both indices were linearly rescaled from 0 to 100 so as to obtain a comparable measurement. Topographic variables included elevation, aspect, slope steepness, solar radiation and snow index. Aspect as circular data (degrees) was transformed to linear data following a method based on the interaction of slope and aspect to indicate the relative solar insolation (Clark, 1990). The snow index was a snow redistribution measure influenced by separate exposure, elevation-slope and aspect effects (Baker and Weisberg, 1995). Solar radiation was a direct clear-sky short-wave radiation measurement based on latitude, season, time interval and a DEM (Kumar et al., 1997; Zimmermann, 2000). Each dataset was relativized by the standard deviate in order to put variables, that were measured in different units, on an equal footing (McCune and Grace, 2002).

A hierarchical cluster analysis was performed on the stand structure data, using Ward’s clustering method based on a Euclidean distance matrix. Sixty-eight sample plots were grouped according to similarity in stand structure using cluster dendrograms and a 50% threshold for total variance explained.

Principal components analysis (PCA) was used to explore the correlation structure of variables and to identify key factors underlying spatial patterns of stand structure variation. Two data matrices for each study site including eleven stand structure variables (Table 1) and seven topographic and anthropogenic variables (elevation, aspect, slope, solar radiation, snow index, distance from roads, distance from buildings) were processed using the statistical package PcOrd (McCune and Mefford, 1999). We used biplots to assess the correlation of environmental variables with the

underlying gradients of stand structure (PCA axes). Moreover PCA was used to extract two synthetic descriptors of stand structure then used as focus variables in the path analysis. Relationships between stand structure, environmental and anthropogenic sets of variables were analyzed using path analysis, a specialized version of Structural Equation Models (Shipley, 2000). With path analysis, the cause-and-effect relationships between the putative causal variables (environmental and anthropogenic) and the hypothesized effect variables (first two stand structure principal components) were tested (Leduc et al., 1992, Cuevas, 2003, Weisberg, 2004, Houle, 2007). This method permits modeling of both directly observed (manifest) and unmeasured (latent) variables. A graphical model is presented using diagrams where arrows symbolize cause-and-effect relationship between variables that are represented by rectangles (manifest) and ellipses (latent). Path analysis allowed us to quantify and graphically illustrate relative influences of topographic, climatic and anthropogenic variables on stand structure, which was our response variable. Several hypothetical models were built under the underlying concept that environmental and anthropogenic variables interact together in shaping forest structure (Figure 2). Alternative models considered subsets of the full model (Figure 2), including the interactive effects of the various combinations of variable groups, Topographic (T), Anthropogenic (A) and Climatic (C): T-C-A, T-A, T-C, T and A only.

#Figure 2 approximately here#

PCA was also employed to reduce the number of stand structure response variables to a smaller subset of integrative, synthetic variables for use in the path models. Quantitative model comparisons used a combination of Akaike's Information Criterion (AIC) statistic and the Root Mean Square Error of Approximation (RMSEA). The latter is a goodness-of-fit index that is relatively independent of sample size. A model with $RMSEA < 0.06$ was considered a good fit

(Hu and Bentler, 1999). All such models were computed and the models with the smallest AIC statistic were selected as the most parsimonious models (Hu and Bentler, 1999). Path analyses were conducted using Mx software that works with covariance matrices as input data and a maximum likelihood (ML) fit function (Neale, 1994).

3. Results

3.1. Multivariate Analyses of Forest Structure and its Environmental Relationships

The cluster analysis performed on the stand structure data set (11 stand descriptors) produced four structural types in each study area (Table 2). Stand structural types varied from sparse, larch-dominated forests with young trees at high elevations (Type 1), to denser stands at lower elevations where spruce is more common (Type 2). Ventina had slightly more sites belonging to Type 1 (28% vs. 21%), while Musella had more sites belonging to Type 3 (29% vs. 18%), mixed-species stands with large trees.

#Table2 approximately here#

PCA was used to relate stand structure to environmental and anthropogenic influences. A biplot of the first two components at Musella showed a strong negative correlation of elevation ($r = -0.74$) with the dominant axis, while distance from roads was weakly and positively correlated with the second axis ($r = 0.23$; Figure 3). The first and second principal components accounted for 40% and 18% of the total variation, respectively. The ordination of plots revealed a clear grouping related to clusters 1 (sparse larch dominated stands with small trees, at higher elevations) and 4 (dense, larch dominated stands with mean size trees, at lower elevations). A perpendicular position of the elevation vector relative to distance from roads indicated that these variables were uncorrelated. The first component (axis 1) reflected variations of diameter, basal area and tree height; the second

was related to density and vertical evenness (Table 3). Sites further from roads tended to have greater tree density and diversity of canopy height layers, although the statistical relationship was weak (Figure 3).

The PCA biplot of 40 plots established at the Ventina study area showed a strong influence of elevation on stand structure (Figure 4). The first and second principal components accounted for 33% and 24% of the total variance, respectively. As for the Musella study area, the first component represented diameter and tree height, while the second represented tree density, canopy cover and vertical evenness (Table 3). The first component was positively influenced by distance from roads and slope, and negatively influenced by elevation. The second component was negatively influenced by elevation and snow and positively influenced by distance to buildings.

The ordination of plots revealed a clear grouping related to cluster 1 (sparse larch dominated stands with small trees) exclusively. Such stands occurred at high-elevation, snowy sites.

#Figures 3, 4 and Table 3 approximately here#

3.2. Causal Models of Forest Structure

We tested alternative path models for two synthetic descriptors of stand structure derived from the PCAs: overall tree size (PC 1) and tree density and vertical complexity (PC 2). Four alternative models emerged as having significant support (Table 4); two each for the Musella and Ventina study areas, respectively. Models for tree size (PC 1) only are shown (Figure 5) because of their better predictive ability and similar behavior to the models based on the second principal component. The two models for Musella differed in that the first included only topographic and anthropogenic factors (Figure 5 – M1), while the second additionally included the explicit influences of climatic variables (Figure 5 – M2). Both models included the interaction of anthropogenic and abiotic influences in that the negative effect of site aspect on proximity to

buildings ($\beta = -0.31$) was explicitly represented. Elevation was strongly negatively associated with tree size in both models ($\beta = -0.83$ and $\beta = -0.90$). A weak negative effect on tree size was also observed for the anthropogenic variables (proximity to buildings and roads). Slope was weakly but positively associated ($\beta = 0.20$ and $\beta = 0.22$) with tree size. In the second model (Figure 5 – M2), a direct positive effect ($\beta = 0.23$) of solar radiation on tree size was slightly reduced by indirect effects mediated by snow depth (total effect = 0.15).

Significant models for Ventina did not include effects of anthropogenic variables. Only topographic variables contributed significantly to the first model (Figure 5 – V1). Elevation was the only variable strongly but negatively related to stand structure ($\beta = -0.42$). The second model for the Ventina site was more complex and included snow as a predictor variable. Snow depth was positively associated with tree size, and in turn mediated the effects of the various topographic variables on tree size, as each of these variables also influenced snow depth (Figure 5 – V2). For example, a direct negative effect ($\beta = -0.49$) of elevation on tree size was slightly reduced by the positive effect of increasing elevation on snow depth (total effect = -0.42).

Elevation had a similar influence on tree size in all the models tested, but it was particularly strong for the Musella study area (models M1 and M2). The anthropogenic variables had a similar, slightly negative effect on tree size in the two Musella models, but were not important predictors of stand structure for the Ventina study area. Aspect had weakly negative direct effects at Ventina, and negative indirect effects through “proximity to buildings” at Musella.

#Figures 5 – M1, M2, V1, V2 and Table 4 approximately here#

4. Discussion

Abiotic environmental gradients, described by simple topographic variables including elevation, slope, and aspect, proved to be good predictors for forest structure even in forests that are still

recovering from recent, intensive human influence. This is likely a result of the extreme environmental gradients that characterize these mountainous landscapes. Furthermore, the effects of the physical environment are largely independent of anthropogenic influences, except for the positive association between building density and aspect observed at the Musella site.

Correlations of environmental variables with components describing dominant trends in stand structure were relatively low, with the exception of elevation (Figure 6 and 7). This indicates that either the effect of elevation is so strong as to swamp other effects, or that the anthropogenic and abiotic variables measured were only weakly related to the true underlying gradients driving forest development processes. Tree size, and especially basal area, emerged as a good, integrative descriptor for forest structure and diversity as exemplified by many other studies (Neumann and Starlinger, 2000; Solomon and Gove, 1999; Staudhammer and LeMay, 2001; Houle, 2007).

#Figure 6 and 7 approximately here#

Predictor variables in vegetation modeling can be classified as representing indirect gradients (topographic variables) or direct or resource gradients (bioclimatic variables; Austin, 1980, 1985; Guisan and Zimmermann, 2000). Our models were constructed using mainly indirect gradients (topographic variables) rather than direct or resource gradients (bioclimatic variables). Use of such indirect predictors to infer complex combinations of resource and direct gradients generally favors model precision instead of generalizability (Guisan et al., 1999; Guisan and Zimmermann, 2000). Path models developed within the present research may not be directly applicable to other study areas and different climatic conditions. However, the general relationships described by the conceptual model should be valid. A certain level of misclassification and uncertainty of the model can be explained by the use of proxy or DEM-derived data (Tappeiner et al., 1998), particularly with regard to snow, which is only weakly predicted by topography (Tappeiner et al., 2001).

Despite the importance of abiotic factors in shaping forest structure, anthropogenic influences play a critical role for most regions of the world. For some areas in the Italian Alps, historical human influences are important for understanding current patterns of forest structure across mountain landscapes (Motta et al., 2006). In these areas culture-landscape interactions are multifaceted and it is essential to consider human impacts on the environment (Naveh, 1995).

The Ventina site has historically been less disturbed by human land uses. Therefore, it is not surprising that anthropogenic variables failed to emerge as important in the most parsimonious path models. Stand structure in this relatively pristine landscape of complex topographic gradients is best explained by abiotic variables. More complex models including the interactive effects of anthropogenic influences and abiotic factors were needed to explain stand structure in the Musella watershed. At Musella, stands that were likely to have experienced historically more intense anthropogenic disturbance currently have smaller trees and sparser, less diverse stand structures. Both watersheds are characterized by a strong dominance of larch stands, but their structure and spatial pattern differ greatly. At Musella, sparse stands are mainly located at higher elevations, but are also common near low-elevation farms. At Ventina, this pattern differs in that sparse forests are exclusively associated with moraines and upper elevations. Vegetation dynamics at Ventina represent primary succession in the path of the retreating glacier, whereas at Musella past and present human actions (grazing and unmanaged thinning) contribute to maintain a more open canopy and even-aged forest at lower and moderate elevations. Another indicator of the relative importance of anthropogenic disturbance in the two study areas is maximum larch age, which is estimated at greater than 1000 years at Ventina and less than 500 years at Musella (Nola, 1994; Nola and Motta, 1996). A high level of anthropogenic disturbance can also explain the scarcity of Swiss stone pine at Musella, where larch trees were likely favored by humans due to their ability to form sparse pastured woodlands. Larch has a light canopy and is resistant to grazing. The stone

337 pine was considered a competing species and an obstacle to traditional agroforestry practices.
338 Seedling removal of stone pines was a common practice in the Western Alps until the first half of
339 the twentieth century (Motta and Lingua, 2005; Motta et al., 2006).
340 Historical human influences are difficult to measure and may play a greater role in determining
341 forest structure than is suggested by these analyses. It would be valuable to develop accurate
342 spatial datasets for such human influences, including more explicit proxy variables for human
343 influences that are independent of abiotic, topographic variables. Spatially explicit data on native
344 and nonnative plant species diversity could be a useful integrated measure of anthropogenic
345 disturbance, especially cattle grazing (Dullinger et al., 2003; Vacher et al., 2007). Remote sensing
346 and GPS tracking could also be used to better understand cattle distribution and behavior for
347 assessment of grazing intensity (Kawamura et al., 2005). Historical data regarding cattle density
348 and their distribution in the pastures, as well as individual-based models of grazing animal
349 behavior (Dumont and Hill, 2001), could be used to estimate historical grazing intensity.
350 Unfortunately, long-term data on historical forest harvesting and density and spatial distribution of
351 cattle were not available for our study areas. For this reason, map-derived data (distances from
352 roads and buildings) were used as proxy indices of potential human impact on the forest structure
353 of the two study areas.
354 Path analysis provides an analytical tool that allowed us to partition the interacting factors
355 involved in shaping forest structure in our complex mountain region with a long history of human
356 impacts. Land abandonment is currently the main cause for forest increase in the Alps. However,
357 over the long term, the importance of climate change in large scale vegetation dynamics will
358 clearly increase because forests will fill anthropogenic gaps below and above the treeline (Walther,
359 1986; Dale, 1997; Gehrig-Fasel et al., 2007).

Our study highlights the importance of landscape context for modeling and interpreting influences of anthropogenic disturbances on forest structure. A model that is able to partition natural and anthropogenic effects on stand structure can provide a robust method for discriminating between anthropogenically disturbed sites and areas where natural disturbances and climate change are the primary driving forces. Such an approach can be informative for developing conservation priorities in relation to those cultural landscapes that are disappearing in many Alpine regions (Margules et al., 1994; Guisan and Zimmermann, 2000; Maurer et al., 2006). It is important to identify cultural landscapes (e.g. the *Larix* forest) in order to calibrate an integrated forest management strategy where some areas are actively restored and managed while others are merely monitored. Cultural landscapes in the Italian Alps are complex mosaics characterized by a greater diversity of forest structural types than may otherwise be found in less disturbed systems (MacDonald et al., 2000). However, the two types of landscapes with different levels of historical human influence can occur in close proximity to each other, as is the case for Ventina and Musella. Future research should attempt to identify and model the two kinds of areas *a priori*, as these are likely to exhibit different stand dynamics and different potential responses to global change.

Acknowledgments

We thank Emanuele Lingua for field assistance and Tiziana Stangoni from the Sondrio Province for providing aerial photography and logistical support. We also thank the Lombardy Region for funding this research through the “Programma regionale di ricerca in campo agricolo 2004-2006” program.

384 **References**

- 385 Antrop, M., 1997. The concept of traditional landscapes as a base for landscape evaluation and
386 planning. The example of Flanders Region. *Landscape and Urban Planning* 38, 105-117.
- 387 Austin, M.P., 1980. Searching for a model for use in vegetation analysis. *Vegetatio* 42, 11-21.
- 388 Austin, M.P., 1985. Continuum concept, ordination methods and niche theory. *Annual Reviews in*
389 *Ecology and Systematics* 16, 39-61.
- 390 Baker, W.L., Weisberg, P.J., 1995. Landscape analysis of the forest-tundra ecotone in Rocky
391 Mountain National Park, Colorado. *Professional Geographer* 47, 361-375.
- 392 Baker, W.L., 2002. Indians and fire in the US Rocky Mountains: the wilderness hypothesis
393 renewed. In: Vale, T.R. (Ed.), *Fire, Native Peoples, and the Natural Landscape*. Island Press,
394 Washington, pp. 41-76.
- 395 Baldock, D., Beaufoy, G., Brouwer, F., Godeschalk, F., 1996. Farming at the margins:
396 abandonment or redeployment of agricultural land in Europe. Institute for European
397 Environmental Policy, London/The Hague.
- 398 Barrett, S.W., Arno, S.F., 1982. Indian fires as an ecological influence in the Northern Rockies.
399 *Journal of Forestry* 80, 647-651.
- 400 Bätzing, W., Perlik, M., Dekleva, M., 1996. Urbanization and depopulation in the Alps: an
401 analysis of current social–economic structural changes. *Mountain Research and Development* 16,
402 335-350.
- 403 Bätzing, W., 2003. *Die Alpen. Geschichte und Zukunft einer europäischen Kulturlandschaft*.
404 Verlag C.H. Beck, München.
- 405 Benz, U.C., Hofmann, P., Willhauck, G., Lingenfelder, I., Heynen, M., 2004. Multi-resolution,
406 object-oriented fuzzy analysis of remote sensing data for GIS-ready information. *Journal of*
407 *Photogrammetry and Remote Sensing* 58, 239-258.

408 Bergeron, Y., Harvey, B., 1997. Basing silviculture on natural ecosystem dynamics: an approach
 409 applied to the southern boreal mixedwood forest of Quebec. *Forest Ecology and Management* 92,
 410 235-242.

411 Bergomi, M., 2006. *Politica e amministrazione in Val Malenco nell'Età Moderna*. Società Storica
 412 Valtellinese, Sondrio.

413 BL/BUWAL (Swiss Federal Office for Agriculture/for Environment Forest and Landscape), 1994.
 414 *Wegleitung für den Gewässerschutz in der Landwirtschaft*, LmZ, Berne, Switzerland, 100 pp.

415 Bolli, J.C., Rigling, A., Bugmann, H., 2007. The influence of changes in climate and land-use on
 416 regeneration dynamics of Norway spruce at the treeline in the Swiss Alps. *Silva Fennica* 41, 55-
 417 70.

418 Bradshaw, R., Gemmel, P., Bjorkman, L., 1994. Development of nature-based silvicultural models
 419 in Southern Sweden: the Scientific background. *Forest and Landscape Research* 1, 1-16.

420 Bray, J.R., Curtis, J.T., 1957. An Ordination of the Upland Forest Communities of Southern
 421 Wisconsin. *Ecological Monographs* 27, 325-349.

422 Carcaillet, C., 1998. A spatially precise study of Holocene fire history, Climate and Human Impact
 423 within the Maurienne Valley, North French Alps. *The Journal of Ecology* 86, 384-396.

424 Carpenter, S., Brock, W., Hanson, P., 1999. Ecological and social dynamics in simple models of
 425 ecosystem management. *Conservation Ecology* 3, 4. URL: [http://www.consecol.org/vol3/](http://www.consecol.org/vol3/iss2/art4)
 426 [iss2/art4](http://www.consecol.org/vol3/iss2/art4)

427 Clark, P.J., Evans, F.C., 1954. Distance to nearest neighbor as a measure of spatial relationships in
 428 populations. *Ecology* 35, 445-453.

429 Clark, J.S., 1990. Fire and climate change during the last 750 years in northwestern Minnesota.
 430 *Ecological Monographs* 60, 135-159.

431 Crookston, N.L., Stage, A.R., 1999. Percent canopy cover and stand structure statistics from the
 432 Forest Vegetation Simulator. US Forest Service General Technical Report RMRS-GTR-24. Fort
 433 Collins, CO.
 434 Cuevas, J.G., 2003. Gap characteristics in relation to forest structure and implications for southern
 435 beech forest dynamics. *Canadian Journal of Forest Research* 33, 1915-1922.
 436 Curtis, J.T., McIntosh, R.P., 1951. An upland forest continuum in the prairie-forest border region
 437 of Wisconsin. *Ecology* 32, 476-496.
 438 Dale, V.H., 1997. The relationship between land-use change and climate change. *Ecological*
 439 *Applications* 7, 753–769
 440 Del Favero, R., 2002. I tipi forestali nella Regione Lombardia. Cierre edizioni, Milano.
 441 Della Marianna, G., Gusmeroli, F., Puccio, C., 2004. Gli alpeggi della Comunità Montana
 442 Valtellina di Sondrio. Fondazione Fojanini, Sondrio.
 443 Didier, L., 2001. Invasion patterns of European larch and Swiss stone pine in subalpine pastures in
 444 the French Alps. *Forest Ecology and Management* 145, 67-77.
 445 Dullinger, S., Dirnböck, T., Greimler, J., Grabherr, G., 2003. A resampling approach for
 446 evaluating effects of pasture abandonment on subalpine plant species diversity. *Journal of*
 447 *Vegetation Science* 14, 243-252.
 448 Dumont, B., Hill, D.R.C., 2001. Multi-agent simulation of group foraging in sheep: effects of
 449 spatial memory, conspecific attraction and plot size. *Ecological Modelling* 141, 201-215.
 450 Ellis, J.E., Swift, D.M., 1988. Stability of African pastoral ecosystem: alternate paradigms and
 451 implications for development. *Journal of Range Management* 41, 450-459.
 452 Foster, D.R., 1992. Land-use history (1730-1990) and vegetation dynamics in central New
 453 England,
 454 USA. *The Journal of Ecology* 80, 753-771.

455 Foster, D.R., Swanson, F., Aber, J., Burke, I., Brokaw, N., Tilman, D., Knapp, A., 2003. The
 456 importance of land-use legacies to ecology and conservation. *BioScience* 53, 77-88.
 457 Gehrig-Fasel, J., Guisan, A., Zimmermann, N.E., 2007. Tree line shifts in the Swiss Alps: Climate
 458 change or land abandonment? *Journal of Vegetation Science* 18, 571-582.
 459 Gellrich, M., Zimmermann, N.E., 2007. Investigating the regional-scale pattern of agricultural
 460 land abandonment in the Swiss mountains: A spatial statistical modelling approach. *Landscape
 461 and Urban Planning* 79, 65–76.
 462 Guisan, A., Weiss, S.B., Weiss, A.D., 1999. GLM versus CCA spatial modeling of plant species
 463 distribution. *Plant Ecology* 143, 107-122.
 464 Guisan, A., Zimmermann, N.E., 2000. Predictive habitat distribution models in ecology.
 465 *Ecological Modelling* 135, 147-186.
 466 Hall, B., Motzkin, G., Foster, D.R., Syfert, m., Burk, J., 2002. Three hundred years of forest and
 467 land-use change in Massachusetts, USA. *Journal of Biogeography* 29, 1319-1335.
 468 Hawkins, B.A., Field, R., Cornell, H.V., Currie, D.J., Guégan, J.F., Kaufman, D.M., Kerr, J.T.,
 469 Mittelbach, G.G., Oberdorff, T., O'Brien, E.M., Porter, E.E., Turner, J.R.G., 2003. Energy, water,
 470 and broad-scale geographic patterns of species richness. *Ecology* 84, 3105-3117.
 471 Hessburg, P.F., Agee, J.K., 2003. An environmental narrative of Inland Northwest United States
 472 forests, 1800–2000. *Forest Ecology and Management* 178, 23-59.
 473 Houle, G., 2007. Determinants of fine scale plant species richness in a deciduous forest of
 474 notheastern North America. *Journal of Vegetation Science* 18, 345-354.
 475 Hu, L., Bentler, P., 1999. Cutoff criteria for fit indexes in covariance structure analysis:
 476 Conventional criteria versus new alternatives. *Structural equation modeling* 6, 1-55.
 477 Hunter, M.L., 1996. *Fundamentals of conservation biology*. Blackwell Science, Cambridge.

478 Kawamura, K., Akiyama, T., Yokota, H., Tsutsumi, M., Yasuda, T., Watanabe, O., Wang, S.,
 479 2005. Quantifying grazing intensities using geographic information systems and satellite remote
 480 sensing in the Xilingol steppe region, Inner Mongolia, China. *Agriculture, Ecosystems and*
 481 *Environment* 107, 83-93.

482 Kumar, L., Skidmore, A.K., Knowles, E., 1997. Modelling topographic variation in solar radiation
 483 in a GIS environment. *International Journal for Geographical Information Science* 11, 475-497.

484 Leduc, A., Drapeau, P., Bergeron, Y., Legendre, P., 1992. Study of Spatial Components of Forest
 485 Cover Using Partial Mantel Tests and Path Analysis. *Journal of Vegetation Science* 3, 69-78.

486 Lehringer, S., Höchtl, F., Konold, W., 2003. Effects of land use changes and depopulation on
 487 landscape, social life and tourism - overview about the results of a case study from the Piedmont
 488 Alps in Italy. *Centralblatt für das gesamte Forstwesen* 120, 1-18.

489 Lingua, E., Cherubini, P., Motta, R., Nola, P., 2007. Spatial structure along an altitudinal gradient
 490 in the Italian Central Alps reveals competition and facilitation processes among different
 491 coniferous species. *Journal of Vegetation Science* 19, 425-436.

492 MacDonald, D., Crabtree, J.R., Wiesinger, G., Dax, T., Stamou, N., Fleury, P., Gutierrez Lazpita,
 493 J., Gibon, A., 2000. Agricultural abandonment in mountain areas of Europe: Environmental
 494 consequences and policy response. *Journal of Environmental Management* 59, 47–69.

495 Margules, C.R., Austin, M.P., Mollison, D., Smith, F., 1994. Biological models for monitoring
 496 species decline: the construction and use of data bases [and discussion]. *Philosophical*
 497 *Transactions: Biological Sciences* 344, 69-75.

498 Marsh, G.P., 1864. *Man and nature; or, physical geography as modified by human action*. Belknap
 499 Press of Harvard University Press, Cambridge.

500 Maurer, K., Weyand, A., Fischer, M., Stöcklin, J., 2006. Old cultural traditions, in addition to land
501 use and topography, are shaping plant diversity of grasslands in the Alps. *Biological Conservation*
502 130, 438-446.

503 McCune, B., Grace, J.B., 2002. *Analysis of Ecological Communities*. MjM Software Design,
504 Gleneden Beach.

505 McCune, B., Mefford, M.J., 1999. *PC-ORD*. MjM Software Design, Gleneden Beach.

506 Motta, R., Lingua, E., 2005. Human impact on size, age, and spatial structure in a mixed European
507 larch and Swiss stone pine forest in the Western Italian Alps. *Canadian Journal of Forest Research*
508 35, 1809-1820.

509 Motta, R., Nola, P., 2001. Growth trends and dynamics in sub-alpine forest stands in the Varaita
510 Valley (Piedmont, Italy) and their relationships with human activities and global change. *Journal*
511 *of Vegetation Science* 12, 219-230.

512 Motta, R., Morales, M., Nola, P., 2006. Human land-use, forest dynamics and tree growth at the
513 treeline in the Western Italian Alps. *Annals of Forest Science* 63, 739-747.

514 Motta, R., Nola, P., Piussi, P., 2002. Long-term investigations in a strict forest reserve in the
515 eastern Italian Alps: spatio-temporal origin and development in two multi-layered subalpine
516 stands. *Journal of Ecology* 90, 495-507.

517 Motzkin, G., Wilson, P., Foster, D.R., Allen, A., 1999. Vegetation patterns in heterogeneous
518 landscapes: the importance of history and environment. *Journal of Vegetation Science* 10, 903-
519 920.

520 Nakamura, T., Go, T., Wuyunna, Hayashi, I., 2000. Effects of grazing on the floristic composition
521 of grasslands in Baiyinxile, Xilingole, Inner Mongolia. *Grassland Science* 45, 324-350.

522 Naveh, Z., 1982. Landscape ecology as an emerging branch of human ecosystem science. In,
523 *Advances in Ecological Research*. Academic press, London, pp. 189-237.

524 Naveh, Z., 1995. Interactions of landscapes and cultures. *Landscape and Urban Planning* 32, 43-
525 54.

526 Neale, M.C., 1994. MxGui 3.2. Department of Psychiatry, Virginia Commonwealth University,
527 Richmond. <http://www.vcu.edu/mx>

528 Neumann, M., Starlinger, F., 2000. The significance of different indices for stand structure and
529 diversity in forests. *Forest Ecology and Management* 145, 91-106.

530 Nola, P., 1994. A dendroecological study of Larch at timberline in the Central Italian Alps.
531 *Dendrochronologia* 12, 77-91.

532 Nola, P., Motta, R., 1996. Una cronologia plurisecolare di larice (*Larix decidua* Mill.) per l'alta
533 Valmalenco (Sondrio, Italia). *Dendrochronologia* 14, 31-42.

534 Parshall, T., Foster, D.R., Faison, E., MacDonald, D., Hansen, B.C.S., 2003. Long-term history of
535 vegetation and fire in pitch pine-oak forests on Cape Cod, Massachusetts. *Ecology* 84, 736-748.

536 Peterken, G.F., 1992. Conservation of old growth: a European perspective. *Natural Areas Journal*
537 12, 10-13.

538 Peterken, G.F., 1996. *Natural Woodland. Ecology and Conservation in Northern Temperate*
539 *Regions*. Cambridge University Press, Cambridge.

540 Piussi, P., 2000. Expansion of European mountain forests. In: Price, M.F., Butt, N. (Eds.), *Forest*
541 *in sustainable mountain development: a state of knowledge report for 2000*. CABI Publishing,
542 Wallingford, pp. 19-25.

543 Roath, L.R., Krueger, W.C., 1982. Cattle grazing and behavior on a forested range. *Journal of*
544 *Range Management* 35, 332-338.

545 Shipley, B., 2000. *Cause and Correlation in Biology: A User's Guide to Path Analysis, Structural*
546 *Equations and Causal Inference*. Cambridge University Press, Cambridge.

547 Società Agraria di Lombardia, 1901. Atti della Commissione d'inchiesta sui pascoli alpini. Società
548 Agraria di Lombardia. Milano.

549 Solomon, D.S., Gove, J.H., 1999. Effects of uneven-age management intensity on structural
550 diversity in two major forest types in New England. *Forest Ecology and Management* 114, 265-
551 274.

552 Spies, T.A., Franklin, J.F., Klopsch, M., 1990. Canopy gaps in Douglas-fir forests of the Cascade
553 Mountains. *Canadian Journal of Forest Research* 20, 649-658.

554 Sprugel, D.G., 1991. Disturbance, equilibrium, and environmental variability: What is "natural"
555 vegetation in a changing environment? *Biological Conservation* 58, 1-18.

556 Staudhammer, C.L., LeMay, V.M., 2001. Introduction and evaluation of possible indices of stand
557 structural diversity. *Canadian Journal of Forest Research* 31, 1105–1115.

558 Stephenson, N.L., 1990. Climatic control of vegetation distribution: the role of the water balance.
559 *The American Naturalist* 135, 649-670.

560 Stern, R., 1983. Human impact on tree borderlines. In: Holzner, W., Werger, M.J.A., Ikusima, I.
561 (Eds.), *Man's impact on vegetation*. Dr W. Junk Publisher, The Hague, pp. 227-236.

562 Swetnam, T.W., Allen, C.D., Betancourt, J.L., 1999. Applied historical ecology: using the past to
563 manage for the future. *Ecological Applications* 9, 1189-1206.

564 Tappeiner, U., Tasser, E., Tappeiner, G., 1998. Modelling vegetation patterns using natural and
565 anthropogenic influence factors: preliminary experience with a GIS based model applied to an
566 Alpine area. *Ecological Modelling* 113, 225-237.

567 Tappeiner, U., Tappeiner, G., Aschenwald, J., Tasser, E., Ostendorf, B., 2001. GIS-based
568 modelling of spatial pattern of snow cover duration in an alpine area. *Ecological Modelling* 138,
569 265-275.

570 Turner, B.L., Meyer, W.B., 1993. Environmental change: the human factor. In: McDonnell, M.J.,
 571 Pickett, S.T.A. (Eds.), Humans as components of ecosystems. Springer-Verlag, New York, pp. 40-
 572 50.
 573 Turner, M.G., Gardner, R.H., O'Neill, R.V., 2001. Landscape ecology in theory and practice.
 574 Springer, New York.
 575 Urban, D.L., Miller, C., Halpin, P.N., Stephenson, N.L., 2000. Forest gradient response in Sierran
 576 landscapes: the physical template. *Landscape Ecology* 15, 603-620.
 577 von Humboldt, A., 1808. Ansichten der Natur mit Wissenschaftlichen Erläuterungen. Cotta J.G.,
 578 Tübingen.
 579 Vacher, K.A., Killingbeck, K.T., August, P.V., 2007. Is the relative abundance of nonnative
 580 species an integrated measure of anthropogenic disturbance? *Landscape Ecology* 22, 821-835.
 581 Zimmermann, N., 2000. Tools for analyzing, summarizing, and mapping of biophysical variables:
 582 Solar radiation routine Shortwavb.aml. In:
 583 http://www.wsl.ch/staff/niklaus.zimmermann/programs/aml1_3.html
 584 Walther, P., 1986. Land abandonment in the Swiss Alps: a new understanding of a land-use
 585 problem. *Mountain Research and Development* 6, 305-314.
 586 Weisberg, P.J., Lingua, E., Pillai, R.B., 2007. Spatial Patterns of Pinyon–Juniper Woodland
 587 Expansion in Central Nevada. *Rangeland Ecology and Management* 60, 115-124.
 588 White, M.A., Mladenoff, D.J., 1994. Old-growth forest landscape transitions from pre-European
 589 settlement to present. *Landscape Ecology* 9, 191-205.
 590 Whitlock, C., Knox, M.A., 2002. Prehistoric Burning in the Pacific Northwest. In: Vale, T.R.
 591 (Ed.), Fire, Native Peoples, and the Natural Landscape. Island Press, Washington, pp. 195-232.
 592 Wu, J., Loucks, O.L., 1995. From balance of nature to hierarchical patch dynamics: a paradigm
 593 shift in ecology. *The Quarterly Review of Biology* 70, 439-466.

594 **Tables**

595 Table 1 - Descriptors and measures used in the stand structure classification.

596

STAND DESCRIPTORS	Symbols	Measures and references
Richness of trees	Ri	n. of trees' species
Dominance	Do	relative density of larch stems
Density	De	n. of stems per hectare
Diameter (mean)	Dbh-Me	diameter at 130 cm (mean value)
Diameter (standard dev.)	Dbh-Sd	diameter at 130 cm (standard deviation)
Basal Area	BA	basal area per hectare
Height	He	mean height
Canopy Cover	CC	canopy cover (following Crookston and Stage 1999)
Nearest neighbor Index	NNI	Clark-Evans Index of aggregation (Clark and Evans, 1954)
Vertical Evenness	VE	vertical evenness following Neumann and Starlinger, 2001
Age of larch	AGE	age estimation of the 3 largest-diameter larches

597

598

599 Table 2 - Mean values of the 11 descriptors for each structure type obtained by cluster analysis at Musella and Ventina sites.

600

	Stand structure types	Plots	Ri	Do	De	DBH-Me	DBH-Sd	BA	He	CC	NNI	VE	AGE
		(n)	(n)	(%)	(n/ha)	(cm)	(cm)	(m2/ha)	(m)	(%)			(yrs)
Musella	1 Sparse; larch dominated; small trees	6	1.50	0.83	301.67	11.75	5.95	4.28	6.05	28.33	0.60	0.65	77.17
	2 Mean density; mixed; mean size trees	4	2.00	0.18	541.00	17.85	12.30	18.83	9.63	43.25	0.64	0.64	123.00
	3 Mean density; mixed; big trees	8	1.88	0.50	480.38	29.23	15.25	38.68	15.79	59.13	0.74	0.62	163.67
	4 Dense; larch dominated; mean size trees	10	2.00	0.84	689.20	15.33	10.80	19.75	8.13	54.90	0.65	0.73	163.00
Ventina	1 Sparse; larch dominated; small trees	11	1.64	0.92	266.73	11.48	5.80	4.02	5.05	18.45	0.60	0.59	84.30
	2 Mean density; mixed; mean size trees	7	2.43	0.43	252.21	19.52	17.96	12.55	7.16	21.14	0.69	0.62	273.29
	3 Mean density; mixed; big trees	7	2.14	0.90	249.33	36.66	20.96	50.93	12.98	42.71	0.79	0.59	298.57
	4 Dense; larch dominated; mean size trees	15	2.33	0.88	640.90	17.76	12.09	20.43	8.54	54.80	0.74	0.74	185.67

601

602

603 Table 3 - Principal component loadings for the first two principal components for both study areas.

604 Loadings greater than 0.4 are indicated in bold.

605

	MUSELLA		VENTINA	
Axis	PC 1	PC 2	PC 1	PC 2
% of variance	40.27	17.88	32.84	24.42
Ri	0.143	0.244	0.127	0.398
Do	-0.253	-0.050	0.016	-0.088
De	0.110	0.637	0.023	0.549
Dbh-Me	0.423	-0.230	0.449	-0.209
Dbh-Sd	0.397	-0.196	0.447	-0.105
BA	0.448	0.062	0.380	0.081
He	0.438	-0.121	0.441	-0.063
CC	0.306	0.345	0.251	0.451
NNI	0.186	0.196	0.230	0.172
VE	-0.171	0.483	-0.027	0.447
AGE	0.130	0.179	0.356	-0.183

606

607

608

609

610

611

612

Table 4 - Path models comparison by means of their fit indices. Akaike's Information Criterion (AIC) statistic and the Root Mean Square Error of Approximation (RMSEA) together with degrees of freedom (df), Maximum Likelihood chi-squared fit function value (ML χ^2) and its probability (P) are reported.

Models	Goodness-of-Fit statistics				
	ML χ^2	df	P	RMSEA	AIC
M1	4.440	9	0.880	<0.001	-13.560
M2	14.847	19	0.732	< 0.001	-23.153
V1	0.981	3	0.806	< 0.001	-5.019
V2	0.927	3	0.819	< 0.001	-5.073

Figure captions

Figure1 - Location of the 70 plots (●) in two watersheds of Malenco valley.

Figure 2 - Conceptual model to be tested for both study areas through path analysis. The full model includes topographic, climatic and anthropogenic variables associated through positive or negative causal paths. “Stand structure” refers to the first principal component (PC 1) defined as tree size.

Figure 3 - Biplot from Principal Components Analysis of 28 plots at Musella. Site scores are shown as triangles; stand structure types are reported as number (1-4). Correlations of environmental variables with PCA axes are shown as linear vectors. Vectors are shown only for correlations > 0.05 . The first and second principal component accounted for 40% and 18% of the total amount of variation, respectively.

Figure 4 - Biplot from Principal Components Analysis of 40 plots at Ventina. Site scores are shown as triangles; stand structure types are reported as number (1-4). Correlations of environmental variables with PCA axes are shown as linear vectors. Vectors are shown only for correlations > 0.05 . The first and second principal component accounted for 33% and 24% of the total amount of variation, respectively.

Figure 5 - Path diagrams for Musella (M1, M2) and Ventina (V1, V2). Continuous lines: positive paths; dotted lines: negative paths; single arrow lines: causal paths; double arrow lines: covariance paths. Thickness of causal path vectors corresponds to the strength of effect. Only significant path coefficients are presented next to each path.

Figure 6 - Bivariate scatter plots of selected environmental variables vs. the stand structure PCs at Musella. Only relationships significant for the SEM analyses are shown.

Figure 7 - Bivariate scatter plots of selected environmental variables vs. the stand structure PCs at Ventina. Only relationships significant for the SEM analyses are shown.

655 Figure 1

656

657

658 Figure 2

659

660 Figure 3

661

662 Figure 4

663

664 Figure 5 (M1, M2, V1, V2)

683 Figure 6

684

685

686

687

688

Figure 7