AperTO - Archivio Istituzionale Open Access dell'Università di Torino

Superpulsed laser irradiation increases osteoblast activity via modulation of bone morphogenetic factors

This is the author's manuscript

Original Citation:

Availability:

This version is available <http://hdl.handle.net/2318/72287> since 2019-04-30T10:20:24Z

Terms of use:

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)

Superpulsed Laser Irradiation Increases Osteoblast Activity Via Modulation of Bone Morphogenetic Factors

Silvia Saracino, BSc,¹ Marco Mozzati, DDS,² Germana Martinasso, PhD,¹ Renato Pol, DMD,² Rosa A. Canuto, MD,¹ and Giuliana Muzio, PhD^{1*}

¹Department of Experimental Medicine and Oncology, University of Turin, Corso Raffaello 30, 10125 Turin, Italy

²Department of Biomedical Sciences and Human Oncology, Section of Prosthetic Dentistry, School of Dentistry, University of Turin, Lingotto, Via Nizza 230, 10126 Turin, Italy

Background and Objective: Laser therapy is a new approach applicable in different medical fields when bone loss occurs, including orthopedics and dentistry. It has also been used to induce soft-tissue healing, for pain relief, bone, and nerve regeneration. With regard to bone synthesis, laser exposure has been shown to increase osteoblast activity and decrease osteoclast number, by inducing alkaline phosphatase (ALP), osteopontin, and bone sialoprotein expression. Studies have investigated the effects of continuous or pulsed laser irradiation, but no data are yet available on the properties of superpulsed laser irradiation. This study thus aimed to investigate the effect of superpulsed laser irradiation on osteogenic activity of human osteoblast-like cells, paying particular attention to investigating the molecular mechanisms underlying the effects of this type of laser radiation.

Study Design/Materials and Methods: Human osteoblast-like MG-63 cells were exposed to 3, 7, or 10 superpulsed laser irradiation (pulse width 200 nanoseconds, minimum peak power 45 W, frequency 30 kHz, total energy 60 J, exposure time 5 minutes). The following parameters were evaluated: cell growth and viability (light microscopy, lactate dehydrogenase release), calcium deposits (Alizarin Red S staining), expression of bone morphogenetic factors (real-time PCR).

Results: Superpulsed laser irradiation decreases cell growth, induces expression of TGF- β 2, BMP-4, and BMP-7, type I collagen, ALP, and osteocalcin, and increases the size and the number of calcium deposits. The stimulatory effect is maximum on day 10, that is, after seven applications.

Conclusions: Reported results show that superpulsed laser irradiation, like the continuous and pulsed counterparts, possesses osteogenic properties, inducing the expression of molecules known to be important mediators of bone formation and, as a consequence, increasing calcium deposits in human MG-63 cells. Moreover, the data suggest a new potential role for PPAR γ as a regulator of osteoblast proliferation. *Lasers Surg. Med.* 41:298–304, 2009.

© 2009 Wiley-Liss, Inc.

Key words: BMPs; calcium deposits; MG-63 cells; superpulsed laser; TGF- β 2

INTRODUCTION

Laser therapy is a new approach in different medical fields, including orthopedics and dentistry, when bone loss occurs, that is, in cases of bone fracture and tooth extraction [1]. Recent studies have reported the benefits of low level laser therapy (LLLT), which has been used to induce soft-tissue healing, for pain relief, bone, and nerve regeneration [2], although the molecular mechanisms triggered are not yet fully clear.

With regard to bone synthesis, *in vivo* experiments on rat femur have shown that pulsed laser irradiation with high peak power stimulates bone formation by increasing osteoblast activity and decreasing osteoclast numbers [3]. Beneficial properties have been ascribed to LLLT's anti-inflammatory effect, postulating that the treatment modulates transcription factors and regulates the expression of pro-inflammatory cytokines [4]. A recent study on the subplantar tissue of rat's paw evidenced that LLLT decreased mRNA content of TNF- α , IL-1 β , and IL-6 [5]; it was suggested that an early target of radiation was TNF- α which, in turn, activates other cytokines. These authors also reported a decreased expression of kinin receptors in the same experimental model [6].

It has also been reported that LLLT induces the formation of small amounts of reactive oxygen species (ROS), which can trigger cell stimulation via increased mitochondrial respiration and ATP formation [7].

The stimulatory effect of LLLT has also been confirmed *in vitro* in different cell lines. In osteoblast-like cells isolated from fetal rat calvariae, LLLT stimulated proliferation and differentiation, inducing alkaline phosphatase (ALP), osteopontin (OP), and bone sialoprotein expression [8]. Similar results have been obtained in cultured human SaOS-2 cells, where early induction of ALP, type I collagen,

Contract grant sponsor: Regione Piemonte, Italy; Contract grant sponsor: University of Turin, Italy; Contract grant sponsor: Heraeus Kulzer GmbH, Division Dentistry, Hanau, Germany.

*Correspondence to: Giuliana Muzio, PhD, Department of Experimental Medicine and Oncology, University of Turin, Corso Raffaello 30, 10125 Turin, Italy. E-mail: giuliana.muzio@unito.it

Accepted 19 February 2009

Published online in Wiley InterScience

(www.interscience.wiley.com).

DOI 10.1002/lsm.20762

and OP was observed in cells irradiated with 670 nm diode laser [9,10]. Moreover, treatment with LLLT of human osteoblast-like cells, cultured on titanium, induced cellular activity in a dose-dependent manner, through increased osteocalcin and TGF- β_1 production [11]. TGF- β is one of the most abundant regulatory factors stored in the bone matrix [12]; it drives the cascade of events leading to new bone formation. In case of bone resorption, TGF- β is released early, and favors bone growth; however, once the process of bone formation is started, it acts as an inhibitor of mineralization and suppresses markers of osteoblast activity; this suggests that TGF- β affects osteoblast precursors but not differentiated osteoblasts [13,14].

Bone morphogenetic proteins (BMPs) are members of the TGF- β superfamily, known to play an important role in bone formation [15]. Of the BMPs, BMP-2, 6, and 9 have been shown to be the most potent agents in inducing osteoblast differentiation of mesenchymal stem cells. On the contrary, BMP-3 exhibits no osteogenic activity and antagonizes the osteogenic activity of BMP-2, 4, 6, 7, and 9 [16].

Recent clinical trials have shown that BMP-7 is effective in stimulating ALP production in relatively mature osteoblasts [17]. Taken together, these findings suggest that interactions among BMPs are fundamental in regulating and driving the bone regeneration process.

Based on the fact that osteoblasts and adipocytes share a common progenitor (multipotential mesenchymal stem cells in bone marrow), and since peroxisome proliferated-activated receptor (PPAR) γ is a key regulator of adipocyte differentiation, the possible role of this nuclear receptor in bone metabolism has been investigated. It has been reported that low PPAR γ expression enhances formation of bone mass by stimulating osteoblastogenesis [18]. In cultured primary osteoblasts, PPAR γ ligands have been shown to decrease mineralized bone nodules and ALP activity, suggesting that activation of PPAR γ inhibits the expression of mediators involved in bone formation, such as BMP-2 and osteocalcin [19], and that the appropriate antagonists of PPAR γ could provide a new approach to treating some bone disorders [20].

Laser irradiation can also be used in the superpulsed modality, although no reports on its use have yet been published. The present study thus investigated the effects of superpulsed laser irradiation on osteogenic activity of human osteoblast-like cells, paying particular attention to investigating the molecular mechanisms underlying those effects.

MATERIALS AND METHODS

Materials

MG-63 human osteoblast-like cells were from ATCC (Rockville, MD); L-glutamine was from (Gibco Life Technologies, Paisley, UK); Trypan Blue (TB) was from Carlo Erba Reagenti (Milano, Italy); all the other reagents were from Sigma (St. Louis, MO).

Cell Culture

Cells were seeded $5 \times 10^4/\text{cm}^2$ in modified eagle medium (MEM) supplemented with 2 mM L-glutamine, 1% (v/v) antibiotic/antimycotic solution, 1 mM sodium pyruvate, and 10% (v/v) fetal bovine serum (FBS). Cells were maintained at 37°C in a 5% CO₂ atmosphere.

Laser Treatment

Twenty-four hours after seeding, cells were exposed to superpulsed laser irradiation, with the following protocol: every 24 hours for the first 5 days, then every 48 hours until day 20. A Lumix 2 HFPL Dental device IR (904–910 nm) Gallium Arsenide laser (Fisioline s.n.c., Verduno, Cuneo, Italy) was used, with the following experimental parameters: pulse width 200 nanoseconds, minimum peak power 33 W, average out power 200 mW, frequency 30 kHz, exposure time 5 minutes, total energy 60 J for each well (illuminated surface size: 9 cm²). The administered dose was 6.7 J/cm². Control cells were not exposed to laser irradiation. Both control and laser-treated cells were harvested by trypsinization at days 4 (3 exposures to laser irradiation), 10 (7 exposures to laser irradiation), and 20 (12 exposures to laser irradiation), and used for the determinations reported below.

Cell Growth and Viability

Cells present in monolayer and culture medium were counted using a Burkner chamber. Cell viability was determined by the TB exclusion test and through lactate dehydrogenase (LDH) activity in the culture medium, as marker of necrotic death. TB exclusion was evaluated on cells suspended in the presence of the dye (0.8 mg/ml); 400 cells were counted for each sample and results expressed as percentages of TB-positive cells. LDH activity was assessed spectrophotometrically on clear culture supernatants after centrifugation at 600g for 10 minutes and expressed as nmoles NADH oxidized/minute/ml of culture medium [21].

Calcium Deposition

Alizarin Red S staining was used to evaluate the formation of calcium deposits and mineral matrix: detached cells were fixed in 70% ethanol, stained with 1% Alizarin Red solution for 2 minutes, washed with Tris-buffered saline solution and observed under a light microscope to evaluate Alizarin positivity.

Real-Time Polymerase Chain Reaction (Real-Time PCR)

Total RNA was isolated from MG-63 cells with a RNEasy[®] Mini Kit (Qiagen, GmbH, Germany) following the manufacturer's protocol.

One microgram of RNA was reverse transcribed using the High Capacity cDNA Archive kit (Applied Biosystems, Foster City, CA). PCR was performed using IQ[™] SYBR-Green Supermix (Bio-Rad, Hercules, CA) in a iCycler system (Bio-Rad).

Each sample was tested three times, and the threshold cycle (C_t) values from each reaction were averaged.

TABLE 1. Human Primer Sequences for Real-Time PCR

Gene	Forward primer 5'-3'	Reverse primer 5'-3'	bp
GAPDH	GTCGGAGTCAACGGATTTGG	GGGTGGAATCATATTGGAACATG	142
ALP	CTCCAGTCTCATCTCCT	AAGACCTCAACTCCCCTGAA	120
BMP-4	CTCGCTCTATGTGGACTT	ATGTTTGTTGAGTTGAGG	130
BMP-7	GTGGAACATGACAAGGAAT	GAAAGATCAAACCGGAAC	65
OCN	GTGACGAGTTGGCTGACC	CAAGGGGAAGAGGAAAGAAGG	129
PPAR α	GGATGTACACAACGCGATT	GCCAGAGATTTGAGATCTGCAGTT	127
PPAR γ	GCCGAGAAGGAGAAGC	TGGTCAGCGGGAAGG	150
TGF- β 2	GAG TAC TAC GCC AAG GAG GTT TAC A	CGA ACA ATT CTG AAG TAG GGT CTG T	104
COLL I	GAG GAA ACT GTA AGA AAG G	GTT CCC ACC GAG ACC	150

GAPDH, glyceraldehyde 3-phosphate dehydrogenase; ALP, alkaline phosphatase; BMP-4, bone morphogenetic protein 4; BMP-7, bone morphogenetic protein 7; OCN, osteocalcin; PPAR α , peroxisome proliferators-activated receptor α ; PPAR γ , peroxisome proliferators-activated receptor γ ; TGF- β 2, transforming growth factor β 2; COLL I, type I collagen; bp, product length.

The change was quantified as the relative expression compared to that at time 0 (before the first exposure to laser irradiation), calculated as $2^{-\Delta\Delta C_t}$, where $\Delta C_t = C_{t\text{sample}} - C_{t\text{GAPDH}}$ and $\Delta\Delta C_t = \Delta C_{t\text{sample}} - \Delta C_{t0}$. Human primer sequences used for real-time PCR are reported in Table 1.

Statistical Analysis

Data are expressed in all cases as means $\pm$ SD. The significance of differences between group means was assessed by variance analysis, followed by the Newman-Keuls test ($P < 0.05$).

RESULTS

The effect of super-pulsed laser irradiation on cell proliferation and viability, and on the expression of some parameters involved in bone synthesis was studied in human osteoblast-like cells MG-63.

Figure 1 reports proliferation, showing a decrease of cell numbers in laser-treated cells versus controls. The

Fig. 1. Growth of cells exposed or not to superpulsed laser irradiation. Data refer to cells present in the monolayer and are means $\pm$ SD of five experiments. Means with different letters are significantly different from one another ($P < 0.05$) as determined by analysis of variance followed by post hoc Newman-Keuls test. C, control cells; L, laser-treated cells.

decrease started from the fourth day and reached a maximum on day 10 after seven applications, at which point the number of laser-treated cells was 75% of the control value. No induction of cell death was shown by the TB exclusion test, nor by counting cells floating in the medium or determining LDH release in the medium, at any time studied (data not reported).

Figures 2–6 show the stimulatory effect of super-pulsed laser irradiation on osteoblast activity. Figure 2 reports the

Fig. 2. mRNA content of alkaline phosphatase (ALP) and osteocalcin in cells exposed or not to superpulsed laser irradiation. Data are means $\pm$ SD of five experiments. Means with different letters are significantly different from one another ($P < 0.05$) as determined by analysis of variance followed by post hoc Newman-Keuls test. C, control cells; L, laser-treated cells.

Fig. 3. mRNA content of TGF- β 2 and type I collagen in cells exposed or not to superpulsed laser irradiation. Data are means $\pm$ SD of five experiments. Means with different letters are significantly different from one another ($P < 0.05$) as determined by analysis of variance followed by post hoc Newman-Keuls test. C, control cells; L, laser-treated cells.

Fig. 4. mRNA content of BMP-4 and BMP-7 in cells exposed or not to superpulsed laser irradiation. Data are means $\pm$ SD of five experiments. Means with different letters are significantly different from one another ($P < 0.05$) as determined by analysis of variance followed by post hoc Newman-Keuls test. C, control cells; L, laser-treated cells.

Fig. 5. mRNA content of PPAR α and γ in cells exposed or not to superpulsed laser irradiation. Data are means $\pm$ SD of five experiments. Means with different letters are significantly different from one another ($P < 0.05$) as determined by analysis of variance followed by post hoc Newman-Keuls test. C, control cells; L, laser-treated cells.

mRNA content of ALP and osteocalcin. In laser-treated cells, the ALP values were higher than those in control cells at days 10 and 20, being the increase versus control 23% and 98%, respectively, whereas no significant difference was present at day 4 between treated and control cells. As regards osteocalcin, in laser-treated cells, mRNA content was higher than in control cells only at day 10 (+60%), and was similar to the other experimental times.

Figure 3 shows the mRNA content of TGF β 2 and type I collagen. In laser-treated cells, the TGF β 2 mRNA content was higher on days 10 and 20 (+30%, +40% vs. controls), whereas it was similar on day 4. Type I collagen showed significant changes on days 10 and 20 (+109%, +58% vs. controls).

As regards BMPs, a significant increase in laser-treated cells was evident in BMP-4 mRNA only on day 20 (+44% vs. control); whereas BMP-7 was induced on both day 10 and day 20, the larger increase being on day 10 (+50% vs. control; Fig. 4).

Figure 5 shows PPAR α and γ mRNA contents; both isotypes increased in laser-treated cells at day 10 in comparison with control cells, although the increase in PPAR γ was larger (90% vs. 25% of PPAR α); at the other times, no significant difference was observed between laser-exposed cells and controls.

Fig. 6. Calcium deposits evidenced with Alizarin S staining in cells exposed or not to superpulsed laser irradiation. MG-63 harvested from the monolayer were stained with Alizarin red S and observed under the light microscope. Cells were analyzed on days 10 and 20. The black arrow shows the presence of calcium deposits. **Panel A:** DAY 10-C, control cells after 10 days of experimental time. **Panel B:** DAY 10-L, laser-

treated cells after 10 days of experimental time. **Panel C:** DAY 20-C, control cells after 20 days of experimental time. **Panel D:** DAY 20-L, laser-treated cells after 20 days of experimental time (magnification of panel **D**). **Panels E and F:** DAY 20-L, laser-treated cells after 20 days of experimental time (magnification of panel **D**).

Calcium deposits were investigated by staining cells with Alizarin red S (Fig. 6). Especially at 20 days (panel D vs. panel C), larger and more numerous calcium deposits were present in laser-treated cells compared to controls. Panels E and F are magnification of nodules present in laser-treated cells at 20 days.

DISCUSSION

The study constitutes the first report investigating the effect of superpulsed laser irradiation on human osteoblastic-like cells. To date, evidence of the osteogenic properties of continuous or pulsed laser irradiation has been reported, both in vitro and in vivo [22–26]. The present results show that repeated super-pulsed laser irradiation inhibits cell proliferation in human osteoblast-like cells and, especially, that it increases the expression of proteins essential for bone formation, that is, TGF- β 2, BMP-4, and BMP-7, ALP, type I collagen, and osteocalcin. The stimulatory effect becomes evident on day 10, after seven laser irradiation exposures.

The inverse correlation between cell growth and osteogenic activity in MG-63 cells is not surprising since it is well known that cell differentiation process is characterized by a reduction of proliferation coupled with an increased tissue-specific gene expression, as also recently confirmed [27].

At present, few studies have addressed the molecular mechanisms underlying the osteogenic effect of continuous or pulsed laser irradiation. It has been suggested that this type of electromagnetic radiation influences biochemical activities, inducing a transient heating of some chromophores [28] and/or changing the redox status, with consequent production of ROS. It is well known that, when produced in large amounts, ROS are cell destructive, but that in small amounts they can act as “secondary messengers” via different pathways [29]. An effect of ROS is to induce lipid peroxidation and, as consequence, to increase production of aldehydes such as 4-hydroxynonenal, which induce cell differentiation at low concentrations [30,31]. Aldehydes derived from lipid peroxidation may also affect the activity of important membrane enzymes, such as ATPase, 5'-nucleotidase, phospholipase C [31].

The possibility that the effects of laser irradiation are in part mediated by changes in plasmamembranes may be supported by observations that shock waves, another kind of mechanical biostimulation, increase substance uptake and delivery via membrane modulation [29,32].

It is likely that, in our experimental conditions, a transient but stimulating heating occurs, since the diode laser equipment used is within the infrared wavelength band (904–910 nm). In turn, the laser-induced transient

heating could increase ROS production and lipid peroxidation [33].

It has been reported very recently that "in vitro" low-power far-infrared laser irradiation activated latent TGF- β 1 immediately, and subsequently TGF- β 3, possibly via induction of a conformational change in the latent complexes. These changes render TGF- β more amenable to the usual activation by altered pH or proteases [34]. These results suggest that the TGF- β pathway might be a key molecular mechanism triggered by laser irradiation. In our research, superpulsed laser-treated cells showed a significant increase of TGF- β 2 mRNA versus controls, on both day 10 and day 20, suggesting the possibility that this signal transduction pathway may also be activated by superpulsed laser irradiation.

Other important changes observed in the superpulsed laser-treated cells concern proteins and transcription factors involved in regulating bone formation. An early increase of BMP-7 was observed, followed by the increase of BMP-4. This temporal sequence could be in agreement with the observation that BMP-7 is an early factor involved in committing mesenchymal stem cells to osteoblast differentiation and, especially, with reports that both BMP-7 and BMP-4 increase expression of ALP and osteocalcin [35].

For the first time, our research showed that laser irradiation affects the expression of PPARs. PPAR α and γ were examined because they are closely connected with the reduction of cell viability and proliferation and with the increase in differentiation [36,37]. In laser-exposed cells, the strongest increase occurred in the isotype γ . The observation regarding the PPAR γ increase is in disagreement with some reports indicating an anti-osteoblastogenic role for this nuclear receptor [38–41]; these studies principally analyzed the role of PPAR γ in determining the destiny of multipotent mesenchymal stem cells in bone marrow.

In the light of these observations we hypothesize that, in differentiated MG-63 cells, the increased PPAR γ is mainly involved in decreasing cell proliferation, as has been reported for other types of both normal and tumor cells [42–44]. The true role of PPAR γ in modulating laser effect is now under investigation in experiments using specific antagonists.

The bulk of the results obtained shows that superpulsed laser irradiation, like the continuous and pulsed counterparts, possesses osteogenic properties, inducing the expression of molecules known to be important mediators of bone formation and, as a consequence, increasing the size and the number of calcium deposits in human MG-63 cells. Moreover, a new potential role may be suggested for the nuclear receptor PPAR γ as a regulator of osteoblast proliferation.

ACKNOWLEDGMENTS

This research was supported by grants from Regione Piemonte, Italy, University of Turin, Italy, and Heraeus Kulzer GmbH, Division Dentistry, Hanau, Germany. We thank Mrs Frances Cooper for revising the English.

REFERENCES

- Ozawa Y, Shimizu N, Kariya G, Abiko Y. Low-energy laser irradiation stimulates bone nodule formation at early stages of cell culture in rat calvarial cells. *Bone* 1998;22:347–354.
- Soares LP, Oliveira MG, Pinheiro AL, Fronza BR, Maciel ME. Effects of laser therapy on experimental wound healing using oxidized regenerated cellulose hemostat. *Photomed Laser Surg* 2008;26:10–13.
- Ninomiya T, Hosoya A, Nakamura H, Sano K, Nishisaka T, Ozawa H. Increase of bone volume by a nanosecond pulsed laser irradiation is caused by a decreased osteoclast number and an activated osteoblasts. *Bone* 2007;40:140–148.
- Albertini R, Aimbire F, Villaverde AB, Silva JA, Jr., Costa MS. COX-2 mRNA expression decreases in the subplantar muscle of rat paw subjected to carrageenan-induced inflammation after low level laser therapy. *Inflamm Res* 2007;56:228–229.
- Albertini R, Villaverde AB, Aimbire F, Bjordal J, Brugnera A, Mittmann J, Silva JA, Costa M. Cytokine mRNA expression is decreased in the subplantar muscle of rat paw subjected to carrageenan-induced inflammation after low-level laser therapy. *Photomed Laser Surg* 2008;26:19–24.
- Bortone F, Santos HA, Albertini R, Pesquero JB, Costa MS, Silva JA, Jr. Low level laser therapy modulates kinin receptors mRNA expression in the subplantar muscle of rat paw subjected to carrageenan-induced inflammation. *Int Immunopharmacol* 2008;8:206–210.
- Morimoto Y, Arai T, Kikuchi M, Nakajima S, Nakamura H. Effect of low-intensity argon laser irradiation on mitochondrial respiration. *Lasers Surg Med* 1994;15:191–199.
- Ueda Y, Shimizu N. Effects of pulse frequency of low-level laser therapy (LLLT) on bone nodule formation in rat calvarial cells. *J Clin Laser Med Surg* 2003;21:271–277.
- Stein E, Koehn J, Sutter W, Wendtlandt G, Wanschitz F, Thurnher D, Baghestanian M, Turhani D. Initial effects of low-level laser therapy on growth and differentiation of human osteoblast-like cells. *Wien Klin Wochenschr* 2008;120:112–117.
- Stein A, Benayahu D, Maltz L, Oron U. Low-level laser irradiation promotes proliferation and differentiation of human osteoblasts in vitro. *Photomed Laser Surg* 2005;23:161–166.
- Khadra M, Lyngstadaas SP, Haanaes HR, Mustafa K. Effect of laser therapy on attachment, proliferation and differentiation of human osteoblast-like cells cultured on titanium implant material. *Biomaterials* 2005;26:3503–3509.
- Harris SE, Bonewald LF, Harris MA, Sabatini M, Dallas S, Feng JQ, Ghosh-Choudhury N, Wozney J, Mundy GR. Effects of transforming growth factor beta on bone nodule formation and expression of bone morphogenetic protein 2, osteocalcin, osteopontin, alkaline phosphatase, and type I collagen mRNA in long-term cultures of fetal rat calvarial osteoblasts. *J Bone Miner Res* 1994;9:855–863.
- Gordon KJ, Blobel GC. Role of transforming growth factor-beta superfamily signaling pathways in human disease. *Biochim Biophys Acta* 2008;1782:197–228.
- Ai-Aql ZS, Alagil AS, Graves DT, Gerstenfeld LC, Einhorn TA. Molecular mechanisms controlling bone formation during fracture healing and distraction osteogenesis. *J Dent Res* 2008;87:107–118.
- Cheng H, Jiang W, Phillips FM, Haydon RC, Peng Y, Zhou L, Luu HH, An N, Breyer B, Vanichakarn P, Szatkowski JP, Park JY, He TC. Osteogenic activity of the fourteen types of human bone morphogenetic proteins (BMPs). *J Bone Joint Surg Am* 2003;85:1544–1552.
- Termaat MF, Den Boer FC, Bakker FC, Patka P, Haarman HJ. Bone morphogenetic proteins. Development and clinical efficacy in the treatment of fractures and bone defects. *J Bone Joint Surg Am* 2005;87:1367–1378.
- White AP, Vaccaro AR, Hall JA, Whang PG, Friel BC, McKee MD. Clinical applications of BMP-7/OP-1 in fractures, nonunions and spinal fusion. *Int Orthop* 2007;31:735–741.
- Akune T, Ohba S, Kamekura S, Yamaguchi M, Chung UI, Kubota N, Terauchi Y, Harada Y, Azuma Y, Nakamura K, Kadowaki T, Kawaguchi HH. PPARgamma insufficiency

- enhances osteogenesis through osteoblast formation from bone marrow progenitors. *J Clin Invest* 2004;113:846–855.
19. Lin TH, Yang RS, Tang CH, Lin CP, Fu WM. PPARgamma inhibits osteogenesis via the down-regulation of the expression of COX-2 and iNOS in rats. *Bone* 2007;41:562–574.
 20. Kawaguchi H, Akune T, Yamaguchi M, Ohba S, Ogata N, Chung UI, Kubota N, Terauchi Y, Kadowaki T, Nakamura K. Distinct effects of PPARgamma insufficiency on bone marrow cells, osteoblasts, and osteoclastic cells. *J Bone Miner Metab* 2005;23:275–279.
 21. Kornberg A. Lactic dehydrogenase of muscle. In: Colowick SP, Kaplan ND, editors. *Methods of enzymology*, Vol. 1. New York, NY, USA: Academic Press 1955:441–443.
 22. Obradovi RR, Kesi LG, Peševska S. Influence of low-level laser therapy on biomaterial osseointegration: A mini-review. *Lasers Med Sci* 2008; Jun 20 [Epub ahead of print].
 23. Saltmarche AE. Low level laser therapy for healing acute and chronic wounds—The extendicare experience. *Int Wound J* 2005;5:351–360.
 24. Ivandic BT, Ivandic T. Low-level laser therapy improves vision in patients with age-related macular degeneration. *Photomed Laser Surg* 2008;26:241–245.
 25. Reis SR, Medrado AP, Marchionni AM, Figueira C, Fracassi LD, Knop LA. Effect of 670-nm Laser Therapy and Dexamethasone on tissue repair: A histological and ultrastructural study. *Photomed Laser Surg* 2008;26:307–313.
 26. De Oliveira RF, Oliveira DA, Monteiro W, Zangaro RA, Magini M, Soares CP. Comparison between the effect of low-level laser therapy and low-intensity pulsed ultrasonic irradiation in vitro. *Photomed Laser Surg* 2008;26:6–9.
 27. Zambuzzi WF, Yano CL, Cavagis AD, Peppelenbosch MP, Granjeiro JM, Ferreira CV. Ascorbate-induced osteoblast differentiation recruits distinct MMP-inhibitors: RECK and TIMP-2. *Mol Cell Biochem* 2009;322:143–150.
 28. Karu TI. Local pulsed heating of adsorbing chromophores as a possible primary mechanism of low-power laser effects. In: Galletti G, Bolognini L, Ussia G, editors. *Laser applications in medicine and surgery*. Bologna: Monduzzi Editore 1992: 253.
 29. Lubart R, Eichler M, Lavi R, Friedman H, Shainberg A. Low-energy laser irradiation promotes cellular redox activity. *Photomed Laser Surg* 2005;23:3–9.
 30. Rinaldi M, Barrera G, Spinsanti P, Pizzimenti S, Ciafrè SA, Parella P, Farace MG, Signori E, Dianzani MU, Fazio VM. Growth inhibition and differentiation induction in murine erythroleukemia cells by 4-hydroxynonenal. *Free Radic Res* 2001;34:629–637.
 31. Dianzani MU, Barrera G, Parola M. 4-Hydroxy-2,3-nonenal as a signal for cell function and differentiation. *Acta Biochim Pol* 1999;46:61–75.
 32. Tschoep K, Hartmann G, Jox R, Thompson S, Eigler A, Krug A, Erhardt S, Adams G, Endres S, Delius M. Shock waves: A novel method for cytoplasmic delivery of antisense oligonucleotides. *J Mol Med* 2001;79:306–313.
 33. Karu T, Tiphlova O, Esenaliev R, Letokhov V. Two different mechanisms of low-intensity laser photobiological effects on *Escherichia coli*. *J Photochem Photobiol B* 1994;24:155–161.
 34. Arany PR, Nayak RS, Hallikerimath S, Limaye AM, Kale AD, Kondaiah P. Activation of latent TGF-beta1 by low-power laser in vitro correlates with increased TGF-beta1 levels in laser-enhanced oral wound healing. *Wound Repair Regen* 2007;15:866–874.
 35. Matsui S, Takeuchi H, Tsujimoto Y, Matsushima K. Effects of Smads and BMPs induced by Ga-Al-As laser irradiation on calcification ability of human dental pulp cells. *J Oral Sci* 2008;50:75–81.
 36. Muzio G, Martinasso G, Trombetta A, Di Simone D, Canuto RA, Maggiora M. HMG-CoA reductase and PPARalpha are involved in clofibrate-induced apoptosis in human keratinocytes. *Apoptosis* 2006;11:265–275.
 37. Muzio G, Trombetta A, Maggiora M, Martinasso G, Vasiliou V, Lassen N, Canuto RA. Arachidonic acid suppresses growth of human lung tumor A549 cells through down-regulation of ALDH3A1 expression. *Free Radic Biol Med* 2006;40:1929–1938.
 38. Hasegawa T, Oizumi K, Yoshiko Y, Tanne K, Maeda N, Aubin JE. The PPAR gamma-selective ligand BRL-49653 differentially regulates the fate choices of rat calvaria versus rat bone marrow stromal cell populations. *BMC Dev Biol* 2008; 8:71.
 39. Wahli W. PPAR gamma: Ally and foe in bone metabolism. *Cell Metab* 2008;7:188–190.
 40. Wan Y, Chong LW, Evans RM. PPAR-gamma regulates osteoclastogenesis in mice. *Nat Med* 2007;13:1496–1503.
 41. Wakino S, Kintscher U, Kim S, Yin F, Hsueh WA, Law RE. Peroxisome proliferator-activated receptor gamma ligands inhibit retinoblastoma phosphorylation and G1 → S transition in vascular smooth muscle cells. *J Biol Chem* 2000;275: 22435–22441.
 42. Martinasso G, Oraldi M, Trombetta A, Maggiora M, Bertetto O, Canuto RA, Muzio G. Involvement of PPARs in cell proliferation and apoptosis in human colon cancer specimens and in normal and cancer cell lines. *PPAR Res* 2007;2007: 93416.
 43. Muzio G, Trombetta A, Maggiora M, Martinasso G, Vasiliou V, Lassen N, Canuto RA. Arachidonic acid suppresses growth of human lung tumor A549 cells through down-regulation of ALDH3A1 expression. *Free Radic Biol Med* 2006;40:1929–1938.
 44. Muzio G, Maggiora M, Trombetta A, Martinasso G, Reffo P, Colombatto S, Canuto RA. Mechanisms involved in growth inhibition induced by clofibrate in hepatoma cells. *Toxicology* 2003;187:149–159.