AperTO - Archivio Istituzionale Open Access dell'Università di Torino

Site and stand effects on coarse woody debris in montane mixed forests of Eastern Italian Alps

This is a pre print version of the following article:

Original Citation:

Availability:

This version is available <http://hdl.handle.net/2318/81368> since 2017-05-25T12:42:42Z

Published version:

DOI:10.1016/j.foreco.2010.08.008

Terms of use:

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)

This is the author's final version of the contribution published as:

[Castagneri D., Garbarino M., Berretti R., Motta R. 2010. Site and stand effects on coarse woody debris in montane mixed forests of Eastern Italian Alps. Forest Ecology and Management 2010. ISSN: 0378-1127, doi: 10.1016/j.foreco.2010.08.008]

The publisher's version is available at:

[<http://www.sciencedirect.com/science/article/pii/S0378112710004548>]

When citing, please refer to the published version.

Link to this full text:

[<http://hdl.handle.net/2318/81368>]

This full text was downloaded from iris-AperTO: <https://iris.unito.it/>

**Legacies of past forest management on present deadwood in montane mixed forests of
Eastern Italian Alps**

Daniele Castagneri ^{a*}, Matteo Garbarino ^a, Roberta Berretti ^a, Renzo Motta ^a

^a Department of AgroSelviTer, University of Turin, I-10095 Grugliasco (TO), Italy

*Corresponding author address: via Leonardo da Vinci 44 10095 Grugliasco (TO)

e-mail: daniele.castagneri@unito.it telephone number: 00390116705549

fax: 00390116705546

M. Garbarino: matteo.garbarino@unito.it

R. Berretti: roberta.berretti@unito.it

R. Motta: renzo.motta@unito.it

35 **Abstract**

36 The role of deadwood on biodiversity conservation of forest ecosystems is widely recognised.
37 Interest on deadwood has increased in the last years, and forest management policy regards
38 deadwood as indicator of sustainable forest management.

39 This study took place in mixed montane forests in Eastern Italian Alps. The objective was to
40 determine how past forest management, topography and forest structure influence deadwood
41 accumulation. 124 sampling points were established in four Forest Reserves, where time of non-
42 intervention ranges from 12 to more than 50 years. A multivariate analysis was performed to
43 investigate the connections between forest stand characteristics and deadwood.

44 Coarse woody debris (CWD) volume in the reserves was similar to other recently-unmanaged
45 forests in central Europe. Both stand characteristics and topographic factors determined CWD
46 distribution. Basal area of living trees and human impact emerged as the most important factors.
47 These aspects are connected with the input (density-dependent mortality) and the output
48 (harvesting) of deadwood in the stand. In the next decades we expect an increase of deadwood, due
49 to density-dependent mortality and disturbances. However, many decades in absence of human
50 interventions are probably required to reach amount of deadwood similar to those in old-growth
51 forests.

52

53

54

55

56

57

58 **Keywords** Alps; coarse woody debris; human impact; mixed forest; montane forest; multivariate
59 analysis.

60

61

62

63

64

65

66

67

68

1. Introduction

Deadwood is an important component in the functioning of the forest ecosystem, as it plays an important role in biodiversity, trophic chains, forest natural regeneration, nutrient cycles and overall carbon storage (Harmon et al., 1986; Franklin et al., 1987; Jonsson and Kruys, 2001; Laiho and Prescott, 2004; Luyssaert et al., 2008). During the last decades research focused on the assessment of deadwood amount in forests have been common in North America (Spies and Franklin, 1988; McCarthy and Bailey, 1994; Sturtevant et al., 1997; Clark et al., 1998) and northern Europe (Sippola et al., 1998; Jonsson, 2000; Siitonen et al., 2000; Krankina et al., 2002). However, in central and southern Europe deadwood has received less attention (Bretz Guby and Dobbertin, 1996; Marage and Lemperiere, 2005; Motta et al., 2006; Lombardi et al., 2008).

In central and southern Europe human pressure has affected forest dynamics since prehistoric times (Farrel et al., 2000; Motta and Nola, 2001; Winter et al., 2010), peaking in the last century. In most regions of the Alps anthropogenic impact has been very severe because forests have been used for timber, fuel wood, forest litter, deadwood, branches and even small twigs collection and were subjected to grazing (Bürge and Gimmi, 2007; Gimmi et al., 2008). During recent decades the anthropogenic impact on forests has considerably diminished, and in eastern Italian Alps a large part of forests have been withdrawn from regular management since the early 1950s (Martinis, 1990; Farrell et al., 2000; Lehringer et al., 2003). Moreover, the public attitude towards forests and forestry has dramatically changed. Past management was concentrated on what was being extracted from the forest, whereas current management emphasizes what is being left (Kohm and Franklin, 1997). In this context, quantity and quality of coarse woody debris (CWD) are regarded as important structural indicators of naturalness and biodiversity (Corona et al., 2003; MCPFE, 2003; Jönsson and Jonsson, 2007), providing information on the intensity of past human disturbances and closeness to old-growth condition (Stokland, 2001; Woodall and Nagel, 2006).

The amount of deadwood is determined by its inputs and outputs in the forest (Siitonen, 2001).

Deadwood natural accumulation is influenced by disturbance regime, climate, tree species composition, tree size (von Oheimb et al., 2007), stand succession stage, stand structure (Siitonen et al., 2000), topography (Rubino and Mc Carty, 2003), decomposer organisms (Harmon, 2009).

Besides, forest management can affect deadwood input. Silvicultural interventions alter tree species composition and forest structure, reducing competition intensity and mortality, and removing weakened trees having the highest potential to die (Siitonen et al., 2000 Rouvinen et al., 2002).

Short rotation loggings interrupt natural stage development, preventing forest ageing and deadwood formation (Duvall and Grigal, 1999; Currie and Nadelhoffer, 2002; Vandekerckhove et al., 2009).

Moreover, a direct removal of dead wood can occur to obtain fire wood and to reduce wildfire and

103 pathogen attack risk (Wolynski, 2001). Therefore, deadwood quantities are normally lower in
 104 managed than in unmanaged forests (Gibb et al., 2005; Müller-Using and Bartsch, 2009;
 105 Vandekerckhove et al., 2009).

106 The present study was carried out in four Forest Reserves of the eastern Italian Alps located in the
 107 montane belt (1000-1600 m a.s.l.) where mature stands have been left unmanaged between 12 and
 108 50 years. The vegetation type is a mixed forest of European beech (*Fagus sylvatica* L.), silver fir
 109 (*Abies alba* Mill.) and Norway spruce (*Picea abies* (L.) Karst.). Even if past anthropogenic impact
 110 has affected forest structure the native species are still present and regenerate naturally.

111 The aim of our analysis was to assess the legacies of past logging and stand characteristics on
 112 deadwood. Main research questions were: (1) what is the current volume of deadwood in mixed
 113 Fagus-Abies-Picea forests withdrawn from regular management in eastern Italian Alps? (2) What is
 114 the influence of former management, topography, and forest structure on quantity and quality of
 115 deadwood?

116

117 **2. Methods**

118 **2.1. Study areas**

119 The study took place in four Forest Reserves located in eastern Italian Alps (Val Novarza, Val
 120 Pontebbana and Col Piova in Friuli-Venezia Giulia Region, Ludrin in Trentino-Alto Adige Region,
 121 Fig. 1) with an elevation range from 1020 to 1630 m a.s.l. The predominant soil type is cambisol,
 122 developed on carbonatic substrate except for Val Pontebbana where a silicic substrate was
 123 dominant. Climate regime is similar among the reserves. Val Novarza and Val Pontebbana belong
 124 to “mesalpic district” with mean annual temperature averaging 8 °C, and annual precipitation
 125 averaging 1600 mm (Del Favero et al., 1998). Ludrin has a similar mean annual temperature, while
 126 annual precipitation differs from the other reserves averaging 1300 mm. Col Piova lies in the
 127 “esalpic district”, with mean annual temperature of 11 °C and precipitation of 1700 mm/year (Del
 128 Favero et al., 1998). In all the study sites the precipitation peaks are in spring (May - June) and
 129 autumn (November).

130 Stands are characterized by Norway spruce (*Picea abies* (L.) Karst.), silver fir (*Abies alba* Mill.)
 131 and European beech (*Fagus sylvatica* L.), representing the main species in Italian Alps montane
 132 forests. Other species (*Larix decidua* Mill., *Acer pseudoplatanus* L., *Sorbus aucuparia* L., and other
 133 montane broadleaf species) occur sporadically.

134 Val Novarza Forest Reserve (37 ha; latitude 46°27' N; longitude 12°46' E) is located at an altitude
 135 from 1300 to 1570 m a.s.l, and predominant aspect is west and north-west. Intensive selection
 136 cutting involving both beech and conifers occurred during 1940s, ending in 1953-1955 (forest

management plan archive). No other activities were recorded in successive period, apart from a 37.5 m³ cut in 1997 in the lower part of the reserve. Val Pontebbana Forest Reserve (37.6 ha; latitude 46°32' N; longitude 13°10' E) is east exposed, and altitude varies from 1240 to 1630 m a.s.l. The upper part was managed with group or individual selection cutting, and the last logging was conducted in the early 1960s (forest management plan archive). In the central part, last cutting was performed in 1982, while in the lower part intense selection cutting have been made until 1996. Col Piova Forest Reserve (36 ha; latitude 46°04' N; longitude 12°26' E) is predominantly west and north-west exposed, and altitude varies from 1020 to 1200 m a.s.l. At higher elevation, cutting brought to an end in 1960, while the rest of the forest was logged until 1989. Ludrin Forest Reserve (26.5 ha; latitude 46°07' N; longitude 10°56' E) occupies a part of a small valley north-south oriented, at an altitude from 1250 to 1350 m a.s.l. Several cutting occurred during 1950s, and the last silvicultural operations were performed in 1962. Comparison of ipsometric curves (tree diameter vs. tree height) of the three principal species showed a similar site productivity among the four reserves (data not shown). Slightly higher dominant height at Col Piova was probably related to lower elevation.

152

153 Fig.1.

154

155 2.2. Field methods

156 Sampling points were located on a 100x100 m regular grid superimposed on each reserve. 33 points
157 were identified within Val Novarza, 33 in Val Pontebbana, 36 in Col Piova, and 22 in Ludrin Forest
158 Reserve (Figure 1). Field data were collected between 2005 and 2007. In correspondence of each
159 sampling point, three type of measurement were applied: (1) a circular sampling plot (12-m radius,
160 at Val Novarza, Val Pontebbana, Col Piova,) or a squared plot (30-m side, at Ludrin) for live trees
161 measurement, (2) two perpendicular rectangular plots (50 x 8 m) for the stumps and the snags, and
162 (3) two perpendicular 50 m long transects for line intersect sampling (LIS) for the logs (Van
163 Wagner, 1968).

164 Snags were defined as standing dead trees having diameter at breast height (dbh) > 7 cm and height
165 > 1.3 m, and stumps were wood pieces with diameter at the top > 7 cm and height < 1.3 m. Logs
166 were stems, pieces of stem or branches laying on the ground having at least 5 cm diameter and
167 length > 1 m.

168 For all live trees and snags with dbh > 7 cm, dbh and height were measured. The number of stumps
169 was recorded. The measurement of logs consisted of the diameter at each intersection point (LIS

method). The decay stage of logs and snags was classified according to a class system from 1 (slightly decayed) to 5 (very decayed) (see Motta et al. 2006 for decay class description).

2.3. Stand and CWD descriptors

Several stand characteristics describing forest structure, human disturbance, and topography have been considered as possible factors (explanatory variables) influencing CWD quantity and quality (focus variables).

In each sampling plot, the following forest structure descriptors were calculated: tree density ($n \cdot ha^{-1}$), basal area (BA) ($m^2 \cdot ha^{-1}$), mean diameter at breast height (dbh) (cm), Shannon's diversity index for tree height (THD), proportion of trees having a BA bigger than the mean BA tree (%), BA proportions of the three principal species (%) (Drobyshev et al., 2008; Smirnova et al., 2008).

Human disturbance was evaluated through number of stumps ($n \cdot ha^{-1}$) (all stumps were considered anthropogenic) and the time elapsed since last intervention derived from forest management plan archives (years). Topography descriptors were percent slope (%) and elevation (m a.s.l.) derived from a digital elevation model (10-m resolution) using ArcGIS 8.2 (ESRI Inc.).

Deadwood constituted by snags and logs was referred as coarse woody debris (CWD). Stumps were not included in CWD, since they were considered as indicators of human impact (Rouvinen et al., 2002). The volume ($m^3 \cdot ha^{-1}$) of logs of each decay class and total was calculated using Van Wagner's formula (1968). The volume of standing dead trees (in classes and total) was estimated from yield tables, while the volume of the broken snags was estimated as a frustum of a cone (Motta et al., 2006). CWD was computed as logs and snags volume (CWD Tot), logs volume, snags volume, 1 and 2 CWD decay classes volume (CWD 1), 3, 4, and 5 CWD decay classes volume (CWD 2). To evaluate the occurrence of recently formed CWD in proportion to the total CWD volume, we calculated CWD ratio as the percent ratio between CWD 1 and CWD Tot.

2.4. Statistical analyses

Relationships between forest structure, human disturbance, topography and CWD were investigated adopting a multivariate approach. Redundancy analysis (RDA) was employed to explore relationships among all stand descriptors (explanatory variables) and CWD descriptors (focus variables) (Wimberly and Spies, 2001). Principal component analysis (PCA) was used to summarize CWD variability in few uncorrelated variables. Afterwards, a path analysis was employed to investigate the cause-and-effect relationships between CWD (expressed as PCs) and the most important explanatory variables (Shipley, 2000; Brais et al., 2005).

Prior to multivariate analyses, normality distribution of parameters was assessed and outlier analysis was performed using PcOrd 5 statistical package (McCune and Mefford, 1999). Each dataset was relativized by the standard deviate in order to put variables, that were measured in different units, on an equal footing (McCune and Grace, 2002).

All explanatory and focus variables calculated for each plot were included in the RDA matrix. This direct gradient analysis was performed using Canoco (Ter-Braak and Šmilauer, 1998) and the statistical significance of the relation between CWD and the explanatory variables was tested by a Monte Carlo test (9999 permutations).

A data matrix including six focus variables was processed to summarize CWD variability in fewer uncorrelated variables. PCA was performed using PcOrd 5 statistical package (McCune and Mefford, 1999) and statistical significance of the axes was tested by a Monte Carlo test (9999 permutations). Moreover, Pearson's correlation between explanatory variables and the principal components (PCs) was calculated to find out explanatory variables more related to CWD variability. Variables with Pearson's r over 0.15 (absolute value) were selected.

A path analysis, which is a specialized version of Structural Equation Models (Shipley, 2000), was employed to develop a model describing the influence of explanatory variables on quantity and quality of deadwood (expressed as PCs). A conceptual path model including variables selected by Pearson's correlation analysis was built under the underlying concept that different stand characteristics interact together to determine CWD in the reserves (Fig. 2). Afterwards, alternative models were tested considering a subset of variables to obtain a statistically significant model (Garbarino et al., 2009). Quantitative model comparisons used a combination of Akaike's Information Criterion (AIC) statistic and the Root Mean Square Error of Approximation (RMSEA). The latter is a goodness-of-fit index that is relatively independent of sample size. A model with $RMSEA < 0.06$ was considered a good fit (Hu and Bentler, 1999). All such models were tested and the models with the smallest AIC statistic were selected as the most parsimonious models (Hu and Bentler, 1999). Path analyses were conducted using Mx software that works with covariance matrices as input data and a maximum likelihood (ML) fit function (Neale, 1994).

Fig. 2.

3. Results

3.1. Amount of CWD in the reserves

Live tree volume was greatest at Val Novarza Reserve ($594.9 \text{ m}^3 \text{ ha}^{-1}$), while CWD volume was greatest at Ludrin ($68.4 \text{ m}^3 \text{ ha}^{-1}$). Col Piova showed the lowest values both for live and dead trees

(Table 1). However, values varied considerably among sample plots into the reserves, and the coefficient of variation ranged from 70% (Val Pontebbana) to 120% (Col Piova). A total absence of CWD was observed on one plot in Ludrin, and two plots in Col Piova. Based on outlier analysis, two plots involved in an uprooting episode in Val Novarza (CWD volume: 787 and 330 m³ ha⁻¹) were excluded from analyses. Considering all reserves as a whole, volume contribution of snags and logs to CWD was similar (46.5 and 53.5% respectively), but snags prevailed (59.1%) at Ludrin and logs prevailed (75.0%) at Col Piova.

Table 1

Stand characteristics and CWD volume in the Forest Reserves.

3.2. Multivariate analyses of CWD and its anthropogenic and environmental relationships

RDA was used to relate deadwood data to anthropogenic and environmental data. Monte Carlo permutations (n = 9999) indicated relations between variables being statistically significant (p = 0.01). The first RDA axis explained 11.7 % of variance in CWD data, and the CWD-environment correlation for the first axis was 0.452. CWD Tot, CWD 1 and CWD 2 emerged to be correlated to each other, and were positively associated to stand density, elevation and basal area (Fig. 3). Volume of snags was highly related to density, while logs volume was related to slope. The number of stumps and management variables, both indicators of anthropogenic disturbance, were correlated to each other, and were negatively related to total volume of CWD, density, BA and elevation. Species BA proportions were not related to CWD variables. Beech proportion was higher at low elevation, and spruce at higher elevation. CWD ratio, a qualitative variable, seemed to be uncorrelated to other parameters.

PCA reduced CWD measures into uncorrelated components that explained most of the variation in the original dataset. The first two principal components explained 87.2% of the variation. PC 1 extracted 59.4% of the variation in the dataset, and was significantly associated with quantitative CWD variables, particularly with CWD Tot (Table 2). PC 2 extracted a lower percentage of variation (27.7%), and it was associated with the qualitative CWD variable, i.e. CWD ratio. Both axes were highly significant (p = 0.0001, Monte Carlo test). BA, density, THD, and elevation were negatively correlated with the first component (PC 1) (Table 3). Management, proportion of beech and number of stumps were negatively correlated to PC 1. Weak correlations (r < 0.15) with the second principal component (PC 2) were found.

Fig. 3.

271 Table 2
272 Principal component loadings for the first five axes for the four reserves.

273
274 Table 3
275 Pearson's correlation coefficients of the explanatory variables with the first 2 ordination axes
276 (principal components).

277

278 **3.3. Causal model for CWD**

279 The conceptual model was used to derive alternative path diagrams for two synthetic descriptors of
280 deadwood derived from the PCAs: CWD quantity (PC 1) and quality (PC 2). Seven explanatory
281 variables on 12 were included in the first path model (Fig. 2). Since PC 2 explained a lower
282 percentage of variation compared to PC 1, and no variables had r value over 0.15 with it, we did not
283 perform a model to predict PC 2. A model emerged as having significant support ($RMSEA < 0.001$;
284 $AIC = -1.755$) and included a topographic (elevation), a forest structure (basal area) and a human
285 disturbance (management) variable (Fig. 4). CWD was positively influenced ($\beta = 0.22$) by basal
286 area of live trees, but was negatively ($\beta = -0.20$) associated with management. The model included
287 the interaction of topographic and anthropogenic influences in that the negative effect ($\beta = -0.58$) of
288 elevation on human disturbance (management) was explicitly represented. Moreover elevation was
289 positively ($\beta = 0.31$) associated with basal area.

290

291 Fig. 4.

292

293 **4. Discussion**

294 **4.1. Disturbance regime and amount of CWD in the reserves**

295 Connections between disturbance history and forest structure are critical for understanding
296 ecological processes in the forest ecosystem (Bellemare et al., 2002; Foster et al., 2003; Gimmi et
297 al., 2008; Fraver et al., 2009). The quantity and quality of deadwood can provide information on
298 mortality processes and disturbance regime. Moreover, they can suggest the degree of forest
299 naturalness, and indicate the proximity to the old-growth stage (Stokland, 2001; Woodall and
300 Nagel, 2006; Winter et al., 2010).

301 In mixed temperate southern European forests, the natural disturbance regime mostly results in
302 individual-tree death or small-scale disturbances caused by wind, insects, and fungi, while large-
303 scale disturbances occur seldom (Nagel and Diaci, 2006; Firm et al., 2009; Kenderes et al., 2009).
304 In the studied reserves, only 2 out of 124 plots had high amounts of deadwood, reflecting the

305 absence of recent catastrophic disturbances. Distribution of deadwood in the reserves was related to
306 single-tree mortality and small-scale wind disturbances.

307 In the eastern Italian Alps many forests have been left unmanaged in last decades due to social and
308 economical changes. Their dynamics are presently influenced by autogenic and allogenic
309 disturbances, but the current structure results from land use management history. Montane mixed
310 forests analysed herein have been recently (from 12 to 50 years ago) withdrawn from regular
311 management, thus an effect of former management on deadwood accumulation was expected.
312 The total CWD volume found in study reserves was comparable or slightly lower than other
313 recently-unmanaged mixed forests in central Europe (Bretz Guby and Dobbertin, 1996;
314 Vandekerkhove et al., 2009). However montane mixed Fagus-Abies-Picea old-growth forests have
315 much more CWD, generally over 200 m³ ha⁻¹ (Vrška et al., 2001; Christensen et al., 2005; Motta et
316 al., 2008). Our results suggest cessation of management for even 50 years is insufficient for
317 accumulation of CWD comparable to old-growth forests.

318

319 **4.2. Relationships between human disturbance, topography characteristics, forest structure** 320 **and CWD components**

321 Deadwood accumulation is influenced by a number of factors, resulting in a complex correlation
322 structure between the involved variables. Thus, few studies have analyzed how different stand
323 characteristics influence deadwood in a forest stand (Hély et al., 2000; Storaunet et al., 2000).
324 In the forests analysed herein, the number of stumps and time of non-intervention were strictly
325 related to each other. In case of a lack of historical information, the number of anthropogenic
326 stumps can be used as a proxy variable of human impact (Storaunet et al., 2005). The number of
327 stumps indicates the intensity of cutting (Siitonen et al., 2000) while historical archives can
328 precisely point out the time span of non-intervention.

329 Elevation was more important than slope percentage in shaping the forest structure and CWD
330 characteristics. At higher elevation, spruce stands had a higher density of live and dead trees.
331 Recently-disturbed stands located at lower elevations had lower BA and CWD volume and were
332 dominated by beech trees.

333 Ordination analyses (RDA) indicated a relationship between forest structure and CWD, since BA
334 and tree density were positively correlated with CWD quantity. This relationship probably reveals
335 the effect of density-dependent mortality. Besides, past logging activities influenced actual forest
336 structure, as BA was negatively related to management and stump density. A few studies have
337 shown tree species composition influence deadwood characteristics (Brassard and Chen, 2008).
338 Nevertheless, type or decay class distribution of CWD in the reserves was not strongly affected by

species composition, although beech proportion was slightly negative related with CWD volume, possibly due to the rapid decomposition rate of its wood (von Oheimb et al., 2007). However, beech proportion was higher at low elevation, where generally smaller CWD amounts occurred, and a direct effect of beech proportion on CWD accumulation can not be ascertain. Decay class distribution showed no pattern, as plots with higher quantity of low-decayed CWD had high quantity of high-decayed CWD as well. Moreover, the proportion of recently formed CWD (CWD ratio) was poorly correlated with other variables. Bretz Guby and Dobbartin (1996) found managed stands had more deadwood in higher decay stages than unmanaged stands in Switzerland. Burrascano et al. (2008) and Lombardi et al. (2008) found an opposite pattern in Central Italy. Inconsistency can be due to differences in species composition, disturbance type and decay processes (Yan et al., 2007). In contrast, the occurrence of logs and snags depended on plot characteristics. Snags were more abundant in stands having higher tree density, probably due to density-dependent mortality (Hély et al., 2000), while logs were more abundant in steep slope stands, where probability of uprooting is generally higher. Excluding anthropogenic stumps, effects of human disturbance on the type of CWD were not observed.

354

355 **4.3. Effects of past forest management on forest structure and CWD accumulation**

Clarify causal relationships that determine the accumulation of deadwood in forest ecosystems is critical for forest ecology and ecosystem management. Consistent with the majority of previous studies (e.g. Christensen et al., 2005), the path model indicated that CWD was related to the time elapsed from human intervention, and to the amount (basal area) of live trees.

The CWD accumulation reflects the cumulative balance between inputs through tree mortality and outputs through decomposition and harvesting (Harmon et al., 1986; Tinker and Knight, 2000). In the studied plots mortality was mainly due to competition or, less frequently, to individual uprooting. Since competitive mortality depends on tree density, the input of CWD was connected with the basal area of live trees.

Past management affected CWD input in the reserves. Logging activities reduced stand basal area, affecting density-dependent mortality. Moreover, suppressed, unhealthy and senescent trees with a high potential for death, representing potential sources of CWD, were generally removed. Besides, past management influenced directly the output, as CWD was generally removed during harvesting activities.

Consistent with the results of Christiansen et al. (2005) we found higher CWD quantities at higher elevation. This trend was probably related to the negative elevation effect on management intensity. Human impact on forest ecosystem is generally stronger at low elevation, due to proximity to

373 human settlement, accessibility and higher forest productivity (Garbarino et al., 2009). Stands at
374 higher elevation had higher basal area and lower human impact, and consequently higher CWD
375 volume.

376

377 **4.4. Future perspectives**

378 The majority of studies on deadwood have been carried out on natural forests influenced by
379 catastrophic disturbances (Harmon, 2009). However, in southern European montane forests,
380 individual, small-scale or, more rarely, intermediate disturbances, are dominant processes driving
381 forest dynamics (Leibundgut, 1987; Kenderes et al., 2009). Besides, past and present presence of
382 man affects all southern European forests (Winter et al., 2010). In mixed montane forests analysed
383 herein, past forest management and stand density seem to be the major aspects influencing CWD
384 accumulation. In the absence of further human interventions, we believe that in the next few
385 decades increasing density-dependent mortality and small disturbances will result a CWD
386 accumulation. However many decades will be required for accumulations of CWD comparable to
387 those in old-growth forests.

388 Quantifying deadwood dynamics is critical for modeling and managing forest ecosystems for the
389 development of old-growth conditions in southern Europe. Future studies of interactions between
390 environmental factors, human disturbance and deadwood are required, especially for those forests,
391 which have been withdrawn from regular management for long periods (e.g. 50-100 years). Such
392 forests will be increasingly common in the next several decades.

393

394

395 **Acknowledgment**

396 Funding for this research was provided by the Autonomous Region Friuli Venezia Giulia,
397 "Direzione centrale risorse agricole, naturali, forestali e montagna - Servizio selvicoltura e
398 antincendio boschivo" and by the Autonomous Province of Trento "Servizio Foreste e Fauna".
399 We are indebted to the staff of both administrations for their support during the field campaign, and
400 to Dr. Raffaello Bertinazzi, Dr. Anna Carpanelli and Dr. Pier Paolo De Biasio (Autonomous Region
401 Friuli Venezia Giulia), Dr. Alessandro Simonetti (IRF di Tolmezzo, UD), Dr. Paolo Barsottini
402 (Stazione forestale di Vittorio Veneto, TV), Dr. Felice Dorna and Dr. Roberto Zoanetti (Ufficio
403 distrettuale forestale di Tione, TN) for archival research.
404 Besides we thank Annamaria Battiston, Giuseppe Dolce, Marta Martinez, Fabio Meloni and Vito
405 for field support and Jim Long (Utah State University) for his comments and advices on the
406 manuscript and for revising language.

407 **References**

- 408 Bellemare, J., Motzkin, G., Foster, D.R., 2002. Legacies of the agricultural past in the forested
 409 present: an assessment of historical land-use effects on rich mesic forests. *J. Biogeogr.* 29(10-11),
 410 1401-1420.
- 411 Brais, S., Sadi, R., Bergeron, Y., Grenier, Y., 2005. Coarse woody debris dynamics in a post-fire
 412 jack pine chronosequence and its relation with site productivity. *For. Ecol. Manage.* 220(1-3), 216-
 413 226.
- 414 Bretz Gubi, N.A., Dobbertin, M., 1996. Quantitative estimates of coarse woody debris and standing
 415 dead trees in selected Swiss forests. *Global Ecol. Biogeogr.* 5(6), 327-341.
- 416 Brassard, B.W., Chen, H.Y.H., 2008. Effects of forest type and disturbance on diversity of coarse
 417 woody debris in boreal forest. *Ecosystems* 11(7), 1078-1090.
- 418 Burrascano, S., Lombardi, F., Marchetti, M., 2008. Old-growth forest structure and deadwood, Are
 419 they indicators of plant species composition? A case study from central Italy. *Plant Biosyst.* 142(2),
 420 313-323.
- 421 Bürgi, M., Gimmi, U., 2007. Three objectives of historical ecology, the case of litter collecting in
 422 Central European forests. *Landscape Ecol.* 22(Supplement 1), 77-87.
- 423 Christensen, M., Hahn, K., et al., 2005. Dead wood in European beech (*Fagus sylvatica*) forest
 424 reserves. *For. Ecol. Manage.* 210(1-3), 267-282.
- 425 Clark, D.F., Kneeshaw, D.D., Burton, P.J., Antos, J.A., 1998. Coarse woody debris in sub-boreal
 426 spruce forest of west- central British Columbia. *Can. J. For. Res.* 28(2), 284-290.
- 427 Corona, P., Köhl, M., Marchetti, M., 2003. Advances in forest inventory for sustainable forest
 428 management and biodiversity monitoring, Kluwer Academic Publishers, Dordrecht.
- 429 Currie, W.S., Nadelhoffer, K.J., 2002. The imprint of land-use history, Patterns of carbon and
 430 nitrogen in downed woody debris at the Harvard Forest. *Ecosystems* 5(5), 446-460.
- 431 Del Favero, R., Poldini, L., Bortoli, P.L., Lasen, C., Dreossi, G., Vanone, G., 1998. La vegetazione
 432 forestale e la selvicoltura nella Regione Friuli - Venezia Giulia. Regione autonoma Friuli-Venezia
 433 Giulia – Servizio della Selvicoltura – Volume 1. In Italian.
- 434 Drobyshev, I., Goebel, P.C., Hix, D.M., Corace, R.G., Semko-Duncan, M.E., 2008. Interactions
 435 among forest composition, structure, fuel loadings and fire history, A case study of red pine-
 436 dominated forests of Seney National Wildlife Refuge, Upper Michigan. *For. Ecol. Manage.*
 437 256(10), 1723-1733.
- 438 Duvall, M.D., Grigal, D.F., 1999. Effects of timber harvesting on coarse woody debris in red pine
 439 forests across the Great Lakes states, USA. *Can. J. For. Res.* 29(12), 1926-1934.

440 Farrell, E.P., Fuhrer, E., Ryan, D., Andersson, F., Huttel, R., Piussi, P., 2000. European forest
 441 ecosystems, building the future on the legacy of the past. *For. Ecol. Manage.* 132(1), 5-20.
 442 Firm, D., Nagel, T.A., Diaci, J., 2009. Disturbance history and dynamics of an old-growth mixed
 443 species mountain forest in the Slovenian Alps. *For. Ecol. Manage.* 257, 1893–1901.
 444 Franklin, J.F., Shugart, H.H., Harmon, M.E., 1987. Tree death as an ecological process. *Bioscience*
 445 37(8), 550-556.
 446 Foster, D., Swanson, F., Aber, I., Burke, N., Brokaw, N., Tilman, D., Knapp, A., 2003. The
 447 importance of land-use legacies to ecology and conservation. *BioScience* 53(1), 77-88.
 448 Fraver, S., White, A.S., Seymour, R.S., 2009. Natural disturbance in an old-growth landscape of
 449 northern Maine, USA. *J. Ecol.* 97, 289-298.
 450 Garbarino, M., Weisberg, P.J., Motta, R., 2009. Interacting effects of physical environment and
 451 anthropogenic disturbances on the structure of European larch (*Larix decidua* Mill.) forests. *For.*
 452 *Ecol. Manage.* 257(8), 1794-1802.
 453 Gibb, H., Ball, J.P., Johansson, T., Atlegrim, O., Hjalten, J., Danell, K., 2005. Effects of
 454 management on coarse woody debris volume and composition in boreal forests in northern Sweden.
 455 *Scand. J. For. Res.* 20(3), 213-222.
 456 Gimmi, U., Burgi, M., Stuber, M., 2008. Reconstructing anthropogenic disturbance regimes in
 457 forest ecosystems. A case study from the Swiss Rhone Valley. *Ecosystems* 11(1), 113-124.
 458 Harmon, M.E., Franklin, J.F., Swanson, F.J., et al., 1986. Ecology of coarse woody debris in
 459 temperate ecosystems. *Adv. Ecol. Res.* 15, 133-302.
 460 Harmon, M.E., 2009. Woody detritus mass and its contribution to carbon dynamics of old-growth
 461 forests. In: Wirth, C., Gleixner, G., Heimann, M. (Eds.), *Old growth forests. Function, fate, value.*
 462 Springer, Berlin. p159-190.
 463 Hély, C., Bergeron, Y., Flannigan, W.D., 2000. Coarse woody debris in the southeastern Canadian
 464 boreal forest, composition and load variations in relation to stand replacement. *Can. J. For. Res.*
 465 30(5), 674-687.
 466 Hu, L., Bentler, P., 1999. Cutoff criteria for fit indexes in covariance structure analysis, conventional
 467 criteria versus new alternatives. *Structural Equation Modeling* 6, 1–55.
 468 Jonsson, B.G., 2000. Availability of coarse woody debris in a boreal old-growth *Picea abies* forest.
 469 *J. Veg. Sci.* 11(1), 51-56.
 470 Jonsson, B.G., Kruys, N., 2001. *Ecology of woody debris in boreal forests.* Wiley-Blackwell,
 471 Hoboken, NJ.
 472 Jönsson, M.T., Jonsson, B.G., 2007. Assessing coarse woody debris in Swedish woodland key
 473 habitats, implications for conservation and management. *For. Ecol. Manage.* 242(2007), 363–373.

474 Kenderes, K., Kral, K., Vrska, T., Standovar, T., 2009. Natural gap dynamics in a Central European
 475 mixed beech-spruce-fir old-growth forest. *Ecoscience* 16, 39-47.
 476 Kohm, K.A., Franklin, J.F., 1997. Creating a forestry for the 21st Century, The science of
 477 ecosystem management, Island Press, Washington D.C.
 478 Krankina, O.N., Harmon, M.E., Kukuev Y.A., et al., 2002. Coarse woody debris in forest regions of
 479 Russia. *Can. J. For. Res.* 32(5), 768-778.
 480 Laiho, R., Prescott, C.E., 2004. Decay and nutrient dynamics of coarse woody debris in northern
 481 coniferous forests, a synthesis. *Can. J. For. Res.* 34(4), 763-777.
 482 Lehringer, S., Höchtl, F., Konold, W., 2003. Effects of land use changes and depopulation on
 483 landscape, social life and tourism - overview about the results of a case study from the Piedmont
 484 Alps in Italy. *Centralblatt für das gesamte Forstwesen* 120(1), 1-18.
 485 Leibundgut, H., 1987. Europäische Urwälder der Bergstufe. Verlag Paul Haupt, Bern, Switzerland.
 486 Lombardi, F., Lasserre, B., Tognetti, R., Marchetti, M., 2008. Deadwood in relation to stand
 487 management and forest type in Central Apennines (Molise, Italy). *Ecosystems* 11(6), 882-894.
 488 Luyssaert, S., Schulze, E.D., Börner, A., Knohl, A., Hessenmöller, M., Law, B.E., Ciais, P., Grace,
 489 J., 2008. Old-growth forests as global carbon sinks. *Nature* 455(7210), 213-215.
 490 Marage, D., Lemperiere, G., 2005. The management of snags. A comparison in managed and
 491 unmanaged ancient forests of the Southern French Alps. *Ann. For. Sci.* 62(2), 135-142.
 492 Martinis, A., 1990. Le utilizzazioni boschive nel passato. In Regione Friuli (Eds.), *Foreste, uomo,*
 493 *economia nel Friuli Venezia Giulia*. In Italian.
 494 McCarthy, B.C., Bailey, R.R., 1994. Distribution and abundance of coarse woody debris in a
 495 managed forest landscape of the central Appalachians. *Can. J. For. Res.* 24, 1317-1329.
 496 McCune, B., Mefford, M.J., 1999. PC-ORD. MjM Software Design, Gleneden Beach.
 497 McCune, B., Grace, J.B., 2002. Analysis of Ecological Communities. MjM Software Design,
 498 Gleneden Beach.
 499 MCPFE, 2003. Improved pan-European indicators for sustainable forest management, MCPFE
 500 Liason Unit, Wien.
 501 Motta, R., Nola, P., 2001. Growth trends and dynamics in sub-alpine forest stands in the Varaita
 502 Valley (Piedmont, Italy) and their relationships with human activities and global change. *J. Veg.*
 503 *Sci.* 12(2), 219-230.
 504 Motta, R., Berretti, R., Lingua, E., Piussi, P., 2006. Coarse woody debris, forest structure and
 505 regeneration in the Valbona Forest Reserve, Paneveggio, Italian Alps. *For. Ecol. Manage.* 235(1-3),
 506 155-163.

507 Motta, R., Maunaga, Z., Berretti, R., Castagneri, D., Lingua, E., Meloni, F., 2008. La Riserva
 508 forestale di Lom (Repubblica di Bosnia Erzegovina), descrizione, caratteristiche, struttura di un
 509 popolamento vetusto e confronto con popolamenti stramaturi delle Alpi italiane. *Forest@* 5(1), 100-
 510 111. In Italian.

511 Muller-Using, S., Bartsch, N., 2009. Decay dynamic of coarse and fine woody debris of a beech
 512 (*Fagus sylvatica* L.) forest in Central Germany. *Eur. J. For. Res.* 128(3), 287-296.

513 Nagel, T.A., Diaci, J., 2006. Intermediate wind disturbance in an old-growth beech–fir forest in
 514 southeastern Slovenia. *Can. J. For. Res.* 36, 629–638.

515 Neale, M.C., 1994. MxGui 3.2. Department of Psychiatry, Virginia Commonwealth University,
 516 Richmond. <http://www.vcu.edu/mx>.

517 Rouvinen, S., Kuuluvainen, T., Karjalainen, L., 2002. Coarse woody debris in old *Pinus sylvestris*
 518 dominated forests along a geographic and human impact gradient in boreal Fennoscandia. *Can. J.*
 519 *For. Res.* 32(12), 2184-2200.

520 Rubino, D.L., McCarthy, B.C., 2003. Evaluation of coarse woody debris and forest vegetation
 521 across topographic gradients in a southern Ohio forest. *For. Ecol. Manage.* 183(1-3), 221-238.

522 Shipley, B., 2000. Cause and correlation in biology, a user's guide to path analysis, structural
 523 equations and causal inference. Cambridge University Press, Cambridge.

524 Siitonen, J., Martikainen, P., Punttila, P., Rauh, J., 2000. Coarse woody debris and stand
 525 characteristics in mature managed and old-growth boreal mesic forests in southern Finland. *For.*
 526 *Ecol. Manage.* 128(3), 211-225.

527 Siitonen, J., 2001. Forest management, coarse woody debris and saprophylic organisms,
 528 Fennoscandian boreal forest as an example. *Ecol. Bull.* 49, 11-41.

529 Sippola, A.L., Siitonen, J., Kallio, R., 1998. Amount and quality of coarse woody debris in natural
 530 and managed coniferous forests near the timberline in Finnish Lapland. *Scand. J. For. Res.* 13(2),
 531 204-214.

532 Spies, T.A., Franklin, J.F., 1988. Coarse woody debris in douglas-fir forest of western Oregon and
 533 Washington. *Ecology* 69(6), 1689-1702.

534 Smirnova, E., Bergeron, Y., Brais, S., 2008. Influence of fire intensity on structure and composition
 535 of jack pine stands in the boreal forest of Quebec, Live trees, understory vegetation and dead wood
 536 dynamics. *For. Ecol. Manage.* 255(7), 2916-2927.

537 Stokland, J.N., 2001. The coarse woody debris profile, an archive of recent forest history and an
 538 important biodiversity indicator. *Ecol. Bull.* 49, 71-83.

539 Storaunet, K.O., Rolstad, J., Groven, R., 2000. Reconstructing 100-150 years of logging history in
540 coastal spruce forest (*Picea abies*) with special conservation values in central Norway. Scand. J.
541 For. Res. 15(6), 591-604.

542 Storaunet, K.O., Rolstad, J., Gjerde, I., Gundersen, V.S., 2005. Historical logging, productivity, and
543 structural characteristics of boreal coniferous forests in Norway. Silva Fenn. 39(3), 429-442.

544 Sturtevant, B.R., Bissonette, J.A., Long, J.N., Roberts, D.W., 1997. Coarse woody debris as a
545 function of age, stand structure, and disturbance in boreal Newfoundland. Ecol. Appl. 7(2), 702-
546 712.

547 Ter-Braak C.J.F., Smilauer, P., 1998. CANOCO 4. Centre for Biometry, Wageningen, the
548 Netherlands.

549 Tinker, D.B., Knight, D.H., 2000. Coarse woody debris following fire and logging in Wyoming
550 lodgepole pine forests. Ecosystems 3(5), 472-483.

551 Vandekerckhove, K., De Keersmaecker, L., Menke, N., Meyer, P., Verschelde, P., 2009. When nature
552 takes over from man. Dead wood accumulation in previously managed oak and beech woodlands in
553 North-western and Central Europe. For. Ecol. Manage. 258(4), 425-435.

554 Van Wagner, C.E., 1968. The line intersect method in forest fuel sampling. For. Sci. 14, 20-26.

555 Von Oheimb, G., Westphal, C., Hardtle, W., 2007. Diversity and spatio-temporal dynamics of dead
556 wood in a temperate near-natural beech forest (*Fagus sylvatica*). Eur. J. For. Res. 126(3), 359-370.

557 Vrška, T., Hort, L., Odehnalova, P., Horal, D., Adam, D., 2001. The Boubín virgin forest after 24
558 years (1972–1996) – development of tree layer. Journal of Forest Science 47, 2001 (10).

559 Wimberly, M.C., Spies, T.A., 2001. Modeling landscape patterns of understory tree regeneration in
560 the Pacific Northwest, USA. App. Veg. Sci. 4, 277-286.

561 Winter, S., Fischer, H.S., Fischer, A., 2010. Relative quantitative reference approach for naturalness
562 assessments of forests. For. Ecol. Manage. 259, 1624-1632.

563 Wolynski, A., 2001. Significato della necromassa legnosa in bosco in un'ottica di gestione forestale
564 sostenibile. Sherwood 7, 5-12. In Italian.

565 Woodall, C.W., Nagel, L.M., 2006. Coarse woody type, a new method for analyzing coarse woody
566 debris and forest change. For. Ecol. Manage. 227(1-2), 115-121.

567 Yan, E.R., Wang, X.H., Huang, J.J., Zeng, F.R., Gong, L., 2007. Long-lasting legacy of forest
568 succession and forest management. Characteristics of coarse woody debris in an evergreen broad-
569 leaved forest of Eastern China. For. Ecol. Manage. 252(1-3), 98-107.

573
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606

Figure captions

Figure 1. Location of the 124 sample plots in the four Forest Reserves in eastern Italian Alps.

Figure 2. Conceptual model tested through path analysis. The model includes forest structure (Basal Area, Density, Beech proportion, THD, vertical diversity), topographic (Elevation), and anthropogenic (Management, Stumps) variables. CWD refers to the first principal component (PC 1) defined as deadwood quantity.

Figure 3. Redundancy analysis ordination biplot of 122 plots in the reserves. Dotted arrows represent the biplot scores of deadwood variables (CWD Tot = total CWD volume; Logs = logs volume; Snags = snags volume; CWD1 = volume of the 1st and 2nd CWD decay classes; CWD2 = volume of 3rd, 4th and 5th CWD decay classes; CWD ratio = CWD1 / CWD Tot). Full-line arrows represent the biplot scores of forest structure, human disturbance and topography variables (BA = basal area; Big Trees = BA proportion of trees larger than the mean BA tree; DBH = mean diameter at breast height; Density = number of live trees; THD = height diversity; Aa = fir basal area proportion; Fs = beech basal area proportion; Pa = spruce basal area proportion; Management = inverse of time of non-intervention; Stumps = number of stumps; Elevation = elevation a.s.l.; Slope = percentage slope).

Figure 4. Path diagram for the studied reserves. Continuous lines, positive paths; dotted lines, negative paths; single arrow lines, causal paths; double arrow lines, covariance paths. Thickness of causal path vectors corresponds to the strength of effect. Only significant path coefficients are presented next to each path.

1

Reserve	Area	Plots	Elevation	Last cutting	Basal area	Vol live trees	Vol CWD	Vol log	Vol snag
	(ha)	(n)	(m a.s.l.)	(year)	(m ² ha ⁻¹)	(m ³ ha ⁻¹)	(m ³ ha ⁻¹)	(m ³ ha ⁻¹)	(m ³ ha ⁻¹)
Val Novarza *	37.0	31	1300-1570	1953	53.1 (17.3)	594.9 (250.3)	45.8 (41.1)	25.1 (28.0)	20.7 (25.3)
Val Pontebbana	37.6	33	1240-1630	1996	43.6 (11.2)	470.7 (130.3)	31.0 (21.7)	16.4 (18.1)	14.6 (16.8)
Col Piova	36.0	36	1020-1200	1989	35.4 (11.3)	435.3 (158.1)	22.6 (27.0)	16.9 (22.5)	5.7 (12.4)
Ludrin	26.5	22	1250-1350	1962	48.3 (12.8)	531.3 (166.0)	68.4 (57.8)	28.0 (32.8)	40.4 (32.3)
All reserves	137.1	122	1020-1630	-	44.4 (14.8)	510.7 (187.6)	39.0 (40.0)	20.9 (25.2)	18.1 (24.5)

2

3 **Table 1**

4 Stand characteristics and CWD volume in the Forest Reserves.

5 Standard deviations are indicated in parentheses. Two plots were considered outliers and excluded
6 from the Val Novarza dataset (*).

7

8

9

10

11

12

Axis	PC 1	PC 2	PC 3	PC 4	PC 5
% of variance	59.45	27.72	10.25	1.32	0.99
Logs	-0.43	0.08	-0.72	0.35	0.42
Snags	-0.43	-0.15	0.68	0.36	0.45
CWD Tot	-0.52	-0.02	0.00	0.20	-0.70
CWD 1	-0.35	-0.56	-0.10	-0.70	0.18
CWD 2	-0.48	0.31	0.08	-0.26	-0.21
CWD ratio	0.11	-0.75	-0.11	0.38	-0.24

13

14 **Table 2**

15 Principal component loadings for the first five axes for the four reserves.

16 Loadings greater than 0.5 (in absolute value) are indicated in bold.

17

	PC 1	PC 2
Density	0.25	0.06
Basal Area (BA)	0.32	0.08
Mean Diameter (DBH)	-0.08	0.01
Spruce (Pa)	0.09	0.03
Beech (Fs)	-0.19	-0.12
Fir (Aa)	0.10	0.05
Height diversity (THD)	0.18	0.09
Big trees	-0.07	0.03
Management	-0.33	-0.11
Stumps	-0.18	-0.01
Elevation	0.22	0.04
Slope	0.12	-0.03

18

19 **Table 3**

20 Pearson’s correlation coefficients of the explanatory variables with the first 2 ordination axes
21 (principal components).

22 Pearson’s *r* values greater than 0.15 (in absolute value) are indicated in bold.

23

24

25

26

Figure1
[Click here to download high resolution image](#)

Figure2

Figure3

Figure4