

AperTO - Archivio Istituzionale Open Access dell'Università di Torino

Influence of grape density and harvest date on the changes in phenolic composition, phenol extractability indices and instrumental texture properties during ripening.

This is the author's manuscript

Original Citation:

Availability:

This version is available <http://hdl.handle.net/2318/83679> since

Published version:

DOI:10.1021/jf201318x

Terms of use:

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)



UNIVERSITÀ DEGLI STUDI DI TORINO

This is an author version of the contribution published on:

Questa è la versione dell'autore dell'opera:

J. Agric. Food Chem. 2011, 59, 8796–8805, [dx.doi.org/10.1021/jf201318x](https://doi.org/10.1021/jf201318x)

The definitive version is available at:

La versione definitiva è disponibile alla URL:

[<http://pubs.acs.org/doi/full/10.1021/jf201318x>]

**Influence of Grape Density and Harvest Date on the Changes in Phenolic Composition,
Phenol Extractability Indices and Instrumental Texture Properties During Ripening**

Running title: Grape Density Effect on Phenolic and Textural Changes

LUCA ROLLE, SUSANA RÍO SEGADÉ*, FABRIZIO TORCHIO, SIMONE GIACOSA,
ENZO CAGNASSO, FABIO MARENGO, VINCENZO GERBI

DIVAPRA - Food Technology sector, University of Turin, Via L. da Vinci 44, 10095
Grugliasco (TO), Italy.

*Author to whom correspondence should be addressed. Telephone: +39 0116708558; fax:
+39 0116708549; email: susana.riosegade@unito.it.

ABSTRACT

Changes in the phenolic composition, phenol extractability indices and mechanical properties occur in grape berries during the ripening process but the heterogeneity of the grapes harvested at different ripening stages affects the reliability of the results obtained. In this work, these changes were studied in Nebbiolo grapes harvested during five consecutive weeks and then separated according to three density classes. The changes observed in chemical and mechanical parameters through the ripening process are more related to berries density than harvest date. Therefore, the winemaker has to select the flotation density according to the objective quality properties of the wine to elaborate. On the other hand, the stiffer grapes were associated with a higher accumulation of proanthocyanidins. The harder grapes provided the higher concentration and extractability of flavanols reactive to vanillin whereas the thicker ones facilitated the extraction of proanthocyanidins.

KEYWORDS: phenolic composition; phenol extractability; anthocyanins; skin hardness; skin thickness; texture analysis; red grapes.

INTRODUCTION

The phenolic composition of grapes is responsible for certain organoleptic properties intimately related to the red wine quality, particularly colour, astringency and bitterness. Taking into account that the colour is one of the most important sensory characteristics in the initial valuation of the red wine quality, anthocyanins play an important role in the consumer acceptance of a wine as they are responsible for the colour of red grapes and young wines (1). On the other hand, proanthocyanidins strongly influence the wine astringency whereas the bitterness is restricted to small flavanol molecules (2, 3). Moreover, anthocyanins can react with other phenolic compounds to produce polymeric pigments resulting in the long-term colour stability of aged red wines and in the decrease of the wine astringency (3, 4).

Anthocyanins are gradually accumulated in berry skins from veraison through the grape ripening (5, 6), malvidin-3-glucoside being the most abundant anthocyanin in almost all red grape varieties (6). However, the anthocyanin concentration may decline just before harvest and/or during over-ripening (6). Instead, proanthocyanidins are mainly accumulated in berry skins before veraison (7). The highest concentration of seed proanthocyanidins is achieved at veraison and, from this moment, they decline slowly until close to the grape ripeness but thereafter remain relatively constant (8).

Phenolic compounds are extracted from berry skins and seeds into the wine during the maceration/fermentation step and, therefore, the assessment of the anthocyanin extractability through the winemaking process is required to predict the wine colour from grape polyphenols (9, 10). Furthermore, the anthocyanin extractability varies through the grape ripening (6), as a consequence of the compositional changes occurring in the skin cell-wall during its degradation by pectolytic enzymes (11). In seeds, the histological and

histochemical modifications that occur during the fruit development also affect the ability to release phenols (8, 12).

Many studies have been performed to define the best indices to evaluate the phenol extractability from berry skins and seeds. Since the assessment of the extractability of phenolic compounds is strongly dependent on the extraction method used, the cellular maturity index (EA) and seed maturity index (MP) seem to provide an adequate robustness to predict those in the resulting wines (13, 14). Instrumental texture analysis parameters also permit the estimation of the anthocyanin extractability because the structural and chemical properties of the skin cell-walls may determine the mechanical resistance, texture and ease of processing berries (11). In particular, the berry skin break force can be considered the best mechanical parameter to estimate anthocyanin extraction kinetics with adequate reliability (15). Recently, Río Segade et al. (16) have proposed the use of the berry skin thickness to predict the anthocyanin extractability. Furthermore, the mechanical methods are inexpensive, which represents an additional advantage since it allows their application as a routine monitoring tool for the grape quality.

Most of studies on the influence of the harvest date on the grape phenolic composition and phenol extractability have been carried out without considering the physiological homogeneity of samples. To reduce the heterogeneity of the physiological characteristics corresponding to the different ripening stages, Fournand et al. (6) calibrated berries according to their density. For the first weeks after veraison, the less dense classes were selected, and for the last weeks, the denser classes were selected, so that the physiological differences between the first and the last sampling date were emphasized.

As the combined effect of both the harvest date and grape densimetric sorting was not previously studied, the aim of this work was first to investigate the changes in berry skin phenolic composition, phenol extractability indices and skin mechanical properties through

the last five ripening weeks at three different grape densities and then to establish a relationship that permits to relate the harvest date to certain chemical and physical parameters for each grape density. This knowledge could be interesting because a new technology, particularly an automatic winery equipment of berry densimetric sorting, has been recently developed and proposed for the oenological company. Since the effectiveness of the grapes selection is strongly dependent on the density of the floating solution, it should be previously optimized on the base of the chemical-physical characteristics of the grape. Finally, this approach aimed to determine whether the skin mechanical attributes may influence the phenol composition and/or extractability, irrespective of the effect of the harvest date and/or the sugar content. The study was carried out on *Vitis vinifera* L. cv Nebbiolo because it is one of the most important and well-known Italian variety whose grapes are usually used for the production of renowned red wines like Barolo and Barbaresco DOCG, which are commercialized in all the world.

MATERIALS AND METHODS

Grape samples. Grape samples of Nebbiolo red cultivar (*Vitis vinifera* L.) were collected at different physiological stages from a vineyard of 0.5 ha located in La Morra within the Cuneo province of Piedmont (North-West Italy) during five consecutive weeks in 2009. About 12 kg of grape berries for each sampling date were randomly picked with attached pedicels from 500 vines by picking the berries one by one and/or for bunch (three or four berries) in each cluster. The berries were separated according to their density, which was estimated by flotation in different saline solutions (from 100 to 190 g/L sodium chloride) (6, 17, 18). These solutions had densities between 1069 and 1125 kg/m³. The berries were introduced into the less dense solution and ‘floating’ berries were considered to have the same density as the solution. These berries were separated from those which sank and they

were counted. Berries which sank were removed and introduced into the next denser solution. The same process was applied for all saline solutions. For each harvest date, the following density classes were studied: A = 1088 kg/m³, B = 1094 kg/m³, C = 1100 kg/m³, as they are the densities to which most of berries belong. The ‘floating’ berries were washed with water, visually inspected before analysis and those with damaged skins were discarded.

For both density class and harvest date, a sub-sample of 30 sorted berries was used for the determination of the physical and mechanical properties. Three sub-samples of 20 sorted berries were used for the determination of the skin phenolic composition and relative extractability. Another two sub-samples of 200 berries were used for the determination of the cellular maturity and the seed maturity indices. The remaining berries, subdivided in three replicates, were used for determining standard physicochemical parameters in the grape must obtained by manual crushing and filtration.

Reagents and standards. Solvents of HPLC-gradient grade and all other chemicals of analytical-reagent grade were purchased from Sigma (Milan, Italy). The solutions were prepared in deionized water produced by a Purelab Classic system (Elga Labwater, Marlow, United Kingdom). Anthocyanin standards (Delphinidin-3-O-glucoside chloride, Malvidin-3-O-glucoside chloride, Petunidin chloride, Peonidin-3-O-glucoside chloride, Cyanidin-3-O-glucoside chloride) were supplied by Extrasynthèse (Genay, France). All the standards were stored at -20 °C away from light before use.

Physical and technological maturity parameters. Reducing sugars, pH and total acidity were determined according to International Organization of Vine and Wine (OIV) methods. The length between top and bottom sides (*L*) and length between both lateral

sides at middle of berry height (\hat{l}) were measured using a calliper which had an accuracy of 0.1 mm. Volume was then calculated comparing the berry form to an ellipsoid, following the equation 1 (16):

$$\text{Volume (cm}^3\text{)} = 4 \pi a b c/3 \quad (1),$$

where $a = b = \hat{l}/2$, $c = L/2$.

Cellular maturity and seed maturity indices (Glories' indices). The phenol extractability indices were assessed in accordance with the procedure proposed by Glories and Saint-Criq (19), which was slightly modified for Nebbiolo grapes (13). Two replicates of 200 grape berries were used. The following parameters were determined in both pH 1 and pH 3.2 solutions: total phenolic content (A_{280}), total anthocyanins (A_1 and $A_{3.2}$), total flavonoids (TF_1 and $TF_{3.2}$) and non-anthocyanin flavonoids (FNA_1 and $FNA_{3.2}$) (13, 20). The cellular maturity index (EA) and the seed maturity index (MP) were calculated following the equations 2 and 3, respectively (9, 13, 14):

$$EA (\%) = (A_1 - A_{3.2}) / A_1 \times 100 \quad (2)$$

$$MP (\%) = (A_{280} - ((A_{3.2} / 1000) \times TAR)) / A_{280} \times 100 \quad (3)$$

The average ratio (TAR) between total phenols (A_{280}) and total anthocyanins in grape skins was 70 for Nebbiolo grapes when $A_{3.2}$ was expressed as g/L (13).

Skin phenolic composition and extractability

Extraction. Three replicates of 20 berries for each density class and harvest date were weighed before phenolic extraction. The berry skins were manually removed from the pulp. Afterwards, they were quickly immersed in 75 mL of a buffer solution containing 12 % (v/v) ethanol to simulate the extraction conditions during industrial production, 100 mg/L sodium metabisulphite to limit the oxidation of phenolic compounds (6), 50 mg/L

sodium azide and 5 g/L tartaric acid. The pH value was adjusted to 3.20 by the addition of 1 mol/L sodium hydroxide (20). They were then introduced in a controlled temperature room at 25 °C for 48 h and the supernatant was used for determining easily extracted phenols (solution A) (21). Residual berry skins were rinsed with the hydroalcoholic solution and quickly immersed in 75 mL of a new hydroalcoholic buffer containing a higher sodium metabisulphite concentration (600 mg/L). After homogenizing at 8000 rpm for 1 min with an Ultraturrax T25 high-speed homogenizer (IKA Labortechnik, Staufen, Germany), the extract was centrifuged in a PK 131 centrifuge (ALC International, MI, Italy) for 10 min at 3000×g at 20 °C. The supernatant was then used for determining non-easily extracted phenols (solution B). The total extractable phenol content in berry skins (for each parameter evaluated) was calculated as A + B and expressed as mg/kg grapes, while the extractability yield was calculated as A / (A+B) and expressed as percentage (%) (21).

Spectrophotometric methods. Phenolic compounds in the berry skin extracts were determined by spectrophotometric methods (20) using a UV-1601PC spectrophotometer (Shimadzu Scientific Instruments Inc., Columbia, MD, USA). Total anthocyanins (TA_{sk}) were expressed as malvidin-3-glucoside chloride while flavanols reactive to vanillin (flavanols vanillin assay, FVA_{sk}) and total flavonoids (TF_{sk}) were expressed as (+)-catechin. Proanthocyanidins (PRO_{sk}) were determined after acid hydrolysis with warming (Bate-Smith reaction) using a ferrous salt (FeSO₄) as catalyst. They were expressed as cyanidin chloride.

HPLC method. Anthocyanin profile was performed after submitting the berry skin extract to phase-solid extraction using a SEP-PAK C₁₈ cartridge (Waters Corporation, Milford,

MA, USA), methanol being the eluent. The chromatography system employed was a P100 pump equipped with an AS3000 autosampler (Spectra Physics Analytical, Inc., San Jose, CA, USA), a 20-mL Reodyne sample loop, a LiChroCART analytical column (25 cm \times 0.4 cm i.d.) purchased from Merck (Darmstadt, Germany) which is packed with LiChrosphere 100 RP-18 (5 μ m) particles supplied by Alltech (Deerfield, IL, USA) and a Spectra Focus Diode Array Detector (DAD, Spectra Physics Analytical, Inc., San Jose, CA, USA) operating at 520 nm. The following mobile phases were used: A = 10 % (v/v) formic acid in water; B = 10 % (v/v) formic acid and 50 % (v/v) methanol in water. All the solvents were filtered through a 0.20 μ m filter. The mobile phase flow-rate was 1 mL/min. The following solvent A proportions were used: from 72 to 55 %, 15 min; to 30 %, 20 min; to 10 %, 10 min; to 1 %, 5 min; to 72 %, 3 min. An equilibrium time of 10 min was selected (20). The data treatment was carried out using the ChromQuestTM chromatography data system (ThermoQuest, Inc., San Jose, CA, USA). The identification of the free forms of anthocyanins in berry skin extracts was performed by comparison with external standards. The acylated forms of anthocyanins were identified by matching the DAD spectrum and retention time of each chromatographic peak, and by comparing these with data available in the literature (22). Individual anthocyanins were expressed in percentages.

Skin mechanical properties. A Universal Testing Machine (UTM) TAxT2i□ Texture Analyzer (Stable Micro System, Godalming, Surrey, UK) equipped with a HDP/90 platform and a 5 kg load cell was used. The operating conditions applied and the mechanical properties measured in skins (sk) are shown in **Table 1**. All the data acquisitions were made at 400 Hz and data were evaluated using the Texture Expert Exceed software version 2.54 for Windows 2000. For each berry weighed and measured, the skin hardness was assessed by a puncture test (23). 30 berries were placed on the metal

plate of the UTM with the pedicel in a horizontal plane in order to be consistently punctured in the lateral face. The measurement of the skin thickness required the manual separation of a piece of skin (ca. 0.25 cm^2) from the lateral side of each berry with a razor blade and its subsequent drying with adsorbent paper. Care was taken in removing the pulp from the skin and in positioning the skin sample on the UTM platform to prevent folds in the skin (16). Furthermore, it was convenient to insert an instrumental trigger threshold equal to 0.05 N that enabled the plane surface of the probe to adhere completely to the skin sample before the acquisition started. This allowed a reduction or elimination of the ‘tail’ effect due to the postponement of the contact point (23). Before each test, the instrument was calibrated for force and distance.

The hardness of the berry skin is assessed by the maximum break force (F_{sk}), by the break energy (W_{sk}) or by the material resistance to the axial deformation (E_{sk}). The first variable corresponds to the resistance to the needle probe penetration while the second variable is represented by the area under the force/time curve, which is limited to between 0 and F_{sk} (23). The third one is defined as the slope of the stress-strain curve in the linear section and measures the stiffness of the skin to a load applied (23, 24). The berry skin thickness (Sp_{sk}) is given by the distance between the point corresponding to the probe contact with the berry skin (trigger) and the platform base during the compression test (16).

Statistical analysis. Statistical analyses were performed using the statistical software package SPSS (version 17.0; SPSS Inc., Chicago, IL, USA). The Tukey-b test for $p < 0.05$ was used in order to establish statistical differences by one-way analysis of variance (ANOVA). Pearson correlation coefficients were calculated to determine significant correlations.

RESULTS AND DISCUSSION

The distribution percentage of Nebbiolo grape berries in different density classes at five harvest dates is reported in **Figure 1**. It is important to note that not all the density classes had the same contribution depending on the grape ripeness stage. As expected, the lower berries density (A) made up the majority in the less ripe grapes (harvest date I) whereas the contribution of the higher one (C) increased with the grape ripeness and, therefore, with the harvest date. The contribution of grapes belonging to the A, B and C density classes ranged as follows 6.1-38.7, 21.4-39.2 and 6.2-47.0 %, respectively, depending on the harvest date. The distribution of the berries in the vineyard based on the density is already present at the beginning of the ripening process and it changes during its advancement. Therefore, a non-negligible heterogeneity occurs through the ripening process. Consequently, this heterogeneity implies that a considerable percentage of unripe grapes are harvested and used to elaborate wine. Since unripe grapes provide a lower sugar content, higher acidity, fewer anthocyanins and in particular more seed tannins, their presence can increase bitterness and astringency affecting the final wine quality adversely (18).

Physical and technological maturity parameters. Table 2 shows the physical and technological maturity parameters for the three density classes at five grape ripeness stages of Nebbiolo variety. Many differences were present in the physical characteristics among both the density classes and the harvest dates. This effect was particularly significant in the first harvest date, the grapes richer in sugars being the smaller and lighter ones. Smaller berries have a relatively higher solute to solvent ratio than larger berries and, therefore, it is widely known that the berry size is a determining factor in the wine grape quality.

The values of the technological maturity parameters obtained are those usually found for the Nebbiolo cultivar in the Piedmont region. At the same density class, the sugar content

agreed among the five harvest dates. Fournand et al. (6) reported that the difference in the total sugar content of the berries belonging to two consecutive density classes was ~17 g/L (i.e., 1 % v/v potential alcohol). For Nebbiolo grapes, these differences ranged from 8-18 g/L. Although the grape berries showed a similar floating behavior among the harvest dates, a significant decrease was found in the total acidity when both the berries density increased and the ripening stage advanced (**Table 2**). On the contrary, the pH values of the floated grapes showed little or no differences among both the density classes and the harvest dates.

Total acidity was the only technological maturity parameter that is statistically correlated with the percentage of grape berries belonging to each density class in the different harvest dates. For A density class, a correlation coefficient of 0.997 ($p = 0.0002$, $n = 5$) was obtained whereas this was -0.930 ($p = 0.022$, $n = 5$) for C berries density. This implies that total acidity is the technological maturity parameter more dependent on the harvest date. Although total acidity diminished through the ripening period, the correlation coefficient was positive for the A density class but negative for the C one. This is due to the fact that proportion of grapes with lower pulp sugar content diminishes in the most advanced ripening stages while that of grapes richer in sugars increases.

Cellular maturity index and seed maturity index (Glories' indices). The modifications found in the phenol extractability indices for the Nebbiolo variety through the grape ripening at three different berries densities can be seen in **Table 3**. EA and MP are considered maturity indices. Although anthocyanins are located in the vacuoles in a free form, the skin cell-wall constitutes a barrier for these compounds. Skin cell-walls undergo compositional and structural changes during the grape ripening that modify the capability to diffuse anthocyanins (11). In seeds, the histological and histochemical modifications

that occur during the grape development also affect the ability to release phenols because the solidification of the cells rich in tannins, before harvest, can negatively affect the aptitude for extraction of these compounds (8).

In the same density class, the results obtained for A_1 and $A_{3.2}$ indicate an increasing tendency to accumulate more anthocyanins in the berry skin when the harvest date is later, excepting the first one. Furthermore, the percentage variation of total anthocyanin concentration ranged from 37.2 % to 40.7 % for both A and B density classes; it varied between 19.7 and 21.0 % for the C one. When the differences in the anthocyanin accumulated in the same harvest date for the three berries densities were studied, an increase in both A_1 and $A_{3.2}$ with the increase in the density was observed. This agrees with the increase previously reported in the anthocyanin accumulation in the berry skin from 235 to 269 g/L sugars for sorted Barbera grapes that were harvested at the same date (17). Nevertheless, the differences among total anthocyanin concentrations corresponding to A and C density classes diminished as the ripening stage advanced. In a previous study performed on sorted berries (by selecting only one class of berries for each harvest date), the total red pigments increased rapidly until 170 g/L sugars in the pulp (6). Afterwards, the amount of total red pigments remained nearly unchanged.

In general, FNA_1 , $FNA_{3.2}$ and MP showed a decreasing tendency with the harvest date for each density class defined by flotation whereas a clear tendency was not observed for TF_1 , $TF_{3.2}$, A_{280} and EA, independently of the berries density measured. For each harvest date, TF_1 , $TF_{3.2}$, FNA_1 , $FNA_{3.2}$ and A_{280} increased with the density class in most of cases but this increase was not always significant. In a previous work, sorted Barbera grapes harvested at the same date also showed an increasing tendency of the above mentioned phenol extractability indices when the sugar content increased from 235 to 269 g/L in three levels (17).

In the last harvest date, the values of the phenol extractability indices (A_1 , $A_{3.2}$, TF_1 , $TF_{3.2}$, A_{280} , EA and MP) agreed with those previously published for Nebbiolo grapes in the same production area (13). On the other hand, the results obtained for A_1 , EA and MP fell within the range reported by Ribéreau-Gayon et al. (25), who consider values of 500-2000 mg/L, 70-20 % and 60-0 %, respectively, as the normal variation range. The values of A_1 and EA depend on the ripeness degree and variety, while those of MP depend also on the number of seeds per berry. Furthermore, values lower than 30 % for both EA and MP are recommended to indicate a good phenolic maturity. Therefore, the values of EA obtained indicate a good skin cell-wall fragility that facilitates the anthocyanin extractability whereas the high values of MP associated with this wine-grape variety involve a high contribution of seed tannins.

The phenol extractability indices are determining factors in the wine-grape quality and have a great impact on the selection of the most suitable winemaking methodology. The elaboration of high quality red wines requires not only a sufficient accumulation of anthocyanins in berry skins through the ripening period but also the anthocyanin extractability has to be also assessed (14). Therefore, high colour intensity requires the management of the winemaking process based on the tendency of the berry skin to yield up anthocyanins (9, 11). It is well-known that the anthocyanin extractability varies through the grape ripening as a consequence of the compositional changes occurring in the skin cell-wall during its degradation by pectolytic enzymes (9, 11). In fact, no clear influence of the density class on the anthocyanin extractability in the same harvest date, or with the ripening stage at the same density class, was found. This agrees with two previously published reports in where no significant difference was found in the anthocyanin extractability for sorted Mencía grapes containing 176, 193 and 210 g/L sugars (16) or for sorted Barbera grapes containing 235, 252 and 269 g/L sugars (17).

A correlation study was performed in an attempt to find a relationship between the percentage of grape berries belonging to each density class in the different ripening stages and the phenol extractability indices. This correlation study permits a better understanding of the effect of the ripening stage on the phenol extractability indices. For A density class, correlation coefficients of -0.983 ($p = 0.017$, $n = 4$) and 0.992 ($p = 0.008$, $n = 4$) were obtained for A_1 and MP, respectively, when the values corresponding to the first harvest date were not considered. This signifies that the anthocyanin accumulation in berry skins increased and the contribution of seed tannins diminished with the harvest date for those grapes with lower density.

Skin phenolic composition and extractability. The total extractable phenolic composition of the berry skin and the extractability yield in a model hydroalcoholic solution for the five grape ripening stages studied at three different density classes are shown in **Table 4**. At the same harvest date, total extractable concentration of TA_{sk} , TF_{sk} , PRO_{sk} and FVA_{sk} increased with the berries density in most of cases. However, the variations were only significant for TA_{sk} concentration in I-IV harvest dates and for TF_{sk} content in the three first ones. In particular, an important increase in TA_{sk} with the density class was observed and the difference between the A and C classes was of 154-204 mg/kg of grapes for the I, II and III harvest dates. Furthermore, TF_{sk} experienced a relevant increase from the density class A to C of 371-522 mg/kg of grapes. Therefore, a useful grape densimetric separation could be performed in winery, by densimetric sorting equipments, at 1094 kg/m^3 to obtain wines with differences in the last two parameters. At the same density class, an irregular tendency was generally found for the total extractable concentration of TA_{sk} , TF_{sk} , PRO_{sk} and FVA_{sk} through the ripening process. In this case, the more significant differences among harvest dates were associated with C density class,

except for FVA_{sk}. The extraction yield ranged as follows 91.1-94.2 %, 53.0-63.3 %, 73.8-80.7 % and 77.5-89.7 % for TA_{sk}, TF_{sk}, PRO_{sk} and FVA_{sk}, respectively, but very few significant changes were found for each phenol with the harvest date or the density of berries. In the last stages of grape ripening, the decline rates for all flavanols slowed and, therefore, the composition changes very little (26).

Other work previously published on Barbera grapes, harvested at the same date and classified into three soluble solid classes defined by flotation, reported a significant increase in the concentrations of TA_{sk} and TF_{sk} but those of PRO_{sk} and FVA_{sk} appeared to be independent of the sugar accumulation in the berry pulp (17). This also agrees with the increase previously reported in the anthocyanin accumulation in the berry skin of sorted Mencía grapes (16).

Regarding the anthocyanin profile, **Table 5** shows few significant differences among harvest dates or among density classes because the anthocyanin profile can be considered as a chemical-taxonomic marker of a certain variety. Nevertheless, some authors evidenced that environmental factors influence the anthocyanin synthesis (27, 28). Nebbiolo variety is characterized by a higher percentage of simple glucosides (85.0-88.9 %), malvidin and peonidin derivative forms being the majority anthocyanin compounds (26.7-39.9 % and 32.8-36.9 %, respectively). The results obtained agreed with others previously reported in the literature (28, 29).

In the same ripening stage, it is important to bear in mind that the higher berries density involved a lower percentage of malvidin derivatives and higher ones of petunidin, cyanidin and delphinidin derivative forms. Hence, the higher proportions of di-substituted anthocyanins and non-acylated anthocyanins contribute to a greater sensitivity to the oxidation reactions and to the colour degradation, the colour of the denser Nebbiolo grapes may be more easily degraded. The scarcity of significant differences also suggests that the

variations observed in the anthocyanin profile are not related to a different sugar accumulation (17, 22).

At the same C density class, significant differences in the anthocyanin profile were obtained in the first harvest date. Higher percentages of simple glucosides, cyanidin derivative and peonidin derivative forms were found in favour of lower percentages of acylated glucosides and malvidin derivative forms. Therefore, the later ripening stages may also involve the red wines production with a less sensitivity to the colour degradation.

A correlation study was performed to establish a relationship between the percentage of Nebbiolo grapes belonging to each density class at the different harvest dates and both total extractable phenolic composition and the easily extractable anthocyanin profile. With regard to total extractable phenolic composition, the TF_{sk} concentration was significantly correlated at B density class when the values corresponding to the last ripening stage were discarded (-0.995 , $p = 0.005$, $n = 4$). On the other hand, the FVA_{sk} content showed a significant correlation at C density class when the values associated with the first harvest date were not considered (-0.959 , $p = 0.041$, $n = 4$). For the higher berries density, a significant correlation was found for delphinidin derivatives (-0.978 , $p = 0.004$, $n = 5$). When the results obtained in the first harvest date were discarded, this correlation factor increased to -1.000 ($p = 0.000$, $n = 4$) and a new statistic correlation was found for petunidin derivatives (-0.973 , $p = 0.027$, $n = 4$). Both delphinidin and petunidin derivatives also showed a good correlation for B density class when the values associated with the last harvest date were not considered (-0.965 , $p = 0.035$, $n = 4$ and -0.976 , $p = 0.024$, $n = 4$, respectively). This confirms that the anthocyanin compounds more prone to oxidation diminished with the increase in the proportion of berries belonging to B and C density classes through the grape ripening, and that the skin flavanols reactivity decreased with the

increase in the proportion of the denser berries. It agreed with the loss in the skin tannins reactivity during ripening (25).

Skin mechanical properties. The mechanical properties of berry skins for Nebbiolo grapes harvested in five different dates and classified in three density classes are shown in **Table 6**. The puncture parameters of berry skins increased with the berries density at each harvest date studied. However, this increment was not always significant. Although very few significant changes were reported in the parameters that characterize the berry skin hardness (F_{sk} and W_{sk}) and the tissue rigidity or stiffness (E_{sk}) of Barbera grapes containing different soluble solid contents, an increasing tendency of Sp_{sk} values with the sugar accumulation was observed (17).

When the modification of the skin mechanical characteristics at a certain density class through the five ripening stages evaluated was studied, **Table 6** shows values of the E_{sk} parameter significantly higher and of Sp_{sk} significantly lower for the three density classes in III harvest date, particularly for the A berries density. Furthermore, a correlation study was then carried out to establish a relationship among the percentage of grape berries belonging to each density class in the different ripening stages and the skin mechanical attributes. Nevertheless, no significant correlation was found.

Several studies suggested that the behaviour of F_{sk} values close to the harvest time could limit the choice of this parameter as a maturity indicator in grape berries. In fact, from veraison to ripeness, an increase in the F_{sk} parameter is observed, particularly in the first ripening phases, with a steady value or a slight decrease close to technological maturity (24). This same behaviour was also observed in Nebbiolo grapes for A and C density classes achieving the higher F_{sk} values in II and III harvest dates.

Recently, instrumental texture analysis has also been used for a rapid estimation of the anthocyanin extractability. In particular, the berry skin break force can be considered the best mechanical parameter to estimate the kinetics of anthocyanin extraction with adequate reliability (15) while the berry skin thickness has been proposed to predict the percentage of extractable anthocyanins (16). The mechanical methods are inexpensive, allowing their application as a routine monitoring tool for the grape quality. Thinner skins seem to be characterized by a greater cellular maturity index (30) but this behaviour was only confirmed when comparing Sp_{sk} values among the three density classes for I, II and III harvest dates. The separation of the grape berries on the basis of the anthocyanin extractability estimated from Sp_{sk} could be possible using flotation with the density of 1094 kg/m^3 , which was already proposed for the determination of the berry skin phenolic composition.

Relationship among mechanical and chemical parameters of berry skins. To summarise, this study evaluated the possible dependence of phenol composition and/or extractability on skin mechanical attributes, irrespective of the effect of the harvest date and/or the density class (**Table 7**). The mechanical properties do not seem to be well related to the red pigments accumulated in berry skins. On the other hand, the puncture parameters have been also evaluated as potential estimators of the accumulation power of berry skins for total and easily extractable concentrations of flavonoids, proanthocyanidins and flavanols reactive to vanillin, as well as of the facility of berry skins to yield them up. Total and easily extractable concentrations of TF_{sk} and FVA_{sk} also showed a low correlation with the E_{sk} parameter whereas the concentrations of PRO_{sk} achieved factors ranging 0.705 and 0.756 ($p < 0.004$, $n = 15$). Furthermore, better relationships for total and easily extractable concentrations of FVA_{sk} were obtained with F_{sk} parameter with

correlation factors varying between 0.766 and 0.774 ($p < 0.001$, $n = 15$). The extractability of FVA_{sk} is also little correlated with the F_{sk} parameter. On the other hand, the extraction yield for TF_{sk} and PRO_{sk} is correlated with Sp_{sk} parameter, correlation factors being 0.567 ($p = 0.028$, $n = 15$) and 0.671 ($p = 0.006$, $n = 15$), respectively. Sp_{sk} parameter facilitates the estimation of the skin cell-wall degradability and, therefore, of the extractability of proanthocyanidins. According to Rolle et al. (29), higher skin hardness probably involves greater cell-wall fragility which agrees with the tendency of the extraction yield of flavanols reactive to vanillin to increase when the F_{sk} parameter increases.

Although further studies are necessary increasing the grape varieties, growing areas and vintages to achieve more robust conclusions, this first approach showed that, for a given harvest date, the denser grapes provide, in general, higher total and easily extractable concentrations of phenolic compounds. Since the heterogeneity of the grapes harvested determines the variability of the results obtained, the relevance of the information provided for the management of the winemaking process diminishes according to the grape heterogeneity. The lack and low reliability of the statistical correlations found between the percentage of grape berries belonging to each density class, in the different ripening stages, and the mechanical/chemical parameters confirmed that the changes observed in the latter through the grape ripening process are more related to the berries density than the harvest date. This suggests the effectiveness of the automatic winery equipment of berry densimetric sorting recently developed whenever the flotation density is selected according to the objective quality properties of the grape like TA_{sk} , TF_{sk} and Sp_{sk} . Therefore, the winemaker has to select the flotation density on the basis of the wine type he wishes to elaborate. This work also highlights the importance of improving knowledge of the physical modifications of the cell tissues through the grape ripening to assess the evolution of the phenol composition and extractability because the influence of the skin mechanical

properties on total flavonoids, proanthocyanidins and flavanols reactive to vanillin has not been previously studied. Stiffer grapes allowed accumulation of more proanthocyanidins, while harder ones provided higher concentration and extractability of flavanols reactive to vanillin. On the other hand, the thicker grapes facilitated the extraction of proanthocyanidins. This first approach demands further research on histological and histochemical changes in berry skins during the grape development.

LITERATURE CITED

- (1) Parpinello, G.P.; Versari, A.; Chinnici, F.; Galassi, S. Relationship among sensory descriptors, consumer preferences and color parameters of Italian Novello red wines. *Food Res. Int.* **2009**, *42*, 1389-1395.
- (2) Vidal, S.; Francis, L.; Noble, A.; Kwiatkowski, M.; Cheynier, V.; Waters, E. Taste and mouth-feel properties of different types of tannin-like polyphenolic compounds and anthocyanins in wine. *Anal. Chim. Acta* **2004**, *513*, 57-65.
- (3) Cheynier, V.; Duenas-Paton, M.; Salas, E.; Maury, C.; Souquet, J.M.; Sarni-Manchado, P.; Fulcrand, H. Structure and properties of wine pigments and tannins. *Am. J. Enol. Vitic.* **2006**, *57*, 298-305.
- (4) Fulcrand, H.; Duenas, M.; Salas, E.; Cheynier, V. Phenolic reactions during winemaking and aging. *Am. J. Enol. Vitic.* **2006**, *57*, 289-297.
- (5) Bautista-Ortín, A.B.; Fernández-Fernández, J.I.; López-Roca, J.M.; Gómez-Plaza, E. The effect of grape ripening stage on red wine color. *J. Int. Sci. Vigne Vin* **2006**, *40*, 15-24.

- (6) Fournand, D.; Vicens, A.; Sidhoum, L.; Souquet, J.M.; Moutounet, M.; Cheynier, V. Accumulation and extractability of grape skin tannins and anthocyanins at different advanced physiological stages. *J. Agric. Food Chem.* **2006**, *54*, 7331-7338.
- (7) Cadot, Y.; Miñana Castelló, M.T.; Chevalier, M. Flavan-3-ol compositional changes in grape berries (*Vitis vinifera* L. cv Cabernet Franc) before veraison, using two complementary analytical approaches, HPLC reversed phase and histochemistry. *Anal. Chim. Acta* **2006**, *563*, 65-75.
- (8) Cadot, Y.; Miñana Castelló, M.T.; Chevalier, M. Anatomical, histological, and histochemical changes in grape seeds from *Vitis vinifera* L. cv Cabernet franc during fruit development. *J. Agric. Food Chem.* **2006**, *54*, 9206-9215.
- (9) Romero-Cascales, I.; Ortega-Regules, A.; López-Roca, J.M.; Fernández-Fernández, J.I.; Gómez-Plaza, E. Differences in anthocyanin extractability from grapes to wines according to variety. *Am. J. Enol. Vitic.* **2005**, *56*, 212-219.
- (10) Kontoudakis, N.; Esteruelas, M.; Fort, F.; Canals, J.M.; Zamora, F. Comparison of methods for estimating phenolic maturity in grapes: correlations between predicted and obtained parameters. *Anal. Chim. Acta* **2010**, *660*, 127-133.
- (11) Ortega-Regules, A.; Romero-Cascales, I.; Ros-García, J.M.; López-Roca, J.M.; Gómez-Plaza, E. A first approach towards the relationship between grape skin cell-wall composition and anthocyanin extractability. *Anal. Chim. Acta* **2006**, *563*, 26-32.
- (12) Mattivi, F.; Vrhovsek, U.; Masuero, D.; Trainotti, D. Differences in the amount and structure of extractable skin and seed tannins amongst red grape varieties. *Aust. J. Grape Wine Res.* **2009**, *15*, 27-35.
- (13) Cagnasso, E.; Rolle, L.; Caudana, A.; Gerbi, V. Relationship between grape phenolic maturity and red wine phenolic composition. *Ital. J. Food Sci.* **2008**, *20*, 365-381.

- (14) González-Neves, G.; Gil, G.; Ferrer, M.; Darwin, C.; Balado, J.; Bochicchio, R.; Gatto, G.; Tessore, A. Prediction of the colour and polyphenolic composition of the young red wines from the phenolic potential of the grapes. *Int. J. Food Sci. Technol.* **2010**, *45*, 1843-1851.
- (15) Rolle, L.; Torchio, F.; Ferrandino, A.; Guidoni, S. Influence of wine-grape skin hardness on the kinetics of anthocyanin extraction. *Int. J. Food Prop.* **2011**, in press (DOI: 10.1080/10942911003778022).
- (16) Río Segade, S.; Giacosa, S.; Gerbi, V.; Rolle, L. Berry skin thickness as main texture parameter to predict anthocyanin extractability in winegrapes. *LWT Food Sci. Technol.* **2011**, *44*, 392-398.
- (17) Torchio, F.; Cagnasso, E.; Gerbi, V.; Rolle, L. Mechanical properties, phenolic composition and extractability indices of Barbera grapes of different soluble solids contents from several growing areas. *Anal. Chim. Acta* **2010**, *660*, 183-189.
- (18) Kontoudakis, N.; Esteruelas, M.; Fort, F.; Canals, J.M.; De Freitas, V.; Zamora, F. Influence of the heterogeneity of grape phenolic maturity on wine composition and quality. *Food Chem.* **2011**, *124*, 767-774.
- (19) Saint-Criq, N.; Vivas, N.; Glories, Y. Maturité phénolique: définition et contrôle. *Rev. Fr. Oenol.* **1998**, *173*, 22-25.
- (20) Rolle, L.; Torchio, F.; Giacosa, S.; Gerbi, V. Modification of mechanical characteristic and phenolic composition in berry skins and seeds of Mondeuse winegrapes throughout the on-vine drying process. *J. Sci. Food Agric.* **2009**, *89*, 1973-1980.
- (21) Rolle, L.; Torchio, F.; Zeppa, G.; Gerbi, V. Relations between break skin force and anthocyanin extractability at different stages of ripening. *Am. J. Enol. Vitic.* **2009**, *60*, 93-97.

- (22) Pomar, F.; Novo, M.; Masa, A. Varietal differences among the anthocyanin profiles of 50 red table grape cultivars studied by high performance liquid chromatography. *J. Chromatogr. A* **2005**, *1094*, 34-41.
- (23) Letaief, H.; Rolle, L.; Gerbi, V. Mechanical behavior of winegrapes under compression tests. *Am. J. Enol. Vitic.* **2008**, *59*, 323-329.
- (24) Maury, C.; Madieta, E.; Le Moigne, M.; Mehinagic, E.; Siret, R.; Jourjon, F. Development of a mechanical texture test to evaluate the ripening process of Cabernet Franc grapes. *J. Text. Stud.* **2009**, *40*, 511-535.
- (25) Ribéreau-Gayon, P.; Glories, Y.; Maujean, A.; Dubourdieu, D. Compuestos fenólicos. In *Tratado de Enología: Química del vino. Estabilización y tratamientos*; Hemisferio Sur: Buenos Aires, 2003; vol. 2.
- (26) Kennedy, J.A.; Troup, G.J.; Pilbrow, J.R.; Hutton, D.R.; Hewitt, D.; Hunter, C.R.; Ristic, R.; Iland, P.G.; Jones, G.P. Development of seed polyphenols in berries from *Vitis vinifera* L. cv. Shiraz. *Aust. J. Grape Wine Res.* **2000**, *6*, 244-254.
- (27) Downey, M.; Dokoozlian, N.K.; Krstic, M.P. Cultural practice and environmental impacts on the flavonoid composition of grapes and wine: a review of recent research. *Am. J. Enol. Vitic.* **2006**, *57*, 257-268.
- (28) Guidoni, S.; Ferrandino, A.; Novello, V. Climate and agronomical practice effects on anthocyanin accumulation in cv. 'Nebbiolo' (*Vitis vinifera* L.) berries. *Am. J. Enol. Vitic.* **2008**, *59*, 22-29.
- (29) Rolle, L.; Torchio, F.; Zeppa, G.; Gerbi, V. Anthocyanin extractability assessment of grape skins by texture analysis. *J. Int. Sci. Vigne Vin* **2008**, *42*, 157-162.
- (30) Río Segade, S.; Soto Vázquez, E.; Orriols, I.; Giacosa, S.; Rolle, L. Possible use of texture characteristics of winegrapes as markers for zoning and their relationship with anthocyanin extractability index. *Int. J. Food Sci. Technol.* **2011**, *46*, 386-394.

Table 1. Operative conditions for the measurement of the berry mechanical parameters

Test	Probe	Test speed (mm/s)	Compression- puncture (mm)	Mechanical property
Skin hardness	P/2N, 2 mm needle	1	3	F_{sk} = berry skin break force (N) W_{sk} = berry skin break energy (mJ) E_{sk} = berry skin Young's modulus (N/mm)
Skin thickness	P/2, Ø 2 mm	0.2	–	Sp_{sk} = berry skin thickness (μm)

Table 2. Physical and technological maturity parameters for Nebbiolo grapes harvested at different ripening stages and sorted according to density

Density class	Harvest date	Berry weight (g)	Volume (cm ³)	Reducing sugars (g/L)	pH	Total acidity (g/L)
A	I	2.04±0.06 ^{a,β}	1.87±0.06 ^{a,β}	212±2 ^{a,α}	2.96±0.06 ^{a,α}	7.2±0.1 ^{e,γ}
	II	2.05±0.07 ^{a,α}	1.88±0.08 ^{a,α}	207±5 ^{a,α}	2.95±0.02 ^{a,α}	7.0±0.2 ^{d,β}
	III	2.00±0.04 ^{a,α}	1.84±0.09 ^{a,α}	212±2 ^{a,α}	2.95±0.05 ^{a,α}	6.4±0.3 ^{c,β}
	IV	1.96±0.02 ^{a,α}	1.80±0.07 ^{a,α}	211±8 ^{a,α}	3.06±0.04 ^{a,α}	6.0±0.6 ^{b,β}
	V	2.05±0.04 ^{a,α}	1.88±0.08 ^{a,α}	212±7 ^{a,α}	3.30±0.04 ^{b,β}	5.8±0.2 ^{a,γ}
Sign ¹		ns	ns	ns	**	***
B	I	2.04±0.06 ^{ab,β}	1.86±0.07 ^{a,β}	224±3 ^{a,β}	3.00±0.07 ^{a,αβ}	6.8±0.3 ^{d,β}
	II	2.15±0.04 ^{bc,α}	1.96±0.05 ^{ab,α}	224±6 ^{a,β}	3.01±0.06 ^{a,α}	6.7±0.1 ^{d,α}
	III	2.03±0.07 ^{ab,α}	1.85±0.05 ^{a,α}	227±4 ^{a,β}	3.01±0.04 ^{a,α}	6.2±0.4 ^{c,α}
	IV	1.97±0.08 ^{a,α}	1.80±0.02 ^{a,α}	224±4 ^{a,β}	3.07±0.04 ^{a,α}	5.7±0.3 ^{b,α}
	V	2.28±0.05 ^{c,β}	2.07±0.09 ^{b,β}	230±8 ^{a,β}	3.10±0.08 ^{a,α}	5.4±0.1 ^{a,β}
Sign ¹		**	**	ns	ns	***
C	I	1.72±0.05 ^{a,α}	1.56±0.04 ^{a,α}	241±5 ^{a,γ}	3.13±0.05 ^{a,β}	6.3±0.3 ^{d,α}
	II	2.16±0.02 ^{c,α}	1.96±0.01 ^{c,α}	235±8 ^{a,γ}	3.03±0.03 ^{a,α}	6.7±0.6 ^{e,α}
	III	1.96±0.05 ^{b,α}	1.78±0.03 ^{b,α}	235±7 ^{a,γ}	3.02±0.03 ^{a,α}	6.1±0.2 ^{c,α}
	IV	1.91±0.04 ^{b,α}	1.74±0.07 ^{b,α}	241±8 ^{a,γ}	3.07±0.06 ^{a,α}	5.7±0.1 ^{b,α}
	V	2.05±0.09 ^{bc,α}	1.86±0.05 ^{b,α}	241±6 ^{a,γ}	3.14±0.05 ^{a,α}	5.1±0.3 ^{a,α}
Sign ¹		***	***	ns	ns	***
Sign ²		***,ns,ns,ns,**	**,ns,ns,ns,*	***,***,***,***,***	*,ns,ns,ns,*	***,**,**,***,***

Data are expressed as average value ± standard deviation: n = 30 for berry weight and volume, n = 3 for technological maturity parameters. Different Latin letters within the same column indicate significant differences (¹) among harvest dates at the same berries density (*Tukey-b test*; $p < 0.05$). Different Greek letters within the same column indicate significant differences (²) among the three density classes at the same harvest date (*Tukey-b test*; $p < 0.05$). ^{1,2}: *, **, *** and ns indicate significance at $p < 0.05$, 0.01, 0.001 and not significant, respectively. A = 1088 kg/m³, B = 1094 kg/m³, C = 1100 kg/m³.

Table 3. Phenol extractability indices for Nebbiolo grapes harvested at different ripening stages and sorted according to density

Density class	Harvest date	A _{3,2} (mg/L malvidin-3-glucoside chloride)	TF _{3,2} (mg/L (+)-catechin)	FNA _{3,2} (mg/L (+)-catechin)	A ₂₈₀	A ₁ (mg/L malvidin-3-glucoside chloride)	TF ₁ (mg/L (+)-catechin)	FNA ₁ (mg/L (+)-catechin)	EA (%)	MP (%)
A	I	245±7 ^{b,α}	2028±60 ^{c,α}	1672±50 ^{b,α}	50.0±1.5 ^{b,α}	388±11 ^{b,α}	3482±104 ^{c,α}	2917±96 ^{b,α}	36.9±1.1 ^{b,α}	65.7±2.1 ^{c,α}
	II	195±5 ^{a,α}	1859±55 ^{ab,α}	1576±47 ^{b,β}	67.0±2.2 ^{c,α}	304±14 ^{a,α}	2662±85 ^{a,α}	2219±62 ^{a,α}	36.0±1.2 ^{ab,α}	79.6±2.7 ^{d,α}
	III	210±9 ^{a,α}	1722±48 ^{a,α}	1416±58 ^{a,α}	45.2±1.8 ^{a,α}	387±15 ^{b,α}	3397±98 ^{c,αβ}	2834±85 ^{b,β}	45.6±1.6 ^{c,γ}	67.4±2.2 ^{c,β}
	IV	267±10 ^{c,α}	1851±78 ^{ab,α}	1462±66 ^{a,α}	47.8±2.2 ^{ab,α}	436±21 ^{c,α}	3032±102 ^{b,α}	2396±99 ^{a,α}	38.8±1.0 ^{b,α}	60.9±2.3 ^{b,α}
	V	329±6 ^{d,α}	1927±84 ^{bc,α}	1447±44 ^{a,α}	50.2±1.9 ^{b,α}	495±12 ^{d,α}	3064±115 ^{b,α}	2344±89 ^{a,α}	33.5±1.8 ^{a,α}	54.1±2.7 ^{a,α}
Sign ¹		***	**	*	**	**	***	***	**	**
B	I	291±9 ^{b,β}	2307±69 ^{c,β}	1883±56 ^{c,β}	59.6±1.8 ^{c,β}	487±17 ^{c,β}	3917±117 ^{d,β}	3209±87 ^{c,β}	40.2±1.6 ^{a,β}	65.8±1.7 ^{c,α}
	II	226±8 ^{a,β}	1775±75 ^{a,α}	1445±38 ^{a,α}	71.8±0.9 ^{d,α}	375±9 ^{a,β}	2829±81 ^{a,α}	2283±69 ^{a,α}	39.7±1.5 ^{a,β}	77.9±1.8 ^{d,α}
	III	278±11 ^{b,β}	2066±85 ^{b,β}	1661±47 ^{b,β}	51.2±2.8 ^{a,β}	444±16 ^{b,β}	3211±105 ^{b,α}	2565±56 ^{b,α}	37.3±1.2 ^{a,β}	62.0±2.0 ^{bc,α}
	IV	313±15 ^{c,β}	2104±63 ^{b,β}	1648±74 ^{b,β}	54.4±1.7 ^{ab,β}	504±13 ^{c,β}	3313±83 ^{bc,β}	2579±88 ^{b,β}	37.8±1.8 ^{a,α}	59.7±1.5 ^{ab,α}
	V	364±11 ^{d,β}	2112±68 ^{b,β}	1583±56 ^{b,β}	57.4±2.0 ^{bc,β}	597±11 ^{d,γ}	3482±101 ^{c,β}	2612±76 ^{b,β}	39.1±2.0 ^{a,β}	55.7±1.5 ^{a,α}
Sign ¹		***	***	***	***	***	***	***	ns	***
C	I	303±9 ^{a,β}	2475±70 ^{d,γ}	1930±45 ^{d,β}	62.1±0.7 ^{b,β}	495±13 ^{ab,β}	4031±119 ^{d,β}	3350±90 ^{d,β}	42.0±1.5 ^{b,β}	66.0±2.3 ^{b,α}
	II	282±10 ^{a,γ}	1876±127 ^{a,α}	1466±52 ^{a,α}	83.8±1.7 ^{c,β}	478±12 ^{a,γ}	3169±95 ^{a,β}	2474±76 ^{a,β}	41.0±0.9 ^{b,β}	76.5±3.2 ^{c,α}
	III	353±11 ^{c,γ}	2233±68 ^{c,γ}	1720±56 ^{c,β}	62.8±3.2 ^{b,γ}	528±19 ^{bc,γ}	3550±111 ^{bc,β}	2781±89 ^{bc,β}	33.2±1.3 ^{a,α}	60.7±2.8 ^{a,α}
	IV	327±8 ^{b,β}	2079±95 ^{b,β}	1602±60 ^{b,β}	55.2±1.3 ^{a,β}	595±17 ^{d,γ}	3708±107 ^{c,γ}	2842±97 ^{c,γ}	44.9±0.8 ^{c,β}	58.5±3.6 ^{a,α}
	V	357±6 ^{c,β}	2248±64 ^{c,β}	1728±44 ^{c,γ}	63.8±1.5 ^{b,γ}	550±9 ^{c,β}	3412±96 ^{b,β}	2611±82 ^{ab,β}	35.1±0.6 ^{a,α}	60.8±2.2 ^{a,β}
Sign ¹		***	***	***	***	***	***	***	***	***
Sign ²		***,***,***,***,*	***,ns,***,*,**	*,*,***,*,**	***,***,***,*,***	***,***,***,***,***	**,*,*,***,***	**,*,*,***,***	**,*,***,***,***	ns,ns,*,ns,***

All data are expressed as average value ± standard deviation (n = 2). Different Latin letters within the same column indicate significant differences (¹) among harvest dates at the same berries density (*Tukey-b test*; $p < 0.05$). Different Greek letters within the same column indicate significant differences (²) among the three density classes at the same harvest date (*Tukey-b test*; $p < 0.05$). ^{1,2}: *, **, *** and ns indicate significance at $p < 0.05$, 0.01, 0.001 and not significant, respectively. A_{3,2} = total anthocyanins extracted at pH 3.2, TF_{3,2} = total flavonoids extracted at pH 3.2, FNA_{3,2} = non-anthocyanin flavonoids extracted at pH 3.2, A₂₈₀ = total phenolic content, A₁ = total anthocyanins extracted at pH 1, TF₁ = total flavonoids extracted at pH 1, FNA₁ = non-anthocyanin flavonoids extracted at pH 1, EA = cellular maturity index, MP = seed maturity index. A = 1088 kg/m³, B = 1094 kg/m³, C = 1100 kg/m³.

Table 4. Skin total extractable phenolic composition and relative extractability in model hydroalcoholic solution for Nebbiolo grapes harvested at different ripening stages and sorted according to density

Density class	Harvest date	TA _{sk} (mg/kg malvidin-3-glucoside chloride)	TA _{sk} (% extraction)	TF _{sk} (mg/kg (+)-catechin)	TF _{sk} (% extraction)	PRO _{sk} (mg/kg cyanidin chloride)	PRO _{sk} (% extraction)	FVA _{sk} (mg/kg (+)-catechin)	FVA _{sk} (% extraction)
A	I	415±9 ^{a,α}	94.1±0.1 ^{a,α}	2435±12 ^{b,α}	59.5±1.8 ^{ab,α}	2385±258 ^{a,α}	79.2±0.9 ^{ab,α}	828±45 ^{ab,α}	85.9±0.6 ^{a,α}
	II	328±52 ^{a,α}	92.3±1.2 ^{a,α}	2040±61 ^{a,α}	54.5±2.2 ^{ab,α}	2249±63 ^{a,α}	74.6±0.6 ^{a,α}	703±142 ^{a,α}	77.5±2.2 ^{a,α}
	III	332±29 ^{a,α}	91.9±0.1 ^{a,α}	2053±28 ^{a,α}	55.2±0.3 ^{ab,α}	2319±104 ^{a,α}	74.7±1.8 ^{a,α}	1334±163 ^{b,α}	88.3±2.5 ^{a,α}
	IV	420±29 ^{a,α}	93.8±0.1 ^{a,α}	2262±134 ^{ab,α}	61.4±2.8 ^{b,α}	2406±397 ^{a,α}	80.7±1.8 ^{b,α}	625±278 ^{a,α}	80.0±9.4 ^{a,α}
	V	418±10 ^{a,α}	91.1±1.2 ^{a,α}	2277±22 ^{ab,α}	53.0±1.2 ^{a,α}	2034±164 ^{a,α}	73.8±1.6 ^{a,α}	786±32 ^{ab,α}	83.7±1.2 ^{a,α}
Sign ¹		ns	ns	**	*	ns	*	*	ns
B	I	511±32 ^{a,β}	94.2±0.0 ^{b,α}	2717±60 ^{b,β}	61.2±1.6 ^{a,α}	2595±346 ^{ab,α}	78.8±4.1 ^{a,α}	1075±225 ^{a,α}	88.0±3.2 ^{a,α}
	II	451±19 ^{a,αβ}	92.9±0.1 ^{ab,α}	2463±154 ^{ab,β}	63.3±0.6 ^{a,β}	2839±36 ^{b,β}	78.0±2.7 ^{a,α}	1296±429 ^{a,α}	86.5±2.0 ^{a,β}
	III	409±12 ^{a,β}	92.4±0.6 ^{ab,α}	2236±22 ^{a,β}	57.0±1.1 ^{a,αβ}	2397±117 ^{ab,α}	76.0±0.8 ^{a,α}	1407±41 ^{a,α}	88.8±1.7 ^{a,α}
	IV	482±25 ^{a,αβ}	92.7±0.0 ^{ab,α}	2396±93 ^{ab,α}	61.8±0.4 ^{a,α}	2640±123 ^{ab,α}	78.7±0.4 ^{a,α}	993±103 ^{a,α}	88.1±1.8 ^{a,α}
	V	457±56 ^{a,α}	91.5±0.7 ^{a,α}	2151±198 ^{a,α}	58.5±4.0 ^{a,α}	2076±116 ^{a,α}	77.5±0.3 ^{a,α}	845±168 ^{a,α}	86.5±3.1 ^{a,α}
Sign ¹		ns	*	*	ns	*	ns	ns	ns
C	I	619±16 ^{b,γ}	94.1±1.1 ^{a,α}	2957±13 ^{d,γ}	63.0±2.6 ^{a,α}	3202±152 ^{c,α}	80.3±4.7 ^{a,α}	1346±251 ^{a,α}	89.5±0.9 ^{a,α}
	II	482±19 ^{a,β}	93.4±0.5 ^{a,α}	2411±1 ^{b,β}	62.9±1.3 ^{a,β}	2642±217 ^{b,αβ}	80.0±2.2 ^{a,α}	1616±382 ^{a,α}	88.7±2.6 ^{a,β}
	III	499±3 ^{a,γ}	92.0±0.0 ^{a,α}	2430±24 ^{b,γ}	59.5±0.5 ^{a,β}	2456±51 ^{b,α}	76.3±1.8 ^{a,α}	1527±209 ^{a,α}	89.7±3.1 ^{a,α}
	IV	533±16 ^{a,β}	93.5±0.3 ^{a,α}	2594±37 ^{c,α}	62.2±0.3 ^{a,α}	2540±20 ^{b,α}	78.2±0.7 ^{a,α}	1002±22 ^{a,α}	87.6±0.0 ^{a,α}
	V	483±49 ^{a,α}	92.2±1.0 ^{a,α}	2224±98 ^{a,α}	60.3±2.6 ^{a,α}	1905±147 ^{a,α}	78.3±2.5 ^{a,α}	713±83 ^{a,α}	86.9±2.2 ^{a,α}
Sign ¹		*	ns	***	ns	**	ns	ns	ns
Sign ²		**, *, **, *, ns	ns, ns, ns, ns, ns	**, *, **, ns, ns	ns, *, *, ns, ns	ns, *, ns, ns, ns	ns, ns, ns, ns, ns	ns, ns, ns, ns, ns	ns, *, ns, ns, ns

All data are expressed as average value ± standard deviation (n = 3). Different Latin letters within the same column indicate significant differences (¹) among harvest dates at the same berries density (*Tukey-b test*; $p < 0.05$). Different Greek letters within the same column indicate significant differences (²) among the three density classes at the same harvest date (*Tukey-b test*; $p < 0.05$). ^{1,2}: *, **, *** and ns indicate significance at $p < 0.05$, 0.01, 0.001 and not significant, respectively. TA_{sk} = skin total anthocyanins, TF_{sk} = skin total flavonoids, PRO_{sk} = skin proanthocyanidins, FVA_{sk} = skin flavanols vanillin assay. A = 1088 kg/m³, B = 1094 kg/m³, C = 1100 kg/m³.

Table 5. Extractable anthocyanin profile for Nebbiolo grapes harvested at different ripening stages and sorted according to density

Density class	Harvest date	Simple glucosides (%)	Acetyl-glucosides (%)	Cinnamoyl-glucosides (%)	Delphinidin derivatives (%)	Cyanidin derivatives (%)	Petunidin derivatives (%)	Peonidin derivatives (%)	Malvidin derivatives (%)
A	I	87.9±0.0 ^{a,α}	4.8±0.1 ^{a,αβ}	7.3±0.1 ^{a,α}	8.5±0.8 ^{a,α}	12.2±1.0 ^{a,α}	7.5±0.5 ^{a,α}	34.0±2.7 ^{a,α}	37.8±2.4 ^{a,β}
	II	86.3±0.9 ^{a,α}	5.1±0.3 ^{a,α}	8.5±0.6 ^{a,α}	7.9±0.1 ^{a,α}	11.9±0.5 ^{a,α}	7.0±0.1 ^{a,α}	34.4±0.2 ^{a,α}	38.9±0.7 ^{a,α}
	III	85.0±1.9 ^{a,α}	5.6±0.6 ^{a,α}	9.4±1.3 ^{a,α}	7.3±0.6 ^{a,α}	11.8±2.9 ^{a,α}	6.7±0.1 ^{a,α}	34.3±2.2 ^{a,α}	39.9±5.6 ^{a,α}
	IV	86.0±0.5 ^{a,α}	5.0±0.3 ^{a,α}	9.0±0.1 ^{a,β}	7.9±0.2 ^{a,α}	12.4±0.2 ^{a,α}	7.1±0.1 ^{a,α}	33.8±0.4 ^{a,α}	38.8±0.8 ^{a,β}
	V	86.1±0.7 ^{a,α}	5.0±0.0 ^{a,α}	8.9±0.7 ^{a,α}	7.7±0.2 ^{a,α}	13.4±1.5 ^{a,α}	6.8±0.1 ^{a,α}	34.5±1.0 ^{a,α}	37.6±2.8 ^{a,α}
Sign ¹		ns	ns	ns	ns	ns	ns	ns	ns
B	I	87.5±0.6 ^{a,α}	5.1±0.1 ^{a,β}	7.4±0.5 ^{a,α}	10.1±0.2 ^{a,α}	14.6±0.7 ^{a,α}	8.2±0.0 ^{a,α}	32.8±0.4 ^{a,α}	34.3±0.5 ^{a,β}
	II	87.3±0.7 ^{a,α}	5.2±0.2 ^{a,α}	7.5±0.5 ^{a,α}	9.6±1.1 ^{a,α}	14.5±1.0 ^{a,α}	7.7±0.6 ^{a,α}	34.4±3.0 ^{a,α}	33.8±2.3 ^{a,α}
	III	85.3±1.6 ^{a,α}	5.9±0.6 ^{a,α}	8.8±1.0 ^{a,α}	8.5±0.3 ^{a,αβ}	12.8±1.0 ^{a,α}	7.3±0.1 ^{a,β}	33.0±0.4 ^{a,α}	38.4±1.8 ^{a,α}
	IV	86.5±0.3 ^{a,α}	5.2±0.2 ^{a,α}	8.3±0.1 ^{a,αβ}	9.0±0.2 ^{a,α}	14.3±1.3 ^{a,α}	7.5±0.0 ^{a,α}	34.7±0.2 ^{a,α}	34.6±1.2 ^{a,α}
	V	86.2±0.5 ^{a,α}	5.2±0.4 ^{a,α}	8.5±0.2 ^{a,α}	8.7±0.5 ^{a,α}	14.6±0.2 ^{a,α}	7.3±0.2 ^{a,αβ}	35.3±1.5 ^{a,α}	34.1±1.0 ^{a,α}
Sign ¹		ns	ns	ns	ns	ns	ns	ns	ns
C	I	88.9±0.5 ^{b,α}	4.7±0.1 ^{a,α}	6.4±0.4 ^{a,α}	10.1±0.7 ^{a,α}	18.5±0.5 ^{b,β}	7.8±0.2 ^{a,α}	36.9±0.8 ^{b,α}	26.7±0.6 ^{a,α}
	II	86.7±0.3 ^{a,α}	5.4±0.1 ^{b,α}	7.9±0.4 ^{b,α}	10.3±0.5 ^{a,α}	15.0±1.7 ^{a,α}	8.1±0.0 ^{a,α}	33.8±0.1 ^{a,α}	32.7±2.2 ^{b,α}
	III	86.6±0.2 ^{a,α}	5.5±0.2 ^{b,α}	7.9±0.0 ^{b,α}	9.7±0.4 ^{a,β}	15.3±0.6 ^{a,α}	7.7±0.0 ^{a,γ}	34.6±0.2 ^{ab,α}	32.8±1.2 ^{b,α}
	IV	86.8±0.6 ^{a,α}	5.4±0.3 ^{b,α}	7.9±0.3 ^{b,α}	9.2±0.8 ^{a,α}	15.0±0.5 ^{a,α}	7.6±0.4 ^{a,α}	35.5±0.8 ^{ab,α}	32.7±0.9 ^{b,α}
	V	86.5±0.2 ^{a,α}	5.5±0.1 ^{b,α}	8.1±0.1 ^{b,α}	8.8±0.1 ^{a,α}	13.3±0.4 ^{a,α}	7.4±0.1 ^{a,β}	34.9±0.7 ^{ab,α}	35.7±1.1 ^{b,α}
Sign ¹		**	*	**	ns	*	ns	*	**
Sign ²		ns,ns,ns,ns,ns	*,ns,ns,ns,ns	ns,ns,ns,*, ns	ns,ns,*,ns, ns	**,ns,ns,ns,ns	ns,ns,**,ns,*	ns,ns,ns,ns,ns	*,ns,ns,*,ns

All data are expressed as average value ± standard deviation (n = 3). Different Latin letters within the same column indicate significant differences (¹) among harvest dates at the same berries density (*Tukey-b test*; $p < 0.05$). Different Greek letters within the same column indicate significant differences (²) among the three density classes at the same harvest date (*Tukey-b test*; $p < 0.05$). ^{1,2}: *,** and ns indicate significance at $p < 0.05$, 0.01 and not significant, respectively. A = 1088 kg/m³, B = 1094 kg/m³, C = 1100 kg/m³.

Table 6. Berry skin mechanical parameters for Nebbiolo grapes harvested at different ripening stages and sorted according to density

Density class	Harvest date	F _{sk} (N)	W _{sk} (mJ)	E _{sk} (N/mm)	Sp _{sk} (μm)
A	I	0.652±0.089 ^{a,α}	0.483±0.100 ^{a,α}	0.402±0.051 ^{a,α}	205±23 ^{b,β}
	II	0.696±0.092 ^{a,α}	0.547±0.161 ^{a,α}	0.409±0.054 ^{a,α}	186±23 ^{b,α}
	III	0.744±0.122 ^{a,α}	0.564±0.158 ^{a,α}	0.457±0.042 ^{b,α}	145±18 ^{a,α}
	IV	0.643±0.136 ^{a,α}	0.511±0.177 ^{a,α}	0.383±0.063 ^{a,α}	202±29 ^{b,α}
	V	0.649±0.145 ^{a,α}	0.521±0.169 ^{a,α}	0.374±0.051 ^{a,α}	192±33 ^{b,α}
Sign ¹		ns	ns	***	***
B	I	0.734±0.111 ^{a,α}	0.563±0.146 ^{a,α}	0.435±0.048 ^{bc,α}	185±24 ^{b,α}
	II	0.714±0.146 ^{a,α}	0.566±0.183 ^{a,α}	0.416±0.066 ^{ab,α}	206±26 ^{b,β}
	III	0.733±0.109 ^{a,α}	0.533±0.119 ^{a,α}	0.468±0.061 ^{c,α}	161±21 ^{a,β}
	IV	0.753±0.164 ^{a,β}	0.593±0.232 ^{a,α}	0.458±0.064 ^{bc,β}	200±32 ^{b,α}
	V	0.657±0.186 ^{a,α}	0.554±0.266 ^{a,α}	0.373±0.054 ^{a,α}	183±36 ^{b,α}
Sign ¹		ns	ns	***	***
C	I	0.728±0.130 ^{a,α}	0.578±0.162 ^{a,α}	0.417±0.076 ^{ab,α}	206±28 ^{bc,β}
	II	0.800±0.126 ^{ab,β}	0.700±0.173 ^{a,β}	0.412±0.047 ^{ab,α}	223±23 ^{c,γ}
	III	0.861±0.117 ^{b,β}	0.714±0.188 ^{a,β}	0.481±0.056 ^{c,α}	164±23 ^{a,β}
	IV	0.740±0.103 ^{a,β}	0.570±0.134 ^{a,α}	0.446±0.059 ^{bc,β}	182±21 ^{ab,α}
	V	0.730±0.125 ^{a,α}	0.650±0.195 ^{a,α}	0.375±0.055 ^{a,α}	200±49 ^{bc,α}
Sign ¹		**	ns	***	***
Sign ²		ns, *, **, *, ns	ns, *, **, ns, ns	ns, ns, ns, ***, ns	*, ***, *, ns, ns

All data are expressed as average value ± standard deviation (n = 30). Different Latin letters within the same column indicate significant differences (¹) among harvest dates at the same berries density (*Tukey-b test*; $p < 0.05$). Different Greek letters within the same column indicate significant differences (²) among the three density classes at the same harvest date (*Tukey-b test*; $p < 0.05$). ^{1,2}: *, **, *** and ns indicate significance at $p < 0.05$, 0.01, 0.001 and not significant, respectively. F_{sk} = berry skin break force, W_{sk} = berry skin break energy, E_{sk} = berry skin Young's modulus, Sp_{sk} = berry skin thickness. A = 1088 kg/m³, B = 1094 kg/m³, C = 1100 kg/m³.

Table 7. Correlation study among mechanical and chemical parameters of berry skins for Nebbiolo grapes

Phenolic composition	F _{sk} (N)	W _{sk} (mJ)	E _{sk} (N/mm)	Sp _{sk} (μm)
TA _{sk} total extractable (mg/kg malvidin-3-glucoside chloride)	ns	ns	0.515*	ns
TA _{sk} easily extractable (mg/kg malvidin-3-glucoside chloride)	ns	ns	0.524*	ns
TF _{sk} total extractable (mg/kg (+)-catechin)	ns	ns	0.621*	ns
TF _{sk} easily extractable (mg/kg (+)-catechin)	ns	ns	0.560*	ns
PRO _{sk} total extractable (mg/kg cyanidin chloride)	ns	ns	0.756**	ns
PRO _{sk} easily extractable (mg/kg cyanidin chloride)	ns	ns	0.705**	ns
FVA _{sk} total extractable (mg/kg (+)-catechin)	0.766**	0.561*	0.615*	ns
FVA _{sk} easily extractable (mg/kg (+)-catechin)	0.774**	0.572*	0.624*	ns
TF _{sk} (% extraction)	ns	ns	ns	0.567*
PRO _{sk} (% extraction)	ns	ns	ns	0.671**
FVA _{sk} (% extraction)	0.618*	ns	0.538*	ns

Significance: *,** and ns indicate significance at $p < 0.05$, 0.01 and not significant, respectively. F_{sk} = berry skin break force, W_{sk} = berry skin break energy, E_{sk} = berry skin Young's modulus, Sp_{sk} = berry skin thickness, TA_{sk} = skin total anthocyanins, TF_{sk} = skin total flavonoids, PRO_{sk} = skin proanthocyanidins, FVA_{sk} = skin flavanols vanillin assay.




Figure 1. Percentage of Nebbiolo grape berries in different density classes, as a function of the ripeness stage.