

AperTO - Archivio Istituzionale Open Access dell'Università di Torino

Glycosaminoglycan Content in Term and Preterm Milk during the First Month of Lactation.

This is the author's manuscript

Original Citation:

Availability:

This version is available <http://hdl.handle.net/2318/89338> since

Published version:

DOI:10.1159/000330848

Terms of use:

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)



UNIVERSITÀ DEGLI STUDI DI TORINO

This is an author version of the contribution published on:

[Neonatology, vol. 101, issue 1 2012, DOI: 10.1159/000330848]

The definitive version is available at:

<http://www.karger.com/Article/FullText/330848>

1 **Glycosaminoglycan content in term and preterm milk during the first month of**
2 **lactation**

3 ¹Giovanni V. Coppa, ¹Orazio Gabrielli, ¹Lucia Zampini, ¹Tiziana Galeazzi, ²Francesca
4 Maccari, ²Dania Buzzega, ²Fabio Galeotti, ³Enrico Bertino, ²Nicola Volpi

5
6 ¹Pediatric Division, Department of Clinical Sciences, Polytechnic University of Marche,
7 Ospedali Riuniti, Presidio Salesi, Ancona, Italy (Prof. G.V. Coppa MD, Prof. O.
8 Gabrielli MD, Dr. L. Zampini PhD. Dr. T. Galeazzi PhD);

9 ²Department of Biology, University of Modena & Reggio Emilia, Modena, Italy (Prof.
10 N. Volpi PhD. Dr. F. Maccari PhD. Dr. D. Buzzega PhD. Dr. F. Galeotti PhD);

11 ³Neonatal Intensive Care Unit, Department of Pediatrics, University of Turin, Turin,
12 Italy (Prof. Bertino MD).

13
14 **Contributors**

15 N.V. developed the applied methodologies. L.Z., T.G., F.M., D.B. and F.B. performed
16 the experimental procedures and analyses. E.B. contributed in milk sample collection.
17 N.V., G.V.C. and O.G. designed and developed the experimental design, performed
18 data analysis and wrote the manuscript.

19 All authors reviewed and approved the study.

20 **Conflicts of interest**

21 We declare that we have no conflicts of interest.

22 **Grant support**

23 None

24 **Running head**

25 Glycosaminoglycans in human milks

26 **Corresponding authors**

27 Prof. Nicola Volpi

28 Department of Biology, Via Campi 213/D

29 University of Modena & Reggio Emilia, 41100 Modena, Italy

30 e-mail: volpi@unimo.it

31 fax number: 0039 59 2055548

32 tel number: 0039 59 2055543

33

34 **Abstract**

35 *Background.* In a recent study, we performed a complete structural characterization
36 of glycosaminoglycans (GAGs) in human mature milk. However, no data are available
37 on the total content of GAGs in human milk from healthy mothers having delivered
38 term or preterm newborns.

39 *Objectives.* In this study, we evaluated the total content of GAGs in pooled milk
40 from healthy mothers having delivered term or preterm newborns during the first month
41 of lactation.

42 *Methods.* Highly specific and sensitive analytical approaches were used to quantify
43 human milk total GAGs.

44 *Results.* Highest GAGs values are present at 4th day (~ 9.3 g/L and ~ 3.8 g/L in
45 preterm and term milk, respectively), followed by a progressive decrease up to day 30th
46 (~ 4.3 g/L and ~ 0.4 g/L). The more remarkable differences are related to the first
47 phases of lactation in which a strong decrease of GAGs was observed between days 4th
48 and 10th (~73% in term and ~50% in preterm).

49 *Conclusions.* During the first month of lactation the absolute amount of
50 polysaccharides was constantly and significantly higher in preterm milk than in term
51 one, with a similar behaviour in the decrease. These data further indicate that human
52 milk GAGs may have an active role in protecting newborns during the first phases of
53 lactation.

54

55 *Keywords.* Human milk; Glycosaminoglycans; Chondroitin sulfate.

56

57 **Introduction**

58 In a recent study [1], we performed a complete structural characterization of
59 glycosaminoglycans (GAGs) in human mature milk in comparison with bovine milk.
60 Quantitative analyses yielded 0.41 g/L of GAGs in human milk, ~7 times more than in
61 bovine milk. In particular, chondroitin sulfate (CS) and dermatan sulfate (DS), were
62 found to differ considerably from one type of milk to the other. In fact, in human milk a
63 low-sulfated CS was the main component (~56%), while DS was observed in very low
64 amount (~2%). On the contrary, bovine milk was demonstrated to be composed of
65 ~66% DS and ~34% CS. Structural analysis also showed the presence of fast-moving

66 heparin (FM-Hep) that account for ~30-40% of total GAGs in both milk samples. The
67 same research [1] offered the first full characterization of GAGs in human milk,
68 providing useful data to gain a better understanding of their physiological role, as well
69 as of their fundamental contribution to the health of the newborn. In fact, several types
70 of microorganisms have also been demonstrated to use CS as receptors for the adhesion
71 to and infection of host cells [1-3]. Furthermore, in human milk, Newburg *et al.* [4]
72 demonstrated that CS or a CS-like moiety is able to inhibit the binding of the HIV
73 envelop glycoprotein gp120 to the cellular CD4 receptor. From these studies it emerges
74 that human milk GAGs, on the contrary of bovine milk, could play a role as soluble
75 receptors. In fact, intact GAG molecules reach the small intestine, as no specific
76 digestive enzyme able to degrade them is present on the intestinal wall.

77 In this study, we evaluated the total content of GAGs in pooled milk from healthy
78 mothers having delivered term or preterm newborns. To include in our study a relatively
79 homogeneous but representative population we selected women who had both the
80 Secretor and Lewis phenotypes, as determined in their saliva by the hemoagglutination-
81 inhibition test. This is the most common phenotype in Europe, as it is present in about
82 70% of the general population [5]. Furthermore, we used pooled milk to have a robust
83 characterization of the total content of GAGs. Additionally, we already observed no
84 great variations between different term healthy mothers at the same time of lactation for
85 the quali/quantitative composition of GAGs that was confirmed quite similar in all
86 human Subjects [1].

87

88 **Results and Discussion**

89 Figure 1 illustrates agarose-gel eletrophoresis utilized to separate and quantify
90 extracted GAGs from different milk samples. Both in term and preterm milk, GAGs
91 showed a constant pattern essentially composed of two main polysaccharides, a low-
92 sulfated CS (~60-70%), FM-Hep (~30-40%) and quite no DS. Quantitative data are
93 shown in Figure 2. As evident, considerable variation of GAGs concentration occurs
94 during the first month of lactation. Highest values are present at 4th day (~ 9.3 g/L and ~
95 3.8 g/L in preterm and term milk, respectively), followed by a progressive decrease up
96 to day 30th (~ 4.3 g/L and ~ 0.4 g/L). The more remarkable differences are related to
97 the first phases of lactation in which a strong decrease of GAGs was observed between
98 days 4th and 10th (~73% in term and ~50% in preterm). During the first month of

99 lactation the absolute amount of polysaccharides was constantly and significantly higher
100 in preterm milk than in term one, with a similar behaviour in the decrease.

101 Comparative analysis of term and preterm milk demonstrate compositional
102 differences in several of their nutritional components [6]. It is well known that several
103 components show different behaviour during lactation. As regard as the carbohydrate
104 component, dynamic variations were observed during the first month. In particular,
105 lactose content increases both in term and preterm milk with higher concentration in the
106 first. On the contrary, oligosaccharides decrease in both types of milk, with higher
107 concentration in preterm milk [7, 8]. Very interestingly, the total amount of GAGs was
108 higher in preterm than term milk, with a strong decrease after the first days of lactation,
109 with the same trend already observed in oligosaccharides.

110 Recent studies showed that some cell surface receptors are constituted by GAGs [2-
111 4], which in this way directly participate in infective and inflammatory processes. As a
112 consequence, along with oligosaccharides, we can suppose that the high concentration
113 of GAGs could be useful for the preterm newborn in defence processes against several
114 pathogens (viruses, bacteria and their toxins) with a receptor-like mechanism preventing
115 the adhesion of pathogens to epithelial cells. Furthermore, CS (and GAGs) is a well
116 known antioxidant and antiinflammatory agent [9] and remarkably, for preterm infants
117 endowed with an immature antioxidant defense system [10], GAGs may have important
118 antioxidant properties. In fact, babies being exclusively fed with mother's milk develop
119 less oxidative stress than babies nourished with formula [11], and colostrum possesses
120 relevant antiinfective and antioxidant properties contributing to the infant's defense
121 against free radicals generated by oxygen administration, infection, or byproducts of
122 nutrient metabolism. In conclusion, further efforts should be addressed to improve our
123 knowledge not only on GAGs composition and structure but also on their role in
124 physiological and pathological conditions.

125

126 **Materials and methods**

127 A morning milk sample was obtained with an electric breast pump at 4, 10, 20, and
128 30 post-partum days from 18 women who delivered at term and from 26 women who
129 delivered between 27th and 35th week of gestation.

130 GAGs were extracted and quantified by analytical procedures reported in details in
131 Coppa *et al.* [1]. 5 mL of milk were defatted with acetone. After centrifugation at

132 10,000 g for 10 min and drying at 60°C for 24 h, the pellet was solubilized in 20 mL of
133 100 mM Na-acetate buffer pH 5.5 containing 5 mM EDTA and 5 mM cysteine. 200 mg
134 of papain were added and the solution incubated for 24 h at 60°C in a stirrer. After
135 boiling for 10 min, the mixture was centrifuged at 5,000 g for 15 min, and three
136 volumes of ethanol saturated with sodium acetate were added to the supernatant. After
137 storing at +4°C for 24 h, the precipitate was recovered by centrifugation at 5,000 g for
138 15 min and dried at 60°C for 6 h. The dried precipitate was dissolved in 20 ml of 50
139 mM NaCl and after centrifugation at 10,000 g for 10 min, the supernatant was applied
140 to a column (2 cm x 7 cm) packed with QAE Sephadex® A-25 anion-exchange resin
141 equilibrated with the same NaCl solution. GAGs were eluted with a linear gradient of
142 NaCl from 50 mM to 1.2 M from 0 to 150 min using low-pressure liquid
143 chromatography (Biologic LP chromatography system from BioRad) at a flow of 1
144 ml/min. Fractions positive to uronic acid assay [1] were collected. Three volumes of
145 ethanol saturated with sodium acetate were added to the pooled fractions and stored at
146 +4°C for 24 h. The precipitate was recovered by centrifugation and dried at 60°C for 12
147 h. The dried precipitate was dissolved in 100 µL distilled water and further analyzed.

148 Quantitative agarose-gel electrophoresis in barium acetate/1,2-diaminopropane was
149 performed essentially as previously reported elsewhere [1].

150

151 **References**

- 152 1. Coppa GV, Gabrielli O, Buzzega D, Zampini L, Galeazzi T, Maccari F, Bertino
153 E, Volpi N. Composition and structure elucidation of human milk
154 glycosaminoglycans. *Glycobiology* 2011;21:295-303.
- 155 2. McClure MO, Moore JP, Blanc DF, Scotting P, Cook GM, Keynes RJ, Weber
156 JN, Davies D, Weiss RA. Investigations into the mechanism by which sulfated
157 polysaccharides inhibit HIV infection in vitro. *AIDS Res Hum Retroviruses*
158 1992;8:19-26.
- 159 3. Wadstrom T, Ljungh L: Glycosaminoglycan-binding microbial proteins on
160 tissue adhesion and invasion: key events in microbial pathogenicity. *J Med*
161 *Microbiol* 1999;48:223-233.
- 162 4. Newburg DS, Linhardt RJ, Ampofo SA, Yolken RH: Human milk
163 glycosaminoglycans inhibit HIV glycoprotein gp120 binding to its host cell CD4
164 receptor. *J Nutr* 1995;125:419-424.

- 165 5. Grimmonprez L : Les glucides du lait. *Ann Nutr Alim* 1971;A39-A79.
- 166 6. Atkinson SA. Effects of gestational stage at delivery on human milk
167 components. In: Jensen RG. Ed. *Handbook of milk composition*. San Diego,
168 Academic Press, 1995, 222-237.
- 169 7. Coppa GV, Gabrielli O, Pierani P, Catassi C, Carlucci A, Giorgi PL: Changes in
170 carbohydrate composition in human milk over 4 months of lactation. *Pediatrics*
171 1993;91:637-641.
- 172 8. Coppa GV, Pierani P, Zampini L, Gabrielli O, Carlucci A, Catassi C, Giorgi PL.
173 Lactose, oligosaccharide and monosaccharide content of milk from mothers
174 delivering preterm newborns over the first month of lactation. *Minerva Pediatr*
175 1997;49:471-475.
- 176 9. Egea J, García AG, Verges J, Montell E, López MG: Antioxidant,
177 antiinflammatory and neuroprotective actions of chondroitin sulfate and
178 proteoglycans. *Osteoarthritis Cartilage* 2010;18 Suppl 1:S24-7.
- 179 10. Rautavaa S, Nanthakumara NN, Dubert-Ferrandona A, Lua L, Rautavaa J,
180 Walkera WA: Breast Milk-Transforming Growth Factor- β 2 Specifically
181 Attenuates IL-1 β -Induced Inflammatory Responses in the Immature Human
182 Intestine via an SMAD6- and ERK-Dependent Mechanism. *Neonatology*
183 2011;99:192-201.
- 184 11. Alberti-Fidanza A, Burini G, Perriello G: Total antioxidant capacity of
185 colostrum, and transitional and mature human milk. *J Matern Fetal Neonatal*
186 *Med* 2002;11:275-279.

187

188 **Legends to Figures**


189

190 **Figure 1.** Agarose-gel electrophoresis of GAGs from human milk samples at different
191 days of lactation in mothers delivering term and preterm newborns. CS: chondroitin
192 sulfte. DS: dermatan sulfate HS: heparan sulfate. St: mix of CS, DS and HS. The
193 suddivision in GAG species CS, DS and HS is just related to the standard.

194


195 **Figure 2.** Total amount of GAGs in human milk during different days of lactation in
196 mothers delivering term and preterm newborns. A curve illustrating the polysaccharides
197 content trend is also reported for each type of milk. Data are reported as mean \pm SD.

198



199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219

Figure 1



220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240

Figure 2