

AperTO - Archivio Istituzionale Open Access dell'Università di Torino

**Review of the paper: Commutator criteria for magnetic pseudodifferential operators**

**This is the author's manuscript**

*Original Citation:*

*Availability:*

This version is available <http://hdl.handle.net/2318/93904> since

*Terms of use:*

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)



## UNIVERSITÀ DEGLI STUDI DI TORINO

**This is an author version of the contribution published on:**

Gianluca Garello.

Review to the paper: Commutator criteria for magnetic pseudodifferential operators. .

*AMS MathSciNet Mathematical Reviews*, Vol. MR2753628 , .

**The definitive version is available at:**

*http : //www.ams.org/mathscinet/search/publdoc.html?pg1 =  
RVRI&pg3 = authreviews&s1 = 343862&vfpref = html&r =  
8&mx - pid = 2753628*

The authors study some commutator techniques in the frame of the twisted pseudodifferential calculus associated to Hamiltonian systems in a bounded, smooth magnetic field. The main goal of the paper is to prove results, analogous to the classic ones, for a generalization of the Weyl calculus adapted to the situation in which a variable magnetic field is also present. In order to take advantage of the gauge invariance the authors formulate the statements about commutators in algebraic frame, using the twisted Moyal algebra.

The main result consists in a generalization of the Beals Criterion [MR0435933 (55 #8884)], in the algebraic setting, which is proved in the case of symbols in the Hörmander class  $S_0^0(\mathbf{R}^{2n})$  and  $S_\rho^m(\mathbf{R}^{2n})$ . A Bony type criterion, see [MR1482829 (98m:35233)], is also proved. Both the criteria are applied to inverses and fractional powers of some twisted pseudo-differential operators. At the end, following the idea of Bony, the authors introduce a class of Twisted Fourier Integral Operators (TFIO) and prove that, for a large class of first order elliptic symbols, the unitary groups they generate (in Hilbert representation) are such TFIO.