

AperTO - Archivio Istituzionale Open Access dell'Università di Torino

Synthesis of water-soluble multidentate aminoalcohol beta-cyclodextrin derivatives via epoxide opening

This is the author's manuscript

Original Citation:

Availability:

This version is available <http://hdl.handle.net/2318/95797> since 2017-12-03T23:33:34Z

Published version:

DOI:10.1016/j.carres.2011.09.018

Terms of use:

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)



UNIVERSITÀ DEGLI STUDI DI TORINO

This Accepted Author Manuscript (AAM) is copyrighted and published by Elsevier. It is posted here by agreement between Elsevier and the University of Turin. Changes resulting from the publishing process - such as editing, corrections, structural formatting, and other quality control mechanisms - may not be reflected in this version of the text. The definitive version of the text was subsequently published in [*insert Carbohydrate research, volume number 346, issue number 17, 2011, doi: 10.1016/j.carres.2011.09.018*].

You may download, copy and otherwise use the AAM for non-commercial purposes provided that your license is limited by the following restrictions:

- (1) You may use this AAM for non-commercial purposes only under the terms of the CC-BY-NC-ND license.
- (2) The integrity of the work and identification of the author, copyright owner, and publisher must be preserved in any copy.
- (3) You must attribute this AAM in the following format: Creative Commons BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/deed.en>), [*doi:10.1016/j.carres.2011.09.018*]

Synthesis of water-soluble multidentate aminoalcohol β -cyclodextrin derivatives via epoxide opening

K. Martina,^[a] M. Caporaso,^[a] S. Tagliapietra,^[a] G. Heropoulos,^[b] O. Rosati,^[c] and G. Cravotto*^[a]

[a] *Dipartimento di Scienza e Tecnologia del Farmaco, University of Torino, Via P. Giuria 9, 10125 Torino, Italy;*

[b] *Institute of Organic and Pharmaceutical Chemistry, National Hellenic Res. Found., Vas. Constantinou Av. 48, Athens, 116 35, Greece;*

[c] *Dipartimento di Chimica e Tecnologia del Farmaco, Università degli Studi, Via del Liceo 1, 06123 Perugia, Italy.*

Abstract

New highly soluble β -aminoalcohol β -cyclodextrin (β -CD) derivatives have been synthesized via nucleophilic epoxide opening reactions with mono-6-amino mono-6-deoxy-permethyl- β -CD and mono-6-amino mono-6-deoxy- β -CD. The binding properties of the β -CD were enhanced by linking aminoalcohol subunits which caused its solubility to improve markedly.

The reaction conditions were optimised using microwave irradiation giving moderate-to-good yields with a series of epoxides. A regioselective epoxide opening reaction was observed in the reaction with styrene oxide while the stereoselectivity was strictly dependent on substrate structure.

Keywords: Cyclodextrin, epoxides, selectivity, microwave activation, solubilization in water

1. Introduction

Cyclodextrins (CDs) possess a well-studied ability to act as hosts for hydrophobic guests in water.¹ Due to their homochirality, CDs may be used as molecular receptors and constitute examples of supramolecular chemistry.² Chromatographic³ and electrophoretic⁴ methods as well as stereoselective reactions mediated by CDs exploit this peculiar property which has been widely explored by calorimetric, X-ray diffraction, NMR, and computational studies.⁵

Because of the regularity of the β -CD ring, the substitution of one or more hydroxyl groups affects the annular asymmetry, increasing solubility and influencing its binding properties.

It has been observed that the substituents on the CD's wider side, exhibit different host properties than those on the narrower, primary hydroxyl side. Therefore the synthesis of monosubstituted CDs has become important over the years and complex procedures are generally required to obtain regioselectively substituted derivatives.⁶ Monosubstituted derivatives have been used in the synthesis of dimers and oligomers.⁷

As has already been reported,⁸ the solubility of β -CD can be markedly improved with the attachment of suitable moieties. Even a random and low degree of substitution with hydrophilic hydroxyethyl- or oligo(ethylene glycol)- groups disrupts the hydrogen bond molecular network and dramatically improves water solubility. This solubilising effect has been observed in the reaction of β -CD with propylene oxide and, as a consequence, 6-O(2-hydroxypropyl) β -CD showed higher solubility than the 2-O substituted derivative.⁹


The ring opening of epoxides by nucleophiles is one of the most versatile and well established reactions of all "click chemistry" protocols. This reaction is typically carried out in the presence of a catalyst and a large excess of amines at elevated temperatures. For this reason, non-conventional techniques, besides the more classical procedures, have been used to promote the reaction.¹⁰ The reaction of epoxides with amines affords β -aminoalcohols which are important intermediates for carbohydrate and nucleoside synthesis. When linked to β -CD, the β -aminoalcohol moiety can enhance the CD's binding property and favourably affects its solubility. These derivatives can efficiently form metal complexes with β -CD which can then be applied as supramolecular catalysts in regio- or stereoselective reactions.¹¹ These chemically reactive subunits interact with the bound guests making a complementary metal coordination effect. Furthermore, aminoalcohol derivatives of β -CD may be very attractive as powerful chiral selectors in capillary electrophoresis, as charged β -CDs they are endowed with a high resolution ability without interfering with the detection of analytes.¹²

To the best of the authors' knowledge, there are very few reports on the synthesis of β -CD aminoalcohols and, in these studies, they are usually obtained by displacement of the tosyl group from the 6-monodeoxy-6-mono-(*p*-tolylsulfonyl)- β -CD with an aminoalcohol.^{10,12c} The direct etherification of β -CD with epoxides was reported by Pitha et al.⁹ and confirmed by Wenz et al.¹³ They demonstrated that in aqueous alkali and the presence of propylene oxide, β -CD is alkylated on the primary or the secondary face which results in a mixture of 2-O, 3-O, 6-O substituted CDs. After long reaction times, the desired product is isolated in very low yield. Furthermore β -CD, native or modified was used as an epoxide opening polymerization initiator with ethylene oxide, so obtain a new family biomimetic synthetic pores.¹⁴ There are no publications that refer to the ring opening of epoxides by monoamino β -CD. In the light of the widespread applications of aminoalcohol β -CD derivatives, we have systematically studied the nucleophilic opening of epoxides with monoamino β -CD: a straightforward way of obtaining a range of multidentate and water-soluble aminoalcohol β -CD derivatives. In this paper the alkylation of monoamino β -CD has been achieved with various epoxides and with the aim of exploring the relationship between structure and reactivity. Mono-6-amino mono-6-deoxy-permethyl- β -CD was compared to the mono-6-amino mono-6-deoxy- β -CD in such a way as to evaluate the reactivity of the hydroxyl group and the influence of epoxide inclusion. The optimization of these reactions was studied under microwave (MW) irradiation. The regioselectivity of the attack was studied when the reaction was performed with styrene oxide, whereas cyclohexene oxide was chosen as a symmetrically substituted epoxide which could report on whether its inclusion into CD yields stereoselective effects.

Given the fact that most terminal epoxides are inexpensive and easily available, we also decided to apply this synthetic approach to the preparation of bifunctional molecules by reacting monoamino β -CD with diethylene glycol diglycidyl ether and octadiene dioxide.

2. Results and discussion

Mono-6-amino mono-6-deoxy-permethyl- β -CD was chosen as a preliminary β -CD substrate and was reacted with styrene oxide in a series of reactions carried out with the aim of optimising conditions. (Scheme 1, Table 1).



Scheme 1

Reaction of mono-6-amino mono-6-deoxy-permethyl- β -CD (**1**) and mono-6-amino mono-6-deoxy- β -CD (**2**) with styrene oxide.

Table 1

Assessment of alkylation of mono-6-amino mono-6-deoxy-permethyl- β -CD (**1**) and mono-6-amino mono-6-deoxy- β -CD (**2**) with styrene oxide.

Entry	Reagent	Eq. Epoxide	Reaction Condition, Time	Yield ^{a,b}
1	1	5	LiBr, DMF, 60°C, 24h	18%
2	1	5	LiBr, DMF, 85°C, 4h, MW	35%
3	1	1	LiBr, DMF, 60°C, 24h	18%
4	1	1	LiBr, DMF, 85°C, 24h	38%
5	1	1	LiBr, DMF, 85°C, 4h, MW	50%
6	1	1	DMF, 85°C, 4h, MW	52%
7	2	1	DMF, 85°C, 4h, MW	-
8	2	5	LiBr, DMF, 85°C, 19h	18%
9	2	5	LiBr DMF, 85°C, 4h, MW	19%
10	2	5	DMF, 85°C, 4h, MW	32%
11	2	5	H ₂ O, 85°C, 4h, MW	traces

^a All the products were characterized by ¹H NMR, 2D NMR(¹H-¹³C HMQC, ¹H-¹H COSY), IR spectroscopy and mass spectrometry; ^b isolated yield obtained after column chromatography;


The reactions were performed at 60 °C or 85 °C in DMF under conventional heating or MW irradiation. The epoxide ratio was varied from 1 to 5 eq. and in some experiments LiBr was added. We monitored reaction conversion by TLC, and the product was isolated by silica gel column chromatography. The optimised procedure was setup at 85 °C, under MW irradiation with 1 eq of styrene oxide, the presence of LiBr points to the catalyst lack of effect on the conversion and yield. As already observed in reported derivatizations of CDs,^{7a,7b,7d,15} MW irradiation promotes the reaction cutting down reaction times from 24 to 4 h. As regards the regiochemistry of the ring opening, styrene oxide displays a general tendency for regioselective attack at the benzylic position when less

nucleophilic aromatic amines are employed. In the presence of aliphatic amines a lower degree of regioselectivity is observed and the reaction is controlled by steric factors. The preferential reaction at the non-benzylic position is achieved only with a catalyst.¹⁶ When permethyl β -CD was reacted with styrene oxide a single regioisomer was isolated. Using a combination of ^1H NMR and 2D NMR experiments (^1H - ^{13}C HMQC, ^1H - ^1H COSY), we were able to discover that the product that resulted from the nucleophilic attack at the less-hindered epoxide carbon was the obtained regioisomer. Likewise, the same reaction with mono-6-amino mono-6-deoxy- β -CD (Entries 7-10, Table 1) selectively attacks the β position (**5**) but differently from the reaction with permethyl β -CD (**1**), 5 eq of epoxide were required to achieve conversion. Turning to stereochemistry, when styrene oxide was reacted with 6-monoamino β -CD derivatives **2**, TLC showed two spots with different intensities and R_f values. Reverse phase separation yielded an inseparable mixture of the two products (same molecular weights by MS spectrometry and same ^1H NMR spectra). For this reason, we assume that the mixture was composed of two epimers with slightly different polarities. The diastereomeric ratio of product **4** was measured by HPLC and was found to be 9:1 for the stereoisomeric secondary alcohol. To evaluate the influence of the solvent on the stereoselectivity, the reaction was repeated in water but only traces of product was observed by TLC and HPLC. By HPLC we could not separate the two epimers of products **3**, the two spots were evidenced only by bidimensional TLC.

In the second part of this study, cyclohexene oxide was taken as a symmetrically substituted epoxide and reacted with permethyl mono-6-amino β -CD in DMF.

As with the experiment above, the influence of LiBr catalysis, the heating conditions and the number of equivalents of epoxides were compared (Scheme 2, Table 2). The reaction was monitored by TLC and the hydroxycyclohexylamino β -CD derivative (**5**) was isolated by flash chromatography.

Differently to styrene oxide, 4 eq of cyclohexene oxide were added to obtain the desired product in satisfactory yields. The reaction required a long reaction time when performed under conventional heating and more than 5 equivalents of epoxide did not provide higher yields (Entry 4, Table 2). The reaction was optimized by heating under MW at 85 °C for 4 h and the final product was obtained in 27% yield.



Scheme 2.

Reaction of mono-6-amino-mono-6-deoxy-permethyl- β -CD (**1**) and mono-6-amino-mono-6-deoxy- β -CD (**2**) with cyclohexene oxide.

Table 2.

Assessment of alkylation of mono-6-amino-mono-6-deoxy-permethyl- β -CD (**1**) and mono-6-amino-mono-6-deoxy- β -CD (**2**) with cyclohexene oxide.

Entry	Reagent	Eq. Epoxide	Reaction condition	Yield ^{b,c}
1	1	1	LiBr, 24h	5%
2	1	4	LiBr, 24h	12%
3	1	4	LiBr, 48h	16.8%
4	1	7	LiBr, 26h	15%
5	1	4	LiBr, 4h, MW	26%
6	1	4	4h, MW	27%
7	2	1	48h	Failed
8	2	10	LiBr, 36h	9.2%
9	2	4	4h, MW	Failed
10	2	10	4h, MW	30%
11	2	10	LiBr, 4h, MW	28%

^a The reaction was performed at 85°C in DMF; ^b all the products were characterized by ¹H NMR, 2D NMR (¹H-¹³C HMQC, ¹H-¹H COSY), and mass spectrometry; ^c isolated yield obtained after column chromatography.

The reaction was repeated on mono-6-amino-mono-6-deoxy- β -CD. The reaction showed the best results when performed with 10 eq of cyclohexene oxide under the optimal conditions found for the permethyl derivatives. The influence of MW in promoting the reaction was confirmed as had been observed for the permethyl derivative.

Regarding the stereochemistry, nucleophilic ring epoxide opening almost certainly involves an S_N2 reaction and occurs with an inversion in configuration at the point of attack. Bearing this in mind, it


has been assumed that this opening method would give trans-diaxial rather than trans-diequatorial products. Trans-diaxial cleavage proceeded via a transition state which is more energetically favourable than in the trans-diequatorial epoxide opening.¹⁷ Assignment of the relative configuration of the β -aminoalcohol derivative is possible via a comparison of the coupling constants 3J .

When cyclohexene oxide was reacted with **1** neither TLC nor HPLC revealed stereospecificity. But when we monitored the synthesis of product **6**, TLC showed two spots as had been previously observed with styrene oxide. Reverse phase separation yielded only the major isomer because the minor was present in a negligible amount. A ^1H NMR spectrum of isolated compound **6** (see Figure 1) was characteristic because the cyclohexyl moiety was found to be strongly influenced by the presence of CD. All the cyclohexanol diastereotopic methylene groups were seen to split each other. 2D-NMR experiments (^1H - ^{13}C HMQC, ^1H - ^1H COSY) were employed to assign the peaks and remove any doubt. From this spectrum it was seen that the complexity of CD derivative **6** renders J coupling measurement impossible so that we were unable to verify whether the final product had been isolated in the trans-diaxial or the trans-diequatorial form. The strong interaction between the CD core and the substituent, as shown in the NMR, has promoted the hypothesis of a stereoselective ring opening.

Figure 1

^1H NMR spectrum of 6-mono(2-hydroxycyclohexylamino)-6-monodeoxy- β -CD (**7**). On the right a particular of 2D HMQC (^1H , ^{13}C).

The third part of this study focused on the application of the epoxide ring opening reaction to terminal epoxides.



Scheme 3.

Reaction of mono-6-amino-mono-6-deoxy- β -CD (**1**) and mono-6-amino-mono-6-deoxy-permethyl- β -CD (**2**) with terminal epoxides **7** and **8**

Table 3.

Results of reaction of mono-6-amino-mono-6-deoxy- β -CD (**1**) and mono-6-amino-mono-6-deoxy-permethyl- β -CD (**2**) with terminal epoxides **7** and **8**

Entry	Reagent	Epoxide	Reaction condition	Yield ^{a,b}
1	1	7	10 eq of 7, DMF, 65-85°C, 4h, MW	4%
2	2	7	10 eq of 7, DMF, 85°C, 4h, MW	9%
3	1	8	10 eq of 8, DMF, 65-85°C, 4h, MW	30%
4	2	8	10 eq of 8, DMF, 65-85°C, 4h, MW	52%

^a all the products were characterized by ¹H NMR, 2D NMR (¹H-¹³C HMQC, ¹H-¹H COSY), and mass spectrometry; ^b isolated yield obtained after column chromatography.

Monoamino CD derivatives **1** and **2** were reacted with diethylene glycol diglycidyl ether (**7**) to give a new highly soluble and functionalized CD derivative. The reaction required 10 eq to obtain satisfactory conversion rates. However, product **9** was only obtained in a 4% yield and derivative **10** was isolated in a slightly better 9% yield. As has already been mentioned, nucleophilic epoxide opening with CD does not afford complete conversion but, differently from the reactions already described, diethylene glycol diglycidyl ether gave a mixture of side products and so the desired products were difficult to isolate.

As expected, product **10** showed higher solubility in water than the starting material. Therefore, to reach our objective, we evaluated a new epoxide substrate from which the desired β -aminoalcohol CD derivative was obtained in a higher yield. 1,7-octadiene was epoxidated with MCPBA and isolated in a 95% yield to obtain 1,7-octadiene dioxide (**8**). The reaction was repeated, with the same conditions as for diethylene glycol diglycidyl ether, and products **11** and **12** were obtained with 30 and 52% yields respectively (Table 3).

After NMR and MS analysis we found that the ring opening reaction with the octadiene dioxide occurred twice so that product **12** is a disubstituted β -CD. When repeated with fewer equivalents of 1,7-octadiene dioxide (**8**) the reaction with β -CD **2** did not afford the desired monosubstituted derivative in satisfactory yields.

3. Conclusions

Overall it can be concluded that nucleophilic epoxide opening with mono-6-amino mono-6-deoxy- β -CD and mono-6-amino mono-6-deoxy-permethyl β -CD, promoted by MW, can be considered a versatile way to obtain β -aminoalcohol derivatives of β -CD in acceptable yields. The selective formation of the β regioisomer was observed when the reaction was performed with styrene oxide and excellent stereoselectivity was achieved in the nucleophilic epoxide cleavage of styrene and cyclohexene oxides with mono-6-amino mono-6-deoxy- β -CD. These newly prepared compounds showed high water solubility and the linked aminoalcohol subunits enhanced the CD binding properties. Using monoamino β -CD and terminal diepoxides this protocol afforded new functionalized β -CD derivatives, versatile intermediates which can be exploited for further modifications.

4. Experimental

Commercially available reagents and solvents were used without further purification. Native CDs were kindly provided by Wacker Chemie. 6-amino-mono-6-deoxy- β -CD (**1**) and mono-6-amino-mono-6-deoxy-permethyl- β -CD (**2**) were prepared in accordance with the well established procedure.^{18,19} Diethylene glycol diglycidyl ether was purified from a commercially available 50% pure solution (column chromatography DCM/MeOH). Reactions were monitored by TLC on Merck 60 F254 (0.25 mm) plates. Spot detection was carried out by staining with 5% H₂SO₄ in ethanol or with phosphomolybdic acid stain. IR spectra were recorded with a Shimadzu FT-IR 8001 spectrophotometer. NMR spectra were recorded with a Bruker 300 Avance (300 MHz and 75 MHz for ¹H and ¹³C, respectively) at 25 °C; chemical shifts are calibrated to the residual proton and carbon resonances of the solvent: DMSO-*d*₆ (δ H = 2.54, δ C = 39.5). Chemical shifts (δ) are given in ppm, coupling constants (J) in Hz. ESI-mass spectra were performed on a Waters Micromass ZQ spectrometer equipped with ESI source.

4.1 6-Mono(2-hydroxyl-2-phenyl)ethylamino-6-monodeoxy-permethyl- β -CD (**3**)

The reaction was carried out under magnetic stirring in a professional MW oven; the temperature was monitored with a fiber-optic thermometer. Mono-6-amino mono-6-deoxy-permethyl- β -CD (130 mg, 0.09 mmol) was dissolved in anhydrous DMF (2 mL) and styrene oxide (10.5 μ L, 0.09 mmol) was added. The mixture was irradiated with MW (160 W) at 85 °C for 4 h. The reaction was monitored by TLC (CHCl₃/MeOH 9:1) spots were visualized with a phosphomolybdic acid stain. The solvent was partially evaporated and the reacted mixture was diluted with CH₂Cl₂, washed with water, and finally dried (Na₂SO₄). The crude residue was purified by CC (CH₂Cl₂/CH₃OH) to yield 73 mg (52% yield) of the desired product.

3 is a white powder; R_f = 0.61 (CHCl₃/CH₃OH 9:1); ¹H NMR (300 MHz, CDCl₃) δ = 7.36-7.27 (m, 5H, H-Ar), 5.12 (m, 7H, H-1), 4.75 (m, 1H, Ph-CH(OH)), 3.84-3.81 (m, 14H, H-5, H-6^A), 3.64-3.502 (m, 63H, 3-OCH₃, 2-OCH₃, H-4, H-6^B, H-3), 3.302 (m, 18H, 6-OCH₃), 3.29 (m, 7H, H-2), 2.95 (m, 1H, CH₂-NH), 2.78 (m, 1H, CH₂-NH) ppm; ¹³C NMR (75 MHz, CDCl₃): δ = 142.6 (C_{ipso}), 128.72-126.2 (C_{orto}, C_{meta}, C_{para}), 99.3 (C1), 82.4-82.06 (C2, C3) 80.81-80.51 (C4), 71.75-71.47 (C6, PhCHOH), 71.31-71.17 (C5), 61.86-61.76 (3-OCH₃), 59.27-59.24 (2-OCH₃), 59.01-58.69 (6-OCH₃), 57.84 (CH₂NH) ppm; MS (ESI): *m/z* calcd. for C₇₀H₁₁₉NO₃₅ [M+H]⁺ 1535.7 found 1535.87, C₇₀H₁₁₉NO₃₅ [M+H+Na]²⁺ 779.35 found 779.25.

4.2 6-Mono(2-hydroxyl-2-phenyl)ethylamino-6-monodeoxy- β -CD (4)

The reaction was carried out under magnetic stirring in a professional MW oven, the temperature was monitored with a fiber-optic thermometer. Mono-6-amino-mono-6-deoxy- β -CD (100 mg, 0.09 mmol) was dissolved in anhydrous DMF (2 mL) and styrene oxide (50.2 μ L, 0.44 mmol) was added. The mixture was irradiated with MW (160 W) at 85 °C for 4 h. It was partially evaporated, dissolved with 800 μ L of water then precipitated with cold acetone. The precipitate was purified by reverse phase column chromatography (H₂O/CH₃OH gradient from 97:3 to methanol 100%) to yield 36 mg (32% yield) of the desired product.

5 is a white powder; $R_f = 0.45$ (iPrOH/H₂O/EtOAc/NH₄OH = 5:3:1:1); ¹H NMR (300 MHz, DMSO-*d*₆) $\delta = 7.37$ - 7.27 (m, 5H, H-Ar), 5.82-5.71 (m, 14H, O(2)H, O(3)H), 4.88 (br.s. 7H, H-1), 4.66 (m, 1H, PhCH(OH)), 4.53 (m, 6H, O(6)H), 3.7-3.6 (m, 28H, H-6, H-3), 3.5-3.2 (m, overlaid with water, H-2, H-4, H-5), 2.61 (m, 2H, CH₂-NH) ppm; ¹³C NMR (75 MHz, DMSO) $\delta = 145.47$ (C_{ipso}), 128.76 (C_{meta}), 126.72 (C_{orto}, C_{para}), 102.84 (C1), 82.4 (C4), 73.3 (C3), 73.28 (C2), 72.88 (C5), 70.48 (PhCH(OH)), 57.5 (C6), 57.35 (CH₂NH) 49.64 (C6') ppm; MS (ESI): *m/z* calcd. for C₅₀H₇₉NO₃₅ [M+H]⁺ 1254.4 found 1255.45.

4.3 6-Mono(2-hydroxycyclohexyl)amino-6-monodeoxy-permethyl- β -CD (5)

The reaction was carried out under magnetic stirring in a professional MW oven, the temperature was monitored with a fiber-optic thermometer. Mono-6-amino mono-6-deoxy-permethyl- β -CD (150 mg, 0.11 mmol) was dissolved in anhydrous DMF (1 mL) and cyclohexene oxide (43 μ L, 0.42 mmol) was added. The mixture was irradiated with MW (160 W) at 85 °C for 4 h. The reaction was monitored by TLC (CHCl₃/MeOH 9:1) spots were visualized with a phosphomolybdic acid stain. The solvent was partially evaporated and the reacted mixture was diluted with CH₂Cl₂, washed with water, and finally dried (Na₂SO₄). The crude residue was purified by CC (CH₂Cl₂/CH₃OH) to yield 50 mg (27% yield) of the desired product.

6 is a white powder; $R_f = 0.54$ (CHCl₃/CH₃OH 9:1); ¹H NMR (CDCl₃, 300 MHz) $\delta = 5.18$ - 5.03 (m, 7H, H-1), 4.05-3.98 (m, 1H, H-6^A), 3.87-3.82 (m, 14H, H-5, H-6^A), 3.64-3.52 (m, 64H, 3-OCH₃, 2-OCH₃, H-4, H-6^B, H-3, H-a), 3.40 (m, 19H, 6-OCH₃, H-6^B), 3.22-3.18 (m, 7H, H-2), 1.27 (m, 4H, H-c, H-f), 0.081 (m, 4H, H-d, H-e) ppm; ¹³C NMR (CDCl₃, 75 MHz): $\delta = 98.7$ (C-1), 82.15, 81.87, 81.66 (C-2, C-3), 80.70, 80.30, 80.14 (C-4), 72.42, 71.64, 71.3 (C-5, C-6), 62.9, 61.84, 61.62 (3-OCH₃), 58.73 (2-OCH₃), 57.73 (6-OCH₃), 42.9 (C-6'), 30.01 (C-c), 23.6 (C-f), 21.7, 21.2 (C-d, C-e) ppm; MS (ESI): *m/z* calcd. for C₆₈H₁₂₁NO₃₅ [M+H]⁺ 1511.77, found 1512.03.

4.4 6-Mono(2-hydroxycyclohexylamino)-6-monodeoxy- β -CD (6):

The reaction was carried out under magnetic stirring in a professional MW oven, the temperature was monitored with a fiber-optic thermometer. Mono-6-amino mono-6-deoxy- β -CD (100 mg, 0.09 mmol) was dissolved in anhydrous DMF (2 mL) and cyclohexene oxide (100 μ L, 0.88 mmol) was added. The mixture was irradiated with MW (160 W) at 85 °C for 4 h. It was partially evaporated, dissolved with 800 μ L of water then precipitated with cold acetone. The precipitate was purified by reverse phase column chromatography (H₂O/CH₃OH gradient from 97:3 to methanol 100%) to yield 33 mg (30% yield) of the desired product.

7 is a white powder; R_f = 0.41 (iPrOH/ H₂O/EtOAc/NH₄OH = 5:3:1:1); ¹H NMR (DMSO-d₆, 300 MHz) δ = 5.72 (m, 14H, O(2)H, O(3)H), 4.83 (m, 7H, H-1), 4.62-4.45 (m, 6H, O(6)H), 3.63-3.48 (m, 28H, H-6, H-3), 3.40-3.2 (m, overlaid with water, H-2, H-4, H-5), 3.04-2.98 (m, 2H, H-b, H-6^A), 2.64 (d, J= 11.4 Hz, H-6^B), 2.11 (m, 1H, H-a), 1.91 (d, 1H, H-f), 1.77 (m, 1H, H-c), 1.56 (m, 2H, H-e, H-d), 1.17 (m, 3H, H-c, H-d, H-e), 0.85 (m, 1H, H-f) ppm; ¹³C NMR (DMSO, 75 MHz): δ = 102.84 (C1), 82.5 (C4), 73.87 (C3), 73.32 (C-b), 73.27(C2), 72.90 (C5), 63.16 (C-a), 60.7 (C-6), 46.23 (C-6'), 34.54 (C-c), 30.03 (C-f), 24.86 (C-d, C-e) ppm; MS (ESI): m/z calcd. for C₄₈H₈₁NO₃₅ [M+H]⁺ 1232.46 found 1233.14.

4.5 1,7-Octadiene dioxide (8)

Octadiene (1.37 mL, 9.07 mmol) was dissolved in 25 mL of DCM and stirred at 0 °C. MCPBA (4.68 g, 18.08 mmol) dissolved in 45 mL of DCM was added dropwise. The solution was left at 0 °C under stirring overnight. The reaction was monitored by TLC (PE/EtOAc 8:2) spots were visualized with a phosphomolybdic acid stain. The solution was washed three times with water and finally dried (Na₂SO₄). 1.39 g (95% yield) of the desired product was recovered and employed without further purifications. Analytical data were in accordance with reported values.²⁰

4.6 6-Mono(2-hydroxy-3-(2-(2-(oxiran-2-ylmethoxy)ethoxy)ethoxy) propylamino)-6-monodeoxy-permethyl- β -CD (9)

The reaction was carried out under magnetic stirring in a professional MW oven, the temperature being monitored with a fiber-optic thermometer. Mono-6-amino mono-6-deoxy-permethyl- β -CD (300 mg, 0.212 mmol) was dissolved in anhydrous DMF (2 mL) and diethylene glycol diglycidyl ether (292 μ L, 2.12 mmol) was added. The mixture was irradiated with MW (160 W) at 65 °C for 1h then

85 °C for 3h. The reaction was monitored by TLC (CHCl₃/MeOH 95:5) spots were visualized with phosphomolybdic acid stain. The solvent was partially evaporated and the reacted mixture was diluted with CH₂Cl₂, washed with brine, and finally dried (Na₂SO₄). The crude residue was purified by column chromatography (CH₂Cl₂/CH₃OH) to yield 14 mg (4% yield) of the desired product.

10 is white powder; $R_f = 0.54$ (CHCl₃/CH₃OH 9:1); ¹H NMR (CDCl₃, 300 MHz) $\delta = 5.14$ (br.s 7H, H-1), 4.01 (m, 1H, H-6^A), 3.86-3.81 (m, 14H, H-5, H-6^A), 3.65-3.51 (m, 64H, 3-OCH₃, 2-OCH₃, H-4, H-6^B, H-3, CH-NH), 3.41 (m, 18H, 6-OCH₃, H-6^B), 3.21-3.18 (m, 23H, H-2, H-e, H-a, H-b, H-c, H-d), 2.81 (m, 1H, H-f), 2.63 (m, 1H, H-f), ppm; ¹³C NMR (CDCl₃, 75 MHz): $\delta = 98.9$ (C-1), 80.04 (C-2), 81.67 (C-3), 80.50 (C-4), 71.45, 71.47 (C-c, C-d), 71.83, 71.51, 71.22 (C-5, C-6), 61.28 (3-OCH₃), 58.92 (2-OCH₃), 58.41 (6-OCH₃), 59.3 (CH₂NH), 52.17 (C-e), 47.35 (C-f); MS (ESI): m/z calcd. for C₇₂H₁₂₉NO₃₉ [M+H]⁺ 1632.81 found 1633.36.

4.7 6-Mono(2-hydroxy-3-(2-(2-(oxiran-2-ylmethoxy)ethoxy)ethoxy) propylamino)-6-monodeoxy- β -CD (**10**)

The reaction was carried out under magnetic stirring in a professional MW oven, the temperature being monitored with a fiber-optic thermometer. Mono-6-amino mono-6-deoxy- β -CD (300 mg, 0.26 mmol) was dissolved in anhydrous DMF (2 mL) and diethylene glycol diglycidyl ether (363 μ L, 2.64 mmol) was added. The mixture was irradiated with MW (160 W) at 85 °C for 4h. It was partially evaporated, dissolved with 800 μ L of water then precipitated with cold acetone. The precipitate was purified by reverse phase column chromatography (H₂O/CH₃OH gradient from 97:3 to methanol 100%) to yield 26 mg (9% yield) of the desired product.

11 is a white powder; $R_f = 0.49$ (iPrOH/H₂O/EtOAc/NH₄OH = 5:3:1:1); ¹H NMR (DMSO-d₆, 300 MHz) $\delta = 5.87$ -5.68 (m, 14H, O(2)H, O(3)H), 4.83 (m, 7H, H-1), 4.48-4.46 (m, 6H, O(6)H), 3.77-3.25 (m, 28H), 3.35-3.31 (m, overlaid with water, H-2, H-4, H-5, H-6, H-3, H-b, H-c, H-d), 2.92 (m, 1H, H-e), 2.75 (m, 1H, H-f), 2.47 (m, 2H, H-f, H-a) 2.35 (m, 1H, H-a), ppm; ¹³C NMR (DMSO-d₆, 75 MHz): $\delta = 102.8$ (C1), 82.38 (C4), 73.88 (C3), 73.27 (C2), 72.89 (C5), 70.15, 70.77 (C-c, C-d), 66.27 (C-b) 62.42 (C-a), 60.76 (C6), 52.42 (C-e), 47.85 (C-f) ppm; MS (ESI): m/z calcd. for C₅₂H₈₉NO₃₉ [M+H]⁺ 1352.50 found 1353.03, [M+Na+H]²⁺ 688.24 found 688.10.

4.7 6-Mono(2-hydroxy-6-(oxyran2-yl)hexylamino)-6-monodeoxy-permethyl- β -CD (**11**)

The reaction was carried out under magnetic stirring in a professional MW oven, the temperature was monitored with a fiber-optic thermometer. Mono-6-amino mono-6-deoxy-permethyl- β -CD (150 mg,

0.11 mmol) was dissolved in anhydrous DMF (1.5 mL) and 1,7-octadiene dioxide (150mg, 1.06 mmol) was added. The mixture was irradiated with MW (160 W) at 85 °C for 4h. The reaction was monitored by TLC (CHCl₃/MeOH 9:1) spots were visualized with a phosphomolybdic acid stain. The solvent was partially evaporated and the reacted mixture was diluted with CH₂Cl₂, washed with water, and finally dried (Na₂SO₄). The crude residue was purified by column chromatography (CH₂Cl₂/CH₃OH) to yield 47 mg (30% yield) of the desired product.

12 is a white powder; $R_f = 0.55$ (CHCl₃/CH₃OH = 9:1); ¹H NMR (CDCl₃, 300 MHz) $\delta = 5.11$ (m, 7H, H-1), 3.99 (m, 1H, H-6^{A'}), 3.87 (m, 14H, H-5, H-6^A), 3.62-3.49 (m, 64H, 3-OCH₃, 2-OCH₃, H-4, H-6^B, H-3, CH-NH), 3.45 (m, 18H, 6-OCH₃), 3.18 (m, 7H, H-2), 2.89 (m, H-g), 2.73 (m, 1H, H-h), 2.44 (m, 1H, H-h), 2.04 (m, 2H, CH₂ aliphatic chain), 1.50 (m, 6H, CH₂ aliphatic chain) ppm; ¹³C NMR (CDCl₃, 75 MHz) $\delta = 98.65, 98.56$ (C-1), 82.08, 81.53 (C-2, C-3), 80.23, 80.11 (C-4), 71.69, 71.27, 71.01 (C-5, C-6), 61.59 (3-OCH₃), 59.15 (2-OCH₃), 58.59 (6-OCH₃), 42.47 (C-6'), 51.85 (C-g), 47.58 (C-h), 33.87 (CH₂ aliphatic chain) 29.8 (CH₂ aliphatic chain) ppm; MS (ESI): m/z calcd. for C₇₀H₁₂₅NO₃₆ [M+H]⁺ 1556.8 found 1557.1.

4.7 6-Mono(di-2-hydroxy-6-(oxyran2-yl) hexylamino)-6-monodeoxy- β -CD (**12**)

The reaction was carried out under magnetic stirring in a professional MW oven, the temperature was monitored with a fiber-optic thermometer. Mono-6-amino mono-6-deoxy- β -CD (400 mg, 0.35 mmol) was dissolved in anhydrous DMF (3 mL) and 1,7-octadiene dioxide (501 mg, 3.52 mmol) was added. The mixture was irradiated with MW (160 W) at 85 °C for 4h. It was partially evaporated, dissolved with 800 μ L of water then precipitated with cold acetone. The precipitate was purified by reverse phase column chromatography (H₂O/CH₃OH gradient from 97:3 to methanol 100%) to yield 234 mg (52% yield) of the desired product.

13 is a white powder; $R_f = 0.46$ (iPrOH/H₂O/EtOAc/NH₄OH = 5:3:1:1); ¹H NMR (DMSO-d₆, 300 MHz) $\delta = 5.87-5.68$ (m, 14H, O(2)H, O(3)H), 4.86 (m, 7H, H-1), 4.64-4.51 (m, 6H, O(6)H), 3.66-3.21 (m, overlaid with water, H-2, H-4, H-5, H-6, H-3, H-b), 2.87 (m, 1H, H-g), 2.72 (m, 1H, H-h), 2.44 (m, 2H, H-h, H-a) 2.32 (m, 1H, H-a) 1.5-1.15 (m, 16H, H-c, H-d, H-e, H-f) ppm; ¹³C NMR (DMSO-d₆, 75 MHz): $\delta = 103.04$ (C1), 82.49 (C4), 74.03 (C3), 73.05 (C2), 72.85 (C5), 66.58 (C-b) 62.16 (C-a), 60.31 (C6), 52.43 (C-g), 47.13 (C-h), 31.9 (C-c, C-f), 26.82, 26.38, 26.47, 25.82 (C-d, C-e) ppm; MS (ESI): m/z calcd. for C₅₈H₉₉NO₃₈ [M+H]⁺ 1418.58 found, [M+Na+H]²⁺ 721.29 found 721.12.

Acknowledgments

Financial support from Regione Piemonte (NanoIGT, Converging Technologies 2007) and MIUR (PRIN 2008 - A Green Approach to Process Intensification in Organic Synthesis) is gratefully acknowledged.

Supplementary Data

Supplementary data associated with this article can be found in the online version at [doi:10.1016/j.carres.2011.09.018](https://doi.org/10.1016/j.carres.2011.09.018)

References

1. a) Szejtli, J.; *Cyclodextrin and their inclusion complexes*, Akademiai Kiado, Budapest, **1982**;
b) Rekharsky, M. V.; Inoue, Y.; *Chem. Rev.*, **1998**, 98, 1875-1917.
2. Easton, C. J.; Lincoln, S. F.; *Modified Cyclodextrins, Scaffolds and Templates for Supramolecular Chemistry*, London, **1999**.
3. a) Wistuba, D.; *J. Chrom. A*, **2010**, 1217, 941-952; b) Bicchi, C.; Cagliero, C.; Liberto, E.; Sgorbini, B.; Martina, K.; Cravotto, G.; Rubiolo, P.; *J. Chrom. A*, **2010**, 1217, 1106-1113; c) Dittman, H.; Scharwächter, K.; König, W.A.; *Carbohydr. Res.*, **2000**, 324, 75-96.
4. Cucinotta, V.; Contino, A.; Giuffrida, A.; Maccarrone, G.; Messina, M.; *J. Chrom A*, **2010**, 1217, 953-967.
5. a) Harata, K.; *Chem. Rev.*, **1998**, 98, 1803-1827; b) Harada, A.; Hashidzume, A.; Takashima, Y.; *Supramolecular Polymers Polymeric Betains Oligomers* (Eds Springer), **2006**, 201, 1-43; c) Schneider, H.J.; Hacket, F.; Rüdiger V.; *Chem. Rev.*, **1998**, 98, 1755-1795; d) Chierotti, M. R.; Gobetto, R.; *Chem. Rev.*, **2008**, 1621-1634; e) Lipkowitz K. B.; *Chem. Rev.*, **1998**, 98, 1829-1873; f) Jaime, C.; de Federico, M.; *Curr. Org. Chem.*, **2006**, 10, 731-743; g) Carmona, T.; Gonzalez-Alvarez, M. J.; Mendicuti, F.; Tagliapietra, S.; Martina, K.; Cravotto, G.; *J. Phys. Chem C*, **2010**, 114, 22431-22440.
6. Bellia, F.; La Mendola, D.; Pedone, C.; Rizzarelli, E.; Saviano, M.; Vecchio G.; *Chem. Soc. Rev.*, **2009**, 38, 2756-2781.
7. a) Cravotto, G.; Mendicuti, F.; Martina, K.; Tagliapietra, S.; Robaldo, B.; Barge A.; *Synlett*, **2008**, 17, 2642-2646; b) Aime, S.; Gianolio, E.; Arena, F.; Barge, A.; Martina, K.; Heropoulos, G.; Cravotto G.; *Org. Biomol. Chem.*, **2009**, 7, 370-379; c) Yamauchi, K.; Takashima, Y.; Hashidzume, A.; Yamaguchi, H.; Harada A.; *J. Am. Chem. Soc.*; **2008**; 130, 5024; d) Aime, S.; Gianolio, E.; Robaldo, B.; Barge, A.; Cravotto, G.; *Org. Biomol. Chem.*, **2006**, 4, 1124; e) Lecourt, T.; Mallet, J.M.; Sinay, P.; *Eur. J. Org. Chem.*, **2003**, 4553; f) Yuan, D.Q.; Liu, J.; Atsumi, M.; Isuka, A.; Kai, M.; Fujita, K.; *Chem. Commun.*, **2002**, 730.
8. Khan, A.R.; Forgo, P.; Stine, K. J.; D'Souza, V.T.; *Chem.Rev.*, **1998**, 98, 1977-1996.
9. Rao, C.T.; Lindberg, B.; Lindberg, J.; Pitha, J.; *J. Org Chem.*, **1991**, 56, 1327-1329.
10. Palmisano, G.; Tagliapietra, S.; Barge, A.; Binello, A.; Boffa, L., Cravotto, G.; *Synlett*, **2007**, 13, 2041-2044.
11. Schlatter, A.; Kundu, M.K.; Woggon, W.D.; *Angew. Chem. Int. Ed.*, **2004**, 43, 6731-6734.

12. a) Lämmerhofer, M.; *J. Chromatogr. A*, **2005**, *1068*, 3-30; b) Lämmerhofer, M.; *J. Chromatogr. A*, **2005**, *1068*, 31-57; c) Xiao, Y.; Ong, T.T.; Tan, T. T. Y.; Ng, S.C.; *J. Chromatogr. A*, **2009**, *1216*, 994-999.
13. Wenz, G.; Höfler, T.; *Carbohydrate Res.*, **1999**, *322*, 153-65.
14. a) Topchieva, I.N.; Mischnick, P.; Kühn, G.; Polyakov, V.A.; Elezkaya, S.V.; Bystryzky, G.I.; Karezin, K.I. *Bioconjugate Chem.*, **1998**, *9*, 676-682; b) Badi, N., Auvray, L., Guégan, P. *Adv. Mater.*, **2009**, *21*, 4054-4057.
15. a) Martina, K.; Trotta, F.; Robaldo, B.; Belliardi, N.; Jicsinszky, L.; Cravotto, G. *Tetrahedron Lett.* **2007**, *48*, 9185-9189 b) Cintas, P.; Barge, A.; Tagliapietra, S.; Boffa, L.; Cravotto, G.; *Nature Prot.* **2010**, *5*, 607-616.
16. Shivani Pujala, B.; Chakraborti, A K.; *J. Org. Chem.*, **2007**, *72*, 3713-3722.
17. Parker, R.E.; Isaac, N.S.; *Chem Rev.*, **1959**, 737-799.
18. Trotta, F.; Martina, K.; Robaldo, B.; Barge, A.; Cravotto, G.; *J. Incl. Phen. and Macrocyclic Chem.*, **2007**, *57*, 3-7.
19. Jicsinszky, L.; Ivanyi, R.; *Carbohydr. Polym.*, **2001**, *45*, 139-145.
20. Meijer, E. W.; Palmans, A. R. A.; van As, B. A. C.; van Buijtenen, J.; Mes, T.; *Chem. Eur. J.*, **2007**, *13*, 8325-8332.