

AperTO - Archivio Istituzionale Open Access dell'Università di Torino

## Meta-Taxonomic Analysis of Poultry and Slaughterhouse Microbiota: A Comprehensive Examination of Resident Microbial Communities

**This is the author's manuscript**

*Original Citation:*

*Availability:*

This version is available <http://hdl.handle.net/2318/2009670> since 2024-09-02T12:42:52Z

*Terms of use:*

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)



# Meta-Taxonomic Analysis of Poultry and Slaughterhouse Microbiota: A Comprehensive Examination of Resident Microbial Communities

E. Chiarini<sup>1</sup>, C. Botta<sup>1</sup>, D. Buzzanca<sup>1</sup>, F. Chiesa<sup>2</sup>, S. Rubiola<sup>2</sup>, I. Ferrocino<sup>1</sup>, E. Fontanella<sup>3</sup>, K. Rantsiou<sup>1</sup>, K. Houf<sup>4</sup>, V. Alessandria<sup>1</sup>




<sup>1</sup> Department of Agricultural, Forest and Food Sciences, University of Torino, Italy  
<sup>2</sup> Department of Veterinary Sciences, University of Turin, Italy  
<sup>3</sup> Local health company, ASL CN1, via Carlo Boggio, 12, Cuneo  
<sup>4</sup> Department of Veterinary and Biosciences, Faculty of Veterinary Medicine, Ghent University, Belgium



## INTRODUCTION & AIMS


Poultry contaminations occur during the processing chain and are found from farm to table. The final microbial profile of poultry meat is influenced by each step of the process. The consumption of contaminated poultry poses a significant threat to public health. It is associated to infections caused by pathogens such as: *Campylobacter*, *Salmonella*, *Listeria monocytogenes* and *Arcobacter*.




The objective is to assess the contribution of broiler skin and gut to the indigenous slaughterhouse microbiota. In addition, the study aims to clarify how the persistence and environmental distribution of *Arcobacter* spp. are influenced by the cross-contamination patterns associated with these specific sources.

## MATERIALS & METHODS


**1. BROILERS AND ENVIRONMENTAL SAMPLING**  
154 samples were collected from a local poultry abattoir



**2. DNA extraction**  
**3. V3-V4 region library** constructed from the 16S rRNA gene region




**4. BIOINFORMATIC ANALYSIS:** taxonomy was assigned with a 99% of sequence similarity by matching ASVs to the Silva prokaryotic reference database




## RESULTS & DISCUSSIONS

The BC microbial community is significantly distinct from the BNS and SE community (Fig. 1, 2).



**Figure 2.** Microbiota composition and distribution in phylum and family taxa. The samples are grouped according to the temporal order of the sampling and according to the order of the slaughter of the flock and the processing stage, BC-BNS and SE. Slaughterhouse sectors and transport lines are indicated: FL = first row; PS = plucking sector; SS = slaughter sector; SL = second row. Taxa are ordered in the legend from most to least abundant (mean >1%).




**Figure 1.** Graphic representation of the network revealing the taxa associated to the three sampling sources. Taxa (colored nodes) are unidirectionally connected with arrows to the sampling sources if significant associations were detected (Indicator Species Analysis: *multipatt* statistics;  $R > 0.4$  and  $P$  value < 0.001).

*Arcobacter butzleri* emerged as one of the most abundant species and was detected throughout the slaughterhouse, showing a higher prevalence compared to other Campylobacterota (Fig. 1, 2). It was uniquely and significantly associated with BNS and SE, while *Helicobacter pullorum* and *Campylobacter jejuni* were indicators of BC (Fig. 1). *A. butzleri* was constantly present on BNS along production runs, but significantly more abundant the first sampling day in SE (Fig.2).

## CONCLUSIONS & IMPORTANCE

Our findings have emphasised the persistence of *A. butzleri* in a modern poultry abattoir and its establishment as part of the resident microbiota in specific environmental niches. The analysis conducted underlines the significance of early monitoring of food pathogens in the production chain, supported by meta-taxonomic analysis. Using these detection approaches, the presence of these pathogens could soon be considered an indicator of food safety and quality in slaughtered poultry.

If you are interested in reading more...




Applied and Environmental Microbiology

Environmental Microbiology | Full-Length Text

Microbial contamination pathways in a poultry abattoir provided clues on the distribution and persistence of *Arcobacter* spp.

Cristian Botta,<sup>1</sup> Davide Buzzanca,<sup>1</sup> Elisabetta Chiarini,<sup>1</sup> Francesco Chiesa,<sup>2</sup> Selene Rubiola,<sup>2</sup> Ilario Ferrocino,<sup>1</sup> Edoardo Fontanella,<sup>3</sup> Kalliope Rantsiou,<sup>1</sup> Kurt Houf,<sup>4</sup> Valentina Alessandria<sup>1</sup>



References:  
Wang H, Qin X, Li X, Wang X, Gao H, Zhang C. 2020. Changes in the microbial communities of air- and water-chilled yellow-feathered broilers during storage at 2°C. Food Microbiology 87:103390.  
Buzzanca D, Kerkhof P-J, Alessandria V, Rantsiou K, Houf K. 2023. *Arcobacteraceae* comparative genome analysis demonstrates genome heterogeneity and reduction in species isolated from animals and associated with human illness. Heliyon 9:e17652.  
Ramees TP, Dhama K, Karthik K, Rathore RS, Kumar A, Saminathan M, Tiwari R, Malik YS, Singh RK. 2017. *Arcobacter*: an emerging food-borne zoonotic pathogen, its public health concerns and advances in diagnosis and control – a comprehensive review. Vet Q 37:136–161.

RESEARCH  
FOUNDED BY



E-mail contact:  
elisabetta.chiarini@unito.it

