

AperTO - Archivio Istituzionale Open Access dell'Università di Torino

**Effects of light quality on carbon assimilation/emission and growth of lettuce: use of an upgraded prototype-platform to unravel mechanisms of photosynthesis/respiration control.**

**This is the author's manuscript**

*Original Citation:*

*Availability:*

This version is available <http://hdl.handle.net/2318/1952034> since 2024-01-16T09:38:23Z

*Publisher:*

SIBV

*Terms of use:*

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)

## Effects of light quality on carbon assimilation/emission and growth of lettuce: use of an upgraded prototype-platform to unravel mechanisms of photosynthesis/respiration control.

DAVIDE L. PATONO (1), FLAVIO MARTINI (2), STEFANO CAZZANIGA (2), MATTEO BALLOTTARI (2), CLAUDIO LOVISOLO (1)

(1) Department of Agriculture, Forest and Food Sciences, University of Torino, Largo Braccini 2, 10095 Grugliasco, Italy

(2) Department of Biotechnology, University of Verona, Strada Le Grazie 15, 37134 Verona, Italy

Responses of plant performances to light quality are focus of research to optimize plant indoor cultivation with artificial light. In a comparative study of lettuce grown with different light spectra, photosynthesis, root and shoot respiration, and growth traits through chlorophyll fluorimetry essays, biochemical and molecular characterization of photosystems, and plant-to-atmosphere gas exchange assessment have been measured. To answer physiological questions at the whole-plant level we tested two phenotyping platforms, one based on 3D and multispectral imaging technology and a prototype based on whole plant gas exchange analyses, capable of differentiating root from canopy gas exchange. Red and blue (RB) light maximized photosynthetic activity, but this advantage did not lead to greater biomass accumulation, which was greater in plants under red, green and blue (RGB) and further under full spectrum (FS) light. Plants subjected to the RB light regime invest less carbon in leaf blade extension and incorporate less carbon than what has been assimilated. RB leaves have a smaller PSII antenna size and the cyclic electron transport around PSI is higher, implying a lower NADPH / ATP ratio. Exposure to RB light affects plant overall metabolic biosynthetic pathways at the expense of growth, while RGB and especially FS light, by not triggering this acclimation, are more suited to constant low light growing conditions, as those usually provided by indoor growing technologies.

We acknowledge financial support from the Italy's National Recovery and Resilience Plan (PNRR), grant agreement No PE0000018 - GRINS – Growing Resilient, INclusive and Sustainable and project VITIMOUNT, RF 2022.1659, CRT Foundation.