

Torus Orbifolds, Slice-Maximal Torus Actions, and Rational Ellipticity

Fernando Galaz-García^{1*}, Martin Kerin², Marco Radeschi², and Michael Wiemeler³

¹Institut für Algebra und Geometrie, Karlsruher Institut für Technologie (KIT), Germany, ²Mathematisches Institut, WWU Münster, Germany, and ³Institut für Mathematik, Universität Augsburg, Germany

*Correspondence to be sent to: e-mail: galazgarcia@kit.edu

In this work, it is shown that a simply connected, rationally elliptic torus orbifold is equivariantly rationally homotopy equivalent to the quotient of a product of spheres by an almost-free, linear torus action, where this torus has rank equal to the number of odd-dimensional spherical factors in the product. As an application, simply connected, rationally elliptic manifolds admitting slice-maximal torus actions are classified up to equivariant rational homotopy. The case where the rational-ellipticity hypothesis is replaced by non-negative curvature is also discussed, and the Bott Conjecture in the presence of a slice-maximal torus action is proved.

1 Introduction

A *torus manifold* is a $2n$ -dimensional, closed, orientable, smooth manifold equipped with a smooth, effective n -torus action which has non-empty fixed-point set. Such spaces have been of long-standing interest, going back, on the one hand, to Orlik and Raymond's work on closed, smooth 4-manifolds equipped with smooth, effective T^2 actions [30, 31] and, on the other hand, to the study of toric varieties in algebraic geometry [9]. Many results on manifolds with torus actions admit generalizations to orbifolds (see, e.g.,

Received October 5, 2016; Revised January 16, 2017; Accepted February 17, 2017
Communicated by Prof. John Lott

[16] for smooth torus actions on orbifolds, [20, 25] for Hamiltonian torus actions on symplectic orbifolds or [33] for quasitoric orbifolds).

Recently, it has been shown in [37] that, if M is simply connected and either a rationally elliptic torus manifold with torsion-free integer cohomology or a torus manifold with non-negative sectional curvature, then M is homeomorphic to the quotient of a product of spheres by a free, linear torus action. In this article, *torus orbifolds* are investigated, and a similar result to that in [37] is proven in this more general context.

Recall that a simply connected topological space X is called *rationally elliptic* if it satisfies $\dim_{\mathbb{Q}} H^*(X; \mathbb{Q}) < \infty$ and $\dim_{\mathbb{Q}}(\pi_*(X) \otimes \mathbb{Q}) < \infty$. Two spaces X and Y are *rationally homotopy equivalent* if their corresponding minimal models are isomorphic. Given a torus T , a rational homotopy equivalence between T -spaces X and Y is *T -equivariant* if the corresponding Borel constructions X_T and Y_T are also rationally homotopy equivalent and there exists a commutative diagram

$$\begin{array}{ccc} H^*(Y; \mathbb{Q}) & \longrightarrow & H^*(X; \mathbb{Q}) \\ \uparrow & & \uparrow \\ H_T^*(Y; \mathbb{Q}) & \longrightarrow & H_T^*(X; \mathbb{Q}) \end{array}$$

where the horizontal arrows are isomorphisms induced by the respective rational homotopy equivalences.

Theorem A. Let (\mathcal{O}, T) be a rationally elliptic, simply connected torus orbifold. Then there is a product \hat{P} of spheres of dimension ≥ 3 , a torus \hat{L} acting linearly and almost freely on \hat{P} , and an effective, linear action of T on $\hat{\mathcal{O}} = \hat{P}/\hat{L}$, such that there is a T -equivariant rational homotopy equivalence $\mathcal{O} \simeq_{\mathbb{Q}} \hat{\mathcal{O}}$.

Moreover, if \mathcal{O} is a manifold, then \hat{L} acts freely on \hat{P} and thus $\hat{\mathcal{O}}$ is a manifold as well. □

The final statement in Theorem A regarding manifolds is closely related to Theorem 1.1 of [37], where a stronger assumption (torsion-free integral cohomology) is required in order to obtain a correspondingly strong conclusion (classification up to homeomorphism).

Torus orbifolds have been studied in [17–19] and arise naturally in the study of smooth torus actions on manifolds, for example, when the action is slice maximal.

Definition. Let M be a closed, orientable, smooth n -manifold on which a torus T^k acts smoothly and effectively, and let m be the minimal dimension of an orbit. The action is *slice maximal* if $2k = n + m$. \square

It is clear from the definition that torus manifolds are an extremal case of slice-maximal actions. For a generic k -torus action on an n -dimensional manifold, it follows from the slice representation at a minimal orbit that $2k \leq n + m$. Thus, if equality holds, the slice representation at a minimal orbit is even dimensional and has maximal symmetry rank, justifying the terminology “slice maximal”. Slice-maximal actions were considered in [21, 36], where they were called *maximal*.

Given an n -manifold M with a slice-maximal T^k action, there exists a subtorus $T^m \subseteq T^k$ acting almost freely on M and the quotient $\mathcal{O} = M/T^m$ is a $2(k - m)$ -dimensional torus orbifold. Moreover, if M is rationally elliptic, so too is the quotient \mathcal{O} .

By applying Theorem A, it turns out that the existence of a slice-maximal torus action has strong implications on the topology of a manifold.

Theorem B. Let M be an n -dimensional, smooth, closed, simply connected, rationally elliptic manifold with a slice-maximal T^k action. Then there is a product \hat{P} of spheres of dimension ≥ 3 , a torus \hat{K} acting linearly and freely on \hat{P} , and an effective, linear action of T^k on $\hat{M} = \hat{P}/\hat{K}$, such that there is a T^k -equivariant rational homotopy equivalence $M \simeq_{\mathbb{Q}} \hat{M}$. \square

It is worth pointing out that, in general, rational homotopy does not behave well with respect to group actions; for example, one cannot “pull back” an action via a rational homotopy equivalence. The difficulties are even more apparent in the case of actions which are not almost free. In particular, while it is not too hard in Theorem B to find some space $\hat{M} = \hat{P}/\hat{K}$ that is rationally homotopy equivalent to M , it is more difficult to prove that such space is a manifold (i.e., that the \hat{K} action is free rather than almost free), and even harder to show that the rational homotopy equivalence is T -equivariant in the sense described above. To prove the latter, some novel approaches are required. In this case, it is shown that the rational homotopy equivalence $\hat{M} \simeq_{\mathbb{Q}} M$ induces a rational homotopy equivalence between the *equivariant 1-skeleta* $\hat{M}^{(1)} \simeq_{\mathbb{Q}} M^{(1)}$ and, moreover, that this rational homotopy equivalence is, in fact, induced by a T^k -equivariant homeomorphism $\hat{M}^{(1)} \rightarrow M^{(1)}$.

As a first application, Theorem B has been used in [10] to obtain a classification of closed, simply connected, rationally elliptic manifolds admitting effective torus actions of maximal rank up to equivariant rational homotopy equivalence.

For another interesting consequence of Theorem B, recall that the largest integer r for which a closed, simply connected space M admits an almost-free T^r -action is called the *toral rank* of M , and is denoted $\text{rk}(M)$. The Toral Rank Conjecture, formulated by S. Halperin, asserts that $\dim H^*(M; \mathbb{Q}) \geq 2^{\text{rk}(M)}$.

Corollary C. Let M be a smooth, closed, simply connected, rationally elliptic, n -dimensional manifold with a slice-maximal torus action. Then M satisfies the Toral Rank Conjecture. \square

Proof. Let T^r act almost freely on M . Given $H^2(M; \mathbb{Q}) = \mathbb{Q}^{b_2(M)}$, there is a principal $T^{b_2(M)}$ -bundle over M with (rationally) 2-connected total space P . As any action by a torus T on M lifts to a $T \times T^{b_2(M)}$ action on P , the slice-maximal action (respectively, the almost-free T^r action) on M lifts to a slice-maximal action (respectively, an almost-free $T^r \times T^{b_2(M)}$ action) on P . By Theorem B and since $H^2(P; \mathbb{Q}) = 0$, P must have the rational cohomology of a product of spheres of dimension ≥ 3 . By [8, Prop. 7.23], P satisfies the Toral Rank Conjecture, that is, $H^*(P; \mathbb{Q}) \geq 2^{r+b_2(M)}$. The result now follows from $\dim H^*(P; \mathbb{Q}) \leq \dim H^*(T^{b_2(M)}; \mathbb{Q}) \cdot \dim H^*(M; \mathbb{Q})$. \blacksquare

Finally, recall that the Bott Conjecture asserts that a closed, simply connected, non-negatively curved Riemannian manifold is rationally elliptic. In [34], W. Spindeler verified the conjecture for simply connected, non-negatively curved torus manifolds. In fact, the conjecture also holds in the slice-maximal setting.

Theorem D. Let M be a closed, simply connected, non-negatively curved Riemannian manifold admitting an isometric, slice-maximal torus action. Then M is rationally elliptic. \square

It is worth noting that non-negatively curved torus manifolds have already been classified up to equivariant diffeomorphism in [37], and a similar classification from a different viewpoint can be found in [6]. If the non-negative-curvature hypothesis in Theorem D were to be replaced by positive curvature, then it would follow from the work of K. Grove and C. Searle [13] that M is equivariantly diffeomorphic to a sphere or complex projective space equipped with a linear action.

The article is organized as follows: In Section 2, some basic definitions and facts about orbifolds are collected, following the presentation in [23], as well as some results on smooth actions on orbifolds. These results have been included to provide a basic

reference for compact Lie group actions on orbifolds, since they seem to be scattered in the literature (see, e.g., [15, 16, 25, 38]). In Section 3, torus orbifolds are introduced and their fundamental properties established. In Section 4, there is a brief review of applied to torus orbifolds, following the work of Goresky, Kottwitz and MacPherson in [11]. The proof of Theorem A is contained in Section 5. In Section 6, an example of a family of rationally elliptic torus orbifolds which are not rationally homotopy equivalent to any rationally elliptic manifold is provided, illustrating that almost-free (rather than free) torus actions on products of spheres are necessary in the conclusion of the Theorem A. Section 7 is devoted to establishing Theorem B. In Section 8, a version of Theorem A for non-negatively curved orbifolds of dimension ≤ 6 is proven. The case of general dimensions remains open. Section 8 concludes with the proof of Theorem D, which is independent of the rest of the article.

The reader is referred to [7] for the basic definitions and results of rational homotopy theory. A brief summary can also be found in [10].

2 Review of Orbifolds

As there are some conflicts in the literature regarding basic notions in the study of orbifolds, it is important to clearly define the notation and terminology which will be used throughout the article. Since the proofs of most of the lemmas in this section use standard arguments, these will only be sketched.

Definition 2.1. A *local model of dimension n* is a pair (\tilde{U}, Γ) , where \tilde{U} is an open, connected subset of a Euclidean space \mathbb{R}^n , and Γ is a finite group acting smoothly and effectively on \tilde{U} .

A *smooth map* $(\tilde{U}_1, \Gamma_1) \rightarrow (\tilde{U}_2, \Gamma_2)$ between local models (\tilde{U}_i, Γ_i) , $i = 1, 2$, is a homomorphism $\varphi_{\#} : \Gamma_1 \rightarrow \Gamma_2$ together with a $\varphi_{\#}$ -equivariant smooth map $\tilde{\varphi} : \tilde{U}_1 \rightarrow \tilde{U}_2$, that is, $\tilde{\varphi}(\gamma \cdot \tilde{u}) = \varphi_{\#}(\gamma) \cdot \tilde{\varphi}(\tilde{u})$, for all $\gamma \in \Gamma_1$, $\tilde{u} \in \tilde{U}_1$. \square

Given a local model (\tilde{U}, Γ) , denote by U the quotient \tilde{U}/Γ . Clearly, a smooth map $\tilde{\varphi} : (\tilde{U}_1, \Gamma_1) \rightarrow (\tilde{U}_2, \Gamma_2)$ induces a map $\varphi : U_1 \rightarrow U_2$. The map φ is called an *embedding* if $\tilde{\varphi}$ is an embedding. In this case, the effectiveness of the actions in the local models implies that $\varphi_{\#}$ is injective.

Definition 2.2. An *n -dimensional local chart* $(U_p, \tilde{U}_p, \Gamma_p, \pi_p)$ around a point p in a topological space X consists of:

- (a) A neighbourhood U_p of p in X ;
- (b) A local model (\tilde{U}_p, Γ_p) of dimension n ;

- (c) A Γ_p -equivariant projection $\pi_p : \tilde{U}_p \rightarrow U_p$, where Γ_p acts trivially on U_p , that induces a homeomorphism $\tilde{U}_p/\Gamma_p \rightarrow U_p$.

If $\pi_p^{-1}(p)$ consists of a single point, \tilde{p} , then $(U_p, \tilde{U}_p, \Gamma_p, \pi_p)$ is called a *good local chart* around p . In particular, \tilde{p} is fixed by the action of Γ_p on \tilde{U}_p . □

Note that, given a good local chart $(U_p, \tilde{U}_p, \Gamma_p, \pi_p)$ around a point p in a topological space X , the 4-tuple $(U_p, \tilde{U}_p, \Gamma_p, \pi_p)$ is also a local chart, not necessarily good, around any other point $q \in U_p$. By abusing notation, a local chart $(U, \tilde{U}, \Gamma, \pi)$ will from now on be denoted simply by U .

Definition 2.3. An n -dimensional (smooth) orbifold, denoted by \mathcal{O}^n or simply \mathcal{O} , is a second-countable, Hausdorff topological space $|\mathcal{O}|$, called the *underlying topological space* of \mathcal{O} , together with a maximal collection of n -dimensional local charts $\mathcal{A} = \{U_\alpha\}_\alpha$ such that:

- (a) The neighbourhoods $U_\alpha \in \mathcal{A}$ give an open covering of $|\mathcal{O}|$, and
- (b) For any $p \in U_\alpha \cap U_\beta$, there is a local chart $U_\gamma \in \mathcal{A}$ with $p \in U_\gamma \subseteq U_\alpha \cap U_\beta$ and embeddings $(\tilde{U}_\gamma, \Gamma_\gamma) \rightarrow (\tilde{U}_\alpha, \Gamma_\alpha), (\tilde{U}_\gamma, \Gamma_\gamma) \rightarrow (\tilde{U}_\beta, \Gamma_\beta)$.

An orbifold is *orientable* if every local model \tilde{U}_α is orientable, and if every Γ_α action and every embedding $\tilde{U}_\gamma \rightarrow \tilde{U}_\alpha$ is orientation preserving. Given an orientable orbifold \mathcal{O} , it is not hard to see that the set of points $p \in \mathcal{O}$ for which Γ_p is non-trivial has codimension at least 2 in \mathcal{O} . An orbifold \mathcal{O} is *connected* (respectively, *closed*) if its underlying topological space $|\mathcal{O}|$ is connected (respectively, compact and without boundary). □

Given an orbifold \mathcal{O} and any point $p \in \mathcal{O}$, one can always find a good local chart U_p around p . Moreover, the corresponding group Γ_p does not depend on the choice of good local chart around p , and is referred to as the *local group at p* . From now on, only good local charts will be considered.

Lemma 2.4. Let \mathcal{O} be an orbifold and U_p a good local chart around $p \in \mathcal{O}$. Let $q \in U_p$, $\tilde{q} \in \pi_p^{-1}(q) \subseteq U_p$ and $(\Gamma_p)_{\tilde{q}} = \{\gamma \in \Gamma_p \mid \gamma \cdot \tilde{q} = \tilde{q}\}$. Then there exists a $(\Gamma_p)_{\tilde{q}}$ -invariant neighbourhood $\tilde{U}_q \subseteq \tilde{U}_p$ of \tilde{q} such that $(\pi_p(\tilde{U}_q), \tilde{U}_q, (\Gamma_p)_{\tilde{q}}, \pi_p|_{\tilde{U}_q})$ is a good local chart around q . □

Proof. Define \tilde{U}_q to be a $(\Gamma_p)_{\tilde{q}}$ -invariant neighbourhood of \tilde{q} such that, for every $\gamma \in \Gamma_p \setminus (\Gamma_p)_{\tilde{q}}$, one has $\tilde{U}_q \cap \tilde{\gamma} \cdot \tilde{U}_q = \emptyset$. ■

In particular, given a good local chart U_p around $p \in \mathcal{O}$, a point $q \in U_p$ and $\tilde{q} \in \pi_p^{-1}(q)$, one can identify the local group Γ_q at q with $(\Gamma_p)_{\tilde{q}}$.

Definition 2.5. A smooth map $\varphi : \mathcal{O}_1 \rightarrow \mathcal{O}_2$ between orbifolds is given by a continuous map $|\varphi| : |\mathcal{O}_1| \rightarrow |\mathcal{O}_2|$ such that, if U_p and $U_{\varphi(p)}$ are (good) local charts around $p \in \mathcal{O}_1$ and $\varphi(p) \in \mathcal{O}_2$, respectively, such that $\varphi(U_p) \subseteq U_{\varphi(p)}$, then there is a (possibly non-unique) smooth lift at $p \in \mathcal{O}_1$, $\tilde{\varphi}_p : (\tilde{U}_p, \Gamma_p) \rightarrow (\tilde{U}_{\varphi(p)}, \Gamma_{\varphi(p)})$, so that $\varphi \circ \pi_p = \pi_{\varphi(p)} \circ \tilde{\varphi}_p$ and there is an induced homomorphism $(\tilde{\varphi}_p)_\# : \Gamma_p \rightarrow \Gamma_{\varphi(p)}$.

A diffeomorphism $\varphi : \mathcal{O}_1 \rightarrow \mathcal{O}_2$ between orbifolds is a smooth map with a smooth inverse. \square

Definition 2.6.

- (a) An orbifold \mathcal{O}_1 is a *suborbifold* of an orbifold \mathcal{O}_2 , if there is a smooth map $\varphi : \mathcal{O}_1 \rightarrow \mathcal{O}_2$ such that $|\varphi|$ maps $|\mathcal{O}_1|$ homeomorphically onto its image in $|\mathcal{O}_2|$ and, for every $p \in \mathcal{O}_1$, some (and, hence, every) smooth lift $\tilde{\varphi}_p : \tilde{U}_p \rightarrow \tilde{U}_{\varphi(p)}$ is an immersion. In this case, \mathcal{O}_1 will be identified with its image.
- (b) A suborbifold $\mathcal{O}_1 \subseteq \mathcal{O}_2$ is a *strong suborbifold* if, for every $p \in \mathcal{O}_1$ and every good local chart U_p , the image of a smooth lift $\tilde{\varphi}_p$ is independent of the choice of lift. \square

The above definition of strong suborbifold is equivalent to Thurston's definition of suborbifold (cf. [35]). Given a strong suborbifold $\mathcal{O}_1 \subseteq \mathcal{O}_2$, let U_p be a good local chart (in \mathcal{O}_2) around $p \in \mathcal{O}_1$ and let $\tilde{T}_p U_p$ be the tangent space to \tilde{U}_p at $\tilde{p} = \pi_p^{-1}(p)$. Denote by $\tilde{T}_p \mathcal{O}_1 \subseteq \tilde{T}_p U_p$ the tangent space to $\pi_p^{-1}(\mathcal{O}_1 \cap U_p)$ at \tilde{p} . Then the space $\tilde{T}_p U_p$ splits as $\tilde{T}_p \mathcal{O}_1 \oplus \tilde{\nu}_p \mathcal{O}_1$, where $\tilde{\nu}_p \mathcal{O}_1$ denotes the normal space to $\tilde{T}_p \mathcal{O}_1 \subseteq \tilde{T}_p U_p$.

If \mathcal{O} is an orbifold, then a *smooth action* of a Lie group G on \mathcal{O} is an action of G on \mathcal{O} such that the map $G \times \mathcal{O} \rightarrow \mathcal{O}$, $(g, p) \mapsto g \cdot p$, is smooth. The set $G(p) = \{g \cdot p \mid g \in G\}$ is the *orbit* of G through $p \in \mathcal{O}$. The *ineffective kernel* of the action is the normal subgroup $\text{Ker} = \{g \in G \mid \varphi(g, \cdot) = \text{id}_{\mathcal{O}}\}$. If the ineffective kernel is trivial, the action is *effective*. The group G/Ker will always act effectively. The *isotropy subgroup* G_p at $p \in \mathcal{O}$ is the subgroup consisting of those elements in G that fix p . Note that, whenever G is compact, one can always find a G_p -invariant good local chart around p . If G_p is trivial (respectively, finite) for every $p \in \mathcal{O}$, the action is *free* (respectively, *almost free*). The orbit space of the action will be denoted by \mathcal{O}/G and the fixed-point set $\{p \in \mathcal{O} \mid G_p = G\}$ by \mathcal{O}^G . The identity component of G is denoted by G° .

Lemma 2.7. Every G -orbit in \mathcal{O} is a manifold, as well as a strong suborbifold of \mathcal{O} . \square

Proof. The fact that $G(p)$, $p \in \mathcal{O}$, is a manifold and a suborbifold follows as in the manifold case. To see that it is a strong suborbifold, one can apply the fact that, since G acts by diffeomorphisms, the local groups at all points in the orbit are isomorphic. \blacksquare

Proposition 2.8. Let \mathcal{O} be an orbifold with a smooth, effective action by a compact Lie group G . Let $p \in \mathcal{O}$ have isotropy subgroup $G_p \subseteq G$ and let U_p be a G_p -invariant good local chart. Then there exists a Lie group \tilde{G}_p such that:

- (a) \tilde{G}_p acts on \tilde{U}_p and $\tilde{U}_p/\tilde{G}_p = U_p/G_p$;
- (b) \tilde{G}_p is an extension of G_p by Γ_p , that is, there exists a short exact sequence

$$\{e\} \rightarrow \Gamma_p \rightarrow \tilde{G}_p \xrightarrow{\rho} G_p \rightarrow \{e\}. \quad \square$$

Proof. Let $g \in G_p$. The action of G_p on U_p gives a smooth map

$$L_g : U_p \rightarrow U_p$$

$$q \mapsto g \cdot q.$$

Then, by definition, there exists a smooth lift $\tilde{L}_g : \tilde{U}_p \rightarrow \tilde{U}_p$ of L_g . Let $\tilde{G}_p = \{F_g : \tilde{U}_p \rightarrow \tilde{U}_p \mid \pi_p \circ F_g = L_g \circ \pi_p, g \in G_p\}$ be the collection of all possible lifts. This is a group and, given that the G_p action is smooth, it is not difficult to see that \tilde{G}_p is a Lie group acting smoothly and effectively on \tilde{U}_p . Note that, since $\Gamma_p = \{F_e : \tilde{U}_p \rightarrow \tilde{U}_p \mid \pi_p \circ F_e = e \circ \pi_p = \pi_p\}$, Γ_p is a normal subgroup of \tilde{G}_p . Moreover, Γ_p acts on \tilde{G}_p via $F_g \mapsto \gamma \cdot F_g$ and $\tilde{G}_p/\Gamma_p = G_p$, that is, the quotient by Γ_p fixes a choice of lift corresponding to L_g . It then follows that $\tilde{U}_p/\tilde{G}_p = U_p/G_p$. \blacksquare

Corollary 2.9. Let \mathcal{O} be an orbifold with a smooth, effective action by a compact Lie group G . Let $p \in \mathcal{O}$ have isotropy subgroup G_p and let U_p be a G_p -invariant good local chart. Then the local group Γ_p commutes with every connected subgroup of the lift \tilde{G}_p . \square

Proof. Let \tilde{H} be a connected subgroup of \tilde{G}_p , $\tilde{g} \in \tilde{H}$, and $\gamma \in \Gamma_p$. From the short exact sequence in Proposition 2.8, the element $\tilde{g}\gamma\tilde{g}^{-1}$ belongs to Γ_p . Since \tilde{H} is connected and Γ_p is discrete, the map $\tilde{g} \mapsto \tilde{g}\gamma\tilde{g}^{-1}$ must be constant and hence $\tilde{g}\gamma\tilde{g}^{-1} = \gamma$ for all $\tilde{g} \in \tilde{H}$. \blacksquare

Corollary 2.10. Let G be a compact, connected Lie group acting smoothly and effectively on an orbifold \mathcal{O} such that the fixed-point set \mathcal{O}^G is non-empty. Then each connected component of \mathcal{O}^G is a strong suborbifold. \square

Proof. Let $p \in \mathcal{O}^G$ and let U_p be a G -invariant good local chart around p . The goal is to prove that $\pi_p^{-1}(\mathcal{O}^G \cap U_p)$ is a submanifold in \tilde{U}_p .

Since G is connected, the map $\tilde{G}^o \rightarrow G$ is a covering and therefore for every $g \in G$ there is a $\tilde{g} \in \tilde{G}^o$ projecting to g . Let $q \in \mathcal{O}^G \cap U_p$ and choose $\tilde{q} \in \pi^{-1}(q) \subseteq \tilde{U}_p$. As $g \cdot q = q$, for every $g \in G$, it follows that for every $\tilde{g} \in \tilde{G}^o$ there is a $\gamma_{\tilde{g}} \in \Gamma_p$ such that $\tilde{g} \cdot \tilde{q} = \gamma_{\tilde{g}} \cdot \tilde{q}$. But \tilde{G}^o is connected, hence $\gamma_{\tilde{g}} = e$ for every $\tilde{g} \in \tilde{G}^o$. Thus \tilde{q} is fixed by \tilde{G}^o and so $\pi_p^{-1}(\mathcal{O}^G \cap U_p) \subseteq \tilde{U}_p^{\tilde{G}^o}$. The other inclusion trivially holds, and therefore

$$\pi_p^{-1}(\mathcal{O}^G \cap U_p) = \tilde{U}_p^{\tilde{G}^o}.$$

By Corollary 2.9, Γ_p commutes with \tilde{G}^o and, in particular, Γ_p preserves the fixed-point set of \tilde{G}^o . Then $\mathcal{O}^G \cap U_p$ is a strong suborbifold and, since p was arbitrary, it follows that \mathcal{O}^G is a strong suborbifold. \blacksquare

Just as for manifolds, one has a notion of Riemannian metric for orbifolds. An *orbifold-Riemannian metric* is given at each point p in the orbifold by the metric on a good local chart U_p around p induced by a Γ_p -invariant Riemannian metric on \tilde{U}_p . An orbifold equipped with an orbifold-Riemannian metric will be referred to as a *Riemannian orbifold*. It is clear that Riemannian notions such as geodesics and completeness carry over to Riemannian orbifolds. Recall that any orbifold on which a compact Lie group G acts smoothly and effectively admits a G -invariant orbifold-Riemannian metric. Kleiner's Isotropy Lemma (cf. [22]) also holds, with the same proof, in the context of complete Riemannian orbifolds.

Lemma 2.11 (Isotropy Lemma). Let \mathcal{O} be a complete Riemannian orbifold and suppose that a compact Lie group G acts effectively and isometrically on \mathcal{O} . Let $c : [0, d] \rightarrow \mathcal{O}$ be a minimal geodesic between the orbits $G(c(0))$ and $G(c(d))$. Then, for any $t \in (0, d)$, $G_{c(t)} = G_c$ does not depend on t and is a subgroup of $G_{c(0)}$ and of $G_{c(d)}$. \square

It turns out that the good local charts given by Lemma 2.4 can be chosen to be compatible with the action of a Lie group.

Lemma 2.12. Let a compact Lie group G act effectively and isometrically on a complete Riemannian orbifold \mathcal{O} and fix $p \in \mathcal{O}$. Then there exists a G_p -invariant good local chart

U_p around p such that, for every $q \in U_p$ and every $\tilde{q} \in \pi_p^{-1}(q)$, there is a G_q -invariant good local chart $U_q \subseteq U_p$ around q and a commutative diagram

$$\begin{array}{ccccccc}
 \{e\} & \longrightarrow & \Gamma_q & \longrightarrow & \tilde{G}_q & \xrightarrow{\rho_q} & G_q & \longrightarrow & \{e\} \\
 & & \downarrow & & \downarrow & & \downarrow & & \\
 \{e\} & \longrightarrow & \Gamma_p & \longrightarrow & \tilde{G}_p & \xrightarrow{\rho_p} & G_p & \longrightarrow & \{e\}
 \end{array} \tag{2.1}$$

of short exact sequences, where the vertical maps identify Γ_q , \tilde{G}_q and G_q with the subgroups $(\Gamma_p)_{\tilde{q}}$, $(\tilde{G}_p)_{\tilde{q}}$ and $(G_p)_q$, respectively. Furthermore, there is a commutative diagram

$$\begin{array}{ccc}
 \tilde{U}_q & \longrightarrow & \tilde{U}_p \\
 \pi_q \downarrow & & \downarrow \pi_p \\
 U_q & \longrightarrow & U_p
 \end{array} \tag{2.2}$$

where each map is equivariant with respect to the appropriate actions of the groups \tilde{G}_q , \tilde{G}_p , G_q and G_p . □

Proof. The proof requires simply checking that everything proceeds as expected and is left to the reader. ■

Recall from Lemma 2.7 that orbits of Lie group actions are strong suborbifolds. Using the notation developed in Section 2, there is a version of the Slice Theorem for orbifolds (for a proof, see, e.g., [38]).

Theorem 2.13 (Slice Theorem). Suppose that a compact Lie group G acts on an orientable orbifold \mathcal{O} equipped with a G -invariant, orbifold-Riemannian metric, and let $G(p)$ be the orbit of G through $p \in \mathcal{O}$. Then a G -invariant neighbourhood of $G(p)$ is equivariantly diffeomorphic to

$$G \times_{G_p} (\tilde{v}_p G(p) / \Gamma_p)$$

and, by Proposition 2.8, this is equivariantly diffeomorphic to

$$G \times_{\tilde{G}_p} \tilde{v}_p G(p). \tag{2.3} \quad \square$$

3 Torus Orbifolds

Definition 3.1. A pair (\mathcal{O}^{2n}, T^n) , $n \geq 1$, is a *torus orbifold* if \mathcal{O}^{2n} is a $2n$ -dimensional, closed, oriented orbifold on which the n -dimensional torus T^n acts smoothly and effectively with non-empty fixed-point set. \square

To avoid confusion, henceforth the notation $G = T^n$ will be used. The identity component of a subgroup $K \subseteq G$ will be denoted by K^o . If the action is clear from the context, a torus orbifold (\mathcal{O}, G) will be denoted simply by \mathcal{O} . It will always be assumed that \mathcal{O} is equipped with an invariant orbifold-Riemannian metric (cf. [2]).

Definition 3.2. Let \mathcal{O} be a torus orbifold and let $p \in \mathcal{O}$. The *stratum* containing p , which will be denoted by Σ_p , is the connected component of the set

$$\{q \in \mathcal{O} \mid G_q^o = G_p^o \text{ as subgroups of } G\}$$

which contains p . The projection $\overline{\Sigma}_p/G \subseteq \mathcal{O}/G$ of the closure, $\overline{\Sigma}_p$, of a stratum Σ_p is called an (*orbifold*) *face* of \mathcal{O}/G . A one-dimensional face of \mathcal{O}/G is called an *edge*. \square

It follows from Definition 3.2 that the closure $\overline{\Sigma}_p$ of the stratum containing p is a connected component of the fixed-point set $\mathcal{O}^{G_p^o}$ and hence, by Corollary 2.10, a strong suborbifold of \mathcal{O} .

Note that the identity component G_p^o of an isotropy group G_p is a connected, compact, abelian Lie group, hence a torus. In particular, \tilde{G}_p^o acts effectively on $\tilde{U}_p \cong \mathbb{R}^{2n}$. This fact implies the following lemma.

Lemma 3.3. The fixed-point set of a torus orbifold (\mathcal{O}, G) consists of finitely many isolated points. Hence $H^{\text{odd}}(\mathcal{O}; \mathbb{Q}) = 0$ if \mathcal{O} is simply connected and rationally elliptic. \square

Proof. Let $p \in \mathcal{O}^G$ be a G -fixed point. As \tilde{G}^o is an n -dimensional torus which acts linearly and effectively on $\tilde{T}_p U_p$, it follows from dimension reasons that this action is (up to automorphisms of \tilde{G}^o) equivalent to the standard action of \tilde{G}^o on \mathbb{C}^n . Hence it follows from the Slice Theorem that p is an isolated fixed point.

It now follows from the Euler characteristic identity $\chi(\mathcal{O}^G) = \chi(\mathcal{O})$ (cf. [2, p. 163], [24]) that $\chi(\mathcal{O}) > 0$. Whenever \mathcal{O} is also simply connected and rationally elliptic, this is equivalent to $H^{\text{odd}}(\mathcal{O}; \mathbb{Q}) = 0$ [7, p. 444]. \blacksquare

Lemma 3.4. Given a torus orbifold (\mathcal{O}, G) , suppose that $p \in \mathcal{O}^G$ and U_p is a G -invariant good local chart around p . Then:

- (a) The action of G on U_p lifts to an action of \tilde{G} on \tilde{U}_p such that the isotropy action $\tilde{G}^\circ \times \tilde{T}_p U_p \rightarrow \tilde{T}_p U_p$ is equivalent to the standard n -torus action on \mathbb{C}^n .
- (b) Γ_p is a subgroup of \tilde{G}° . In particular, $\tilde{G} = \tilde{G}^\circ$, i.e. \tilde{G} is connected, hence a torus. □

Proof. Part (a) was already proven in the proof of Lemma 3.3.

There is an \tilde{G} -invariant scalar product on $\tilde{T}_p \mathcal{O}$. Hence, \tilde{G} is a closed subgroup of $SO(\tilde{T}_p \mathcal{O})$. By part (a), \tilde{G}° is a maximal torus of $SO(\tilde{T}_p \mathcal{O})$. Hence, the centralizer of \tilde{G}° in $SO(\tilde{T}_p \mathcal{O})$ is \tilde{G}° itself. Now part (b) follows from Corollary 2.9. ■

Corollary 3.5. Let \mathcal{O}^{2n} be a $2n$ -dimensional torus orbifold. Fix $p \in \mathcal{O}^G$ and let U_p be a G -invariant good local chart around p . Then $U_p/G = \tilde{U}_p/\tilde{G}$ is face-preserving diffeomorphic to $\mathbb{R}_+^n = \{(x_1, \dots, x_n) \in \mathbb{R}^n \mid x_i \geq 0, i = 1, \dots, n\}$. □

Lemma 3.6 (cf. [29, Lemma 2.2]). Let \mathcal{O} be a closed n -orbifold with a smooth, effective action by a k -torus G , $k \leq n$. Let $H \subseteq G$ be a subtorus and $\mathcal{N} \subseteq \mathcal{O}^H$ a connected component of its fixed-point set. If $H^{\text{odd}}(\mathcal{O}; \mathbb{Q}) = 0$, then $H^{\text{odd}}(\mathcal{N}; \mathbb{Q}) = 0$ and $\mathcal{N}^G \neq \emptyset$ (i.e. $\mathcal{N} \cap \mathcal{O}^G \neq \emptyset$). □

Proof. Since $H^{\text{odd}}(\mathcal{O}; \mathbb{Q}) = 0$, it follows from [1, Lemma 4.2.1] and [1, Lemma 3.10.13] that $H^{\text{odd}}(\mathcal{N}; \mathbb{Q}) = 0$. Because \mathcal{N} is a G -invariant strong suborbifold of \mathcal{O} , it follows that $\chi(\mathcal{N}^G) = \chi(\mathcal{N}) > 0$ and, hence, that there is a G -fixed point in \mathcal{N} . ■

Proposition 3.7. Let \mathcal{O}^{2n} be a torus orbifold with $H^{\text{odd}}(\mathcal{O}; \mathbb{Q}) = 0$. Fix $p \in \mathcal{O}^{2n}$ and let $\bar{\Sigma}_p$ be the closure of the stratum Σ_p in \mathcal{O} . Then:

- (a) $\bar{\Sigma}_p$ is a codimension- $(2 \dim G_p)$ torus orbifold with $H^{\text{odd}}(\bar{\Sigma}_p; \mathbb{Q}) = 0$.
- (b) The linear, effective action of \tilde{G}_p° on $\tilde{T}_p U_p = \tilde{T}_p \bar{\Sigma}_p \oplus \tilde{\nu}_p \bar{\Sigma}_p$ is trivial on the first summand and equivalent to the standard $(\dim G_p)$ -torus action on $\mathbb{C}^{\dim G_p}$ on the second. □

Proof. By Lemma 3.6, $\bar{\Sigma}_p \subseteq \mathcal{O}^{G_p}$ contains some fixed point p_0 of the G action. Since $\bar{\Sigma}_p$ is a strong G -invariant suborbifold of \mathcal{O} , it follows that $\tilde{T}_{p_0} \bar{\Sigma}_p$ is an \tilde{G} -invariant subspace of $\tilde{T}_{p_0} \mathcal{O}$. As \tilde{G} acts in the standard way on $\tilde{T}_{p_0} \mathcal{O}$, it follows that $\bar{\Sigma}_p$ has codimension $2 \dim G_p$ and, hence, is a torus orbifold with $H^{\text{odd}}(\bar{\Sigma}_p; \mathbb{Q}) = 0$.

The second claim follows from dimension reasons in a similar way as in the proof of Lemma 3.4. ■

Proposition 3.8. Let \mathcal{O}^{2n} be a torus orbifold with $H^{odd}(\mathcal{O}; \mathbb{Q}) = 0$. Then every point $p \in \mathcal{O}^{2n}$ lies in the closures of exactly $\dim(G_p)$ strata of codimension 2. Equivalently, a point $[p] \in \mathcal{O}/G$ in the (relative) interior of a face of codimension k , lies in exactly k faces of codimension 1 in \mathcal{O}/G . □

Proof. This follows from part (b) of Proposition 3.7 and the Slice Theorem. ■

Lemma 3.9. Let \mathcal{O} be a torus orbifold with $H^{odd}(\mathcal{O}; \mathbb{Q}) = 0$. Then the closure of each two-dimensional stratum of \mathcal{O} is homeomorphic to a two-sphere and each one-dimensional face (edge) in the quotient \mathcal{O}/G contains exactly two fixed points. □

Proof. Recall that the closure $\overline{\Sigma}_i^2$ of each two-dimensional stratum Σ_i^2 in a torus orbifold \mathcal{O} projects down to a one-dimensional face of \mathcal{O}/G . By Proposition 3.7, each $\overline{\Sigma}_i^2$ contains a fixed point of the G action and is a two-dimensional torus orbifold with $H^{odd}(\overline{\Sigma}_i^2; \mathbb{Q}) = 0$. By [35, Chap. 13] the $\overline{\Sigma}_i^2$ are closed, orientable, topological 2-manifolds with positive Euler characteristic, hence each must be homeomorphic to a two-dimensional sphere. Therefore each $\overline{\Sigma}_i^2$ has Euler characteristic 2 and contains exactly two fixed points of the G action. ■

4 Weights, GKM-Graphs and the Moment-Angle Complex

Let \mathcal{O} be a $2n$ -dimensional torus orbifold with $H^{odd}(\mathcal{O}; \mathbb{Q}) = 0$. A *facet* of the orbit space $Q = \mathcal{O}/G$ is a face of codimension one. Recall that, by Proposition 3.8, this corresponds, in \mathcal{O} , to the closure $\overline{\Sigma}_p$ of a stratum Σ_p defined by a one-dimensional isotropy group G_p .

Given a facet F , let $p \in \mathcal{O}$ be a point with $\dim(G_p) = 1$ such that $\overline{\Sigma}_p$ is the pre-image of F , and let $U_p \subset \mathcal{O}$ be a G_p -invariant good local chart around p .

Formally assign a circle S_F^1 to F and let the *label*

$$\lambda_F : S_F^1 \rightarrow G$$

denote the composition (covering)

$$S_F^1 \xrightarrow{\cong} \tilde{G}_p^o \xrightarrow{\rho_p} G_p^o \subseteq G,$$

where the map $\rho_p : \tilde{G}_p \rightarrow G_p$ is that arising in Proposition 2.8. Set now $T_Q = \prod_F S_F^1$ and define the *label map*

$$\lambda = \prod_F \lambda_F : T_Q \rightarrow G.$$

Lemma 4.1. The label map $\lambda : T_Q \rightarrow G$ is well defined. □

Proof. In order to verify that the map λ is well defined, it need only be demonstrated that the labels λ_F do not depend on the choice of the point in the pre-image $\bar{\Sigma}_p$ of a facet F . By Definition 3.2, if $q \in \bar{\Sigma}_p$ is another point with $\dim(G_q) = 1$, then $G_p^o = G_q^o$ as subgroups of G . It suffices to show that there is an isomorphism $\alpha : \tilde{G}_q^o \rightarrow \tilde{G}_p^o$ such that the following diagram commutes:

$$\begin{array}{ccccc}
 S_F^1 & \xrightarrow{\cong} & \tilde{G}_p^o & \xrightarrow{\rho_p} & G_p^o \\
 & \searrow \cong & \uparrow \alpha & & \uparrow = \\
 & & \tilde{G}_q^o & \xrightarrow{\rho_q} & G_q^o
 \end{array}$$

As each of $G_p^o, G_q^o, \tilde{G}_p^o$ and \tilde{G}_q^o is a circle, if the kernels of ρ_p and ρ_q have the same order, then it is possible to lift the identity $G_q^o \xrightarrow{=} G_p^o$ to such an isomorphism α .

The kernels of ρ_p and ρ_q are given by $\Gamma_p \cap \tilde{G}_p^o$ and $\Gamma_q \cap \tilde{G}_q^o$, respectively. Since the stratum Σ_p is connected, it is enough to show that the order of $\Gamma_p \cap \tilde{G}_p^o$ is locally constant.

Let U_p be a sufficiently small G_p -invariant good local chart around p such that \tilde{U}_p is a linear \tilde{G}_p -representation. Then \tilde{U}_p is of the form $V \oplus W$ such that \tilde{G}_p^o acts non-trivially on V and trivially on W (see Proposition 3.7).

Moreover, since Σ_p has codimension 2, it follows that V is two dimensional and $\pi_p(W) = \Sigma_p \cap U_p$. Therefore, the subgroup $\Gamma_p \cap \tilde{G}_p^o$ of Γ_p acts trivially on W . For any $q \in \Sigma_p \cap U_p$ and $\tilde{q} \in \pi_p^{-1}(q) \cap W$ one has $\Gamma_p \cap \tilde{G}_p^o \subseteq (\Gamma_p)_{\tilde{q}}$. By Lemmas 2.4 and 2.12, $\Gamma_q = (\Gamma_p)_{\tilde{q}}$ and $\tilde{G}_q^o = \tilde{G}_p^o$, hence $\Gamma_p \cap \tilde{G}_p^o \subseteq \Gamma_q \cap \tilde{G}_q^o$. On the other hand, the same lemmas yield $\Gamma_q \subseteq \Gamma_p$ and $\tilde{G}_q^o \subseteq \tilde{G}_p^o$, hence $\Gamma_q \cap \tilde{G}_q^o \subseteq \Gamma_p \cap \tilde{G}_p^o$. Therefore, $\Gamma_p \cap \tilde{G}_p^o$ is locally constant. ■

The labels of the facets can be used to define weights on the edges of the orbit space. By Proposition 3.8, any edge E is the intersection of $n - 1$ facets $\{F_1, \dots, F_{n-1}\}$ and, by restricting the label map to $T_E = \prod_{i=1}^{n-1} S_{F_i}^1$, one obtains a homomorphism $\lambda_E : T_E \rightarrow G$. Let p_i be a generic point in the stratum corresponding to F_i . As $S_{F_i}^1 \rightarrow G_{p_i}^o \subseteq G$ is a covering, for all facets F_i , the map λ_E induces an injective map $t_E \rightarrow \mathfrak{g}$ on Lie algebras,

hence a surjective map $\mathfrak{g}^* \rightarrow \mathfrak{t}_E$ on the corresponding dual spaces. Since the dual \mathfrak{l}^* of the Lie algebra of a Lie group L is canonically isomorphic to $H^2(BL; \mathbb{R})$, one concludes that the induced map $\lambda_E^* : H^2(BG; \mathbb{Z}) = \mathbb{Z}^n \rightarrow H^2(BT_E; \mathbb{Z}) = \mathbb{Z}^{n-1}$ has full rank. Define the *weight* $\mu(E) \in H^2(BG; \mathbb{Z})$ of E to be a generator of the kernel of λ_E^* .

In this way, one obtains a system of weights on the vertex-edge graph of the orbit space \mathcal{O} , that is, on the union of edges and vertices. This is the well-known GKM-graph associated to the torus orbifold \mathcal{O} . In an analogous manner to the manifold case (cf. [28]), this graph determines the rational equivariant cohomology ring $H_G^*(\mathcal{O}; \mathbb{Q}) = H^*(\mathcal{O}_G; \mathbb{Q})$ of \mathcal{O} , where $\mathcal{O}_G = \mathcal{O} \times_G EG$ is the Borel construction, in the following way: Since $H^{\text{odd}}(\mathcal{O}; \mathbb{Q}) = 0$ by assumption, it follows that $H_G^*(\mathcal{O}; \mathbb{Q})$ is a free $H^*(BG; \mathbb{Q})$ -module, which is easily seen from the spectral sequence of the homotopy fibration $\mathcal{O} \rightarrow \mathcal{O}_G \rightarrow BG$. In particular, the induced homomorphism $H_G^*(\mathcal{O}; \mathbb{Q}) \rightarrow H^*(\mathcal{O}; \mathbb{Q})$ is surjective.

If $\mathcal{O}^{(1)} \subseteq \mathcal{O}$ is the union of all G -orbits of dimension at most one, that is, the preimage of the vertex-edge graph of \mathcal{O} , the respective inclusion maps induce a commutative diagram

$$\begin{array}{ccc} H_G^*(\mathcal{O}^{(1)}; \mathbb{Q}) & \longrightarrow & H_G^*(\mathcal{O}^G; \mathbb{Q}) \\ \uparrow & \nearrow & \\ H_G^*(\mathcal{O}; \mathbb{Q}) & & \end{array}$$

It follows from Lemma 2.3 and Proposition 2.4 of [5] that the homomorphisms $H_G^*(\mathcal{O}; \mathbb{Q}) \rightarrow H_G^*(\mathcal{O}^G; \mathbb{Q})$ and $H_G^*(\mathcal{O}^{(1)}; \mathbb{Q}) \rightarrow H_G^*(\mathcal{O}^G; \mathbb{Q})$ have the same image, and the former homomorphism is injective. Furthermore, the homomorphism $H_G^*(\mathcal{O}; \mathbb{Q}) \rightarrow H_G^*(\mathcal{O}^{(1)}; \mathbb{Q})$ must, therefore, also be injective.

By Lemma 3.9, $\mathcal{O}^{(1)}$ is a union of two-dimensional spheres (intersecting only in the fixed points of the G action). Therefore, the following theorem follows as in the manifold case [11, Theorem 7.2]:

Theorem 4.2. Let \mathcal{O} be a torus orbifold with fixed points $\{p_1, \dots, p_N\}$ and $H^{\text{odd}}(\mathcal{O}; \mathbb{Q}) = 0$. Then, via the natural restriction map

$$H_G^*(\mathcal{O}; \mathbb{Q}) \rightarrow H_G^*(\mathcal{O}^G; \mathbb{Q}) = \bigoplus_{i=1}^N H^*(BG; \mathbb{Q}),$$

the equivariant cohomology algebra $H_G^*(\mathcal{O}; \mathbb{Q})$ is isomorphic to the set of N -tuples $(f_1, \dots, f_N) \in H_G^*(\mathcal{O}^G; \mathbb{Q})$, with the property that if the vertices p_i and p_j in the associated

GKM-graph are joined by an edge with weight $\mu_{ij} \in H^2(BG; \mathbb{Q})$, then $f_i - f_j$ lies in the ideal of $H^*(BG; \mathbb{Q})$ generated by μ_{ij} . □

Remark 4.3. The process by which one obtains $H_G^*(\mathcal{O}; \mathbb{Q})$ from the GKM-graph is functorial in the following sense: Suppose that \mathcal{O} (respectively, $\hat{\mathcal{O}}$) is a $2n$ -dimensional torus orbifold with fixed points p_1, \dots, p_N (respectively, $\hat{p}_1, \dots, \hat{p}_{\hat{N}}$) and weights μ (respectively, $\hat{\mu}$). Suppose, further, that there is a weight-preserving, injective map φ between the GKM-graphs of \mathcal{O} and $\hat{\mathcal{O}}$, i.e. $\hat{\mu}(\varphi(E)) = \mu(E)$, for each edge E of \mathcal{O}/G .

The map φ induces an injective homomorphism

$$\varphi_{\#} : \bigoplus_{i=1}^N H^*(BG; \mathbb{Q}) \rightarrow \bigoplus_{i=1}^{\hat{N}} H^*(BG; \mathbb{Q}),$$

where, for each $i_0 \in \{1, \dots, N\}$ and given $\varphi(p_{i_0}) = \hat{p}_{j_0}$, the restriction of $\varphi_{\#}$ to the i_0 -th summand of $\bigoplus_{i=1}^N H^*(BG; \mathbb{Q})$ is given by the identity map onto the j_0 -th summand of the target space $\bigoplus_{i=1}^{\hat{N}} H^*(BG; \mathbb{Q})$.

By Theorem 4.2, $H_G^*(\mathcal{O}; \mathbb{Q})$ and $H_G^*(\hat{\mathcal{O}}; \mathbb{Q})$ embed into $\bigoplus_{i=1}^N H^*(BG; \mathbb{Q})$ and $\bigoplus_{i=1}^{\hat{N}} H^*(BG; \mathbb{Q})$, respectively. Since φ is weight-preserving, $\varphi_{\#}$ maps $H_G^*(\mathcal{O}; \mathbb{Q})$ into $H_G^*(\hat{\mathcal{O}}; \mathbb{Q})$. It then follows that there is an induced $H^*(BG; \mathbb{Q})$ -module homomorphism

$$H_G^*(\mathcal{O}; \mathbb{Q}) \rightarrow H_G^*(\hat{\mathcal{O}}; \mathbb{Q}),$$

which, by abuse of language, will be denoted also by $\varphi_{\#}$. Moreover, if $H^{\text{odd}}(\mathcal{O}; \mathbb{Q}) = H^{\text{odd}}(\hat{\mathcal{O}}; \mathbb{Q}) = 0$, then

$$H^*(\mathcal{O}; \mathbb{Q}) = H_G^*(\mathcal{O}; \mathbb{Q})/H^{>0}(BG; \mathbb{Q}) \cdot H_G^*(\mathcal{O}; \mathbb{Q})$$

and similarly for $H^*(\hat{\mathcal{O}}; \mathbb{Q})$. Hence, there is an induced homomorphism

$$\bar{\varphi}_{\#} : H^*(\mathcal{O}; \mathbb{Q}) \rightarrow H^*(\hat{\mathcal{O}}; \mathbb{Q})$$

such that the diagram

$$\begin{array}{ccc} H_G^*(\mathcal{O}; \mathbb{Q}) & \xrightarrow{\varphi_{\#}} & H_G^*(\hat{\mathcal{O}}; \mathbb{Q}) \\ \downarrow & & \downarrow \\ H^*(\mathcal{O}; \mathbb{Q}) & \xrightarrow{\bar{\varphi}_{\#}} & H^*(\hat{\mathcal{O}}; \mathbb{Q}) \end{array} \tag{4.1}$$

commutes. □

Recall now that an n -dimensional manifold with corners Q , that is, a manifold locally modelled on \mathbb{R}_+^n , is called *nice* if each one of its codimension- k faces is contained in exactly k facets, that is, codimension-1 faces, of Q .

Formally assign a copy S_F^1 of the circle to each facet F of Q and let $T_Q = \prod_F S_F^1$ be the torus given by their product.

For any $q \in Q$, let $T(q) = \prod_{F \ni q} S_F^1 \subseteq T_Q$ denote the subtorus generated by the circles corresponding to the facet of Q which contain q . The *moment-angle complex* is defined by $Z_Q = (Q \times T_Q)/\sim$, where $(q_1, t_1) \sim (q_2, t_2)$ if $q_1 = q_2$ and $t_1 t_2^{-1} \in T(q_1)$.

As Q is a nice manifold with corners, it follows that Z_Q is a topological manifold with a continuous T_Q action, such that Z_Q/T_Q is homeomorphic to Q .

Suppose that, in addition, Q has zero-dimensional faces. Consider a torus $G = T^n$ and a homomorphism

$$\hat{\lambda} : T_Q \rightarrow G$$

such that, for every $q \in Q$, the restriction $\hat{\lambda}|_{T(q)} : T(q) \rightarrow G$ has finite kernel. This condition ensures that the kernel K of $\hat{\lambda}$ acts almost freely on Z_Q . The group G then acts on the quotient $\mathcal{O}_Q = Z_Q/K$ such that (\mathcal{O}_Q, G) is a $2n$ -dimensional torus orbifold whose orbit space \mathcal{O}_Q/G has labels induced by the assignment $\hat{\lambda}$, and there is a face-preserving homeomorphism $\mathcal{O}_Q/G \rightarrow Q$.

The following three standard examples will be needed in the proof of Theorem A.

Example 4.4. [4, Ex. 6.7] If $Q = \Delta^n$ is an n -dimensional simplex, then T_Q is an $(n+1)$ -dimensional torus. Moreover, Z_Q is equivariantly homeomorphic to $\mathbb{S}^{2n+1} \subseteq \mathbb{C}^{n+1}$ with the standard linear torus action. \square

Example 4.5. [29, Ex. 4.3] If $Q = \Sigma^n$ is the suspension of the simplex Δ^{n-1} , then T_Q is n -dimensional. Moreover, Z_Q is equivariantly homeomorphic to $\mathbb{S}^{2n} \subseteq \mathbb{C}^n \times \mathbb{R}$ with the standard linear torus action. \square

Example 4.6. [4, Prop. 6.4] Let Q_1 and Q_2 be two nice manifolds with corners. If $Q = Q_1 \times Q_2$, then $T_Q = T_{Q_1} \times T_{Q_2}$ and Z_Q is equivariantly homeomorphic to $Z_{Q_1} \times Z_{Q_2}$. \square

5 Equivariant Classification of Torus Orbifolds

In order to prove Theorem A, it is necessary to first understand the combinatorial properties of the face poset of the orbit space \mathcal{O}/G .

Proposition 5.1. Let \mathcal{O} be a simply connected, rationally elliptic torus orbifold. Then the face poset of \mathcal{O}/G satisfies:

- (a) The vertex-edge graph of each face is connected.
- (b) Each face of \mathcal{O}/G contains at least one vertex.
- (c) Each face of \mathcal{O}/G of codimension k is contained in exactly k faces of codimension 1.
- (d) Each one-dimensional face of \mathcal{O}/G contains exactly two fixed points of the G action.
- (e) Every two-dimensional face of \mathcal{O}/G contains at most four vertices.
- (f) For $d \geq 3$, no d -dimensional face is combinatorially equivalent to the face poset of $[-1, 1]^d/\{\pm \text{id}\}$. \square

Proof. Property (a) follows from [5, Prop. 2.5]. Indeed, $H^{\text{odd}}(\mathcal{O}; \mathbb{Q}) = 0$ and $H^{\text{odd}}(BG; \mathbb{Q}) = 0$ imply that the differentials in the spectral sequence of the homotopy fibration $\mathcal{O} \rightarrow \mathcal{O}_G \rightarrow BG$ are trivial. Therefore $H_G^*(\mathcal{O}; \mathbb{Q}) = H^*(\mathcal{O}_G; \mathbb{Q}) = H^*(\mathcal{O}; \mathbb{Q}) \otimes H^*(BG; \mathbb{Q})$, thus fulfilling the hypotheses of the aforementioned proposition. Properties (b), (c) and (d) have been verified in Lemmas 3.6, 3.8, and 3.9, respectively.

To see that property (e) holds, one must modify the proof of Lemma 4.2 of [37] for the case of torus orbifolds only slightly. The original proof invokes [1, Corollary 3.3.11] which, although stated only for rationally elliptic G -CW-complexes, also holds for compact spaces with finitely many orbit types, for example, torus orbifolds, as indicated on page 160 of [1].

Finally, suppose that there is a d -dimensional face F combinatorially equivalent to the face poset of $X := [-1, 1]^d/\{\pm \text{id}\}$. Notice first that the standard linear, effective T^d action on $(\mathbb{S}^2)^d$ commutes with the diagonal antipodal map and, therefore, induces an effective T^d action on $N = (\mathbb{S}^2)^d/\mathbb{Z}_2$ with orbit space X . Thus, the quotient of the T^d action on the preimage of F is combinatorially equivalent to the quotient of the T^d action on N and, in particular, the corresponding GKM-graphs are isomorphic. By the discussion before Theorem 4.2, their rational cohomology rings are the same. However, the pre-image of F is rationally elliptic by [1, Cor. 3.3.11], while, on the other hand, N is not: Indeed, by [2, Thm. 2.4], $H^*(N; \mathbb{Q}) = H^*((\mathbb{S}^2)^d; \mathbb{Q})^{\mathbb{Z}_2}$ and therefore the Betti numbers of N satisfy $b_1(N) = b_2(N) = b_3(N) = 0$, $b_4(N) = d(d-1)/2$. In particular $\dim_{\mathbb{Q}}(\pi_4(N) \otimes \mathbb{Q}) = b_4(N)$ and, if N were rationally elliptic, Theorem 32.6 in [7] would yield

$$2d(d-1) = 4 \dim_{\mathbb{Q}}(\pi_4(N) \otimes \mathbb{Q}) \leq \sum_j 2j \dim_{\mathbb{Q}}(\pi_{2j}(N) \otimes \mathbb{Q}) \leq 2d$$

which is not possible for $d > 2$. ■

Proof of Theorem A. Following the arguments involved in proving [37, Prop. 4.5], the properties established in Proposition 5.1 are precisely those required to prove that the face poset of \mathcal{O}/G is combinatorially equivalent to the face poset of $Q = \prod_i \Delta^{n_i} \times \prod_j \Sigma^{n_j}$ as in Examples 4.4–4.6, that is, there is an isomorphism of face posets $\varphi : \mathcal{P}(\mathcal{O}/G) \rightarrow \mathcal{P}(Q)$. For each facet $F \in \mathcal{P}(Q)$, fix an isomorphism $\iota_F : S_F^1 \rightarrow S_{\varphi^{-1}(F)}^1$.

With Q as above, the moment-angle complex Z_Q of Q , together with the action of T_Q as discussed in Section 4, is equivariantly homeomorphic to a product of spheres $\prod \mathbb{S}^{n_i}$ equipped with a linear action.

The isomorphism φ induces a label map $\varphi_*\lambda : T_Q \rightarrow G$, such that the restriction to each factor S_F^1 is given by $\lambda_{\varphi^{-1}(F)} \circ \iota_F$. By setting $\hat{\lambda} = \varphi_*\lambda$, one can construct, as before, a torus orbifold (\mathcal{O}_Q, G) , where \mathcal{O}_Q is the quotient of $\prod \mathbb{S}^{n_i}$ by the linear and almost-free action of a subtorus of T_Q complementary to G .

This is achieved as follows: The kernel \hat{L} of $\hat{\lambda}$ acts almost freely on $\prod \mathbb{S}^{n_i}$, although it may not be connected. Therefore, $\prod \mathbb{S}^{n_i}/\hat{L}$ is a torus orbifold. Moreover, the identity component \hat{L}° of \hat{L} is a subtorus of T_Q .

Since the natural action of the finite group \hat{L}/\hat{L}° on $\prod \mathbb{S}^{n_i}/\hat{L}^\circ$ extends to an action of the connected group T_Q/\hat{L}° , the induced action on cohomology is trivial. Hence, by [2, Thm. 2.4], $\prod \mathbb{S}^{n_i}/\hat{L}$ and $\prod \mathbb{S}^{n_i}/\hat{L}^\circ$ have isomorphic rational cohomology rings. Moreover, by Proposition 32.16 of [7] and Corollary 2.7.9 of [1], the minimal models of these spaces are formal and, therefore, isomorphic. Hence, it may be assumed that \hat{L} is connected. In this case, define $\mathcal{O}_Q = \prod \mathbb{S}^{n_i}/\hat{L}$.

By construction, the torus orbifolds (\mathcal{O}, G) and (\mathcal{O}_Q, G) have isomorphic labelled face posets, hence isomorphic GKM-graphs. Therefore the rational cohomology rings of \mathcal{O} and \mathcal{O}_Q are isomorphic, as discussed after Theorem 4.2. But once again, the minimal models of these spaces are formal by Proposition 32.16 of [7] and Corollary 2.7.9 of [1]. Since their cohomology rings are isomorphic, this implies that the spaces are rationally homotopy equivalent.

Since $H^*(\mathcal{O}_G; \mathbb{Q}) = H^*(\mathcal{O}; \mathbb{Q}) \otimes H^*(BG; \mathbb{Q})$ as modules over $H^*(BG; \mathbb{Q})$, BT , \mathcal{O} and \mathcal{O}_Q are formal and \mathcal{O} and \mathcal{O}_Q are rationally elliptic, it follows from Proposition 3.2 of [26] that the minimal models of \mathcal{O}_G and $(\mathcal{O}_Q)_G$ are formal. As $H^*(\mathcal{O}_G; \mathbb{Q})$ is isomorphic to $H^*((\mathcal{O}_Q)_G; \mathbb{Q})$, this ensures that the minimal models of the Borel constructions \mathcal{O}_G and $(\mathcal{O}_Q)_G$ are isomorphic, hence $\mathcal{O}_G \simeq_{\mathbb{Q}} (\mathcal{O}_Q)_G$.

Furthermore, from the face-poset isomorphism φ , one obtains a commutative diagram as in (4.1), where the horizontal arrows are clearly the isomorphisms induced by the rational homotopy equivalence $\mathcal{O} \simeq_{\mathbb{Q}} \mathcal{O}_Q$. Therefore, \mathcal{O} is G -equivariantly rationally homotopy equivalent to \mathcal{O}_Q .

Finally, if \mathcal{O} is a (torus) manifold, then all local groups are trivial and one can identify \tilde{U}_p with U_p , \tilde{G}_p with G_p , and so on. Given any $p \in \mathcal{O}$ with $\dim(G_p) = l$, Proposition 3.8 states that p belongs to the closures $\bar{\Sigma}_i$, $i = 1, \dots, l$, of l codimension-2 strata. By Proposition 3.8 again, each $\bar{\Sigma}_i$ is fixed by a different factor S_i^1 of $G_p^o = T^l$, and the $\bar{\Sigma}_i$ project to distinct facets F_i of \mathcal{O}/G .

By definition, $\lambda_{F_i} : S_{F_i}^1 \rightarrow G$ sends $S_{F_i}^1$ isomorphically into $S_i^1 \subseteq G_p^o$ and, therefore, the label map λ sends $T([p]) = \prod_i S_{F_i}^1$ isomorphically into $G_p^o = \prod_i S_i^1$, where $[p] \in \mathcal{O}/G$ is the image of p .

In particular, the restriction of λ to $T([p])$ has trivial kernel. It then follows that the kernel of λ has trivial intersection with each such torus $T([p])$. Since the label map $\varphi_*\lambda$ has the same properties as λ , the kernel of $\varphi_*\lambda$ has trivial intersection with all isotropy subgroups of the T_Q action on Z_Q and, therefore, acts freely on Z_Q . Hence, \mathcal{O}_Q is also a manifold. ■

6 A Family of Examples

The family of examples in this section shows the necessity of including almost-free actions in the conclusion of Theorem A, and also gives an explicit demonstration of how to apply the GKM algorithm discussed in Section 4.

Consider a 2-torus G and a two-dimensional nice manifold with corners, Q , whose boundary consists of four segments labelled as in the square on the left of Figure 1, where $a, b, c, d \in \mathbb{Z}$. In this example, the four edges and facets of Q coincide. The labels of the facets of Q are the slopes $\in \mathbb{Z}^2$ corresponding to circle subgroups (tori of codimension one) in G . By the discussion following Example 4.6, in order to construct




Fig. 1. (l) Labelled orbit space Q ; (r) GKM-graph Γ_Q .

a four-dimensional torus orbifold with orbit space Q the corresponding labels must be linearly independent whenever two facets intersect. Assume therefore that

$$a, d, \det \begin{pmatrix} a & b \\ c & d \end{pmatrix} \neq 0.$$

Since $Q = [0, 1]^2$, it follows from Examples 4.4 and 4.6 that the moment angle complex Z_Q is equivariantly homeomorphic to $\mathbb{S}^3 \times \mathbb{S}^3$ equipped with the standard linear T^4 action. There is, moreover, a surjective homomorphism $T^4 \rightarrow G$ whose kernel is a 2-torus K acting almost freely on $Z_Q \cong \mathbb{S}^3 \times \mathbb{S}^3$. The resulting orbifold $\mathcal{O}_Q = Z_Q/K \cong (\mathbb{S}^3 \times \mathbb{S}^3)/K$ is a simply connected, rationally elliptic, four-dimensional torus orbifold whose labelled orbit space Q under the action of G is as on the left of Figure 1. Indeed, in this case the action of K on $\mathbb{S}^3 \times \mathbb{S}^3$ can be written explicitly as

$$K \times (\mathrm{SU}(2) \times \mathrm{SU}(2)) \rightarrow \mathrm{SU}(2) \times \mathrm{SU}(2)$$

$$((z, w), (A, B)) \mapsto \begin{pmatrix} \mathrm{diag}(z^{1-a}\bar{w}^c, 1)A \mathrm{diag}(z^a w^c, \bar{z}) \\ \mathrm{diag}(\bar{z}^b w^{1-d}, 1)B \mathrm{diag}(z^b w^d, \bar{w}) \end{pmatrix}.$$

It remains to demonstrate that not all such labelled orbit spaces Q can be realised by torus manifolds, hence that one cannot always find a torus manifold which is rationally homotopy equivalent to a given torus orbifold. This will be achieved by computing the cohomology ring and intersection form of the torus orbifold \mathcal{O}_Q .

As discussed in Section 4, each edge E of the labelled orbit space Q can be assigned a weight $\mu(E) \in H^2(BG; \mathbb{Z})$. The resulting GKM-graph Γ_Q is shown on the right of Figure 1. As there are four vertices (corresponding to the fixed points of the G action on \mathcal{O}_Q), Theorem 4.2 implies that $H_G^*(\mathcal{O}_Q; \mathbb{Q})$, the equivariant cohomology algebra of \mathcal{O}_Q , is isomorphic to the set of all 4-tuples $(f_1, f_2, f_3, f_4) \in \bigoplus_{i=1}^4 H^*(BG; \mathbb{Q}) = \bigoplus_{i=1}^4 \mathbb{Q}[x, y]$ satisfying the relations

$$f_1 - f_2 = m_1 y,$$

$$f_2 - f_3 = m_2 x,$$

$$f_3 - f_4 = m_3 (bx - ay), \text{ and}$$

$$f_4 - f_1 = m_4 (dx - cy),$$

where $m_1, m_2, m_3, m_4 \in \mathbb{Q}[x, y]$, and the ring structure is given by coordinate-wise multiplication. It is straightforward to check that the equivariant cohomology of \mathcal{O}_Q

is then generated as a $\mathbb{Q}[x, y]$ -module by $\mathbf{1} = (1, 1, 1, 1)$, $u = (0, -ay, bx - ay, 0)$, $v = (0, -cy, dx - cy, dx - cy)$ and $w = (0, xy, 0, 0)$, of degree 0, 2, 2 and 4 (in $H_G^*(\mathcal{O}_a; \mathbb{Q})$), respectively. Clearly $\mathbf{1}$ is the unit element.

As $H_G^{\text{odd}}(\mathcal{O}_a; \mathbb{Q}) = 0$ (Lemma 3.3) and, hence, $H_G^*(\mathcal{O}_a; \mathbb{Q})$ is a free $H^*(BG; \mathbb{Q})$ -module, it follows that the rational cohomology of \mathcal{O}_a is given by

$$H^*(\mathcal{O}_a; \mathbb{Q}) = H_G^*(\mathcal{O}_a; \mathbb{Q}) / (R^+ \cdot H_G^*(\mathcal{O}_a; \mathbb{Q})),$$

where $R^+ = H^{>0}(BG; \mathbb{Q})$ and $R^+ \cdot H_G^*(\mathcal{O}_a; \mathbb{Q})$ is the set of all 4-tuples of the form $m_1 \mathbf{1} + m_2 u + m_3 v + m_4 w$, for polynomials $m_1, \dots, m_4 \in \mathbb{Q}[x, y]$ with zero constant term. Therefore, letting α, β and γ in $H^*(\mathcal{O}_a; \mathbb{Q})$ be the classes represented by u, v and w , respectively, $H^2(\mathcal{O}_a; \mathbb{Q})$ is generated (over \mathbb{Q}) by α and β , $H^4(\mathcal{O}_a; \mathbb{Q})$ by γ , and $H^i(\mathcal{O}_a; \mathbb{Q}) = 0, i \neq 0, 2, 4$. Moreover, the ring structure is given by the relations

$$\alpha^2 = ab\gamma, \beta^2 = cd\gamma, \text{ and } \alpha\beta = ad\gamma.$$

Indeed, this implies that $\alpha(d\alpha - b\beta) = 0$ and $\beta(a\beta - c\alpha) = 0$.

Whenever either $b = 0$ or $c = 0$, it is easy to see that one can find generators $\tilde{\alpha}, \tilde{\beta} \in H^2(\mathcal{O}_a; \mathbb{Q})$ such that $\tilde{\alpha}^2 = 0, \tilde{\beta}^2 = 0$ and $\gamma = \tilde{\alpha}\tilde{\beta}$.

On the other hand, if $bc \neq 0$ (by assumption $ad(ad - bc) \neq 0$), then the generators $\tilde{\alpha} = \frac{1}{b}\alpha$ and $\tilde{\beta} = \frac{a}{b}(d\alpha - b\beta)$ satisfy

$$\tilde{\alpha}\tilde{\beta} = 0 \text{ and } \tilde{\beta}^2 + ad(ad - bc)\tilde{\alpha}^2 = 0.$$

Furthermore, $\tilde{\alpha}^2$ generates $H^4(\mathcal{O}_a; \mathbb{Q})$ and the intersection form is given by $\text{diag}(1, -ad(ad - bc))$.

However, $\mathbb{S}^4, \mathbb{C}P^2, \mathbb{C}P^2 \# \pm \mathbb{C}P^2$ and $\mathbb{S}^2 \times \mathbb{S}^2$ are the only closed, simply connected, smooth, rationally elliptic 4-manifolds. Therefore, if \mathcal{O}_a were to be rationally homotopy equivalent to such a manifold, it would have intersection form either $\text{diag}(1, \pm 1)$ or $\begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$, corresponding to $\mathbb{C}P^2 \# \pm \mathbb{C}P^2$ or $\mathbb{S}^2 \times \mathbb{S}^2$. This is clearly not true for generic $a, b, c, d \in \mathbb{Z}$.

7 Slice-Maximal Torus Actions

The goal of this section is to prove Theorem B. To that end, let M be a closed, smooth, simply connected, rationally elliptic, n -dimensional manifold admitting a slice-maximal action by a torus T_M of rank k . If s denotes the maximal dimension of an isotropy subgroup, the action being slice maximal is equivalent to the identity $n = k + s$.

Under these hypotheses, there is a torus $K_M \subseteq T_M$ acting almost freely on M , with $\dim K_M = k - s$. Since the action of K_M on M is almost free, the orbit space M/K_M is an orbifold \mathcal{O} . Moreover, \mathcal{O} is rationally elliptic and has an induced action of the torus $T_{\mathcal{O}} = T_M/K_M$ of rank $s = \frac{1}{2} \dim \mathcal{O}$. The images of the T_M -orbits of (minimal) dimension $k - s$ under the quotient map $M \rightarrow \mathcal{O}$ correspond to fixed points of the $T_{\mathcal{O}}$ action. Hence, $(\mathcal{O}, T_{\mathcal{O}})$ is a simply connected, rationally elliptic torus orbifold.

By Theorem A, \mathcal{O} is $T_{\mathcal{O}}$ -equivariantly rationally homotopy equivalent to a simply connected torus orbifold $(\hat{\mathcal{O}} = \hat{P}/\hat{L}, T_{\mathcal{O}})$, where \hat{P} is a product of spheres of dimension ≥ 3 and \hat{L} is a compact abelian Lie group acting linearly and almost freely on \hat{P} . Recall from Section 5 that \hat{L} is defined as the kernel of the label map $\lambda : T_{\mathcal{O}} \rightarrow T_{\mathcal{O}}$, where $T_{\mathcal{O}} = \prod_F S_F^1$ is the product of a copy of S^1 for each facet of the orbit space $Q = \mathcal{O}/T_{\mathcal{O}} = M/T_M$. Since λ is onto, this yields an isomorphism $T_{\mathcal{O}} = T_{\mathcal{O}}/\hat{L}$.

Consider the map $\pi : M \rightarrow M/K_M = \mathcal{O}$, $p \in M$ and $p^* = \pi(p) \in \mathcal{O}$. A $(T_{\mathcal{O}})_{p^*}$ -invariant good local chart around p^* is given by $\tilde{U}_{p^*} = \nu_p(K_M(p))$ with map $\tilde{U}_{p^*} \rightarrow \mathcal{O}$ given by the composition $\nu_p(K_M(p)) \xrightarrow{\text{exp}} M \xrightarrow{\pi} \mathcal{O}$. The local group at p^* is given by $\Gamma_{p^*} = K_M \cap (T_M)_p$. Thus, following the notation of Proposition 2.8, one has $(\tilde{T}_{\mathcal{O}})_{p^*} \subset T_M$ and

$$(\tilde{T}_{\mathcal{O}})_{p^*}^o = (T_M)_p^o.$$

In particular, the slice representation of $(T_M)_p^o$ on $\nu_p(T_M(p))$ coincides with the slice representation of $(\tilde{T}_{\mathcal{O}})_{p^*}$ on $\tilde{\nu}_{p^*}(T_{\mathcal{O}}(p^*))$ as in the Slice Theorem (Theorem 2.13). From Proposition 3.7(b), this action is a sum of a trivial summand and a maximal-rank summand. Such actions belong to a class called *polar actions* and, since every slice representation of T_M is polar, the T_M action on M is *infinitesimally polar* (see [14, 32]). Here an isometric action on a Riemannian manifold N is called polar if there is a submanifold of N which intersects every orbit orthogonally. A group action is called infinitesimally polar if all slice representations are polar.

By definition of the label map, each $\lambda_F : S_F^1 \rightarrow (T_{\mathcal{O}})_{q^*}^o \subseteq T_{\mathcal{O}}$, with $q^* \in \mathcal{O}$ a point projecting to F , factors through $(\lambda_M)_F : S_F^1 \rightarrow (\tilde{T}_{\mathcal{O}})_{q^*}^o = (T_M)_q^o$ and, therefore, the map $\lambda : T_{\mathcal{O}} \rightarrow T_{\mathcal{O}}$ naturally admits a lift to a map $\lambda_M : T_{\mathcal{O}} \rightarrow T_M$.

Lemma 7.1. Let $\lambda_M : T_{\mathcal{O}} \rightarrow T_M$ be the above-defined lift of the label map. Then:

- (a) λ_M is surjective.
- (b) For every $p \in M$ projecting to $q \in Q$, the torus $T(q) \subseteq T_{\mathcal{O}}$ is mapped isomorphically onto $(T_M)_p^o$. □

Proof. Part (a). Let M_{reg} denote the collection of principal orbits, and $O_{\text{reg}} = M_{\text{reg}}/T_M$. Since the T_M action on M_{reg} is free, there is a principal bundle

$$T_M \rightarrow M_{\text{reg}} \rightarrow O_{\text{reg}}.$$

Since M is simply connected and the T_M -action on M is infinitesimally polar, by Theorem 1.8 of [27] there are no orbits with finite isotropy and, therefore, the set O_{reg} consists precisely of the orbits of maximal dimension. On the other hand, O_{reg} is the quotient $O_{\text{reg}}/T_{\mathcal{O}}$, where O_{reg} also consists of the orbits of \mathcal{O} of maximal dimension. Because \mathcal{O} is a rationally elliptic torus orbifold, $H^{\text{odd}}(\mathcal{O}; \mathbb{Q}) = 0$. Therefore, Corollary 1 of [3] can be applied to conclude that O_{reg} is rationally acyclic. Since $\pi_1(O_{\text{reg}}, [p_0]) = \pi_1(O, [p_0]) = 0$, by Hurewicz $\pi_2(O_{\text{reg}}, [p_0]) \otimes \mathbb{Q} = H_2(O_{\text{reg}}; \mathbb{Q}) = 0$ and, in particular, $\pi_2(O_{\text{reg}}, [p_0])$ is torsion. From the long exact sequence in homotopy for $M_{\text{reg}} \rightarrow O_{\text{reg}}$, it follows that the kernel of $\pi_1(T_M) \rightarrow \pi_1(M_{\text{reg}}, p_0)$ must be torsion as well, but since $\pi_1(T_M)$ is free abelian, the kernel must be trivial. Therefore, the map $\pi_1(T_M) \rightarrow \pi_1(M_{\text{reg}}, p_0)$ is injective.

In order to prove that λ_M is surjective, it is enough to show that it induces a surjective map $(\lambda_M)_* : \pi_1(T_{\mathcal{O}}) \rightarrow \pi_1(T_M)$. Letting $\Omega \subseteq \pi_1(T_M)$ denote the image of $(\lambda_M)_*$, from the discussion above it is enough to prove that Ω has the same image as $\pi_1(T_M)$ in $\pi_1(M_{\text{reg}}, p_0)$.

For any $\alpha \in \pi_1(T_M)$, its image in $\pi_1(M_{\text{reg}}, p_0)$ is represented by some loop C in a principal T_M -orbit in M and, since M is simply connected, hence bounds a two-dimensional disk D in M . The pre-images of the facets of Q are codimension-2 submanifolds of M . Hence, by performing a suitable deformation, it may be assumed without loss of generality that D intersects only finitely many of these codimension-2 submanifolds, in only finitely many points x_1, \dots, x_N , and that these intersections are transversal. As D is simply connected, C is homotopy equivalent (within D) to a concatenation of lassos based at $p_0 \in C$, each of which has a noose γ_i which is a circle around a single intersection point x_i , $i \in \{1, \dots, N\}$.

For each $i \in \{1, \dots, N\}$, in a sufficiently small neighbourhood of the intersection point x_i , the disk D can be assumed to coincide with the normal slice to the T_M -orbit through x_i . By the Slice Theorem, a noose γ_i around x_i can be assumed to lie in an orbit of the slice action of the one-dimensional isotropy subgroup $(T_M)_{x_i}$, hence, to be some (positive or negative) iterate of the circle $(T_M)_{x_i}^o$.

Together with the isomorphisms arising via change of base points, the above discussion ensures that C is homotopic to the concatenation of the γ_i , each of which represents an element in $\pi_1(M_{\text{reg}}, p_0)$ in the image of Ω .

Part (b). This follows closely the last part of the proof of Theorem A. Given any $p \in M$ with $\dim((T_M)_p) = l$, Proposition 3.8 states that the image $p^* \in \mathcal{O}$ of p belongs to the closures $\overline{\Sigma}_i$ of codimension-2 strata Σ_i , $i = 1, \dots, l$. By Proposition 3.8 again, each $\overline{\Sigma}_i$ projects to a different facet F_i of Q , and it is fixed by a different factor S_i^1 of $(T_{\mathcal{O}})_{p^*}^o = T^l$, which lifts to a factor \tilde{S}_i^1 of $(\tilde{T}_{\mathcal{O}})_{p^*}^o = (T_M)_p^o$.


By definition, $(\lambda_M)_{F_i} : S_{F_i}^1 \rightarrow T_M$ sends $S_{F_i}^1$ isomorphically into $\tilde{S}_i^1 \subseteq (T_M)_p^o$ and, therefore, the label map λ sends $T([p]) = \prod_i S_{F_i}^1$ isomorphically into $(T_M)_p^o$, where $[p] \in Q$ is the image of p . ■

Let $K_{\hat{P}} \subseteq T_Q$ denote the kernel of λ_M and let \hat{M} be the quotient $\hat{P}/K_{\hat{P}}$. The group $K_{\hat{M}} = \hat{L}/K_{\hat{P}}$ acts almost freely on \hat{M} , with quotient $\hat{O} = \hat{P}/\hat{L}$. Recall, furthermore, that there is an isomorphism

$$\varphi : \mathcal{P}(Q) \rightarrow \mathcal{P}(\hat{Q})$$

of face posets of the quotients $Q = \mathcal{O}/T_{\mathcal{O}} = M/T_M$ and $\hat{Q} = \hat{O}/T_{\mathcal{O}} = \hat{P}/T_Q$, such that, for every face F of $\mathcal{P}(Q)$, F and $\varphi(F)$ have the same isotropy.

All of the above information is contained in the following diagram, where the label on each arrow denotes the quotient by the given torus, the dashed line indicates rational homotopy equivalence (not a map!), and the dotted line indicates that there is an isomorphism of face posets $\varphi : \mathcal{P}(Q) \rightarrow \mathcal{P}(\hat{Q})$.



It remains to show that the space \hat{M} is a manifold and that M is T_M -equivariantly rationally homotopy equivalent to \hat{M} equipped with the induced action of the torus $T_{\hat{M}} = T_Q/K_{\hat{P}}$.

Lemma 7.2. The group $K_{\hat{P}}$ acts freely on \hat{P} and, hence, the quotient $\hat{M} = \hat{P}/K_{\hat{P}}$ is a (topological) manifold. \square

Proof. Recall that \hat{P} is defined by $\hat{O} \times T_O / \sim$, where $(q, t) \sim (q', t')$ if and only if $q = q'$ and $tt'^{-1} \in T(q) = \prod_{F \ni q} S_F^1 \subseteq T_O$. The action of T_O on \hat{P} is given by left multiplication on the second factor. The action of $K_{\hat{P}}$ on \hat{P} is simply the restriction of the T_O action to $K_{\hat{P}}$ and, therefore, the isotropy subgroup of the $K_{\hat{P}}$ action at a point $[(q, t)] \in \hat{P}$ is given by $T(q) \cap K_{\hat{P}}$.

Let \hat{F}_q denote the face of \hat{O} of minimal dimension containing q , and $F_q = \varphi^{-1}(\hat{F}_q)$ the corresponding face in O , given by the face-poset isomorphism $\varphi : \mathcal{P}(O) \rightarrow \mathcal{P}(\hat{O})$. Since M is a manifold, $T(q)$ maps injectively via λ_M into the isotropy of T_M at a point $x \in M$ in the preimage of F_q . Thus $T(q) \cap \ker(\lambda_M) = T(q) \cap K_{\hat{P}}$ is trivial, as desired. \blacksquare

Lemma 7.3. The manifolds $\hat{P}/K_{\hat{P}}$ and $\hat{P}/K_{\hat{P}}^o$ are rationally homotopy equivalent. Therefore, in the following it may be assumed that $K_{\hat{P}}$ is connected and \hat{M} is simply connected. \square

Proof. It will be shown that the orbit map of the $\Gamma = K_{\hat{P}}/K_{\hat{P}}^o$ -action on $\hat{P}/K_{\hat{P}}^o$ induces a rational homotopy equivalence $\hat{P}/K_{\hat{P}}^o \rightarrow \hat{P}/K_{\hat{P}}$. The Γ action commutes with the $K_{\hat{M}}$ -action on $\hat{P}/K_{\hat{P}}^o$ and, therefore, induces a Γ action on the orbifold $\hat{O}' = (\hat{P}/K_{\hat{P}}^o)/K_{\hat{M}}$ with orbit space $\hat{O} = (\hat{P}/K_{\hat{P}})/K_{\hat{M}}$. Moreover, there is a commutative diagram

$$\begin{array}{ccc}
 K_{\hat{M}} & \longrightarrow & K_{\hat{M}}/(\Gamma \cap K_{\hat{M}}) \\
 \downarrow & & \downarrow \\
 \hat{P}/K_{\hat{P}}^o & \longrightarrow & \hat{P}/K_{\hat{P}} \\
 \downarrow & & \downarrow \\
 \hat{O}' & \longrightarrow & \hat{O}.
 \end{array}$$

Here the top and bottom maps are rational homotopy equivalences, since the Γ -actions on $K_{\hat{M}}$ and \hat{O}' induce trivial actions on cohomology and the spaces in the corners of the diagram are formal. Because a model for the spaces in the middle is given by the tensor product of the models for the corresponding top and bottom spaces, it follows that the map in the middle is a rational homotopy equivalence. Hence, it may be assumed that $K_{\hat{P}}$ is connected. \blacksquare

Observe now that, since the torus $K_{\hat{M}} = \hat{L}/K_{\hat{P}}$ acts almost freely on \hat{M} with $\hat{M}/K_{\hat{M}} = \hat{P}/\hat{L} = \hat{O}$, the projection $\hat{M} \rightarrow \hat{O}$ is, up to rational homotopy, a principal $K_{\hat{M}}$ -bundle.

The label map $\lambda_M : T_O \rightarrow T_M$ described above descends to an isomorphism $\underline{\lambda}_M : T_{\hat{M}} \rightarrow T_M$ with inverse $\mu_M : T_M \rightarrow T_{\hat{M}}$. Since $\lambda : T_O \rightarrow T_{\mathcal{O}}$ factors through λ_M , there is an induced map $\hat{\pi} : T_{\hat{M}} \rightarrow T_{\mathcal{O}}$ with kernel $K_{\hat{M}}$. Then the following diagram commutes

$$\begin{array}{ccc}
 T_M & \xrightarrow{\pi} & T_{\mathcal{O}} \\
 \mu_M \downarrow & & \downarrow = \\
 T_{\hat{M}} & \xrightarrow{\hat{\pi}} & T_{\mathcal{O}}
 \end{array} \tag{7.1}$$

where the vertical maps are isomorphisms. Moreover, there is an induced isomorphism $\mu_K : K_M \rightarrow K_{\hat{M}}$ given by the restriction of μ_M to K_M . Therefore, $M \rightarrow \mathcal{O}$ and $\hat{M} \rightarrow \hat{\mathcal{O}}$ can be thought of as rational homotopy principal K_M -bundles, and the goal is to show that M and \hat{M} are rationally homotopy equivalent.

Theorem 7.4. Let X, Y be rationally homotopy equivalent spaces, and let $\phi : H^2(Y; \mathbb{Q}) \rightarrow H^2(X; \mathbb{Q})$ be the isomorphism induced by a rational equivalence. Moreover, let T be a k -torus and let $\xi_X : E_X \rightarrow X, \xi_Y : E_Y \rightarrow Y$ be rational homotopy principal T -bundles with classifying maps $\rho_X : X \rightarrow BT, \rho_Y : Y \rightarrow BT$. Suppose, finally, that there is a map $\beta : H^2(BT; \mathbb{Q}) \rightarrow H^2(BT; \mathbb{Q})$ such that the diagram

$$\begin{array}{ccc}
 H^2(X; \mathbb{Q}) & \xleftarrow{\phi} & H^2(Y; \mathbb{Q}) \\
 \rho_X^* \uparrow & & \rho_Y^* \uparrow \\
 H^2(BT; \mathbb{Q}) & \xleftarrow{\beta} & H^2(BT; \mathbb{Q})
 \end{array} \tag{7.2}$$

commutes. Then E_X is rationally homotopy equivalent to E_Y . □

Proof. Let $(\wedge V_X, d_X)$ and $(\wedge V_Y, d_Y)$ be the minimal models of X and Y respectively. Let $\phi : (\wedge V_Y, d_Y) \rightarrow (\wedge V_X, d_X)$ be an isomorphism inducing $\phi : H^2(Y; \mathbb{Q}) \rightarrow H^2(X; \mathbb{Q})$.

The minimal model of T is $(\wedge(t_1, \dots, t_k), 0)$ with $|t_i| = 1$. The minimal model of BT is $\mathbb{Q}[\bar{t}_1, \dots, \bar{t}_k]$, where $|\bar{t}_i| = 2$. The t_i are mapped to \bar{t}_i via the isomorphism

$$\delta : W = \text{Hom}(\pi_1(T), \mathbb{Q}) \rightarrow \overline{W} = \text{Hom}(\pi_2(BT), \mathbb{Q})$$

induced by the long exact homotopy sequence of the fibration $T \rightarrow ET \rightarrow BT$. It's clear that $H^2(BT; \mathbb{Q})$ can now be identified with the vector space $\overline{W} = \text{span}_{\mathbb{Q}}\{\bar{t}_1, \dots, \bar{t}_k\}$. Using δ , the map $\beta : \overline{W} \rightarrow \overline{W}$ induces a map

$$\check{\beta} = \delta^{-1} \circ \beta \circ \delta : W \rightarrow W.$$

A model for E_X is $(\wedge V_X \otimes \wedge(t_1, \dots, t_k), D_X)$, where $D_X|_{\wedge V_X} = d_X$ and $D_X|_W = \rho_X^* \circ \delta$. Similarly, a model for E_Y is $(\wedge V_Y \otimes \wedge(t_1, \dots, t_k), D_Y)$, where $D_Y|_{\wedge V_Y} = d_Y$ and $D_Y|_W = \rho_Y^* \circ \delta$. Define now an isomorphism

$$\psi : (\wedge V_Y \otimes \wedge(t_1, \dots, t_k), D_Y) \longrightarrow (\wedge V_X \otimes \wedge(t_1, \dots, t_k), D_X)$$

by letting $\psi|_{\wedge V_Y} = \varphi$ and $\psi(1 \otimes t_i) = 1 \otimes \check{\beta}(t_i)$. It is clear that ψ preserves the grading and $(\psi \circ D_Y)|_{\wedge V_Y} = (D_X \circ \psi)|_{\wedge V_Y}$. Moreover, using the commutativity of diagram (7.2) (and Hurewicz),

$$\begin{aligned} \psi \circ D_Y|_W &= \psi \circ \rho_Y^* \circ \delta \\ &= \varphi \circ \rho_Y^* \circ \delta \\ &= \rho_X^* \circ \beta \circ \delta \\ &= \rho_X^* \circ \delta \circ \check{\beta} \\ &= D_X|_W \circ \check{\beta} \\ &= D_X \circ \psi|_W. \end{aligned}$$

Then ψ is an isomorphism between the models of E_X and E_Y . Consequently, there is an isomorphism between the corresponding minimal models and $E_X \simeq_{\mathbb{Q}} E_Y$. ■

It is now apparent that, in order to show that M and \hat{M} are rationally homotopy equivalent, it suffices to show that the hypotheses of Theorem 7.4 are satisfied by the rational homotopy principal K_M -bundles $M \rightarrow \mathcal{O}$ and $\hat{M} \rightarrow \hat{\mathcal{O}}$.

Proposition 7.5. The diagram

$$\begin{array}{ccc} H^2(\mathcal{O}; \mathbb{Q}) & \xleftarrow{f} & H^2(\hat{\mathcal{O}}; \mathbb{Q}) \\ \uparrow & & \uparrow \\ H^2(BK_M; \mathbb{Q}) & \xleftarrow{(B\mu_K)^*} & H^2(BK_{\hat{M}}; \mathbb{Q}) \end{array} \tag{7.3}$$

is commutative, where the vertical arrows are induced by the bundles $M \rightarrow \mathcal{O}$ and $\hat{M} \rightarrow \hat{\mathcal{O}}$, and f is the isomorphism given by $\mathcal{O} \simeq_{\mathbb{Q}} \hat{\mathcal{O}}$. □

As a first step towards proving Proposition 7.5, the following lemma is necessary.

Lemma 7.6. Let $M^{(1)} \subset M$, $\hat{M}^{(1)} \subset \hat{M}$ be the pre-images of the vertex-edge graphs of \mathcal{O} and $\hat{\mathcal{O}}$, respectively. Then there is a T_M -equivariant homeomorphism $\tilde{h} : M^{(1)} \rightarrow \hat{M}^{(1)}$.

Moreover, \tilde{h} induces a $T_{\mathcal{O}}$ -equivariant homeomorphism $h : \mathcal{O}^{(1)} \rightarrow \hat{\mathcal{O}}^{(1)}$ whose induced map in cohomology completes the commutative diagram

$$\begin{array}{ccc}
 H^2(\mathcal{O}; \mathbb{Q}) & \xleftarrow{f} & H^2(\hat{\mathcal{O}}; \mathbb{Q}) \\
 \downarrow i^* & & \downarrow \hat{i}^* \\
 H^2(\mathcal{O}^{(1)}; \mathbb{Q}) & \xleftarrow{h^*} & H^2(\hat{\mathcal{O}}^{(1)}; \mathbb{Q})
 \end{array}$$

where the vertical maps are induced by the respective inclusions. □

Proof. $M^{(1)}$ and $\hat{M}^{(1)}$ are unions of cohomogeneity-one manifolds N_{ij} and \hat{N}_{ij} , respectively. Each of these cohomogeneity-one manifolds is the pre-image of an edge in \mathcal{O} or $\hat{\mathcal{O}}$, respectively. Moreover, the indices run over the edges e_{ij} of the vertex-edge graph of \mathcal{O} .

Since the isotropy subgroups of the T_M -action on each N_{ij} and \hat{N}_{ij} are the same, there are equivariant homeomorphisms $N_{ij} \rightarrow \hat{N}_{ij}$. Since T_M is a compact, connected, abelian Lie group, these homeomorphisms can be chosen in such a way that they extend to an equivariant homeomorphism $\tilde{h} : M^{(1)} \rightarrow \hat{M}^{(1)}$.

Because $\mathcal{O}^{(1)} = M^{(1)}/K_M$ and $\hat{\mathcal{O}}^{(1)} = \hat{M}^{(1)}/K_M$, it follows that there is an $T_{\mathcal{O}}$ -equivariant homeomorphism $h : \mathcal{O}^{(1)} \rightarrow \hat{\mathcal{O}}^{(1)}$.

It remains to show that the induced map in cohomology completes a commutative diagram as in the statement of the lemma. Consider the diagram below, where the back (by equivariant rational homotopy equivalence), base and sides are each commutative. The goal is to show that the dotted arrow in the diagram below makes the top of the

cube into a commutative diagram.

$$\begin{array}{ccccc}
 H^2(\mathcal{O}; \mathbb{Q}) & \xleftarrow{f} & H^2(\hat{\mathcal{O}}; \mathbb{Q}) & & \\
 \uparrow & \searrow^{i^*} & \uparrow & \searrow^{\hat{i}^*} & \\
 & & H^2(\mathcal{O}^{(1)}; \mathbb{Q}) & \xleftarrow{h^*} & H^2(\hat{\mathcal{O}}^{(1)}; \mathbb{Q}) \\
 & & \uparrow & & \uparrow \\
 H_{T_{\mathcal{O}}}^2(\mathcal{O}; \mathbb{Q}) & \xleftarrow{\quad} & H_{T_{\mathcal{O}}}^2(\hat{\mathcal{O}}; \mathbb{Q}) & \xrightarrow{\hat{i}^*} & H_{T_{\mathcal{O}}}^2(\hat{\mathcal{O}}^{(1)}; \mathbb{Q}) \\
 & \searrow^{i^*} & & \searrow^{i^*} & \\
 & & H_{T_{\mathcal{O}}}^2(\mathcal{O}^{(1)}; \mathbb{Q}) & \xleftarrow{\quad} & H_{T_{\mathcal{O}}}^2(\hat{\mathcal{O}}^{(1)}; \mathbb{Q})
 \end{array}$$

Here the bottom maps are induced by functoriality from the isomorphism of face posets $\varphi : \mathcal{P}(\mathcal{O}) \rightarrow \mathcal{P}(\hat{\mathcal{O}})$, see Remark 4.3.

By a diagram chase, one readily sees that it suffices to show both that the vertical map $H_{T_{\mathcal{O}}}^2(\mathcal{O}; \mathbb{Q}) \rightarrow H^2(\mathcal{O}; \mathbb{Q})$ is surjective and that the front of the cube is commutative.

Since $H^1(\mathcal{O}, \mathbb{Q}) = 0$ the natural map $H_{T_{\mathcal{O}}}^2(\mathcal{O}; \mathbb{Q}) \rightarrow H^2(\mathcal{O}; \mathbb{Q})$ is surjective.

By using an inductive Mayer–Vietoris sequence argument, one sees that $H^2(\mathcal{O}^{(1)}; \mathbb{Q})$ is generated by the duals α_{ij} of the fundamental classes $[\mathbb{S}_{ij}^2] \in H_2(\mathcal{O}^{(1)}; \mathbb{Q})$. Similarly, $H^2(\hat{\mathcal{O}}^{(1)}; \mathbb{Q})$ is generated by the duals $\hat{\alpha}_{ij}$ of the classes $[\hat{\mathbb{S}}_{ij}^2] \in H_2(\hat{\mathcal{O}}^{(1)}; \mathbb{Q})$, and $h^*(\hat{\alpha}_{ij}) = \alpha_{ij}$.

On the other hand, by using the Mayer–Vietoris sequence on the Borel construction $(\mathbb{S}_{ij}^2)_T$ (using the decomposition $(\mathbb{S}_{ij}^2)_T = U \cup V$, with $U = (\mathbb{S}_{ij}^2 \setminus p_i)_T$ and $V = (\mathbb{S}_{ij}^2 \setminus p_j)_T$)

$$H_T^2(\mathbb{S}_{ij}^2; \mathbb{Z}) = \{(g_i, g_j) \in H_T^2(p_i; \mathbb{Z}) \oplus H_T^2(p_j; \mathbb{Z}) \mid g_i - g_j \in \mathbb{Z} \cdot \mu(e_{ij})\},$$

where $\mu(e_{ij})$ is the weight of e_{ij} in the GKM-graph associated to \mathcal{O} . From the Serre spectral sequence of the fibration $\mathbb{S}_{ij}^2 \rightarrow (\mathbb{S}_{ij}^2)_T \rightarrow BT$, which degenerates in the E_2 -page, it follows that the map $H_T^2(\mathbb{S}_{ij}^2; \mathbb{Z}) \rightarrow H^2(\mathbb{S}_{ij}^2; \mathbb{Z})$ is surjective, and sends $(\mu(e_{ij}), 0)$ to α_{ij} . The same discussion carries over identically to the spheres $\hat{\mathbb{S}}_{ij}^2$. The map $h^* : H^2(\hat{\mathcal{O}}^{(1)}; \mathbb{Q}) \rightarrow H^2(\mathcal{O}^{(1)}; \mathbb{Q})$ can now be factored as

$$\begin{array}{ccccc}
 H^2(\hat{\mathcal{O}}^{(1)}; \mathbb{Q}) & \longrightarrow & H_{T_{\mathcal{O}}}^2(\hat{\mathcal{O}}^{(1)}; \mathbb{Q}) & \longrightarrow & H_{T_{\mathcal{O}}}^2(\mathcal{O}^{(1)}; \mathbb{Q}) \longrightarrow H^2(\mathcal{O}^{(1)}; \mathbb{Q}) \\
 \hat{\alpha}_{ij} \longmapsto & & (\hat{\mu}(\hat{e}_{ij}), 0) & \longmapsto & (\mu(e_{ij}), 0) \longmapsto \alpha_{ij}
 \end{array}$$

and, therefore, there is a commutative diagram, as desired. ■

Lemma 7.7. The inclusion map $i : \mathcal{O}^{(1)} \rightarrow \mathcal{O}$ induces an injection $i^* : H^2(\mathcal{O}; \mathbb{Q}) \rightarrow H^2(\mathcal{O}^{(1)}; \mathbb{Q})$. □

Proof. Recall that there is a map of fibrations

$$\begin{array}{ccccc}
 \mathcal{O}^{(1)} & \longrightarrow & \mathcal{O}_{T\mathcal{O}}^{(1)} & \longrightarrow & BT_{\mathcal{O}} \\
 i \downarrow & & i^* \downarrow & & \parallel \\
 \mathcal{O} & \longrightarrow & \mathcal{O}_{T\mathcal{O}} & \longrightarrow & BT_{\mathcal{O}}
 \end{array} \tag{7.4}$$

which induces a map between the corresponding Serre spectral sequences with rational coefficients. Both spectral sequences have the property that $E_2^{1j} = E_2^{3j} = 0$ for all $j \geq 0$. Therefore, for $X = \mathcal{O}^{(1)}$ or $X = \mathcal{O}$ there are exact sequences

$$0 \rightarrow E_{\infty}^{2,0} \rightarrow H^2(X_{T\mathcal{O}}; \mathbb{Q}) \rightarrow E_{\infty}^{0,2} \rightarrow 0.$$

The natural map $H^2(BT_{\mathcal{O}}; \mathbb{Q}) \rightarrow H^2(X_{T\mathcal{O}}; \mathbb{Q})$ is injective, since there are $T_{\mathcal{O}}$ -fixed points in X . It then follows that $E_{\infty}^{2,0} = H^2(BT_{\mathcal{O}}; \mathbb{Q})$. Moreover, $E_{\infty}^{0,2} \subseteq H^2(X; \mathbb{Q})$, with equality holding if $b_1(X) = 0$. This last condition holds if $X = \mathcal{O}$

The map i^* , in particular, induces a row-exact, commutative diagram

$$\begin{array}{ccccccc}
 0 & \longrightarrow & H^2(BT_{\mathcal{O}}; \mathbb{Q}) & \longrightarrow & H^2(\mathcal{O}_{T\mathcal{O}}^{(1)}; \mathbb{Q}) & \longrightarrow & H^2(\mathcal{O}^{(1)}; \mathbb{Q}) \\
 & & \parallel & & (i_{T\mathcal{O}})^* \uparrow & & i^* \uparrow \\
 0 & \longrightarrow & H^2(BT_{\mathcal{O}}; \mathbb{Q}) & \longrightarrow & H^2(\mathcal{O}_{T\mathcal{O}}; \mathbb{Q}) & \longrightarrow & H^2(\mathcal{O}; \mathbb{Q}) \longrightarrow 0.
 \end{array}$$

By [5, Proof of Prop. 2.4], $(i_{T\mathcal{O}})^*$ is injective and, by diagram chasing it follows that $i^* : H^2(\mathcal{O}; \mathbb{Q}) \rightarrow H^2(\mathcal{O}^{(1)}; \mathbb{Q})$ is injective as well. ■

Proof of Proposition 7.5. Diagram (7.3) is part of the larger diagram

$$\begin{array}{ccc}
 H^2(\mathcal{O}^{(1)}; \mathbb{Q}) & \xleftarrow{h^*} & H^2(\hat{\mathcal{O}}^{(1)}; \mathbb{Q}) \\
 i^* \uparrow & & \hat{i}^* \uparrow \\
 H^2(\mathcal{O}; \mathbb{Q}) & \xleftarrow{f} & H^2(\hat{\mathcal{O}}; \mathbb{Q}) \\
 \uparrow & & \uparrow \\
 H^2(BK_M; \mathbb{Q}) & \xleftarrow{(B\mu_K)^*} & H^2(BK_{\hat{M}}; \mathbb{Q})
 \end{array} \tag{7.5}$$

where, $i : \mathcal{O}^{(1)} \rightarrow \mathcal{O}$, $\hat{i} : \hat{\mathcal{O}}^{(1)} \rightarrow \hat{\mathcal{O}}$ denote the inclusions. The upper square comes from Lemma 7.6 and hence commutes. By Lemma 7.7, the map i^* is injective. In order to prove the proposition it now suffices to show that the outer square commutes.

By Lemma 7.6, the map $h : \mathcal{O}^{(1)} \rightarrow \hat{\mathcal{O}}^{(1)}$ can be lifted to a μ_K -equivariant map $\tilde{h} : M^{(1)} \rightarrow \hat{M}^{(1)}$ such that the diagram

$$\begin{array}{ccc}
 K_M & \xrightarrow{\mu_K} & K_{\hat{M}} \\
 \downarrow & & \downarrow \\
 M^{(1)} & \xrightarrow{\tilde{h}} & \hat{M}^{(1)} \\
 \downarrow & & \downarrow \\
 \mathcal{O}^{(1)} & \xrightarrow{h} & \hat{\mathcal{O}}^{(1)}
 \end{array}$$

is a pull-back diagram between the (rational homotopy) principal torus bundles $M^{(1)} \rightarrow \mathcal{O}^{(1)}$ and $\hat{M}^{(1)} \rightarrow \hat{\mathcal{O}}^{(1)}$. This induces a (rational homotopy) commutative diagram

$$\begin{array}{ccc}
 \mathcal{O}^{(1)} & \xrightarrow{h} & \hat{\mathcal{O}}^{(1)} \\
 \downarrow & & \downarrow \\
 BK_M & \xrightarrow{B\mu_K} & BK_{\hat{M}}
 \end{array}$$

from which the commutativity of the outer square in diagram (7.5) follows. ■

Theorem 7.8. The manifolds M and \hat{M} are T_M -equivariantly rationally homotopy equivalent. □

Proof. Recall that $M \rightarrow \mathcal{O}$ (respectively, $\hat{M} \rightarrow \hat{\mathcal{O}}$) is a rational principal K_M -bundle (respectively, $K_{\hat{M}}$ -bundle), with an isomorphism $\mu_K : K_M \rightarrow K_{\hat{M}}$. With respect to the induced identification $(B\mu_K)^* : H^*(BK_{\hat{M}}; \mathbb{Q}) \rightarrow H^*(BK_M; \mathbb{Q})$, Proposition 7.5 yields a commutative diagram

$$\begin{array}{ccc}
 H^2(\mathcal{O}; \mathbb{Q}) & \xleftarrow{f} & H^2(\hat{\mathcal{O}}; \mathbb{Q}) \\
 \uparrow & & \uparrow \\
 H^2(BK_M; \mathbb{Q}) & \xlongequal{\quad} & H^2(BK_M; \mathbb{Q})
 \end{array}$$

where f is the isomorphism induced by the rational homotopy equivalence $\mathcal{O} \simeq_{\mathbb{Q}} \hat{\mathcal{O}}$ and the vertical arrows are induced by the rational principal K_M -bundles. By Theorem 7.4, this implies that the total spaces M and \hat{M} are rationally homotopy equivalent.

Consider now the diagram

$$\begin{array}{ccc}
 H^*(M; \mathbb{Q}) & \longleftarrow & H^*(\hat{M}; \mathbb{Q}) \\
 \uparrow & & \uparrow \\
 H^*(\mathcal{O}; \mathbb{Q}) & \xleftarrow{f} & H^*(\hat{\mathcal{O}}; \mathbb{Q}) \\
 \uparrow & & \uparrow \\
 H_{T_{\mathcal{O}}}^*(\mathcal{O}; \mathbb{Q}) & \longleftarrow & H_{T_{\hat{\mathcal{O}}}}^*(\hat{\mathcal{O}}; \mathbb{Q})
 \end{array}$$

where the uppermost map is the isomorphism induced by the rational homotopy equivalence $M \simeq_{\mathbb{Q}} \hat{M}$ constructed in Theorem 7.4. From that construction, it is clear that the upper square commutes. On the other hand, the lower square commutes because of the equivariance of the rational homotopy equivalence $\mathcal{O} \simeq_{\mathbb{Q}} \hat{\mathcal{O}}$. Since $\mathcal{O}/T_{\mathcal{O}} = M/T_M$ and $\hat{\mathcal{O}}/T_{\hat{\mathcal{O}}} = \hat{M}/T_{\hat{M}}$, it follows from the commutativity of (7.1) that M and \hat{M} are T_M -equivariantly rationally homotopy equivalent.

Indeed, since K_M acts almost freely on M with orbit space \mathcal{O} , there is an isomorphism

$$H_{T_M}^*(M; \mathbb{Q}) \cong H_{T_{\mathcal{O}}}^*(\mathcal{O}; \mathbb{Q}).$$

and similarly for \hat{M} and $\hat{\mathcal{O}}$. Therefore, since the above diagram commutes, there is a commutative diagram

$$\begin{array}{ccc}
 H^*(M; \mathbb{Q}) & \longleftarrow & H^*(\hat{M}; \mathbb{Q}) \\
 \uparrow & & \uparrow \\
 H_{T_M}^*(M; \mathbb{Q}) & \longleftarrow & H_{T_M}^*(\hat{M}; \mathbb{Q})
 \end{array}$$

as desired. ■

8 Torus Actions in Non-Negative Curvature

To begin this section, a version of Theorem A for non-negatively curved torus orbifolds of dimension at most six will be established.

Theorem 8.1. Let (\mathcal{O}, G) be a non-negatively curved and simply connected torus orbifold of dimension at most six such that $H^{\text{odd}}(\mathcal{O}; \mathbb{Q}) = 0$. Then there is a product \hat{P} of spheres of dimension ≥ 3 , a torus \hat{L} acting linearly and almost freely on \hat{P} , a linear action of G on $\hat{\mathcal{O}} = \hat{P}/\hat{L}$ and a G -equivariant rational homotopy equivalence $\mathcal{O} \simeq_{\mathbb{Q}} \hat{\mathcal{O}}$. \square

To prove this theorem, it suffices to show that \mathcal{O}/G satisfies all the properties listed in Proposition 5.1. All of these, except for Property (f), can be proved as in the rationally elliptic case.

Note that, as \mathcal{O}/G is being viewed as a face of itself of codimension zero, in order to prove Theorem 8.1, Property (f) needs to be discussed in dimension six. Since the rational cohomology of \mathcal{O} is concentrated in even degrees, it follows that all faces of \mathcal{O}/G are acyclic over the rationals [3, Corollary 3]. Hence, the following lemma implies that Property (f) holds for \mathcal{O}/G .

Lemma 8.2. There is no simply connected, six-dimensional torus orbifold (\mathcal{O}, G) such that each face of \mathcal{O}/G is acyclic over the rationals, each facet of \mathcal{O}/G is combinatorially equivalent to a square, and the intersection of any two facets has two components. \square

Proof. Assume that there is a torus orbifold whose orbit space contradicts the conclusion of the lemma. First note that all two-dimensional orbifolds are homeomorphic to two-dimensional topological manifolds. Therefore, since the facets of \mathcal{O}/G are acyclic over the rationals and orientable, they are all homeomorphic to two-dimensional discs. Hence, with the same argument as in the proof of Lemma 4.4 of [37], one sees that the boundary of \mathcal{O}/G is homeomorphic to $\mathbb{R}P^2$. However, \mathcal{O}/G is an orientable orbifold with boundary, while $\mathbb{R}P^2$ is non-orientable, yielding a contradiction, as desired. \blacksquare

Proof of Theorem 8.1. Since $H^{\text{odd}}(\mathcal{O}; \mathbb{Q}) = 0$ and \mathcal{O} admits an invariant metric with non-negative sectional curvature, the conclusion of Proposition 5.1 holds for \mathcal{O}/G as discussed above. Therefore the same arguments as in the proof of Theorem A can be carried out to prove Theorem 8.1. \blacksquare

To conclude the article, a proof of Theorem D is provided, that is, the Bott Conjecture in the presence of an isometric, slice-maximal torus action is verified. This is a generalisation of Theorem 1.2 of [34].

Proof of Theorem D. Let T denote the torus whose action on M is slice maximal. It is sufficient to show that M is *rationally Ω -elliptic*, that is, the pointed loop space ΩM

of M satisfies $\sum_r \dim(\pi_r(\Omega M) \otimes \mathbb{Q}) < \infty$, since, M being simply connected, this property implies that M is rationally elliptic. The proof will proceed by induction on the dimension $d = \dim(M/T)$ and no longer assumes that M is simply connected.

When $d = 0$, M consists of one orbit and is, therefore, a torus, hence rationally Ω -elliptic. Suppose now that every non-negatively curved, closed manifold admitting an isometric, slice-maximal torus action with quotient of dimension $d - 1$ is rationally Ω -elliptic.

From the introduction, the action of T on M being slice maximal ensures that, at every point on a (fixed) minimal orbit, the normal slice is even dimensional and the identity component G of the isotropy subgroup acts on it with maximal rank, that is, the action is equivalent to the standard linear, effective action of G on $\mathbb{C}^{\dim(G)}$. Hence, one can find a circle subgroup $S \subseteq G \subseteq T$ such that some component M' of its fixed-point set M^S is of codimension two and contains the minimal orbit. Consequently, the induced action of $T' = T/S$ on M' is slice-maximal. Moreover, since M' is totally geodesic, hence non-negatively curved, and $\dim(M'/T') = d - 1$, the induction hypothesis yields that M' is rationally Ω -elliptic.

As the action of S on M has a fixed-point component of codimension 2, meaning that it is fixed-point homogeneous, by Theorem 4.1 of [34] there exists a submanifold $N \subseteq M$ such that M is diffeomorphic to the union of the normal disc bundles $D(M')$ and $D(N)$ of M' and N along their common boundary E :

$$M = D(M') \cup_E D(N).$$

The foot-point projection $D(M') \rightarrow M'$ induces an S^1 -bundle $E \rightarrow M'$. Since M' is rationally Ω -elliptic, it follows from the homotopy long exact sequence that E is also rationally Ω -elliptic. Moreover, by Theorem D of [12], the homotopy fibre F of the inclusion $\iota : E \hookrightarrow M$ is rationally Ω -elliptic. Therefore, from the homotopy long exact sequence for ι and the fact that E is rationally Ω -elliptic, it follows that M is rationally Ω -elliptic as well, as desired. ■

Funding

F.G.-G., M.K., and M.R. received support from SFB 878: *Groups, Geometry & Actions* at WWU Münster. M.W. was partially supported by DFG Grant HA 3160/6-1.

Acknowledgements

We would like to thank Matthias Franz for helpful comments on a previous version of this article, and Wolfgang Spindeler for discussions regarding his work in [34].

References

- [1] Allday, C. and V. Puppe. *Cohomological Methods in Transformation Groups*. Cambridge Studies in Advanced Mathematics 32. Cambridge: Cambridge University Press, 1993.
- [2] Bredon, G. E. *Introduction to Compact Transformation Groups*. Pure and Applied Mathematics, vol. 46. Academic Press, New York, London, 1972.
- [3] Bredon, G. E. "The free part of a torus action and related numerical equalities." *Duke Mathematical Journal* 41 (1974): 843–54.
- [4] Buchstaber, V. and T. Panov. *Torus Actions and Their Applications in Topology and Combinatorics*. University Lecture Series 24. Providence, RI: American Mathematical Society, 2002.
- [5] Chang, T. and T. Skjelbred. "The topological Schur lemma and related results." *Annals of Mathematics* 100, no. 2 (1974): 307–21.
- [6] Escher, C. and C. Searle. "Non-negative curvature and torus actions." preprint 2015, arXiv:1506.08685.
- [7] Félix, Y., S. Halperin, and Thomas, J.-C. *Rational Homotopy Theory*. Graduate Texts in Mathematics 205. New York: Springer, 2001.
- [8] Félix, Y., J. Oprea, and Tanré, D. *Algebraic Models in Geometry*. Oxford Graduate Texts in Mathematics 17. Oxford: Oxford University Press, 2008.
- [9] Fulton, W. *Introduction to Toric Varieties*. Annals of Mathematics Studies 131. The William H. Roever Lectures in Geometry. Princeton, NJ: Princeton University Press, 1993.
- [10] Galaz-García, F., M. Kerin, and M. Radeschi. "Torus actions on rationally elliptic manifolds." preprint 2015, arXiv:1511.08383.
- [11] Goresky, M., R. Kottwitz, and R. MacPherson. "Equivariant cohomology, Koszul duality, and the localization theorem." *Inventiones Mathematicae* 131, no. 1 (1998): 25–83.
- [12] Grove, K. and S. Halperin "Dupin hypersurfaces, group actions and the double mapping cylinder." *Journal of Differential Geometry* 26 (1987): 429–59.
- [13] Grove, K. and C. Searle. "Positively curved manifolds with maximal symmetry-rank." *Journal of Pure and Applied Algebra* 91, no. 1–3 (1994): 137–42.
- [14] Grove, K. and W. Ziller. "Polar manifolds and actions." *Journal of Fixed Point Theory and Applications* 11, no. 2 (2012): 279–313.
- [15] Haefliger, A. and É. Salem. "Riemannian foliations on simply connected manifolds and actions of tori on orbifolds." *Illinois Journal of Mathematics* 34, no. 4 (1990): 706–30.
- [16] Haefliger, A. and É. Salem. "Actions of tori on orbifolds." *Annals of Global Analysis and Geometry* 9, no. 1 (1991): 37–59.
- [17] Hattori, A. "Elliptic genera, torus orbifolds and multi-fans. II." *International Journal of Mathematics* 17, no. 6 (2006): 707–35.
- [18] Hattori, A. and M. Masuda. "Theory of multi-fans." *Osaka Journal of Mathematics* 40, no. 1 (2003): 1–68.
- [19] Hattori, A. and M. Masuda. "Elliptic genera, torus orbifolds and multi-fans." *International Journal of Mathematics* 16, no. 9 (2005): 957–98.

- [20] Holm, T. S. and T. Matsumura. "Equivariant cohomology for Hamiltonian torus actions on symplectic orbifolds." *Transformation Groups* 17, no. 3 (2012): 717–46.
- [21] Ishida, H. "Complex manifolds with maximal torus actions." *J. Reine Angew. Math.*, doi: <https://doi.org/10.1515/crelle-2016-0023>, July 2016.
- [22] Kleiner, B. "Riemannian four-manifolds with nonnegative curvature and continuous symmetry." PhD Thesis, U.C. Berkeley, 1989.
- [23] Kleiner, B. and J. Lott. "Geometrization of Three-Dimensional Orbifolds via Ricci Flow." *Astérisque* 365 (2014): 101–77.
- [24] Kobayashi, S. "Fixed points of isometries." *Nagoya Mathematical Journal* 13 (1958): 63–8.
- [25] Lerman, E. and S. Tolman. "Hamiltonian torus actions on symplectic orbifolds and toric varieties." *Transactions of the American Mathematical Society* 349, no. 10 (1997): 4201–30.
- [26] Lupton, G. "Variations on a conjecture of Halperin." In *Homotopy and Geometry (Warsaw, 1997)*, edited by J. Oprea and A. Tralle, 115–35. Banach Center Publ. 45. Warsaw: Polish Acad. Sci., 1998.
- [27] Lytchak, A. "Geometric resolution of singular Riemannian foliations." *Geometriae Dedicata* 149 (2010): 379–95.
- [28] Maeda, H., M. Masuda, and T. Panov. "Torus graphs and simplicial posets." *Advances in Mathematics* 212 (2007): 458–83.
- [29] Masuda, M. and T. Panov. "On the cohomology of torus manifolds." *Osaka Journal of Mathematics* 43, no. 3 (2006): 711–46.
- [30] Orlik, P. and F. Raymond. "Actions of the torus on 4-manifolds. I." *Transactions of the American Mathematical Society* 152 (1970): 531–59.
- [31] Orlik, P. and F. Raymond. "Actions of the torus on 4-manifolds. II." *Topology* 13 (1974): 89–112.
- [32] Palais, R. S. and C.-L. Terng. "A general theory of canonical forms." *Transactions of the American Mathematical Society* 300, no. 2 (1987): 771–89.
- [33] Poddar, M. and S. Sarkar. "On quasitoric orbifolds." *Osaka Journal of Mathematics* 47, no. 4 (2010): 1055–76.
- [34] Spindeler, W. "Nonnegatively curved quotient spaces with boundary." preprint 2015, arXiv:1510.01908.
- [35] Thurston, W. *The Geometry and Topology of 3-Manifolds*. Lecture Notes. Princeton, NJ: Princeton University Mathematics Department, 1978.
- [36] Ustinovsky, Yu. M. "Geometry of compact complex manifolds with maximal torus action." *Trudy Matematicheskogo Instituta imeni V. A. Steklova* 286 (2014): 219–30. Translation in *Proceedings of the Steklov Institute of Mathematics* 286 (2014): 198–208.
- [37] Wiemeler, M. "Torus manifolds and non-negative curvature." *Journal of the London Mathematical Society* (2) 91, no. 3 (2015): 667–92.
- [38] Yeroshkin, D. "Riemannian orbifolds with non-negative curvature." PhD Thesis, University of Pennsylvania, 2014.