

AperTO - Archivio Istituzionale Open Access dell'Università di Torino

Primary aldosteronism in pregnancy

This is a pre print version of the following article:

Original Citation:

Availability:

This version is available <http://hdl.handle.net/2318/1889332> since 2023-02-01T18:24:51Z

Published version:

DOI:10.1007/s11154-022-09729-6

Terms of use:

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)

PA SUSPECTED DURING PREGNANCY
(high BP +/- hypokaliemia)



Screening Test (ARR)

PA excluded



Consider re-screening in the post-partum in case of persistent hypertension

PA confirmed (high PAC + low PRA/DRC)



Postpone confirmatory test and subtype differentiation
Start pregnancy approved anti-hypertensive drugs +/- potassium

NO change

BP controlled

BP NOT controlled or persistent hypokaliemia



MRI-scan

Bilateral alterations or no alterations

Unilateral adenoma

Consider MRA (eplerenone) or amiloride

Consider MRA (eplerenone) or amiloride
OR
Adrenalectomy (preferably 2° trimester)