



High-resolution temporal variations of nitrate in a high-elevation pond in alpine tundra (NW Italian Alps)

N. Colombo^{a,b,c}, R. Balestrini^d, D. Godone^{e,c,*}, D. Vione^{f,c}, D. Said-Pullicino^a, G. Viviano^d, M. Martin^a, C.A. Delconte^d, S. Fratianni^{g,c}, A.G. Capodaglio^h, E. Pintaldi^a, M. Freppaz^{a,c}, F. Salerno^{d,i}

^a University of Turin, Department of Agricultural, Forest and Food Sciences, Grugliasco, Italy

^b CNR-IRSA (National Research Council - Water Research Institute), Rome, Italy

^c Research Center on Natural Risk in Mountain and Hilly Environments, NatRisk, University of Turin, Grugliasco, Italy

^d CNR-IRSA (National Research Council - Water Research Institute), Brughiero, Italy

^e CNR-IRPI (National Research Council - Research Institute for Geo-Hydrological Protection), Turin, Italy

^f University of Turin, Department of Chemistry, Turin, Italy

^g University of Turin, Department of Earth Sciences, Turin, Italy

^h University of Pavia, Department of Civil Engineering and Architecture, Pavia, Italy

ⁱ CNR-ISP (National Research Council - Institute of Polar Sciences), Venezia Mestre, VE, Italy

ARTICLE INFO

Keywords:

NO₃⁻
Surface water
Mountains
LTER
Turbidity
Nitrogen retention

ABSTRACT

High-resolution temporal measurements in remote, high-elevation surface waters are required to better understand the dynamics of nitrate (NO₃⁻) in response to changes in meteorological conditions. This study reports on the first use of a UV-Vis submersible spectrophotometric probe (UV-Vis probe) to measure the hourly concentration of nitrate nitrogen (NO₃⁻-N) in a pond located at 2722 m a.s.l. in an alpine tundra area (NW Italian Alps), during two snow-free seasons (July–October) in 2014 and 2015. Weekly analyses of NO₃⁻-N and stable isotopes of water (δ¹⁸O and δ²H), together with continuous meteorological, water temperature, and turbidity measurements, were performed over the same period. The integration of in-situ UV-Vis spectrophotometric measurements with weekly samples allowed depicting the role of summer precipitation, snow melt, and temperature (air and water) in influencing NO₃⁻ dynamics. Short-duration meteorological events (e.g., summer storms and rain-on-snow events) produced rapid variations of in-pond NO₃⁻ concentration, i.e., fivefold increase in 18 h, that would not be detectable using the traditional manual collection of discrete samples. The observed seasonal variability of NO₃⁻ concentration, negatively correlated with water temperature, highlighted the important role of in-pond biological processes leading to an enhanced N uptake and to the lowest NO₃⁻ concentration in the warmer periods. The occurrence of heavy rainfall events critically altered the expected seasonal NO₃⁻ trends, increasing the N supply to the pond. The comparison of N dynamics in two years characterised by extremely different meteorological conditions allowed us to obtain insights on the potential effects of climate changes (e.g., high air temperature, heavy rainfalls, and rain-on-snow events) on sensitive aquatic ecosystems as high-elevation ponds.

1. Introduction

Changes in climate and nutrient input can have strong effects on mountain ecosystems, in particular on those located above the treeline, in alpine tundra (Balestrini et al., 2013). These ecosystems are susceptible to alterations that affect their physical structure and biological communities, because of their complex topography, harsh climate, long-lasting snow cover, and a short growing season (Williams et al., 2002;

Balestrini et al., 2013; Barnes et al., 2014). These extreme environments are key components of the water cycle and are integral parts of the so called “water towers”, referring to the role of mountains as providers of essential freshwater to lowland areas (Viviroli et al., 2007, Viviroli et al., 2020).

In this context, high-elevation lakes play an important role in the hydrological and chemical dynamics of mountain watersheds (Catalan et al., 2006; Tartari et al., 2008; Tolotti et al., 2009; Salerno et al.,

* Corresponding author at: CNR-IRPI (National Research Council - Research Institute for Geo-Hydrological Protection), Turin, Italy.

E-mail address: daniilo.godone@irpi.cnr.it (D. Godone).

<https://doi.org/10.1016/j.catena.2023.107635>

Received 1 June 2023; Received in revised form 10 October 2023; Accepted 24 October 2023

Available online 5 November 2023

0341-8162/© 2023 The Authors. Published by Elsevier B.V. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).

2016a). The general lack of direct human influence on these lakes and the fact that their physical, chemical, and biological properties respond rapidly to climate-related changes make them key freshwater reference sites for global scale processes (Adrian et al., 2009; Mladenov et al., 2009; Salerno et al., 2016b; Rogora et al., 2020). In addition, water bodies located in high-elevation environments are generally characterised by small size, and can be defined as ponds (area < 2×10^4 m²; Hamerlík et al., 2014). The relatively low water volumes and high surface area to depth ratios make these water bodies even more fragile and sensitive to environmental changes (Hamerlík et al., 2014).

Chronic, relatively high, inputs of nitrogen (N) from atmospheric deposition can, over time, saturate the N assimilation capacity of biological processes in remote and commonly N-limited ecosystems (N saturation, see Aber et al., 1998). The effects of N-saturation are complex with impact on both terrestrial and aquatic ecosystems, and include the enhanced leaching of inorganic N from soils to surface waters (Balestrini et al., 2006; Rogora et al., 2012). In this regard, nitrate (NO₃) has been investigated due to its role in affecting the productivity and species diversity in remote mountain waters (Elser et al., 2009; Slemmons et al., 2017). In mountain catchments, NO₃ dynamics in surface waters depend on several drivers, such as land-cover and topographic characteristics (slope and presence of soil, bedrock, and cryospheric elements; e.g., Kopáček et al., 2005; Balestrini et al., 2013; Colombo et al., 2019a), climatic conditions (e.g., precipitation, snow-cover duration, and air temperature; e.g., Kopáček et al., 2005; Williams et al., 2015a,b; Freppaz et al., 2019), and anthropogenic activities (e.g., N deposition from atmospheric pollution; Elser et al., 2009; Rogora et al., 2012). Meteo-hydrological events such as snowmelt and summer storms, able to generate rapid changes in water flow paths, nutrient source areas, and biogeochemical processes, can strongly affect NO₃ concentrations and fluxes in mountain surface waters (Williams et al., 2002; Clow et al., 2003; Sickman et al., 2003; Williams et al., 2007; Sebestyen et al., 2008). However, little work has been done to characterise the hydrochemical response, with particular reference to NO₃, to meteorological events in ponds and lakes located above the treeline. Furthermore, the biological in-stream/lake processes are often ignored in alpine catchments, although a growing body of literature demonstrates that they are important regulators of nutrient retention and export in headwater catchments (Mulholland et al., 2006; Roberts & Mulholland, 2007; Rusjan & Mikos, 2010).

Studies focusing on NO₃ dynamics in remote surface waters generally rely on manual collection of discrete samples and subsequent laboratory analysis (e.g., Williams et al., 2007; Vione et al., 2021). This approach is expensive, time-consuming, and is often affected by intrinsic risks associated with extreme weather conditions and location (e.g., orographic thunderstorms, rockfalls, snow avalanches, ice falls, etc.), resulting in sporadic and/or low temporal-resolution data sets. For this reason, essential insights into the processes occurring in these ecosystems are lacking or incomplete, especially the ones occurring on short time scales (hours, days, or even weeks).

High-resolution in-situ measurements could refine the assessment of nutrient fluxes and temporal dynamics of high-elevation aquatic ecosystems, in turn improving the understanding of how weather- and climate-driven modifications could impact these fragile and rapidly changing ecosystems. To do this, in situ sensors can be deployed in remote and hardly accessible locations where repeated grab sampling operations would be logistically difficult and potentially dangerous (e.g., Beaton et al., 2017). Ultraviolet-visible light (UV-Vis) spectrophotometers are currently available to evaluate variations in NO₃ concentrations in surface waters with high temporal resolution. Research has focused on the use of similar sensors, especially in running waters in lowland areas (e.g., Pellerin et al., 2012; Burns et al., 2019); however, applications of in-situ UV-Vis spectrophotometers in remote, high-elevation water bodies have not yet been reported.

In the present work, a UV-Vis spectrophotometer was installed and used to monitor nitrate nitrogen (NO₃-N) concentration and turbidity in

the Col d'Olen Rock Glacier Pond (2722 m a.s.l.) located in the NW Italian Alps, during the summers 2014 and 2015. In addition, grab samples were collected on a weekly basis and analysed for NO₃-N concentration and isotopic composition of water ($\delta^{18}\text{O}$ and $\delta^2\text{H}$). Finally, meteorological parameters (air temperature, rainfall, and snow depth) and water temperature were continuously measured during the investigated period. The Col d'Olen Rock Glacier Pond was selected as a model system for this investigation since the hydrochemical features of the pond have been deeply studied in the last years to understand the influence of a rock glacier flowing into the pond (Colombo et al., 2018a, Colombo et al., 2018b; Colombo et al., 2019a; Colombo et al., 2020). This provided a solid knowledge baseline for interpreting the high-temporal-resolution data obtained in this work.

The aim of this study was to unravel the mechanisms underlying the dynamics of nitrate during seasonal transitions and short-lived meteorological events in a remote, high-elevation pond. The implications of our findings on the effect of climate change on N cycle were also discussed.

2. Materials and methods

2.1. Study area

The research site (Angelo Mosso Scientific Institute site) is a node of the Long-Term Ecological Research (LTER) network in Italy, situated in the North-Western Italian Alps, at the boundary between Valle d'Aosta and Piemonte regions (Fig. 1a). The Col d'Olen Rock Glacier Pond is situated at the Col d'Olen Rock Glacier terminus (Fig. 1b,c), at an elevation of 2722 m a.s.l. Its catchment area is approximately 206,000 m² (Fig. 1b). The pond has an area of ca. 1600 m² (covering 0.8 % of the catchment), with maximum length and width of ca. 60 × 40 m, and reaches a maximum depth of about 3 m (average depth: 1.8 m). It is a low-turbidity water pond, characterised by ultraoligotrophic conditions and by the lack of macroalgal and macrophyte cover at the bottom (Mania et al., 2019). Two water temperature profiles were performed in the pond on 12 July and 6 September 2015, showing a slight temperature decrease toward the bottom (300 cm depth) of 0.9 °C and 0.8 °C, respectively (Colombo et al., 2018a). A thick layer (up to several decimetres) of fine-grained sediments covers the bottom of the pond (Sam-buelli et al., 2015; Colombo et al., 2020). Coarse sediment constitutes the main land cover in the pond catchment, followed by bedrock outcrops; vegetated soil is also present, especially in the pond surroundings (Colombo et al., 2019a). More details on the catchment structural setting, hydrological and chemical dynamics, and ecosystem features of the pond can be found in Colombo et al. (2018a, Colombo et al. 2018b, Colombo et al. 2019a, Colombo et al. 2020) and Mania et al. (2019).

According to recent climate data series (2008–2015) obtained by the Col d'Olen AWS (Automatic Weather Station, Meteomont Service, Italian Army, 2900 m a.s.l., located approx. 900 m from the pond), the area is characterised by 400 mm of rainfall (on average) during the summer season, a mean annual air temperature of -2.6 °C, and a mean cumulative snowfall of 850 cm. The snowpack generally develops by late October to early November, and melt out usually occurs in July. At the site, heavy (>10 mm) and very heavy (>20 mm) rainfall events are relatively frequent during the snow-free season (Freppaz et al., 2019). At the site, mean annual nitrate concentrations in snow and rain are 5 and 10 μmol L⁻¹, respectively (year 2015, Colombo et al., 2019a).

2.2. Meteorological measurements

Air temperature and snow depth were measured at the Col d'Olen AWS. Rain data were obtained from the Gressoney-La-Trinité - Lago Gabiet AWS (2379 m a.s.l., managed by Regione Autonoma Valle d'Aosta), located ca. 2.5 km from the pond; the use of the Gressoney-La-Trinité - Lago Gabiet AWS was due to some gaps in the precipitation data of the Col d'Olen AWS during the analysed time-span. Data were




Fig. 1. (a) Location of the study area in Italy. (b) Elevation map of the study area showing the extent of the catchment, the Col d'Olen Rock Glacier, and the Col d'Olen Rock Glacier Pond. (c) Aerial view of the Rock Glacier Pond system and location of the UV-Vis probe (green polygon) on the southern side of the pond (cyan polygon) (aerial image year 2006, coordinate system WGS 84 / UTM zone 32 N). (d) Image of the probe used in this study. Details of the probe installation site: (e) installation in the pond, (f) on-shore installation with electrical and solar panels, and (g) internal view of the electrical panel, with pressurised air system and battery.

acquired on an hourly basis.

2.3. Spectroscopic and water temperature measurements

A UV-Vis submersible spectrophotometric probe (s::can sprectro::lyser™, s::can Messtechnik GmbH, Austria; Fig. 1d) was used for high-resolution spectroscopic measurements in the pond. The deployed probe had a measuring path length of 35 mm, suggested by the manufacturer for use in natural water. The probe measured absorbance over a 200–730 nm range at 2.5 nm intervals, thus it is potentially able to monitor concentrations of all compounds which generate an absorption. In particular, nitrate nitrogen (NO_3^- -N) induces an absorption between 210 and 240 nm (UV range) and turbidity does so between 400 and 700 nm (Visible range). Although NO_3^- -N can be estimated by the probe using the manufacturer's default global calibration (Langergraber et al., 2003), a local calibration algorithm can also be used to adapt the global calibration to the local conditions (Fleischmann et al., 2001). Local calibration uses laboratory analysis on actual samples from the investigated medium (Fleischmann et al., 2001), to account for matrix effects. To properly perform the local calibration, a sufficient number of water samples should be collected onsite prior to the deployment of the instrument, to cover the entire expected range of possible NO_3^- -N concentrations. In our case, considering the remoteness of the study area and the scarce availability of enough existing grab samples covering the whole range of nitrate concentrations (which is a typical condition encountered by researchers in these kinds of high-elevation settings), the default global calibration was used (detection range: $0.35\text{--}715 \mu\text{mol L}^{-1}$; Snazelle, 2015).

The reduced size of the probe (length: 58 cm, diameter: 4.4 cm, weight: ca. 2 kg) made this instrument particularly suitable for installation in remote areas. The probe was installed at the southern side of the pond (Fig. 1c), approximately 3-m from the shoreline (Fig. 1e) and at ca. 1-m depth. The sensor was installed in a zone of the pond not directly influenced by the rock glacier, thus representative of the processes ongoing in the entire pond and its catchment (Colombo et al., 2018a, Colombo et al., 2018b). The sensor was placed in horizontal position close to the pond bottom, with bottom-facing measuring path to avoid direct solar radiation incidence, prevent particles sedimentation in the measuring window, and avoid adhesion of gas bubbles. Power was provided by a 12 V/18 Ah battery charged by a solar panel (Fig. 1f) that ensured continuous measurements during the monitoring period. In addition, the measuring window was automatically cleaned at 3-hour intervals using pressurised air (Fig. 1g). Data were acquired at 3-hour intervals over the periods 14 July–9 October 2014 (88 days) and 29 June–31 July 2015 (33 days). The probe was removed on 9 October 2014 to prevent damages to the installation due to ice formation on the pond surface. On 1 August 2015, a sensor malfunction occurred due to a thunderstorm, preventing its recovery and thus further measurements during the remaining summer 2015 campaign. Raw data were saved in an internal datalogger. Every two weeks the probe was checked for its operational conditions and no technical issues were found.

Several processes are capable of influencing the performance of UV-Vis probes in estimating nitrate concentrations. For instance, interference of pH and salinity on nitrate quantification might occur (cf., Edwards et al., 2001), however, pH and electrical conductivity at the Rock Glacier Pond during the investigated period were between 7 and 7.5 and 32 and $45 \mu\text{S cm}^{-1}$, respectively (Colombo et al., 2018b), thus any interference was excluded. Also, since NO_2^- -N concentrations in pond water were extremely low, generally below the detection limits (Colombo et al., 2018b), an interference effect of nitrite could be excluded too (Huebsch et al., 2015). Furthermore, the probe internal algorithm compensates for the effect of turbidity, thus from this point of view any relevant interference could be discarded. Finally, water temperature might also have an effect on nitrate concentration estimation (Pellerin et al., 2012), however, during a comparison test between several probes for measuring nitrate, no relevant drift was observed for a

probe like the one used in the present study (Snazelle, 2015). Thus, the conditions of the pond and the characteristics of the employed UV-Vis probe were considered suitable for estimating the NO_3^- -N temporal variations in our setting.

Water temperature in the pond was measured continuously (at 3-hour intervals) from 28 July to 9 October 2014 and from 29 June to 12 October 2015 by means of a miniature temperature Onset HOBO® TidbiT v2 Temp logger (accuracy $\pm 0.21 \text{ }^\circ\text{C}$, resolution $0.02 \text{ }^\circ\text{C}$) installed at the pond bottom close to the probe.

2.4. Water level and grab samples collection and analysis

Water level was measured approximately on a weekly basis using a hydrometric station with direct observations from 14 July to 9 October 2014 and from 29 June to 12 October 2015. In conjunction with the water level measurements, the pond was sampled using a telescopic sampling pole, collecting samples close to the probe site (13 samples in 2014 and 15 samples in 2015), approximately at the same depth as the probe. To validate the probe during its deployment period, sample collection was performed at the exact time of the probe measurement to compare sample analyses with the instantaneous probe data (13 samples in 2014 and 5 samples in 2015).

Water samples for NO_3^- -N analyses were collected in new polyethylene tubes (volume 50 mL). To properly choose the tubes used for the sampling, they were preliminarily tested by storing Milli-Q water in conditions similar to those of the samples; no release of NO_3^- -N was detected. The content of the tubes was immediately filtered in the field through a $0.2 \mu\text{m}$ nylon membrane filter. Given the expected low NO_3^- -N concentrations, the suitability of the filters for our analyses was also tested; several blank filtration experiments were performed and no modifications in NO_3^- -N occurred. In addition, blank filtration experiments were conducted on different filters lots and filters were tested also on calibration standards, and no anomalies were detected. Samples were stored in an ice-packed cooler during transport from the field site, and then immediately transferred to the laboratory where they were refrigerated (at $+4 \text{ }^\circ\text{C}$) until analysis. The concentration of NO_3^- -N was determined by ion chromatography (Dionex DX-500, Sunnyvale, California, USA). Analysis quality was determined by including method blanks and repeated measurements of standard reference samples. Analytical precision was $< 10 \%$ and LOD was $1 \mu\text{mol L}^{-1}$.

Water samples for isotopic analyses were collected using new polyethylene tubes with airtight caps (volume 50 mL), completely filled to avoid head space. Furthermore, a 300 cm-deep snow profile was sampled before the melting season near the Col d'Olen AWS on 14 April 2015; six snow samples were collected at 50-cm intervals. A bulk rain-water sampler was also installed on 28 July 2014 close to the pond and sampled weekly (if precipitation occurred). Eight and four rain samples were collected in 2014 and 2015, respectively. Analysis was performed at the INSTAAR (Institute of Arctic and Alpine Research) Kiowa Environmental Chemistry Laboratory of the University of Colorado at Boulder (USA), by means of a cavity ring-down spectroscopy analyzer - Picarro L2130-i (Picarro Inc., Sunnyvale, California, USA). Isotopic composition was expressed as a δ (per mil) ratio of the sample to the Vienna Standard Mean Ocean Water (VSMOW), where δ is the ratio of $^{18}\text{O}/^{16}\text{O}$ and $^2\text{H}/^1\text{H}$. Analytical precision was $0.1 \text{ }^\circ\text{‰}$ and $1 \text{ }^\circ\text{‰}$ for $\delta^{18}\text{O}$ and $\delta^2\text{H}$, respectively.

2.5. Data elaboration and analysis

To assess the performance of the probe, NO_3^- -N concentrations estimated by the probe were compared to the ones determined in 18 grab samples; then, grab sample data were used to post-calibrate the probe data, through a simple linear regression approach. In addition, the degree of correlation among selected parameters was verified through the Pearson's correlation coefficient (r), if data were normally distributed (Kolmogorov-Smirnov test; Carvalho, 2015). The non-parametric

Spearman's test was used if data did not follow the normal distribution, even after the application of log-transformation. All analyses were performed in R environment (R Core Team, 2022).

3. Results

3.1. Meteorological and hydrological conditions

The two investigated years showed a marked difference in the snow-cover duration. Indeed, the melt-out date of snow occurred at the beginning of August in 2014 (Fig. 2a), while it occurred one month earlier in 2015 (Fig. 2b). The two periods differed also considering the rainfall, as 2014 was characterised by a lower cumulated amount (351 mm) with respect to 2015 (567 mm). In 2014, heavy rain events mostly occurred from mid-July, when snowpack was still present, to mid-August (Fig. 2a). July was the wettest month in 2014, with 154 mm of cumulated rain. Differently, in 2015, most of the heavy rain events occurred in the snow-free August (Fig. 2b), which was also the wettest month (303 mm). The mean daily air temperature during the investigated periods was +2.9 °C and +4 °C in 2014 and 2015, respectively.

The range of variation was also different between the two years. Indeed, daily air temperature variation in 2014 was more reduced (Fig. 2a), ranging from -3.0 °C (6 October) to $+8.1$ °C (17 July), while in 2015 it was larger (Fig. 2b), ranging from -5.8 °C (1 October) to $+12.1$ °C (5 July).

Regarding the daily water temperature, in 2014, an absolute minimum value of $+3.5$ °C was registered at the beginning of the measurement period (28 July), which was followed by a progressive increase until the absolute maximum value of $+10.7$ °C, that was reached at the end of August (31 August, Fig. 2a). Then, a decrease in water temperature occurred until the end of the investigated period, reaching a minimum value of $+5.7$ °C (10 October). It is also worth noting a short period (10 days), from the end of September to the beginning of October, when a water temperature increase occurred (maximum value: $+8.8$ °C, 4 October, Fig. 2a). In 2015, the monitored period began with a sharp water temperature increase, from $+4.0$ °C to $+13.2$ °C in 18 days (29 June–17 July, Fig. 2b). The temperature remained relatively high until 6 August (absolute maximum value: $+13.7$ °C), and then decreased to $+3.9$ °C in correspondence with an intense rainy period (8–24 August). Another water temperature increase occurred at the beginning of




Fig. 2. Daily air temperature, water temperature, rain, and snow depth in (a) 2014 and (b) 2015. Water level and $\delta^{18}\text{O}$ values in grab samples in (c) 2014 and (d) 2015 (only $\delta^{18}\text{O}$ is shown given the high correlation with $\delta^2\text{H}$). $\text{NO}_3^- \text{-N}$ concentrations in grab samples in (e) 2014 and (f) 2015.

September, after which the temperature finally declined toward the end of the season (absolute minimum value: $+2.3\text{ }^{\circ}\text{C}$, 3 October), when the snowpack started building up (Fig. 2b). During the deployment period of the probe, the water temperature exhibited daily cycles with the lowest mean values at 09:00 AM and the highest mean values at 6:00 PM, during both years; the mean difference between the highest and lowest diel values was $1.2\text{ }^{\circ}\text{C}$ in 2014 and $0.9\text{ }^{\circ}\text{C}$ in 2015 (not shown).

Regarding the water level, starting from the same initial value (about 100 cm), a level decline was weekly recorded in both seasons with a minimum value measured in October, which led to an overall decrease of about 30 cm (Fig. 2c,d). Despite the decreasing trend, some increases due to precipitation events occurred, for instance in September 2014 (4 cm increase) and August 2015 (10 cm increase).

The $\delta^{18}\text{O}$ and $\delta^2\text{H}$ values of water molecule were used as hydrologic tracers to detect the main water sources of the pond, e.g., water from snow melt and precipitation events (cf., Colombo et al., 2018b, 2019b; Brighenti et al., 2021). The most depleted values were shown by snow for $\delta^{18}\text{O}$ (mean $-18.9\text{ }‰$, range: -22.9 to $-16.2\text{ }‰$) and $\delta^2\text{H}$ (mean: $-139.2\text{ }‰$, range: -171.2 to $-116.2\text{ }‰$). On the opposite, rain had the most enriched values of $\delta^{18}\text{O}$ (mean: $-10.0\text{ }‰$, range: -12.3 to $-7.8\text{ }‰$) and $\delta^2\text{H}$ (mean: $-63.9\text{ }‰$, range: -89.3 to $-46.8\text{ }‰$). Pond water values of $\delta^{18}\text{O}$ (mean: $-12.8\text{ }‰$, range: -14.8 to $-10.4\text{ }‰$) and $\delta^2\text{H}$ (mean: $-90.3\text{ }‰$, range: -106.9 to $-70.8\text{ }‰$) arranged in between snow and rain ones. A progressive isotopic enrichment occurred during the analysed periods, with some evident sharp enrichments after heavy rain events in July 2014, July 2015, and August 2015 (Fig. 2c,d, 3). The isotope values of snow, rain, and water pond, shown in a dual isotope diagram (Fig. 3), fell close to the Local Meteoric Water Line (LMWL) calculated for Northern Italy by Longinelli and Selmo (2003).

3.2. NO_3^- -N concentrations

3.2.1. Grab samples

In 2014, the highest NO_3^- -N concentrations occurred at the beginning of the sampling season, in July (6.6 – $8.9\text{ }\mu\text{mol L}^{-1}$), then they decreased in August (2.7 – $4.1\text{ }\mu\text{mol L}^{-1}$, Fig. 2e). A slight, progressive NO_3^- -N increase occurred toward the end of the sampling period, in September and October (ca. $5\text{ }\mu\text{mol L}^{-1}$). In 2015, NO_3^- -N concentrations declined from the end of June ($5.6\text{ }\mu\text{mol L}^{-1}$) toward the minimum values measured in second half of July (ca. $2\text{ }\mu\text{mol L}^{-1}$), which were then followed by a sharp increase between the end of July and mid-August (6.2 – $13.8\text{ }\mu\text{mol L}^{-1}$, Fig. 2f). After a concentration decline toward

mid-September ($3.7\text{ }\mu\text{mol L}^{-1}$), NO_3^- -N concentrations increased again between the end of September and the beginning of October (5.0 – $6.6\text{ }\mu\text{mol L}^{-1}$).

3.2.2. UV-Vis probe measurements

The probe provided a good estimate of NO_3^- -N temporal variations; indeed, the correlation coefficient (r) between grab sample and probe NO_3^- -N concentrations was 0.83 ($p < 0.01$) (Fig. 4). Simple linear regression was used to post-calibrate the probe, obtaining a mean absolute percentage error of $\pm 18\%$. The residuals of the probe vs. grab sample regression were normally distributed (Fig. 4).

The daily probe data showed that, in 2014, after a 6-day period of relatively dry conditions (14–19 July 2014; cumulated rain: 7.8 mm), during which NO_3^- -N concentrations slightly declined (from 7.0 to 5.5




Fig. 4. Scatterplot of grab samples against the values predicted by the linear regression model with confidence levels at 95 % ($n = 18$, $r = 0.83$, $p < 0.01$). The inset shows the normal quantile-quantile plot of residuals of the final regression model (x axis: theoretical quantiles; y axis: residual quantiles).




Fig. 3. Dual-plot isotope distribution showing pond, snow and rain data. For pond samples, different colours identify the sampling months (from June to October). NIMWL: Northern Italian Meteoric Water Line.

$\mu\text{mol L}^{-1}$), a sharp NO_3^- -N increase was measured on 20–21 July 2014 (Fig. 5a,c). During these days, 3-hour data showed that 39.8 mm of rain fell in 33 h and NO_3^- -N concentrations increased from 5.6 to $11.2 \mu\text{mol L}^{-1}$ (Fig. 6a). Then, NO_3^- -N remained relatively stable (ca. 8–10 $\mu\text{mol L}^{-1}$) for 8 days even though other abundant rainy events occurred (22–29 July 2014; cumulated rain: 107 mm). The following period (1–22 August) was characterised by a gradual NO_3^- -N decline, until reaching a plateau that lasted approximately two weeks (23 August–5 September), with minimum concentrations of ca. $3.0 \mu\text{mol L}^{-1}$ (Fig. 5c, 6b). After this period, NO_3^- -N concentrations slowly increased until 16 September (ca. $+1 \mu\text{mol L}^{-1}$), when a faster increase occurred (ca. $+2 \mu\text{mol L}^{-1}$ in 4 days; Fig. 5c, 6c). Again, NO_3^- -N concentrations reached a plateau around $5 \mu\text{mol L}^{-1}$ and then increased at the end of the season ($6.5 \mu\text{mol L}^{-1}$). Temporal variations of turbidity showed similarities with NO_3^- -N ones, such as higher values in July (mean \pm st.dev.: 3.3 ± 2.1 Formazine Turbidity Unit - FTU; Fig. 5c). From 1 August to the third week of September, turbidity remained low and stable (mean \pm st.dev.: 0.8 ± 0.2 FTU), while it slightly increased in October (mean \pm st.dev.: 1.2 ± 0.6 FTU), similarly to NO_3^- -N (Fig. 5c). Turbidity was significantly correlated to NO_3^- -N ($r = 0.71$, $p < 0.001$, $n = 697$, 3-hour data).

In 2015, a dry period characterised the first investigated weeks (29 June–21 July, cumulated rain: 13.2 mm), during which daily NO_3^- -N concentrations progressively decreased from $7.1 \mu\text{mol L}^{-1}$ (29 June) to $3.1 \mu\text{mol L}^{-1}$ (21 July; Fig. 5b,d). After the initial dry period, a rain event occurred on 22 July, with 32.2 mm of rain cumulated in 6 h; a fivefold NO_3^- -N increase occurred, with 3-hour concentrations that increased from 2.9 to $15 \mu\text{mol L}^{-1}$ (Fig. 5b,d, 6d). Then, a progressive decrease in NO_3^- -N occurred until the end of the studied period (Fig. 5d). Turbidity and NO_3^- -N showed similar temporal variations (Fig. 5d, 6d), indeed they were significantly correlated ($r = 0.86$, $p < 0.001$, $n = 260$, 3-hour data).

In 2014, an inverse correlation was found between 3-hour water temperatures and NO_3^- -N concentrations estimated by the probe ($r =$

-0.89 , $p < 0.001$, $n = 589$). A correlation analysis was also performed between daily water temperatures and NO_3^- -N concentrations in grab samples: also in this case, a significant, negative correlation was found ($r = -0.74$, $p < 0.01$; $n = 11$). In 2015, no significant correlation was found between 3-hour water temperatures and NO_3^- -N concentrations estimated by the probe ($r = -0.01$, $p = 0.93$; $n = 260$) and neither between daily water temperatures and NO_3^- -N concentrations in grab samples ($r = -0.57$, $p = 0.31$; $n = 5$).

Finally, no evident and temporally consistent daily cycles in NO_3^- -N concentrations were found during both investigated years; the mean difference between the highest and lowest diel values was $0.3 \mu\text{mol L}^{-1}$ in 2014 and $0.6 \mu\text{mol L}^{-1}$ in 2015 (not shown). However, in the first days of 2015 (29 June–5 July), during the snow-melt phase, daily NO_3^- -N amplitude reached values up to $4.1 \mu\text{mol L}^{-1}$, with the highest concentrations ($7\text{--}9 \mu\text{mol L}^{-1}$) occurring in the evening (generally at 09:00 PM; not shown).

4. Discussion

4.1. Early summer: Snow-melt period

The different temperature and snow regimes at the beginning of the summer seasons have greatly influenced the snowmelt period in the analysed years. In 2014, the sampling campaign started when snow depth was 120 cm (14 July) corresponding to ca. 30 % of the maximum snow depth while, in 2015, snow depth was 70 cm, thus 90 % of snowpack was already melted.

In the early summer of both years, the higher NO_3^- concentrations followed by a progressive decline during the first days, before the occurrence of rain events, likely reflected the nitrate regression phase following the nitrate pulse (Pellerin et al., 2012; Beaton et al., 2017). Indeed, although the main nitrate pulse was not captured in this study, since it generally occurs during the early snow-melt period (e.g.,




Fig. 5. Daily air temperature, water temperature, rain, and snow depth in (a) 2014 and (b) 2015 (meteorological conditions are shown in Fig. 2 and here to enhance the interpretation of the NO_3^- -N and turbidity variations measured by the probe). Daily NO_3^- -N concentrations and turbidity estimated by the probe in (c) 2014 and (d) 2015. Red windows in panels c and d refer to panels a–d in Fig. 6.




Fig. 6. Three-hourly rain, water temperature (when available), and NO₃⁻-N concentrations and turbidity estimated by the probe in four selected periods: (a) 14 – 22 July 2014 (water temperature was not measured in this period), (b) 20 August – 3 September 2014, (c) 15 – 23 September 2014, and (d) 20 – 25 July 2015.

Sebestyen et al., 2008; Pellerin et al., 2012), the depleted isotope values in the pond water and low water temperature suggested a contribution from snow melt (cf., Hayashi, 2020; Marchina et al., 2020; Bearzot et al., 2023). Several studies have shown that the nitrate (and other solute) peak, often observed in montane surface waters during the early phases of snow melt (Johannessen & Henriksen, 1978), originates from the preferential elution of nitrate from the snowpack during the early melting phase (atmospheric origin, Sebestyen et al., 2008), and/or from the flushing of nitrate, produced by microbial nitrification, from catchment soils (terrestrial origin, Campbell et al., 2002; Sickman et al., 2003).

In 2014, 5 main rain-on-snow events occurred during the late snow-melt phase, contributing to 54 % of the cumulated rain during the entire 2014 monitoring period. During the first event (ca. 40 mm), NO_3^- concentrations doubled in 33 h (from 5.6 to $11.2 \mu\text{mol L}^{-1}$) while, for instance, the concentration increase after the third event (ca. 50 mm) was rather reduced (from 8 to $10 \mu\text{mol L}^{-1}$). These variations likely reflect the occurrence of new sources of N made available by an increase in the wetness during rain events. Indeed, these events could have increased the fraction of the catchment hydrologically connected to the pond through shallow subsurface and overland flows. At the end of July, an enrichment in the isotopic values of pond water with respect to the preceding days (from -14.8 to -13.6 ‰ for $\delta^{18}\text{O}$ and from -106.9 to -96.3 ‰ for $\delta^2\text{H}$) seems to indicate a relevant hydrological contribution from rain water. Enhanced infiltration could also have favoured the rising of groundwater and saturation of soil, resulting in the release of nitrate originated by nitrification. In addition, NO_3^- could also have derived from the flushing of other debris deposits in the catchment, such as talus, where it could be released by microbial pools (Sickman et al., 2003; Ley et al., 2004; Nemergut et al., 2005). The expansion of the hydrological network, enhancing the interaction between water and soil/debris deposits, might also explain the increase in turbidity, due to the transport of suspended particles to the pond.

In 2015, the snow-melt phase lasted few days (29 June–9 July) however, in this period, pronounced NO_3^- diel cycles (amplitude up to $4.1 \mu\text{mol L}^{-1}$) might indicate a connection to the diel variations of the hydrological fluxes originated from snow melt in the catchment. Indeed, previous research using in-situ high-frequency sensors in alpine streams (Pellerin et al., 2012) and proglacial meltwater rivers (Beaton et al., 2017) reported a diurnal nitrate variability caused by variations in diurnal discharge (inverse relationship), during the early and middle stage of the snow melt phase. However, it is not clear why these diel cycles were larger in 2015 with respect to 2014. An explanation might be related to the warmer atmospheric conditions during the snow-melt period in 2015, which caused a higher daily snow melt (ca. 6.5 cm day^{-1}) with respect to the one in 2014 (ca. 4.5 cm day^{-1}), resulting in potentially larger daily snowmelt fluxes. Unfortunately, the lack of continuous measurements of the pond water level does not allow us to deepen the analysis on this process.

4.2. Mid-summer: Snow-free season

In 2014, after the July peaks, a decline in NO_3^- concentrations was observed until they reached the minimum values at the end of August (ca. $3 \mu\text{mol L}^{-1}$). Concurrently with the NO_3^- decline, a gradual increase in water temperature was measured. The significant negative correlation between water temperature and NO_3^- in 2014 suggests that the biological processes consuming N (e.g., plant uptake and microbial immobilisation), and taking place in the pond and in the catchment, played a fundamental role in the nitrate dynamics. A number of studies reported relatively low values of NO_3^- in alpine forest streams during summer, highlighting the importance of soil biological community in the retention and loss of N and, therefore, the strict connection between soil and waters in mountain ecosystems (e.g., Balestrini et al., 2006; Helliwell et al., 2007; Curtis et al., 2011). In agreement with these findings, Balestrini et al. (2013) showed a strong positive relationship between

the areal extension of developed soils and the retention of N in running waters above the treeline.

A growing number of studies (e.g., Roberts & Mulholland, 2007; Pellerin et al., 2012; Beaton et al., 2017) have shown that in-stream and in-lake processes can also be important drivers of water nutrient concentrations. For instance, Pellerin et al. (2012) and Beaton et al. (2017) attributed the nitrate diel cycles (amplitude $1\text{--}2 \mu\text{mol L}^{-1}$), detected in a forest stream and in a proglacial lake, to the biological uptake directly linked to autotrophic production. Photosynthesis provides supplementary energy that can be used by biological communities to lessen nitrate, for use in metabolism and biosynthesis (Roberts & Mulholland, 2007). In lentic systems like ponds, the biological processes occurring within the water column and in the sediments should be even more important compared to lotic waters. In addition, the lack of tree shading and the high radiation intensity should be able to enhance the photosynthesis in aquatic ecosystems over the treeline. The fact that, in our case, evident nitrate diel cycles were not recorded might be attributable to the low sensitivity of the probe at very low concentrations, which did not allow us to detect NO_3^- variations like those expected as a result of autotrophic assimilation. Although the pond lacks the most evident examples of benthic primary producers (e.g., submerged macrophytes and macroalgae), the findings of Mania et al. (2019, Mania et al. 2021) demonstrated the presence of higher proportions of cyanobacterial sequences in the deepest area of the pond, and attested that photo- and chemolithoautotrophic bacteria may represent an important component of the total benthic microbial community. Finally, during the entire mid-summer period, rainfall events (cumulated rain: 110 mm) did not strongly affect turbidity (and NO_3^- concentrations), which remained low and stable, possibly indicating a poor hydrological connection between the catchment and the pond. Thus, it is possible to hypothesise that bacterial processes in the pond played a major role in driving the nitrate temporal evolution during the snow-free season, despite a water temperature lower than $+11$ °C. Another process potentially responsible for the NO_3^- removal is denitrification in pond sediments. However, the quantification of ambient denitrification rate is critical and a relevant gap of knowledge exists concerning this process in small oligotrophic lakes/ponds with very low N concentrations (Seitzinger et al., 2006). Some studies performed in cold and oligotrophic environments (Myrstener et al., 2016; Vila-Costa et al., 2016; Palacin-Lizarbe et al., 2018) reported denitrification rates falling in the low range of freshwater sediments (Seitzinger, 1988; Piña-Ochoa & Alvarez-Cobelas, 2006).

The snow-free season 2015 started one month earlier with respect to the 2014's one. During the first two weeks of July, the water temperature increased rapidly (up to ca. $+13$ °C), while NO_3^- concentrations reached their minimum values (ca. $3 \mu\text{mol L}^{-1}$) approximately a month and a half earlier compared to the previous year. The first rain event, after several consecutive dry days, resulted in a higher and faster NO_3^- ($15 \mu\text{mol L}^{-1}$) and turbidity (29.7 FTU) peak with respect to the one recorded in July 2014. The rapid decline of both parameters in the following days, when the water temperature was relatively stable, could be attributed to the exhaustion of the N source, such as the organic soils (Schlesinger, 1997; Campbell et al., 2002; Burns et al., 2019) located around the pond (Colombo et al., 2020).

The NO_3^- temporal evolution after the probe malfunction (31 July) was investigated by considering the weekly grab samples. In particular, one more NO_3^- peak ($14 \mu\text{mol L}^{-1}$) was measured on 13 August, after a heavy rain event that occurred between 8 and 10 August (cumulated rain: 115 mm). The higher soil temperatures measured in 2015 close to the pond (Freppaz et al., 2019) might have enhanced the microbial activity and, therefore, both mineralisation of organic matter and nitrification (Rogora et al., 2008; Dawes et al., 2017; Donhauser et al., 2021), favouring the release of nitrate during the precipitation events. Enhanced mobilisation due to stronger evapoconcentration in the soil under previous drier conditions might have also played a role (Knapp et al., 2020). Moreover, in addition to the meteorological conditions during the summer season, also the previous snow-covered season might

have contributed in influencing the nutrient dynamics in soil and surface water in the area (Freppaz et al., 2019). Specifically, Magnani et al. (2017) demonstrated that a short snow-cover duration, like the one recorded in 2015, may increase soil temperature and substrate availability during the subsequent growing season, favouring soil microbial biomass.

Other processes might have contributed in supplying NO_3^- to the pond during and after the rainfall events, although they were considered of minor importance. For instance, Colombo et al. (2018b, Colombo et al., 2019a) showed the role of rainfall in enhancing the export of NO_3^- from the rock glacier. However, the authors also estimated that the hydrochemical influence of the rock glacier on the overall pond was limited in frequency and magnitude, and it was not likely to affect the entire pond in terms of hydrology and water chemistry. The contribution of a direct atmospheric input of NO_3^- to the pond water cannot be excluded either. However, taking the mid-August 2015 event as an example, the NO_3^- peak concentration in pond water ($14 \mu\text{mol L}^{-1}$) was above the mean NO_3^- concentration measured in rain in the area ($10 \mu\text{mol L}^{-1}$) and during the rain event itself ($12.7 \mu\text{mol L}^{-1}$), as reported by Colombo et al. (2019a). In addition, previous studies in this area have shown that physical and chemical processes occurring in the catchments, rather than direct precipitation contributions, are likely the predominant drivers in determining nitrate concentrations, and their seasonal variations, in surface waters (Magnani et al., 2017; Colombo et al., 2019a,b; Freppaz et al., 2019).

4.3. Late summer-fall transition

The behaviour of NO_3^- during September and October 2014 was the opposite of that observed in the previous two months. Indeed, increasing NO_3^- concentrations, in conjunction with decreasing water temperatures, were observed. Some rain events interrupted the gradual evolution of both parameters with sharper changes depending on rainfall intensity and duration. Turbidity also showed a response to the latest rain events, although its response occurred later and was weaker compared to the one of NO_3^- . The available weekly data exhibited a similar NO_3^- pattern in 2015. Indeed, the NO_3^- concentration increased from $3.7 \mu\text{mol L}^{-1}$, at the beginning of September, to $6.4 \mu\text{mol L}^{-1}$ after a month. The NO_3^- concentration at the end of the season was comparable to that recorded in 2014, suggesting a sort of stability in the processes characterising this aquatic ecosystem. This nitrate behaviour during the late summer-fall transition phase is coherent with the findings of Freppaz et al. (2019) in a nearby pond, and with similar reports from alpine tundra (Sickman et al., 2003; Balestrini et al., 2013; Williams et al., 2015a) and alpine forest areas (Balestrini et al., 2006; Rusjan & Mikoš, 2010). The common hypothesis explaining the increase in nitrate is the slowdown of the biologically-mediated immobilisation processes in soil, due to colder soil temperature at the end of summer. In these conditions, the possible transfer of nitrate to the pond depends on rainfall occurrence. In addition, the temperature decrease can affect the in-pond nutrient uptake (cf., Roberts & Mulholland, 2007), also lowering the photosynthetic activity. Consequently, the autotrophic community might play a role in N retention during this season (Rusjan & Mikoš, 2010; Oleksy et al., 2021).

Nitrate increase due to simple concentration as a result of evaporation might be also considered. However, in this phase of the season, the increases of NO_3^- were observed during colder atmospheric periods. In addition, the distribution of the stable water isotopic data was consistent with the NIMWL. Thus, results suggest that evaporation can be considered a negligible process in isotope fractionation in the pond, since it would have caused an enrichment in $\delta^{18}\text{O}$ and a subsequent decrease in the distribution slope with respect to the NIMWL. It is also true that, in both years, the water level of the pond as measured on a weekly basis by Colombo et al. (2018a) progressively decreased from June/July to October, thus this might seem to point towards a role of evaporation, considering the lack of stable pond outflows. However, this water level

decrease was found to be mostly driven by a sub-surface seepage at the pond bottom, where a minor fault zone in bedrock is located, characterised by altered and highly-fractured rocks (Colombo et al., 2018a). Finally, isotopically-enriched groundwater (cf., Fan et al., 2022), possibly nitrate-concentrated, might also have provided a higher contribution in late summer-early fall (cf., Hayashi, 2020). Unfortunately, the contribution of groundwater, that was not sampled in this study since no groundwater springs are present in the analysed catchment, is not easy to disentangle when dealing with high-elevation lakes and ponds, especially in the absence of stable surface inflows/outflows (cf., Langston et al., 2013), like in the study site. Thus, the role of groundwater in supplying NO_3^- in this high-elevated setting must be further investigated.

4.4. Environmental implications and research/operative perspectives

The complex interplay between rainfall, snow melt, and temperature (water and air) during the summer season showed the capability to drive daily and seasonal NO_3^- concentrations in the investigated pond. In the European Alps, significant increases in air temperature (Gobiet et al., 2014) and snow line elevation (Koehler et al., 2022), as well as reductions in snow cover duration (Klein et al., 2016), snow depth (Matiu et al., 2021), and snow water equivalent (Marty et al., 2017; Colombo et al., 2022; Colombo et al., 2023), have occurred in the last decades. These occurrences are expected to become even more dramatic in the future (Gobiet et al., 2014; Beniston et al., 2018). Regional climate model simulations also indicate that summer rainfall at Alpine high elevations will increase due to global warming, despite the expected large-scale precipitation reduction (Giorgi et al., 2016) and drought event increases (Spinoni et al., 2018), together with potential increases in heavy precipitation and hot temperature extremes (Scherrer et al., 2016). Furthermore, rain-on-snow events are also predicted to increase in the next decades (Beniston & Stoffel, 2016).

The alteration of the hydrological cycle induced by climate change (e.g., changes in runoff peak and timing together with modifications in the water residence time) could have a great influence on both N input and removal to/from aquatic ecosystems (e.g., Baron et al., 2013). In addition, during dry periods, leaching of NO_3^- from soils is reduced/absent due to low hydraulic conductivity in the soil profile. As a consequence, a residence time of potentially leachable N in soils (and its plant uptake) as well as microbial immobilisation could increase. Then, this N temporally stored in soils could be mobilised (NO_3^-)/mineralised (organic N) and leached in the following wet periods. In the present study, the N dynamics were analysed and compared in two years characterised by extremely different meteorological conditions and features that might be typical under the climate change impacts, such as very high air temperature, heavy rainfalls, and rain-on-snow events. In 2015, a higher N supply to the pond was found, in the form of peaks due to rapid and intense hydrological flows. These events occurred in the mid-summer phase when the biological N uptake is commonly maximum and thus minimal NO_3^- values are expected. This is even more evident if compared to the NO_3^- dynamics in 2014, when the relatively stable meteorological conditions during mid-summer led to limited variations and low concentrations of NO_3^- . Moreover, this study further highlights the role of air and water temperature in controlling the N production and retention at daily and seasonal time scale. In the snow-free season, when dry conditions occur and the pond is, therefore, hydrologically disconnected from the catchment, water temperature could be used as a robust predictor of pond water NO_3^- concentration. Only by using an approach based on in-situ high-frequency measurements it was possible to fully grasp both the effect of hydrological processes and the role of in-lake nutrient biological retention processes in the N dynamics in the Col d'Olen Rock Glacier Pond, representative of ultraoligotrophic water bodies in alpine tundra. Finally, the high-resolution temporal assessment of nitrate dynamics could also avoid potentially misleading comparisons of water chemistry collected through synoptic surveys

during different seasons and years, and improve the understanding of the connections between surface water, soil water and groundwater in these remote environments.

Future work could take into consideration the use of longer measuring path lengths to increase the accuracy and sensitivity of the UV-Vis probe at these low concentrations, which are at the lower end of the probe measurement range. In addition, performing local calibrations using samples taken from the actual water investigated will help account for the matrix effect and, thus, further increase the measurement accuracy. In this context, a consistent number of samples (for instance, 40–50, half for calibration and half for validation) could be collected in order to improve the probe performance assessment, also taking into account a possible wider range of nitrate concentrations. However, considering a weekly sampling plan and the fact that ponds like the one analysed in this study are typically ice-free for a limited amount of time (maximum 3/4 months), the collection of a large number of samples would require the in-situ maintenance of the probe for three or more ice-free seasons, which could be hard to perform in such highly elevated remote areas. Thus, alternatively, a focus on shorter periods could be recommended, such as during the snow-ionic pulse (if the ice conditions in the pond allow for the installation of the probe), late snow melt, in-lake retention conditions during periods with poor hydrological connection, or single months for the monitoring of possible heavy rain events; in turn, a larger number of samples could be collected, even on a daily basis. Finally, performing measurements with a higher sensitivity is considered necessary to properly investigate the diel variability of nitrate in order to verify the occurrence of a retention process strictly connected to photosynthesis (e.g., autotrophic assimilation). In this regard, the high-frequency measurement of O₂ to explore the extent of its diel variations and to record the groundwater inflow would be beneficial for supporting the interpretation of the NO₃⁻ analysis.

5. Conclusions

This study reports on the first use of a UV-Vis submersible spectrophotometric probe in association with weekly analyses of NO₃⁻-N and stable isotopes of water ($\delta^{18}\text{O}$ and $\delta^2\text{H}$), together with continuous meteorological, water temperature, and turbidity measurements, in a high-elevation pond in alpine tundra. The proposed approach allowed for disentangling the complex effects, and their interplay, of snow melt, temperature (air and water), and summer rainfall on nitrate dynamics. In particular, snow-melt duration and temperature fluctuations drove the nitrate variations on a seasonal basis, also determining the timing of the seasonal transitions. However, short-duration meteorological events (lasting even just a few hours), such as heavy rainfalls and rain-on-snow events, deeply disrupted these dynamics, in the form of NO₃⁻ peaks due to rapid and intense hydrological flows, which lasted up to few days/weeks. The effects of these hydrological events were properly assessed thanks to the use of in-situ high-frequency measurements, which also allowed us to better define the role of in-lake nutrient biological retention processes in the N dynamics in the investigated ultra-oligotrophic pond. Therefore, high-resolution temporal monitoring of nitrate dynamics may contribute to a better understanding of the biogeochemical processes occurring in these remote, yet highly sensitive environments. Ultimately, this could help predicting how the quality of high-elevation surface waters will respond to changing climate and related climate extremes (e.g., reduction in snow-cover duration and increases in magnitude and frequency of heavy rainfall events).

CRedit authorship contribution statement

N. Colombo: Conceptualization, Investigation, Methodology, Formal analysis, Data curation, Software, Visualization, Writing – original draft, Writing – review & editing. **R. Balestrini:** Methodology, Formal analysis, Writing – original draft, Writing – review & editing. **D. Godone:** Investigation, Methodology, Formal analysis, Data curation,

Software, Visualization, Writing – review & editing. **D. Vione:** Writing – review & editing. **D. Said-Pullicino:** Writing – review & editing. **G. Viviano:** Investigation, Writing – review & editing. **M. Martin:** Writing – review & editing. **C.A. Delconte:** Writing – review & editing. **S. Fratianni:** Conceptualization, Funding acquisition, Writing – review & editing. **A.G. Capodaglio:** Writing – review & editing. **E. Pintaldi:** Writing – review & editing. **M. Freppaz:** Writing – review & editing. **F. Salerno:** Investigation, Methodology, Formal analysis, Funding acquisition, Funding acquisition, Writing – review & editing.

Declaration of Competing Interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Data availability

Data will be made available on request.

Acknowledgments

We thank Elena Serra, Elisa Giaccone, Andrea Magnani, Diego Guenzi, Maria Chiara Caschetto, and Davide Viglietti for their help in fieldwork and laboratory activities. We give special thanks to Mark Williams and Holly Hughes (Department of Geography, University of Colorado, Boulder, USA), Consorzio di Miglioramento Fondiario di Gressoney (Aosta), Monterosa Spa and Monterosa 2000 (Monterosa ski). This research was supported by the PRIN 2010-11 (funded project of the Italian Ministry for Education and Research) named “Response of morphoclimatic system dynamics to global changes and related geomorphological hazards” (coord. Prof. Carlo Baroni). Nicola Colombo and Michele Freppaz were partially supported by the project NODES, which has received funding from the MUR – M4C2 1.5 of PNRR funded by the European Union - NextGeneration EU (Grant agreement no. ECS00000036). Davide Vione acknowledges financial support by Project CH4.0 under the MUR program “Dipartimenti di Eccellenza 2023-2027” (CUP: D13C22003520001), as well as Next Generation EU - PNRR project GRINS (Growing Resilient, INclusive, and Sustainable), PE9 - spoke 6. We are grateful to the editor and two reviewers who provided valuable feedback and input during the review of this manuscript. We would like to dedicate this paper to the memory of our colleague Simona Fratianni, who passed away prematurely.

References

- Aber, J., McDowell, W., Nadelhoffer, K., Magill, A., Berntson, G., Kamakea, M., McNulty, S., Currie, W., Rustad, L., Fernandez, I., 1998. Nitrogen saturation in temperate forest ecosystems: hypotheses revisited. *Bioscience* 48 (11), 921–934. <https://doi.org/10.2307/1313296>.
- Adrian, R., O'Reilly, C.M., Zagarese, H., Baines, S.B., Hessen, D.O., Keller, W., Livingstone, D.M., Sommaruga, R., Straile, D., Van Donk, E., Weyhenmeyer, G.A., Winder, M., 2009. Lakes as sentinels of climate change. *Limnol. Oceanogr.* 54 (6), 2283–2297. <https://doi.org/10.4319/lo.2009.54.6.part.2.2283>.
- Balestrini, R., Di Martino, N., Van Mieghroet, H., 2006. Nitrogen cycling and mass balance for a forested catchment in the Italian Alps. *Biogeochemistry* 78, 97–123. <https://doi.org/10.1007/s10533-005-3429-7>.
- Balestrini, R., Arese, C., Freppaz, M., Buffagni, A., 2013. Catchment features controlling nitrogen dynamics in running waters above the tree line (central Italian Alps). *Hydrol. Earth Syst. Sci.* 17 (3), 989–1001. <https://doi.org/10.5194/hess-17-989-2013>.
- Barnes, R.T., Williams, M.W., Parman, J.N., Hill, K., Caine, N., 2014. Thawing glacial and permafrost features contribute to nitrogen export from Green Lakes Valley, Colorado Front Range, USA. *Biogeochemistry* 117, 413–430. <https://doi.org/10.1007/s10533-013-9886-5>.
- Baron, J.S., Hall, E.K., Nolan, B.T., Finlay, J.C., Bernhardt, E.S., Harrison, J.A., Chan, F., Boyer, E.W., 2013. The interactive effects of excess reactive nitrogen and climate change on aquatic ecosystems and water resources of the United States. *Biogeochemistry* 114, 71–92. <https://doi.org/10.1007/s10533-012-9788-y>.
- Beazot, F., Colombo, N., Cremonese, E., Morra di Cella, U., Drigo, E., Caschetto, M., Basirico, S., Crosta, G.B., Frattini, P., Freppaz, M., Pogliotti, P., Salerno, F., Brunier, A., Rossini, M., 2023. Hydrological, thermal and chemical influence of an

- intact rock glacier discharge on mountain stream water. *Sci. Total Environ.* 876, 162777 <https://doi.org/10.1016/j.scitotenv.2023.162777>.
- Beaton, A.D., Wadhwa, J.L., Hawkings, J., Bagshaw, E.A., Lamarche-Gagnon, G., Mowlem, M.C., Tranter, M.C., 2017. High-resolution in situ measurement of nitrate in runoff from the Greenland Ice Sheet. *Environ. Sci. Tech.* 51 (21), 12518–12527. <https://doi.org/10.1021/acs.est.7b03121>.
- Beniston, M., Stoffel, M., 2016. Rain-on-snow events, floods and climate change in the Alps: Events may increase with warming up to 4 °C and decrease thereafter. *Sci. Total Environ.* 571, 228–236. <https://doi.org/10.1016/j.scitotenv.2016.07.146>.
- Beniston, M., Farinotti, D., Stoffel, M., Andressen, L.M., Coppola, E., Eckert, N., Fantini, A., Giacoma, F., Hauck, C., Huss, M., Huwald, H., Lehning, M., López-Moreno, J.-L., Magnusson, J., Marty, C., Morán-Tejeda, E., Morin, S., Naaim, M., Provenzale, A., Rabatel, A., Six, D., Stötter, J., Strasser, U., Terzagio, S., Vincent, C., 2018. The European mountain cryosphere: a review of its current state, trends, and future challenges. *Cryosphere* 12, 759–794. <https://doi.org/10.5194/tc-12-759-2018>.
- Brighenti, S., Engel, M., Tolotti, M., Bruno, M.C., Wharton, G., Comiti, F., Tirlir, W., Cerasino, L., Bertoldi, W., 2021. Contrasting physical and chemical conditions of two rock glacier springs. *Hydrol. Process.* 35, e14159.
- Burns, D.A., Pellerin, B.A., Miller, M.P., Capel, P.D., Tesorieri, A.J., Duncan, J.M., 2019. Monitoring the riverine pulse: Applying high-frequency nitrate data to advance integrative understanding of biogeochemical and hydrological processes. *WIREs Water* 6 (4), e1348.
- Campbell, D.H., Kendall, C., Chang, C.C.Y., Silva, S.R., Tonnessen, K.A., 2002. Pathways for nitrate release from an alpine watershed: Determination using $\delta^{15}N$ and $\delta^{18}O$. *Water Resour. Res.* 38 (5), 8. <https://doi.org/10.1029/2001WR000294>, 10–1–10–9.
- Carvalho, L., 2015. An improved evaluation of Kolmogorov's distribution. *J. Stat. Softw.* 65, 1–8. <https://doi.org/10.18637/jss.v065.c03>.
- Catalan, J., Camarero, L., Felip, M., Pla, S., Ventura, M., Buchaca, T., Bartumeus, F., De Mendoza, G., Miró, A., Casamayor, E.O., Medina-Sánchez, J.M., Bacardit, M., Altuna, M., Bartrons, M., Díaz De Quijano, D., 2006. High mountain lakes: Extreme habitats and witnesses of environmental changes. *Limnologia* 25 (1–2), 551–584. <https://doi.org/10.23818/limn.25.38>.
- Clow, D.W., Sickman, J.O., Striegl, R.G., Krabbenhoft, D.P., Elliott, J.G., Dornblaser, M., Roth, D.A., Campbell, D.H., 2003. Changes in the chemistry of lakes and precipitation in high-elevation national parks in the western United States, 1985–1999. *Water Resour. Res.* 39 (6), 1171–1184. <https://doi.org/10.1029/2002WR001533>.
- Colombo, N., Sambuelli, L., Comina, C., Colombero, C., Giardino, M., Gruber, S., Viviano, G., Vittori Antisari, L., Salerno, F., 2018a. Mechanisms linking active rock glaciers and impounded surface water formation in high-mountain areas. *Earth Surf. Proc. Land* 43 (2), 417–431. <https://doi.org/10.1002/esp.4257>.
- Colombo, N., Gruber, S., Martin, M., Malandrino, M., Magnani, A., Godone, D., Freppaz, M., Fratianni, S., Salerno, F., 2018b. Rainfall as primary driver of discharge and solute export from rock glaciers: the Col d'Olen Rock Glacier in the NW Italian Alps. *Sci. Total Environ.* 639, 316–330. <https://doi.org/10.1016/j.scitotenv.2018.05.098>.
- Colombo, N., Salerno, F., Martin, M., Malandrino, M., Giardino, M., Serra, E., Godone, D., Said-Pullicino, D., Fratianni, S., Paro, L., Tartari, G., Freppaz, M., 2019a. Influence of permafrost, rock and ice glaciers on chemistry of high-elevation ponds (NW Italian Alps). *Sci. Total Environ.* 685, 886–901. <https://doi.org/10.1016/j.scitotenv.2019.06.233>.
- Colombo, N., Bocchiola, D., Martin, M., Confortola, G., Salerno, F., Godone, D., D'Amico, M.E., Freppaz, M., 2019b. High export of nitrogen and dissolved organic carbon from an Alpine glacier (Indren Glacier, NW Italian Alps). *Aquat. Sci.* 81, 74. <https://doi.org/10.1007/s00027-019-0670-z>.
- Colombo, N., Ferronato, C., Vittori Antisari, L., Marziali, L., Salerno, F., Fratianni, S., D'Amico, M.E., Ribolini, A., Godone, D., Sartini, S., Paro, L., Morra di Cella, U., Freppaz, M., 2020. A rock-glacier–pond system (NW Italian Alps): Soil and sediment properties, geochemistry, and trace-metal bioavailability. *Catena* 194, 104700. <https://doi.org/10.1016/j.catena.2020.104700>.
- Colombo, N., Valt, M., Romano, E., Salerno, F., Godone, D., Cianfarra, P., Freppaz, M., Maugeri, M., Guyennon, N., 2022. Long-term trend of snow water equivalent in the Italian Alps. *J. Hydrol.* 614 (A) <https://doi.org/10.1016/j.jhydrol.2022.128532>.
- Colombo, N., Guyennon, N., Valt, M., Salerno, F., Godone, D., Cianfarra, P., Freppaz, M., Maugeri, M., Manara, V., Acquafotta, F., Pietrangeli, A.B., Romano, E., 2023. Unprecedented snow-drought conditions in the Italian Alps during the early 2020s. *Environ. Res. Lett.* 18, 074014 <https://doi.org/10.1088/1748-9326/acdb88>.
- Curtis, C., Evans, C., Goodale, C., Heaton, T., 2011. What have stable isotope studies revealed about the nature and mechanisms of N Saturation and nitrate leaching from semi-natural catchments? *Ecosystems* 14, 1021–1037. <https://www.jstor.org/stable/41505929>.
- Dawes, M.A., Schleppli, P., Hättenschwiler, S., Rixen, C., Hagedorn, F., 2017. Soil warming opens the nitrogen cycle at the alpine treeline. *Glob. Chang. Biol.* 23 (1), 421–434. <https://doi.org/10.1111/gcb.13365>.
- Donhauser, J., Qi, W., Bergk-Pinto, B., Frey, B., 2021. High temperatures enhance the microbial genetic potential to recycle C and N from necromass in high-mountain soils. *Glob. Chang. Biol.* 27 (7), 1365–1386. <https://doi.org/10.1111/gcb.15492>.
- Edwards, A.C., Hooda, P.S., Cook, Y., 2001. Determination of nitrate in water containing dissolved organic carbon by ultraviolet spectroscopy. *Int. J. Environ. Anal. Chem.* 80 (1), 49–59. <https://doi.org/10.1080/03067310108044385>.
- Elser, J.J., Kyle, M., Steger, L., Nydick, K.R., Baron, J.S., 2009. Nutrient availability and phytoplankton nutrient limitation across a gradient of atmospheric nitrogen deposition. *Ecology* 90 (11), 3062–3073. <https://doi.org/10.1890/08-1742.1>.
- Fan, Y., Wu, Y., Wang, Y., Jiang, S., Yu, S., Shang, H., 2022. An analysis of surface water-groundwater interactions based on isotopic data from the Kaidu River Basin. *South Tianshan Mountain. Water* 14, 2259. <https://doi.org/10.3390/w14142259>.
- Fleischmann, N., Langergraber, G., Weingartner, A., Hofstädter, F., Nusch, S., Maurer, P., 2001. On-line and in-situ measurement of turbidity and COD in wastewater using UV/VIS spectrometry. SCAN Media Library. https://www.s-can.at/wp-content/uploads/2021/09/p_2001_06.pdf.
- Freppaz, M., Viglietti, D., Balestrini, R., Lonati, M., Colombo, N., 2019. Climatic and pedoclimatic factors driving C and N dynamics in soil and surface water in the alpine tundra (NW-Italian Alps). *Nat. Conserv.* 34, 67–90. <https://doi.org/10.3897/natureconservation.34.30737>.
- Giorgi, F., Torma, C., Coppola, E., Ban, N., Schär, C., Somot, S., 2016. Enhanced summer convective rainfall at alpine high elevations in response to climate warming. *Nat. Geosci.* 9, 584–589. <https://doi.org/10.1038/ngeo2761>.
- Gobiet, A., Kotlarski, S., Beniston, M., Heinrich, G., Rajczak, J., Stoffel, M., 2014. 21st century climate change in the European Alps: a review. *Sci. Total Environ.* 493, 1138–1151. <https://doi.org/10.1016/j.scitotenv.2013.07.050>.
- Hamerlík, L., Svitok, M., Novikmec, M., Očadlík, M., Bitušík, P., 2014. Local, among-site, and regional diversity patterns of benthic macroinvertebrates in high altitude waterbodies: Do ponds differ from lakes? *Hydrobiologia* 723 (1), 41–52. <https://doi.org/10.1007/s10750-013-1621-7>.
- Hayashi, M., 2020. Alpine hydrogeology: The critical role of groundwater in sourcing the headwaters of the world. *Groundwater* 58, 498–510. <https://doi.org/10.1111/gwat.12965>.
- Helliwell, R.C., Coull, M.C., Davies, J.J.L., Evans, C.D., Norris, D., Ferrier, R.C., Jenkins, A., Reynolds, B., 2007. The role of catchment characteristics in determining surface water nitrogen in four upland regions in the UK. *Hydrol. Earth Syst. Sci.* 11 (1), 356–371. <https://doi.org/10.5194/hess-11-356-2007>.
- Huebsch, M., Grimmeisen, F., Zemann, M., Fenton, O., Richards, K.G., Jordan, P., Sawarieh, A., Blum, P., Goldscheider, N., 2015. Technical Note: Field experiences using UV/VIS sensors for high-resolution monitoring of nitrate in groundwater. *Hydrol. Earth Syst. Sci.* 19, 1589–1598. <https://doi.org/10.5194/hess-19-1589-2015>.
- Johannessen, M., Henriksen, A., 1978. Chemistry of snow meltwater: Changes in concentration during melting. *Water Resour. Res.* 14 (4), 615–619. <https://doi.org/10.1029/WR014i004p0615>.
- Klein, G., Vitasse, Y., Rixen, C., Marty, C., Rebetez, M., 2016. Shorter snow cover duration since 1970 in the Swiss alps due to earlier snowmelt more than to later snow onset. *Clim. Chang.* 139 (3), 637–649. <https://doi.org/10.1007/s10584-016-1806-y>.
- Knapp, J.L.A., Freyberg, J., Studer, B., Kiewiet, L., Kirchner, J.W., 2020. Concentration–discharge relationships vary among hydrological events, reflecting differences in event characteristics. *Hydrol. Earth Syst. Sci.* 24, 2561–2576. <https://doi.org/10.5194/hess-24-2561-2020>.
- Koehler, J., Bauer, A., Dietz, A.J., Kuenzer, C., 2022. Towards forecasting future snow cover dynamics in the European Alps—The potential of long optical remote-sensing time series. *Remote Sens.* 14, 4461. <https://doi.org/10.3390/rs14184461>.
- Kopáček, J., Stuchlík, E., Wright, R.F., 2005. Long-term trends and spatial variability in nitrate leaching from alpine catchment–lake ecosystems in the Tatras Mountains (Slovakia–Poland). *Environ. Pollut.* 135 (1), 89–101. <https://doi.org/10.1016/j.envpol.2004.12.012>.
- Langergraber, G., Fleischmann, N., Hofstädter, F., 2003. A multivariate calibration procedure for UV/VIS spectrometric quantification of organic matter and nitrate in wastewater. *Water Sci. Technol.* 47, 63–71. <https://doi.org/10.2166/wst.2003.0086>.
- Langston, G., Hayashi, M., Roy, J.W., 2013. Quantifying groundwater-surface water interactions in a proglacial moraine using heat and solute tracers. *Water Resour. Res.* 49 (9), 5411–5426. <https://doi.org/10.1002/wrcr.20372>.
- Ley, R.E., Williams, M.W., Schmidt, S.K., 2004. Microbial population dynamics in an extreme environment: Controlling factors in talus soils at 3750 m in the Colorado Rocky Mountains. *Biogeochemistry* 68, 297–311. <https://doi.org/10.1023/B:BIOG.0000031032.58611.d0>.
- Longinelli, A., Selmo, E., 2003. Isotopic composition of precipitation in Italy: a first overall map. *J. Hydrol.* 270, 75–88. [https://doi.org/10.1016/S0022-1694\(02\)00281-0](https://doi.org/10.1016/S0022-1694(02)00281-0).
- Magnani, A., Viglietti, D., Balestrini, R., Williams, M.W., Freppaz, M., 2017. Contribution of deeper soil horizons to N and C cycling during the snow-free season in alpine tundra, NW Italy. *Catena* 155, 75–85. <https://doi.org/10.1016/j.catena.2017.03.007>.
- Mania, I., Gorra, R., Colombo, N., Freppaz, M., Martin, M., Anesio, A.M., 2019. Prokaryotic diversity and distribution in different habitats of an alpine rock glacier-pond system. *Microb. Ecol.* 78 (1), 70–84. <https://doi.org/10.1007/s00248-018-1272-3>.
- Mania, I., Pellicciaro, M., Gorra, R., 2021. Insights into the microbial autotrophic potential of a shallow oligotrophic alpine pond. *Mar. Freshw. Res.* 72 (6), 899–903. <https://doi.org/10.1071/MF20241>.
- Marchina, C., Lencioni, V., Paoli, F., Rizzo, M., Bianchini, G., 2020. Headwaters' isotopic signature as a tracer of stream origins and climatic anomalies: evidence from the Italian Alps in summer 2018. *Water* 12 (2), 390. <https://doi.org/10.3390/w12020390>.
- Marty, C., Tilg, A.-M., Jonas, T., 2017. Recent evidence of large scale receding snow water equivalents in the European Alps. *J. Hydrometeorol.* 18, 1021–1031. <https://doi.org/10.1175/JHM-D-16-0188.1>.
- Matiu, M., Crespi, A., Bertoldi, G., Carmagnola, C.M., Marty, C., Morin, S., Schöner, W., Cat Berro, D., Chiogna, G., De Gregorio, L., Kotlarski, S., Majone, B., Resch, G., Terzagio, S., Valt, M., Beozzo, W., Cianfarra, P., Gouttevin, I., Marcolini, G.,

- Notarnicola, C., Petitta, M., Scherrer, S.C., Strasser, U., Winkler, M., Zebisch, M., Cicogna, A., Cremonini, R., Debernardi, A., Falletto, M., Gaddo, M., Giovannini, L., Mercalli, L., Soubeyrou, J.-M., Sušnik, A., Trenti, A., Urbani, S., Weigluni, V., 2021. Observed snow depth trends in the European Alps: 1971 to 2019. *Cryosphere* 15, 1343–1382. <https://doi.org/10.5194/tc-15-1343-2021>.
- Mladenov, N., Lipez-Ramos, J., McKnight, D.M., Rechea, L., 2009. Alpine lake optical properties as sentinels of dust deposition and global change. *Limnol. Oceanogr.* 54, 2386–2400. https://doi.org/10.4319/lo.2009.54.6_part.2.2386.
- Mulholland, P.J., Thomas, S.A., Valett, H.M., Webster, J.R., Beaulieu, J., 2006. Effects of light on NO₃ uptake in small forested streams: diurnal and day-to-day variations. *J. N. Am. Benthol. Soc.* 25, 583–595. [https://doi.org/10.1899/0887-3593\(2006\)25\[583:EOLONU\]2.0.CO;2](https://doi.org/10.1899/0887-3593(2006)25[583:EOLONU]2.0.CO;2).
- Myrsten, M., Jonsson, A., Bergstrom, A.K., 2016. The effects of temperature and resource availability on denitrification and relative N₂O production in boreal lake sediments. *J. Environ. Sci.* 47, 82–90. <https://doi.org/10.1016/j.jes.2016.03.003>.
- Nemergut, D.R., Costello, E.K., Meyer, A.F., Pescador, M.Y., Weintraub, M.N., Schmidt, S. K., 2005. Structure and function of alpine and arctic soil microbial communities. *Res. Microbiol.* 156, 775–784. <https://doi.org/10.1016/j.resmic.2005.03.004>.
- Oleksy, I.A., Baron, J.S., Beck, W.S., 2021. Nutrients and warming alter mountain lake benthic algal structure and function. *Freshw. Sci.* 40 (1), 88–102. <https://doi.org/10.1086/713068>.
- Palacin-Lizarbe, C., Camarero, L., Catalan, J., 2018. Denitrification temperature dependence in remote, cold, and N-poor lake sediments. *Water Resour. Res.* 54, 1161–1173. <https://doi.org/10.1002/2017WR021680>.
- Pellerin, B.A., Saraceno, J.F., Shanley, J.B., Sebestyen, S.D., Aiken, G.R., Wollheim, W. M., Bergamaschi, B.A., 2012. Taking the pulse of snowmelt: in situ sensors reveal seasonal, event and diurnal patterns of nitrate and dissolved organic matter variability in an upland forest stream. *Biogeochemistry* 108, 183–198. <https://doi.org/10.1007/s10533-011-9589-8>.
- Piña-Ochoa, E., Alvarez-Cobelas, M., 2006. Denitrification in aquatic environments: A cross-system analysis. *Biogeochemistry* 81 (1), 111–130. <https://www.jstor.org/stable/20456415>.
- R Core Team, 2022. *R: A Language and Environment for Statistical Computing*. R Foundation for Statistical Computing, Vienna, Austria URL.
- Roberts, B.J., Mulholland, P.J., 2007. In-stream biotic control on nutrient biogeochemistry in a forested stream, West Fork of Walker Branch. *J. Geophys. Res.* Biogeosci. 112, G04002. <https://doi.org/10.1029/2007JG000422>.
- Rogora, M., Aresè, C., Balestrini, R., Marchetto, A., 2008. Climate control on sulphate and nitrate concentrations in alpine streams of Northern Italy along a nitrogen saturation gradient. *Hydrol. Earth Syst. Sci.* 12 (2), 371–381. <https://doi.org/10.5194/hess-12-371-2008>.
- Rogora, M., Arisci, S., Marchetto, A., 2012. The role of nitrogen deposition in the recent nitrate decline in lakes and rivers in Northern Italy. *Sci. Total Environ.* 417–418, 214–223. <https://doi.org/10.1016/j.scitotenv.2011.12.067>.
- Rogora, M., Somaschini, L., Marchetto, A., Mosello, R., Tartari, G.A., Paro, L., 2020. Decadal trends in water chemistry of Alpine lakes in calcareous catchments driven by climate change. *Sci. Total Environ.* 708, 135180. <https://doi.org/10.1016/j.scitotenv.2019.135180>.
- Rusjan, B., Mikoš, M., 2010. Seasonal variability of diurnal instream nitrate concentration oscillations under hydrologically stable conditions. *Biogeochemistry* 97, 123–140. <https://doi.org/10.1007/s10533-009-9361-5>.
- Salerno, F., Thakuri, S., Guyennon, N., Viviano, G., Tartari, G., 2016a. Glacier melting and precipitation trends detected by surface area changes in Himalayan ponds. *Cryosphere* 10 (4), 1433–1448. <https://doi.org/10.5194/tc-10-1433-2016>.
- Salerno, F., Rogora, M., Balestrini, R., Lami, A., Tartari, G.A., Thakuri, S., Godone, D., Freppaz, M., Tartari, G., 2016b. Glacier melting increases the solute concentrations of Himalayan glacial lakes. *Environ. Sci. Tech.* 50 (17), 9150–9160. <https://doi.org/10.1021/acs.est.6b02735>.
- Sambuell, L., Colombo N., Giardino M., Godone D., 2015. A Waterborne GPR Survey to Estimate Fine Sediments Volume and Find Optimum Core Location in a Rockglacier Lake. Near Surface Geoscience 2015. In: 21st European Meeting of Environmental and Engineering Geophysics. 10.3997/2214-4609.201413826.
- Scherrer, S.C., Fischer, E.M., Posselt, R., Liniger, M.A., Croci-Maspoli, M., Knutti, R., 2016. Emerging trends in heavy precipitation and hot temperature extremes in Switzerland. *J. Geophys. Res. Atmos.* 121, 2626–2637. <https://doi.org/10.1002/2015JD024634>.
- Schlesinger, W.H., 1997. *Biogeochemistry: An Analysis of Global Change*, third ed. Academic Press, San Diego.
- Sebestyen, S.D., Boyer, E.W., Shanley, J.B., Kendall, C., Doctor, D.H., Aiken, G.R., Ohte, N., 2008. Sources, transformations, and hydrological processes that control stream nitrate and dissolved organic matter concentrations during snowmelt in an upland forest. *Water Resour. Res.* 44 (12) <https://doi.org/10.1029/2008WR006983>.
- Seitzinger, S.P., 1988. Denitrification in fresh-water and coastal marine ecosystems: Ecological and geochemical significance. *Limnol. Oceanogr.* 33 (4), 702–724. <https://doi.org/10.4319/lo.1988.33.4part2.0702>.
- Seitzinger, S., Harrison, J.A., Böhlke, J.K., Bouwman, A.F., Lowrance, R., Peterson, B., Tobias, C., Van Drecht, G., 2006. Denitrification across landscapes and waterscapes: a synthesis. *Ecol. Appl.* 16, 2064–2090. [https://doi.org/10.1890/1051-0761\(2006\)016\[2064:DALAWA\]2.0.CO;2](https://doi.org/10.1890/1051-0761(2006)016[2064:DALAWA]2.0.CO;2).
- Sickman, J.O., Leydecker, A., Chang, C.C.Y., Kendall, C., Melack, J.M., Lucero, D.M., Schimel, J., 2003. Mechanisms underlying export of N from high-elevation catchments during seasonal transition. *Biogeochemistry* 64 (1), 1–24. <https://doi.org/10.1023/A:1024928317057>.
- Slemmons, K.E., Rodgers, M.L., Stone, J.R., Saros, J.E., 2017. Nitrogen subsidies in glacial meltwaters have altered planktonic diatom communities in lakes of the US Rocky Mountains for at least a century. *Hydrobiologia* 800 (1), 129–144. <https://doi.org/10.1007/s10750-017-3187-2>.
- Snazelle TT, 2015. Results from laboratory and field testing of nitrate measuring spectrophotometers: U.S. Geological Survey Open-File Report 2015–1065, 35p. doi: 10.3133/ofr20151065.
- Spinoni, J., Vogt, J.V., Naumann, G., Barbosa, P., Dosio, A., 2018. Will drought events become more frequent and severe in Europe? *Int. J. Climatol.* 38 (4), 1718–1736. <https://doi.org/10.1002/joc.5291>.
- Tartari, G., Salerno, F., Buraschi, E., Bruccoleri, G., Smiraglia, C., 2008. Lake surface area variations in the North-Eastern sector of Sagarmatha National Park (Nepal) at the end of the 20th Century by comparison of historical maps. *J. Limnol.* 67 (2), 139. <https://doi.org/10.4081/jlimnol.2008.139>.
- Tolotti, M., Forsström, L., Morabito, G., Thaler, B., Stoyneva, M., Cantonati, M., Šiško, M., Lotter, A., 2009. Biogeographical characterisation of phytoplankton assemblages in high altitude, and high latitude European lakes. *Adv. Limnol.* 62, 55–75. <https://doi.org/10.1127/advlim/62/2009/55>.
- Vila-Costa, M., Pulido, C., Chappuis, E., Calviño, A., Casamayor, E.O., Gacia, E., 2016. Macrophyte landscape modulates lake ecosystem level nitrogen losses through tightly coupled plant-microbe interactions. *Limnol. Oceanogr.* 61, 78–88. <https://doi.org/10.1002/lno.10209>.
- Vione, D., Colombo, N., Said-Pullicino, D., Bocchiola, D., Confortola, G., Salerno, F., Viviano, G., Fratianni, S., Martin, M., Godone, D., Freppaz, M., 2021. Seasonal variations in the optical characteristics of dissolved organic matter in glacial pond water. *Sci. Total Environ.* 759, 143464. <https://doi.org/10.1016/j.scitotenv.2020.143464>.
- Viviroli, D., Dürr, H.H., Messerli, B., Meybeck, M., Weingartner, R., 2007. Mountains of the world, water towers for humanity: Typology, mapping, and global significance. *Water Resour. Res.* 43, 1–13. <https://doi.org/10.1029/2006WR005653>.
- Viviroli, D., Kumm, M., Meybeck, M., Kallio, M., Wada, Y., 2020. Increasing dependence of lowland populations on mountain water resources. *Nat. Sustain.* 3, 917–928. <https://doi.org/10.1038/s41893-020-0559-9>.
- Williams, M.W., Losleben, M.V., Hamann, H.B., 2002. Alpine areas in the Colorado Front Range as monitors of climate change and ecosystem response. *Geogr. Rev.* 92 (2), 180–191. <https://doi.org/10.2307/4140969>.
- Williams, M.W., Knauf, M., Cory, R., Caine, N., Liu, F., 2007. Nitrate content and potential microbial signature of rock glacier outflow, Colorado Front Range. *Earth Surf. Proc. Land.* 32, 1032–1047. <https://doi.org/10.1002/esp.1455>.
- Williams, M.W., Seastedt, T.R., Bowman, W.D., McKnight, D.M., Suding, K.N., 2015a. An overview of research from a high elevation landscape: the Niwot Ridge, Colorado long term ecological research programme. *Plant Ecol. Divers.* 8 (5–6), 597–605. <https://doi.org/10.1080/17550874.2015.1123320>.
- Williams, M.W., Hood, E., Molotch, N.P., Caine, N., Cowie, R., Liu, F., 2015b. The 'teflon basin' myth: hydrology and hydrochemistry of a seasonally snow-covered catchment. *Plant Ecol. Divers.* 8 (5–6), 639–661. <https://doi.org/10.1080/17550874.2015.1123318>.