




## Article

# Quali–Quantitative Fingerprinting of the Fruit Extract of *Uapaca bojeri* Bail. (*Euphorbiaceae*) and Its Antioxidant, Analgesic, Anti-Inflammatory, and Antihyperglycemic Effects: An Example of Biodiversity Conservation and Sustainable Use of Natural Resources in Madagascar

Finiavana Mihary Valisoa Rakotonirina <sup>1,2</sup>, Dario Donno <sup>3,\*</sup>, Zoarilala Rinah Razafindrakoto <sup>4</sup>, Nantenaina Tombozara <sup>4</sup>, Roger Marie Rafanomezantsoa <sup>2</sup>, Charles Andrianjara <sup>4</sup>, David Ramanitrahambola <sup>4,5</sup> and Gabriele Loris Beccaro <sup>3</sup>

- <sup>1</sup> Centre Hospitalier Universitaire Andrainjato Fianarantsoa, Faculté de Médecine, Université de Fianarantsoa, Antananarivo 101, Madagascar
  - <sup>2</sup> Ecole Doctorale de Géochimie et Chimie Médicinale, Université de Fianarantsoa, Antananarivo 101, Madagascar
  - <sup>3</sup> Dipartimento di Scienze Agrarie, Forestali e Alimentari, Università degli Studi di Torino, 10095 Grugliasco, Italy
  - <sup>4</sup> Institut Malgache de Recherches Appliquées, Antananarivo 101, Madagascar
  - <sup>5</sup> Mention Pharmacie, Faculté de Médecine, Université d'Antananarivo, Antananarivo 101, Madagascar
- \* Correspondence: dario.donno@unit.it; Tel.: +39-0116-708751



**Citation:** Rakotonirina, F.M.V.; Donno, D.; Razafindrakoto, Z.R.; Tombozara, N.; Rafanomezantsoa, R.M.; Andrianjara, C.; Ramanitrahambola, D.; Beccaro, G.L. Quali–Quantitative Fingerprinting of the Fruit Extract of *Uapaca bojeri* Bail. (*Euphorbiaceae*) and Its Antioxidant, Analgesic, Anti-Inflammatory, and Antihyperglycemic Effects: An Example of Biodiversity Conservation and Sustainable Use of Natural Resources in Madagascar. *Plants* **2023**, *12*, 475. <https://doi.org/10.3390/plants12030475>

Academic Editors: Jana Šic Žlabur, Sanja Radman and Martina Skendrović Babojelić

Received: 24 November 2022

Revised: 16 January 2023

Accepted: 17 January 2023

Published: 19 January 2023



**Copyright:** © 2023 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

**Abstract:** Antioxidants are important supplements for the human body for their roles in human life for the maintenance of homeostasis. *Uapaca bojeri* fruits are used by the riverain population of the *Uapaca* forests in Madagascar as complementary foods. This study aims to quantify the main antioxidants in the *U. bojeri* fruits to verify their contribution to the enhancement of their anti-inflammatory and antihyperglycemic effects. Standard phytochemical screening was used for qualitative analysis, while spectrophotometric (TPC, TAC, and TFC) and chromatographic analyses (HPLC) were used to quantify several phytochemicals in *U. bojeri* fruits. The antioxidant activity was evaluated using DPPH and FRAP assays. The writhing test was used for the analgesic effects, the carrageenan-induced paw edema was used for the anti-inflammatory activity, and OGTT was used to test the anti-hyperglycemia property of the MEUB in mice. Several phytochemicals were detected and quantified in the fruits, including succinic acid (67.73%) as the main quantified compound. Fruits exerted a good antioxidant capacity and showed analgesic, anti-inflammatory, and antihyperglycemic activities in mice. Isolation of the bioactive compounds should be carried out to confirm these pharmacological properties and develop health-promoting food products or medicinal applications derived from this species.

**Keywords:** *Uapaca bojeri*; phytochemicals; antioxidant capacity; analgesic activity; anti-inflammatory activity; antidiabetic activity; endemism; Madagascar

## 1. Introduction

Antioxidant compounds play important roles in human life for the maintenance of homeostasis. The ingestion of exogenous antioxidants could help our organism to stabilize reactive oxygen species (ROS), which are harmful to cells and tissues [1,2]. An unbalance between ROS and natural antioxidants is the main cause of oxidative stress; in this case, ROS, mainly produced by the mitochondria in cells, cause several problems to other cells and tissues [3,4]. ROS are often composed of free radical atoms, molecules, or ions with unpaired electrons that are extremely unstable and susceptible to active chemical reactions with other molecules. Oxidative stress impairs several endothelial functions, vascular

smooth muscle, and adventitial cells involved in several cardiovascular disorders, including hypertension, atherosclerosis, hypercholesterolemia, and diabetes (Cai and Harrison, 2000; Harrison et al., 2003) [5,6]. Moreover, during their respiration, leukocytes and activated macrophages produce free radicals, which damage epithelial and stromal cells leading to carcinogenesis by modifying targets and pathways essential for normal tissue homeostasis (Hussain et al., 2003) [7]. Antioxidant supplements are necessary in the case of pathologic excessive production of ROS. These compounds are often molecules possessing labile hydrogen that may interact with the ROS and produce stable radicals (e.g., phenolic and organic acid compounds). These compounds can be extracted from edible plants, including fruits and vegetables used in diets [8–10]. Moreover, alternative treatment with fruits showed a growing interest in a broad spectrum of diseases, including diabetes, obesity, neurodegenerative, gastric, inflammation and cardiovascular disorders, and certain types of cancers [11–15].

Tapia, also known by its botanical name *Uapaca bojeri* Bail. (*Euphorbiaceae*), is an endemic tree species from the Tapia forests of Madagascar (Figure 1) localized in the Imamo massif, near Arivonimamo and Miarinarivo, the Hill of Tapia in the region of Manandona between Antsirabe and Ambositra, the Itremo massif in the west of Ambatofinandrahana, and the Isalo massif near Ranohira [16]. The Tapia fruits have a drupe shape with a fleshy, sweet, sticky mesocarp; woody endocarp protecting the three seeds at maturity. The Tapia produces large quantities of small, juicy, oval, and edible fruits, called Voapaka or Voatapia in the Malagasy language. Its flowering period is between March and September, while its fruiting period is from mid-September to early December [17]. This species is known as a medicinal plant and possesses several health-promoting properties. The local population uses this plant in the treatment of diabetes, infectious diseases, and hypertension [18], and the rape fruits are used as a complementary food and source of income for the riverain population during the fructification period. Recently, several compounds have been quantified from the *U. bojeri* leaves and the stems, including carotenoids, organic acids, and phenolics which contribute to the antioxidant, antidiabetic, and anti-inflammatory properties of this species [19]. Leaves, stems, and fruits are the main storage of secondary metabolites in plants. Nowadays, the trend in the research of natural treatments is growing due to the side effects caused by long-term treatment with synthetic drugs [20–23]. As discussed in the study of Razafindrakoto et al. [19], the quantified antioxidant secondary metabolites in the methanol extracts of the *U. bojeri* leaves and stems contribute to the antidiabetic, analgic, and anti-inflammatory activities of this species. However, few studies are reported in the literature on the phytochemical and biological characterization of *U. bojeri* fruits. For this reason, the main aim of this study was to identify and quantify the main bioactive compounds susceptible as antioxidants in the methanol extract of the *U. bojeri* fruits to verify their contribution to the enhancement of their potential therapeutic effects toward inflammatory diseases and diabetes.




Figure 1. Tapia forests distribution in Madagascar.

## 2. Results and Discussion

### 2.1. Phytochemicals and Antioxidant Properties

The methanol maceration produced an extract of *U. bojeri* fruits (MEUB) with a yield of 12.45% relative to the fresh powder. Classes of phytochemicals detected in the fruits of *U. bojeri* were reported in Table 1.

Table 1. Classes of secondary metabolites in the fruit of *U. bojeri*.

Class of Secondary Metabolites	<i>U. Bojeri</i> Fruits
Flavonoids	+
Anthocyanins	+
Leucoanthocyanins	+
Phenolic compounds	+
Tannins	+
Terpenoids	–
Alkaloids	–
Steroids	+
Lactonic steroids	–
Unsaturated sterols	+
Cardiac glycosides	–
Quinones	–
Polysaccharides	–
Saponins	+

+ as the presence of phytochemicals; – as the absence of phytochemicals.

Most of the main antioxidant phytochemicals, including flavonoids, anthocyanins, phenolics, and tannins, were identified in the fruits. These phytochemicals were quantitatively determined as total flavonoid content (TFC), total phenolic content (TPC), and total anthocyanin content (TAC). They showed values of  $118.05 \pm 31.05$  mg<sub>QE</sub>/100 g fresh weight (FW),  $883.60 \pm 186.62$  mg<sub>GAE</sub>/100 g FW, and  $423.51 \pm 14.89$  mg<sub>C3GE</sub>/100 g FW (Table 2), respectively.

**Table 2.** TPC, TFC, TFC, and FRAP result from the fruit of *U. bojeri* and the IC<sub>50</sub> of MEUB on the radical DPPH.

Assay/Sample	Fruit Extract	Gallic Acid
TPC (mgGAE/100 g FW)	$883.60 \pm 186.62$	-
TAC (mgC3GE/100 g FW)	$423.51 \pm 14.89$	-
TFC (mgQE/100 g FW)	$118.05 \pm 31.05$	-
FRAP (mmol Fe <sup>2+</sup> /kg of FW)	$39.72 \pm 0.34$	-
DPPH: IC <sub>50</sub> (μg/mL)	$301.85 \pm 3.62$ <sup>a</sup>	$20.73 \pm 1.35$

Values are expressed as mean  $\pm$  SD ( $n = 3$ ); a:  $p < 0.05$  vs. gallic acid (student *t*-test); GAE: gallic acid equivalent; C3GE: cyanidin-3-glycoside equivalent; QE: quercetin equivalent.

The amounts of the main bioactive compounds are reported in Table 3. Among the 28 biomarkers considered in this study, 14 molecules were detected and quantified; succinic acid was the main compound, with a percentage of 67.73% in relation to the total bioactive compound content (TBCC) that was represented by the sum of all the quantified compounds. Organic acids were the main class of detected phytochemicals, with a percentage of 79.56% in relation to TBCC, in particular succinic, quinic, and oxalic acids. Succinic acid was also the main compound quantified in leaves and the stems with values of  $533.74 \pm 340.08$  and  $1275.65 \pm 434.99$  mg/100 g dry weight, respectively [19], showing that this plant may be considered a good source of succinic acid for the growing interest in pharmaceutical, agricultural, food, and chemical industry applications [24]. Cinnamic acids were represented by caffeic and chlorogenic acids (0.04% and 1.09%, respectively); moreover, four flavonols were detected, including hyperoside (0.03%), isoquercitrin (0.05%), quercetin (0.73%), and quercitrin (3.57%). Ellagic acid (8.34%) represented the class of benzoic acids, and epicatechin (0.84%) represented the class of catechins, while castalagin (4.20%) and vescalagin (0.94%) were detected in the class of tannins. Vitamin C (0.60%) was also identified and quantified in the samples. Most of these quantified compounds show labile hydrogens and could stabilize ROS.

**Table 3.** Phytochemical contents in the fruit of *U. bojeri*.

Class	Standards	Rt (min)	λ (nm)	Value (mg/100 g FW)	Rate <sup>a</sup> (%)
Cinnamic acids	Caffeic acid	4.54	330	$0.71 \pm 0.13$	0.04
	Chlorogenic acid	3.89	330	$19.18 \pm 0.38$	1.09
	Coumaric acid	6.74	330	ND	0
	Ferulic acid	7.99	330	ND	0
	Hyperoside	10.89	330	$0.59 \pm 0.23$	0.03
Flavonols	Isoquercitrin	11.24	330	$0.90 \pm 0.06$	0.05
	Quercetin	17.67	330	$13.08 \pm 2.62$	0.74
	Quercitrin	13.28	330	$62.96 \pm 18.78$	3.57
	Rutin	12.95	330	ND	0
Benzoic acids	Ellagic acid	18.65	280	$147.14 \pm 6.53$	8.34
	Gallic acid	4.26	280	ND	0
Catechins	Catechin	10.31	280	ND	0
	Epicatechin	14.30	280	$14.86 \pm 2.91$	0.84
Tannins	Castalagin	16.35	280	$74.02 \pm 4.21$	4.20
	Vescalagin	17.25	280	$16.65 \pm 7.62$	0.94

Table 3. Cont.

Class	Standards	Rt (min)	$\lambda$ (nm)	Value (mg/100 g FW)	Rate <sup>a</sup> (%)
Vitamins	Ascorbic acid	4.14	261	ND	0
	Dehydroascorbic acid	3.41	348	10.50 ± 3.22	0.60
	Citric acid	5.30	214	ND	0
	Malic acid	4.04	214	ND	0
Organic acids	Oxalic acid	7.85	214	70.49 ± 17.17	4.00
	Quinic acid	3.21	214	138.03 ± 23.02	7.83
	Succinic acid	3.46	214	1194.38 ± 269.07	67.73
	Tartaric acid	5.69	214	ND	0
	Limonene	3.35	250	ND	0
Monoterpenes	Phellandrene	3.57	210	ND	0
	Sabinene	3.45	220	ND	0
	$\gamma$ -terpinene	3.28	235	ND	0
	Terpinolene	4.83	220	ND	0
TBCC				1763.50 ± 355.95	100

Values are expressed as mean ± SD of the amount of phytochemicals ( $n = 3$ ). Rt: Retention time;  $\lambda$ : Wavelength; ND: Not detected; a: Relative to TBCC (Total bioactive compound content).

The antioxidant capacity of the *U. bojeri* fruits was evaluated using the free radical DPPH assay on the MEUB and confirmed with the FRAP assay. The DPPH assay is based on the ability of the compounds in the extracts to donate hydrogen to the radical DPPH (used as the ROS model) and form stable DPPH-H and stable radicals, while the FRAP assay is based on the capacity of the compounds in the plants to give electrons to the ferric ions and form ferrous ions and stable radicals [25]. The results are reported in Table 2. The fruit extracts averagely inhibited the radical DPPH with an IC<sub>50</sub> of 301.85 ± 3.62 µg/mL in relation to the gallic acid ( $p < 0.05$ ), higher values than leaf (47.36 ± 3.00 µg/mL;  $p < 0.05$ ) and stem (33.32 ± 0.69 µg/mL;  $p < 0.05$ ) extracts [19].

The FRAP results showed that the fruits were two-fold less active than the leaves and stems. Effectively, its capacity to reduce Fe<sup>3+</sup>-TPTZ into Fe<sup>2+</sup>-TPTZ is two-fold less important (39.72 ± 0.34 mmol Fe<sup>2+</sup>/kg FW) than that of leaves (69.20 ± 1.41 mmol DW [19]) or stems (70.17 ± 9.53 mmol Fe<sup>2+</sup>/kg DW [19]). It may be due to the TPC content. Indeed, fruits contain less TPC than leaves (3624.72 ± 268.07 mg<sub>GAE</sub>/100 g FW) or stems (5854.17 ± 1247.67 mg<sub>GAE</sub>/100 g FW). Moreover, several studies showed the relationship between the phenolics, organic acids, flavonoids, and vitamin content and the antioxidant capacity that could influence the AOC values in several plant materials [16,19,26,27].

## 2.2. Analgesic Effect of MEUB

The analgesic activity of MEUB was evaluated using the writhing method, one of the most common peripheral analgesic animal models for the screening of analgesic drugs [28]. The number of writhing and the percentage of inhibition exerted by the animals after the treatment with MEUB were reported in Table 4. The intraperitoneal injection of 1% acetic acid solution into the control group caused 24.6 ± 4.3 writhing between the intervals of 25 min after the 5th min of acetic acid induction. The treatment with MEUB at each dose significantly decreased the number of writhing ( $p < 0.01$ ) in a dose-dependent manner ( $p < 0.001$  with ANOVA). The paracetamol at the dose of 100 mg/kg inhibited the pain caused by the acetic acid induction at 82.11%, showing the effectiveness of the protocol. MEUB at the dose of 200 and 400 mg/kg exerted similar analgesic activity when compared to the paracetamol 100 mg/kg ( $p > 0.05$ ). The peripheral analgesic effect may be mediated by the inhibition of cyclo-oxygenases and/or lipoxygenases (and other inflammatory mediators), while the central analgesic action may be mediated by the inhibition of central pain receptors. Several classes of phytochemicals identified in this species exerted these properties (e.g., steroids, flavonoids such as quercetin and quercitrin, and tannins such as castalagin and vescalgin) [29–33], showing their contribution to the pain reduction effect of the MEUB. The pain inhibition effect of the ripe fruit is better than the effect of leaves



(49.40% vs. 70.73% at the dose of 400 mg/kg) and stems (57.43 vs. 70.73% at the dose of 400 mg/kg), showing the importance of the other compounds detected in the fruits (e.g., steroids) [19].

**Table 4.** Effects of MEUB on acetic acid-induced mice by *i.p.* route ( $n = 5$ ).

Sample	Dose (mg/kg)	Writhing Number	Inhibition (%)
Vehicle	-	24.6 ± 4.3	-
Paracetamol	100	4.4 ± 0.7 <sup>a</sup>	82.11
	100	8.6 ± 1.3 <sup>a,b</sup>	65.04
MEUB	200	7.4 ± 1.2 <sup>a</sup>	69.92
	400	7.0 ± 0.9 <sup>a</sup>	71.54

Writhing numbers were expressed as mean ± SEM; a:  $p < 0.01$  vs. vehicle; b:  $p < 0.05$  vs. paracetamol. Student *t*-test and ANOVA, followed by Tukey posthoc test, were performed.

### 2.3. Anti-Inflammatory Effect of MEUB

The anti-inflammatory effect of the MEUB was evaluated using carrageenan-induced mice paw edema, one of the most used *in vivo* protocols for the evaluation of the anti-inflammatory activity of natural products [33]. The injection of carrageenan (2%) caused inflammation at the mice's paw level that was progressively reduced after the treatment. The inflammatory response was detected in two phases; the first phase occurred between the 0 and 120th min post carrageenan injection due to the release of histamine, serotonin, and bradykinin. These mediators increased vascular permeability in the surrounding damaged tissues [34]. The second phase occurred after the 120th min and was identified together with the biosynthesis of prostaglandins and infiltration of neutrophils [35]. The treatment with MEUB at the dose of 100, 200, and 400 mg/kg significantly reduced the inflammation at 60, 120, 180, and 240 min ( $p < 0.05$ ,  $p < 0.01$ , and  $p < 0.001$ ; Table 5) compared with the negative control. The effect of MEUB was in a dose-dependent manner after 180 and 240 min ( $p < 0.001$ , respectively). The effect of the MEUB was comparable to the effect of the indomethacin at the dose of 10 mg/kg at the 120th, 180th, and 240th min post carrageenan injection ( $p > 0.05$ ), showing that the MEUB was more effective during the second phase (however, the inhibition of the first phase should not be unnoticed during the release of prostaglandins). These observations showed that the secondary metabolites in the MEUB could antagonize the release and/or the activity of the chemical mediators during both phases. Compared to the leaf and stem extracts, as reported by Razafindrakoto et al. [19], the MEUB was more effective at higher doses (200 and 400 mg/kg) after the 60th, 120th, and 180th min carrageenan injection. It may be due to the high rate of succinic acid in the fruit, with a value of 67.73% vs. 32.68% in the leaves and 41.64% in the stems. Succinic acid and derivatives are reported as a potent anti-inflammatory by increasing TNF- $\alpha$  secretion and suppressing IL-6 production in LPS-stimulated cells [36]. Moreover, Chang et al. [37] reported the contribution of vescalagin as an antioxidant in its anti-inflammatory properties; on the other hand, Al-Sayed and Abdel-Daim [31] also demonstrated the antioxidant and anti-inflammatory activity of epicatechin.

**Table 5.** Effects of MEUB on carrageenan-induced mice hind paw edema ( $n = 5$ ).

Sample	Dose (mg/kg)	Inhibition of Inflammation (%)				
		30 min	60 min	120 min	180 min	240 min
Vehicle	-	4.17 ± 2.04	11.50 ± 2.05	15.39 ± 2.88	20.32 ± 3.96	27.77 ± 4.35
Indomethacin	10	34.78 ± 6.00 <sup>b</sup>	59.98 ± 6.00 <sup>a</sup>	74.63 ± 5.83 <sup>a</sup>	88.69 ± 5.02 <sup>a</sup>	91.80 ± 2.02 <sup>a</sup>
	100	15.90 ± 6.89	36.83 ± 4.54 <sup>b</sup>	44.33 ± 6.01 <sup>b</sup>	51.60 ± 5.41 <sup>b</sup>	72.81 ± 8.13 <sup>b</sup>
MEUB	200	10.87 ± 5.11	30.31 ± 9.09	67.00 ± 7.55 <sup>a</sup>	81.77 ± 3.25 <sup>a</sup>	91.08 ± 2.98 <sup>a</sup>
	400	15.88 ± 5.81	36.04 ± 6.96 <sup>c</sup>	62.85 ± 11.02 <sup>b</sup>	91.40 ± 8.05 <sup>a</sup>	93.99 ± 5.46 <sup>a</sup>

a:  $p < 0.001$ ; b:  $p < 0.01$ ; c:  $p < 0.05$  vs. vehicle. Student *t*-test and ANOVA, followed by Tukey posthoc test, were performed.

#### 2.4. Antihyperglycemic Activity

The effects of MEUB on the blood glucose level were evaluated using an oral glucose tolerance test (OGTT) in mice. The results are reported in Table 6. As shown in Table 6, the glycemia increased from the administration of glucose (t = 0 min) to the 30th min, then it gradually decreased from the 60th min for each treatment showing that the body releases insulin to regulate the rate of glucose and keep the homeostatic state and demonstrating the effectiveness of the protocol. The glycemia dose-dependently decreased after the 60th, 90th, 120th, and 150th min ( $p < 0.01$ ) glucose loading, and already after 30 min from the glucose administration, mice fed with MEUB at 200 and 400 mg/kg doses already showed lower glucose values compared to those of negative control. Compared to the GBC, after the 60th min glucose loading, MEUB at the dose of 200 mg/kg showed a silent, low effect in terms of variation of glycemia, but the glycemia values did not show any statistical difference compared to GBC; moreover, at the dose of 400 mg/kg, the glycemia variation was higher for MEUB than glycemia variation of GBC, but the glucose level did show any statistical difference. Benzoic acid-related molecules, castalagin, vescalagin, and other antioxidant compounds enhanced insulin effects and reduced insulin resistance [37,38] and the improvement of liver; cardiovascular and metabolic parameters in a rat model of human diseases [39] could be at the origin of these effects on the glycemia decreasing. Additionally, epicatechin, a potent antioxidant that lowers blood pressure [40], plays an important role as insulin-like, contributing to the antihyperglycemic activity of the *U. bojeri* fruits [41].

**Table 6.** Effect of MEUB and GBC on blood glucose level in mice (mg/dL).

Sample (n = 5)	Dose (mg/kg)	Time (min)						
		−60	0	30	60	90	120	150
Negative control	0	102.20 ± 2.37	99.80 ± 2.23	316.80 ± 37.32 0%	285.20 ± 37.04 14.56%	204.20 ± 28.09 51.89%	134.40 ± 6.03 84.06%	122.60 ± 6.24 89.49%
	100	101.60 ± 5.64	93.50 ± 5.52	333.20 ± 26.35 0%	282.20 ± 29.60 <sup>d</sup> 21.27%	202.20 ± 17.43 <sup>d</sup> 54.65%	141.60 ± 7.02 <sup>d</sup> 79.93%	122.60 ± 5.30 <sup>d</sup> 87.86%
MEUB	200	105.00 ± 4.27	101.25 ± 4.17	287.80 ± 22.63 0%	190.80 ± 16.23 <sup>b</sup> 52.00%	133.80 ± 5.02 <sup>c,d</sup> 82.55%	105.80 ± 4.37 <sup>b</sup> 97.56%	97.60 ± 7.07 <sup>c</sup> 101.96%
	400	90.60 ± 7.17	93.75 ± 6.68	274.40 ± 10.81 0%	143.60 ± 15.55 <sup>b</sup> 72.41%	104.40 ± 4.93 <sup>b</sup> 94.10%	95.60 ± 4.46 <sup>b</sup> 98.98%	89.80 ± 8.33 <sup>c</sup> 102.19%
GBC	10	106.00 ± 3.71	103.00 ± 3.52	302.00 ± 20.38 0%	171.20 ± 7.30 <sup>c</sup> 65.73%	101.60 ± 6.50 <sup>b</sup> 100.70%	91.40 ± 5.25 <sup>a</sup> 105.83%	85.20 ± 5.58 <sup>b</sup> 108.94%

The above values are expressed as mean ± SEM of glycemia (n = 5). Below values are the percentage of glycemia variation relative to the glycemia variation at the 30th min. MEUB: Methanol extract of *U. bojeri*; GBC: Glibenclamide a:  $p < 0.001$  vs. negative control; b:  $p < 0.01$  vs. negative control; c:  $p < 0.05$  vs. negative control; d:  $p < 0.05$  vs. GBC. Student *t*-test and ANOVA, followed by Tukey posthoc test, were performed.

### 3. Materials and Methods

#### 3.1. Plant Materials

The ripe fruits of *U. bojeri* were collected in November 2019 in the Imamo forests in the district of Miarinarivo, about 70 km from Antananarivo. A specimen was identified by Mr. Benja Rakotonirina, the botanist of the *Institut Malgache de Recherches Appliquées* (IMRA), and the voucher specimen was compared to the previous reference TN-021/LPA at the IMRA Botanical Department. The ripe fruits were stored in a cool and airy place away from sunlight before being ground.

#### 3.2. Animals

Swiss albino male and female mice (weight: 25 ± 5 g; age: 4–5 months), kept under controlled conditions (12 h dark and 12 h light cycle, 25 ± 2 °C temperature, and 50 ± 10% humidity) at the IMRA animal house, were used. The animals received standard food pellets (1420, Livestock Feed Ltd., Port Louis, Mauritius), and they remained fasting for one night before the experiment. All experiments were carried out following the DIRECTIVE 2010/63/EU and approved by the local ethic committee (number: 06/CEA-IMRA/2020).

### 3.3. Chemicals and Reagents

The chemicals and reagents used during the study are reported in Supplementary Material.

### 3.4. Qualitative Analysis

The classes of secondary metabolites were detected by the traditional methods for phytochemical screening described in the work of Tombozara et al. [42]. Their principle is based on the formation of colored soluble or precipitated compounds by the specific reactive reagent used.

### 3.5. Quantitative Analysis

#### 3.5.1. Total Phenolic Content (TPC), Total Anthocyanin Content (TAC), and Total Flavonoid Content (TFC)

The protocol described by Slinkard and Singleton [43] was used to evaluate the TPC in triplicate using the Folin–Ciocalteu reagent. TAC was determined using the pH differential method in triplicate described in the protocol of Lee et al. [44]. The method is based on the coloration change of the colored monomeric anthocyanin in oxonium form when diluted at pH 1.0 to the colorless hemiketal form at pH 4.5. The aluminum chloride (AlCl<sub>3</sub>) method was used for the determination of the TFC in triplicate, according to Matic et al. [45]. The details are reported in Supplementary Material.

#### 3.5.2. HPLC Analysis

The samples for the HPLC analysis of phytoconstituents were prepared in triplicate according to the method described by Razafindrakoto et al. [19]. The extraction protocol was reported in Supplementary Material. Twenty-eight standards (Table 3) were selected and manually injected (20 µL) in triplicate for their quantification in this study. The quantitation of these compounds was performed using an Agilent 1200 High-Performance Liquid Chromatograph coupled to an Agilent UV-Vis diode array detector (Agilent Technologies, Santa Clara, CA, USA) according to the protocol described by Razafindrakoto et al. [19].

### 3.6. Antioxidant Activity Evaluation

The antioxidant activity of the methanolic extract of *U. bojeri* (MEUB) was determined using the free radical DPPH assay along with the FRAP assay that is based on the capacity of the sample to reduce the ferric ions Fe<sup>3+</sup> into ferrous ions Fe<sup>2+</sup> in the 2,4,6-tripyridyl-s-triazine (TPTZ) complex [46]. Both protocols have been described by Razafindrakoto et al. [19], and they are detailed in Supplementary Material.

### 3.7. Acetic-Acid-Induced Writhing Test

The protocol of Olajide et al. [47], slightly modified by Razafindrakoto et al. [19], was used to determine the analgesic property of MEUB (Supplementary Material).

### 3.8. Carrageenan-Induced in Paw Oedema Test

In vivo anti-inflammatory activity was evaluated based on the inhibition of carrageenan-induced mouse hind paw edema using a plethysmometer as previously described by Buisseret et al. [48], slightly modified by Razafindrakoto et al. [49] (Supplementary Material).

### 3.9. Oral Glucose Tolerance Test (OGTT)

OGTT, described by Tombozara et al. [16] with slight modifications, was applied to determine the hypoglycemia property of MEUB (Supplementary Material).

### 3.10. Statistical Analysis

The results were expressed as mean ± standard error means (S.E.M.), and the data were statistically analyzed using the Student's *t*-test and one-way analysis of variance (ANOVA) followed by the HSD Tukey multiple range test using SPSS 20.0 software. All the differences showing a *p* < 0.05 were accepted as statistically significant.



#### 4. Conclusions

This work was performed to determine the antioxidant compounds in the fruit of *U. bojeri*, consumed by the local people living around the Tapia forests, and their contribution to the management of metabolic diseases, including inflammatory and related diseases and diabetes. Overall, 14 compounds belonging to phenolic, flavonoid, vitamin, and organic acid classes have been quantified in the fruit of *U. bojeri*, contributing to its antioxidant, analgesic, anti-inflammatory, and antihyperglycemic activities. The tested product is a methanolic crude extract (MEIC) of the *U. bojeri* fruit, which is a mixture of more secondary metabolites. Some of them could contribute to the pharmacological activity. In the case of the mixture of active compounds, the synergy of action can be envisaged, but all molecules have their pharmacokinetic profile. It is therefore not entirely obvious to have a dose-dependent response as in the case of an isolated bioactive molecule. This study contributed to the valorization of this species as a health promotor, mostly as complementary medicine in the treatment of diabetes and inflammatory and related diseases, but further studies, including the isolation of the bioactive compounds, should be performed to confirm the properties of this plant. It is very important to investigate the local use of biodiversity to identify new bioactive compounds for supporting biodiversity conservation and sustainable development projects in Madagascar.

**Supplementary Materials:** The following supporting information can be downloaded at: <https://www.mdpi.com/article/10.3390/plants12030475/s1>.

**Author Contributions:** Conceptualization, D.R. and G.L.B.; methodology, F.M.V.R., D.D., Z.R.R. and N.T.; software, D.D. and N.T.; validation, D.D. and D.R.; formal analysis, D.D., Z.R.R. and N.T.; investigation, F.M.V.R., D.D., Z.R.R. and N.T.; resources, C.A. and G.L.B.; data curation, F.M.V.R., Z.R.R. and N.T.; writing—original draft preparation, F.M.V.R. and N.T.; writing—review and editing, D.D., Z.R.R., R.M.R., C.A., D.R. and G.L.B.; visualization, C.A. and G.L.B.; supervision, R.M.R., D.R. and G.L.B.; project administration, D.R. and G.L.B. All authors have read and agreed to the published version of the manuscript.

**Funding:** This research received no external funding.

**Institutional Review Board Statement:** The study was conducted following the DIRECTIVE 2010/63/EU and approved by the local ethic committee of the *Institut Malgache de Recherches Appliquées* (IMRA) (number: 06/CEA-IMRA/2020, approved on 19 November 2020).

**Informed Consent Statement:** Not applicable.

**Data Availability Statement:** Not applicable.

**Conflicts of Interest:** The authors declare no conflict of interest.

#### References

1. Vitaglione, P.; Morisco, F.; Caporaso, N.; Fogliano, V. Dietary Antioxidant Compounds and Liver Health. *Crit. Rev. Food Sci. Nutr.* **2005**, *44*, 575–586. [CrossRef]
2. Gulcin, A. Antioxidants and antioxidant methods: An updated overview. *Arch. Toxicol.* **2020**, *94*, 651–715. [CrossRef]
3. Lenaz, G. The Mitochondrial Production of Reactive Oxygen Species: Mechanisms and Implications in Human Pathology. *IUBMB Life* **2001**, *52*, 159–164. [CrossRef]
4. Chen, Q.; Vazquez, E.J.; Moghaddas, S.; Hoppel, C.L.; Lesnfsky, E.J. Production of Reactive Oxygen Species by Mitochondria: Central role of complex III. *J. Biol. Chem.* **2003**, *278*, 36027–36031. [CrossRef]
5. Cai, H.; Harrison, D.G. Endothelial dysfunction in cardiovascular diseases: The role of oxidant stress. *Circ. Res.* **2000**, *87*, 840–844. [CrossRef]
6. Harrison, D.; Griendling, K.K.; Landmesser, U.; Hornig, B.; Drexler, H. Role of oxidative stress in atherosclerosis. *Am. J. Cardiol.* **2003**, *91*, 7–11. [CrossRef]
7. Hussain, S.; Hofseth, L.; Harris, C. Radical causes of cancer. *Nat. Rev. Cancer* **2003**, *3*, 276–285. [CrossRef]
8. Kaur, C.; Kapoor, H.C. Antioxidants in fruits and vegetables—The millennium’s health. *Int. J. Food Sci. Technol.* **2001**, *36*, 703–725. [CrossRef]
9. Donno, D.; Mellano, M.G.; Riondato, I.; De Biaggi, M.; Andriamaniraka, H.; Gamba, G.; Beccaro, G.L. Traditional and Unconventional Dried Fruit Snacks as a Source of Health-Promoting Compounds. *Antioxidants* **2019**, *8*, 396. [CrossRef]

10. Kim, S. Antioxidant Compounds for the Inhibition of Enzymatic Browning by Polyphenol Oxidases in the Fruiting Body Extract of the Edible Mushroom *Hericium. Erinaceus*. *Foods* **2020**, *9*, 951. [[CrossRef](#)]
11. Afrin, S.; Giampieri, F.; Gasparrini, M.; Forbes-Hernandez, T.Y.; Varela-López, A.; Quiles, J.L.; Mezzetti, B.; Battino, M. Chemopreventive and Therapeutic Effects of Edible Berries: A Focus on Colon Cancer Prevention and Treatment. *Molecules* **2016**, *21*, 169. [[CrossRef](#)]
12. Forino, M.; Tartaglione, L.; Dell'Aversano, C.; Ciminiello, P. NMR-based identification of the phenolic profile of fruits of *Lycium barbarum* (goji berries). Isolation and structural determination of a novel N-feruloyl tyramine dimer as the most abundant antioxidant polyphenol of goji berries. *Food Chem.* **2016**, *194*, 1254–1259. [[CrossRef](#)]
13. Musumeci, L.; Maugeri, A.; Cirmi, S.; Lombardo, G.E.; Russo, C.; Gangemi, S.; Calapai, G.; Navarra, M. Citrus fruits and their flavonoids in inflammatory bowel disease: An overview. *Nat. Prod. Res.* **2019**, *34*, 1–15. [[CrossRef](#)]
14. Areche, C.; Hernandez, M.; Cano, T.; Ticona, J.; Cortes, C.; Simirgiotis, M.; Caceres, F.; Borquez, J.; Echeverría, J.; Sepulveda, B. *Corryocactus brevistylus* (K. Schum. ex Vaupel) Britton & Rose (Cactaceae): Antioxidant, gastroprotective effects, and metabolomic profiling by ultrahigh-pressure liquid chromatography and electrospray high resolution orbitrap tandem mass spectrometry. *Front. Pharmacol.* **2020**, *11*, 417. [[CrossRef](#)]
15. Belwal, T.; Bisht, A.; Devkota, H.P.; Ullah, H.; Khan, H.; Pandey, A.; Bhatt, I.D.; Echeverría, J. Phytopharmacology and clinical updates of Berberis species against diabetes and other metabolic diseases. *Front. Pharmacol.* **2020**, *11*, 41. [[CrossRef](#)]
16. Tombozara, N.; Donno, D.; Razafindrakoto, Z.R.; Randriamampionona, D.; Ramanitrahasimbola, D.; Andrianjara, C.; Ramilison-Razafimahefa, R.D.; Rakotondramanana, A.D.; Beccaro, G.L. The first assessment on antioxidant and antidiabetic activities of leaves and stems of *Vaccinium secundiflorum* Hook. (Ericaceae), an endemic plant of Madagascar. *S. Afr. J. Bot.* **2020**, *130*, 422–429. [[CrossRef](#)]
17. Direction des Eaux et Forêts (DEF). *Inventaire Ecologique Forestier National: Recueil Botanique de 200 Espèces Forestières*; Ministère des Eaux et Forêts: Antananarivo, Madagascar, 1996; 503p.
18. Pernet, R.; Meyer, G. *Pharmacopée de Madagascar*; Institut de recherche scientifique de Tananarive-Tsimbazaza: Antananarivo, Madagascar, 1957.
19. Razafindrakoto, Z.R.; Donno, D.; Tombozara, N.; Andriamaniraka, H.; Andrianjara, C.; Ramanitrahasimbola, D.; Beccaro, G.L. Activities of Leaves and Stems of *Uapaca bojeri* Bail. (EUPHORBIACEAE), an Endemic Plant of Madagascar. *Pharmaceuticals* **2020**, *13*, 71. [[CrossRef](#)]
20. Cohen, K.; Weinstein, A.M. Synthetic and Non-synthetic Cannabinoid Drugs and Their Adverse Effects—A Review From Public Health Prospective. *Front. Public Health* **2018**, *6*, 1–8. [[CrossRef](#)]
21. Khursheed, R.; Singh, S.K.; Wadhwa, S.; Kapoor, B.; Gulati, M.; Kumar, R.; Ramanunni, A.K.; Awasthi, A.; Dua, K. Treatment strategies against diabetes: Success so far and challenges ahead. *Eur. J. Pharmacol.* **2019**, *862*, 172625. [[CrossRef](#)]
22. Atkinson, T.J.; Fudin, J. Nonsteroidal Antiinflammatory Drugs for Acute and Chronic Pain. *Phys. Med. Rehabil. Clin.* **2020**, *31*, 219–231. [[CrossRef](#)]
23. Gupta, M.; Singh, N.; Gulati, M.; Gupta, R.; Sudhakar, K.; Kapoor, B. Herbal bioactives in treatment of inflammation: An overview. *S. Afr. J. Bot.* **2021**, *143*, 205–225. [[CrossRef](#)]
24. Pateraki, C.; Almqvist, H.; Ladakis, D.; Liden, G.; Koutinas, A.A.; Vlysidis, A. Modelling succinic acid fermentation using a xylose-based substrate. *Biochem. Eng.* **2016**, *114*, 26–41. [[CrossRef](#)]
25. Prior, R.L.; Wu, X.; Schaich, K. Standardized methods for the determination of antioxidant capacity and phenolics in foods and dietary supplements. *J. Agric. Food Chem.* **2005**, *53*, 4290–4302. [[CrossRef](#)]
26. Abeysuriya, H.I.; Bulugahapitiya, V.P.; Loku Pulukkuttige, J. Total Vitamin C, Ascorbic Acid, Dehydroascorbic Acid, Antioxidant Properties, and Iron Content of Underutilized and Commonly Consumed Fruits in Sri Lanka. *Int. J. Food Sci.* **2020**, *2020*, 4783029. [[CrossRef](#)]
27. Singh, B.; Singh, J.P.; Kaur, A.; Singh, N. Phenolic composition, antioxidant potential and health benefits of citrus peel. *Food Res. Int.* **2020**, *132*, 109114. [[CrossRef](#)]
28. Negus, S.S.; Vanderah, T.W.; Brandt, M.R.; Bilsky, E.J.; Becerra, L.; Borsook, D. Preclinical assessment of candidate analgesic drugs: Recent advances and future challenges. *J. Pharmacol. Experiment. Ther.* **2006**, *319*, 507–514. [[CrossRef](#)] [[PubMed](#)]
29. Viana, G.S.B.; Bandeira, M.A.M.; Moura, L.C.; Souza-Filho, M.V.P.; Matos, F.J.A.; Ribeiro, R.A. Analgesic and Antiinflammatory Effects of the Tannin Fraction from *Myracrodruon urundeuva* Fr. All. *Phytother. Res.* **1997**, *11*, 118–122. [[CrossRef](#)]
30. Afzal, M.; Gupta, G.; Kazmi, I.; Rahman, M.; Afzal, O.; Alam, J.; Hakeem, K.R.; Pravez, M.; Gupta, R.; Anwar, F. Anti-inflammatory and analgesic potential of a novel steroidal derivative from *Bryophyllum pinnatum*. *Fitoterapia* **2012**, *83*, 853–858. [[CrossRef](#)]
31. Al-Sayed, E.; Abdel-Daim, M.M. Analgesic and anti-inflammatory activities of epicatechin gallate from *Bauhinia hookeri*. *Drug Dev. Res.* **2018**, *79*, 157–164. [[CrossRef](#)]
32. Lichota, A.; Gwozdziński, L.; Gwozdziński, K. Therapeutic potential of natural compounds in inflammation and chronic venous insufficiency. *Eur. J. Med. Chem.* **2019**, *176*, 68–91. [[CrossRef](#)]
33. Razafin-drabazo, F.; Donno, D.; Tombozara, N.; Razafindrakoto, Z.R.; Rajaonarison, J.F.; Andrianjara, C.; Ramanitrahasimbola, D.; Beccaro, G.L. Phyto-compounds and pharmacological activities of *Lygodium lanceolatum* Desv. (Schizaeaceae). *S. Afr. J. Bot.* **2020**, *135*, 225–232. [[CrossRef](#)]
34. Zhu, Z.; Ma, K.; Ran, X.; Zhang, H.; Zheng, C.; Han, T. Analgesic, anti-inflammatory and antipyretic activities of the petroleum ether fraction from the ethanol extract of *Desmodium podocarpum*. *J. Ethnopharmacol.* **2021**, *133*, 1126–1131. [[CrossRef](#)] [[PubMed](#)]

35. Tamrat, Y.; Nedi, T.; Assefa, S.; Teklehaymanot, T.; Shibeshi, W. Anti-inflammatory and analgesic activities of solvent fractions of the leaves of *Moringa stenopetala* Bak. (Moringaceae) in mice models. *BMC Complement. Altern. Med.* **2017**, *17*, 473. [[CrossRef](#)]
36. Chien, S.C.; Chen, M.L.; Kuo, H.T.; Tsai, Y.C.; Lin, B.F.; Kuo, Y.H. Anti-inflammatory activities of new succinic and maleic derivatives from the fruiting body of *antrodia camphorate*. *J. Agric. Food Chem.* **2008**, *56*, 7017–7022. [[CrossRef](#)] [[PubMed](#)]
37. Chang, W.C.; Shen, S.C.; Wu, J.S.B. Protective effects of vescalagin from pink wax apple [*Syzygium samarangense* (blume) merrill and perry] fruit against methylglyoxal-induced inflammation and carbohydrate metabolic disorder in rats. *J. Agric. Food Chem.* **2013**, *61*, 7102–7109. [[CrossRef](#)] [[PubMed](#)]
38. Marles, R.J.; Farnsworth, N.R. Antidiabetic plants and their active constituents. *Phytomedicine* **1995**, *2*, 137–189. [[CrossRef](#)]
39. Panchal, S.K.; Brown, L. Cardioprotective and hepatoprotective effects of ellagitannins from European oak bark (*Quercus petraea* L.) extract in rats. *Eur. J. Nutr.* **2013**, *52*, 397–408. [[CrossRef](#)]
40. Iacopini, P.; Baldi, M.; Storch, P.; Sebastiani, L. Catechin, epicatechin, quercetin, rutin and resveratrol in red grape: Content, in vitro antioxidant activity and interactions. *J. Food Compos. Anal.* **2008**, *21*, 589–598. [[CrossRef](#)]
41. Ahmad, F.; Khalid, P.; Khan, M.M.; Rastogi, A.K.; Kidwai, J.R. Insulin like activity in (–) epicatechin. *Acta Diabetol. Lat.* **1989**, *26*, 291–300. [[CrossRef](#)]
42. Tombozara, N.; Razafindrakoto, Z.R.; Ramanitrahasimbola, D.; Razafimahefa-Ramilison, R.D.; Marchioni, E.; Rakotondramanana, A.D. Isolation of the gallic acid in the butanolic fraction of *crassula ovata* (Mill.) Druce (CRASSULACEAE) leaves and its vaso-relaxing effect. *Am. J. Innov. Res. Appl. Sci.* **2017**, *4*, 200–207.
43. Slinkard, K.; Singleton, V.L. Total phenol analysis: Automation and comparison with manual methods. *Am. J. Enol. Vitic.* **1977**, *28*, 49–55.
44. Lee, J.; Durst, R.W.; Wrolstad, R.E. Determination of Total Monomeric Anthocyanin Pigment Content of Fruit Juices, Beverages, Natural Colorants, and Wines by the pH Differential Method: Collaborative Study. *J. AOAC Int.* **2005**, *88*, 1269–1278. [[CrossRef](#)] [[PubMed](#)]
45. Matic, P.; Sabljic, M.; Jakobek, L. Validation of Spectrophotometric Methods for the Determination of Total Polyphenol and Total Flavonoid Content. *J. AOAC Int.* **2017**, *100*, 1795–1803. [[CrossRef](#)] [[PubMed](#)]
46. Benzie, I.F.; Strain, J.J. Ferric reducing/antioxidant power assay: Direct measure of total antioxidant activity of biological fluids and modified version for simultaneous measurement of total antioxidant power and ascorbic acid concentration. *Meth. Enzymol.* **1999**, *299*, 15–27.
47. Olajide, O.A.; Awe, S.O.; Makinde, J.M.; Ekhelar, A.I.; Olusola, A.; Morebise, O.; Okpako, D.T. Studies on the anti-inflammatory, antipyretic and analgesic properties of *Alstonia boonei* stem bark. *J. Ethnopharmacol.* **2000**, *71*, 179–186. [[CrossRef](#)]
48. Buisseret, B.; Guillemot-Legris, O.; Muccioli, G.G.; Alhouayek, M. Prostaglandin D2-glycerol ester decreases carrageenan-induced inflammation and hyperalgesia in mice. *Biochim. Biophys. Acta—Mol. Cell Biol. Lipids* **2019**, *1864*, 609–618. [[CrossRef](#)]
49. Razafindrakoto, Z.R.; Tombozara, N.; Donno, D.; Gamba, G.; Nalimanana, N.R.; Rakotondramanana, A.D.; Andrianjara, C.; Beccaro, G.L.; Ramanitrahasimbola, D. Antioxidant, analgesic, anti-inflammatory and antipyretic properties, and toxicity studies of the aerial parts of *Imperata cylindrica* (L.) Beauv. *S. Afr. J. Bot.* **2021**, *142*, 222–229. [[CrossRef](#)]

**Disclaimer/Publisher’s Note:** The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.