

AperTO - Archivio Istituzionale Open Access dell'Università di Torino

Detection of fentanyl, synthetic opioids, and ketamine in hair specimens from purposive samples of American and Italian populations

This is a pre print version of the following article:

Original Citation:

Availability:

This version is available <http://hdl.handle.net/2318/1945832> since 2023-12-04T11:08:29Z

Published version:

DOI:10.1111/1556-4029.15348

Terms of use:

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)



Detection of fentanyl, synthetic opioids, and ketamine in hair specimens from purposive samples of American and Italian populations

Journal:	<i>Journal of Forensic Sciences</i>
Manuscript ID	JOFS-23-385.R1
Manuscript Type:	Original Paper
Keywords:	hair analysis, synthetic opioids, fentanyl, ketamine, NPS, MDMA

SCHOLARONE™
Manuscripts

Detection of fentanyl, synthetic opioids, and ketamine in hair specimens from purposive samples of American and Italian populations

ABSTRACT

With the current crisis related to the diffusion of fentanyl and other novel opioids in several countries and populations, new and effective approaches are needed to better elucidate the phenomenon. In this context, hair testing offers a unique perspective in the investigation of drug consumption, producing useful information in terms of exposure to psychoactive substances. In this research, we applied targeted ultra-high performance liquid chromatography tandem mass spectrometry (UHPLC-MS/MS) analytical methods to detect novel synthetic and prescription opioids and other common controlled psychoactive drugs in the keratin matrix. A total of 120 hair samples were analyzed from the United States (US) and Italy, segmented when longer than 6 cm, and then analyzed. In the 60 samples (83 segments in total) analyzed from a purposive sample of data collected in [the US](#), fentanyl was detected in 14 cases (16.9%), with no detection of nitazens or bromphine. [We also detected](#) fentanyl metabolites, despropionyl-*p*-fluorofentanyl, and prescription opioids. In the 60 samples collected in Italy (91 segments in total), ketamine was the most prevalent compound detected (in 41 cases; [45.1%](#)), with ketamine demonstrating a strong correlation with detection of amphetamines and MDMA, likely due to co-use of these substances in recreational contexts. Several common drugs were also detected, but no exposure to fentanyl or its analogs were detected. Results of this retrospective exploration of drug use add to increasing evidence that hair testing can serve as a useful adjunct to epidemiology studies that seek to determine biologically confirmed use and exposure in high-risk populations.

Keywords: hair analysis; fentanyl; synthetic opioids; ketamine; [new psychoactive substances](#); [MDMA](#)

Highlights

- Results from real samples are crucial to understand the molecules used by at-risk populations
- In the US, fentanyl was detected in 16.9% of cases, with no detection of nitazens or bropphine
- In Italy, ketamine was the most prevalent compound detected, detected in 45.1% of hair segments
- Detection of ketamine was strongly correlated with detection of amphetamines and MDMA
- Hair analysis is effective in investigating the diffusion of new psychoactive drugs

1. Introduction

In the past decade, illicit opioid use has progressed from nonmedical use of legal analgesic drugs such as hydrocodone, oxycodone, and tramadol, to the diffusion of illicitly manufactured fentanyl and its analogs, often referred to as “fentalogs” (1, 2). In addition, a new class of synthetic opioids referred to as nitazenes has been recently reported in several illegal drug markets (3). Many of these novel synthetic opioids (NSOs) are considered particularly risky due to their high potency and because they are often introduced into the market as cutting/adulterant agents of drugs such as heroin or simply as cheaper substitutes for other drugs (4, 5).

While heroin was the first drug to become frequently replaced or cut with NSO, these compounds began to appear to counterfeit pills representing common prescription drugs (e.g., oxycodone, alprazolam) and other illegal powder drugs such as cocaine (5-10), raising a major health concern for unaware users. In parallel, however, preference for fentanyl has increased among some populations, leaving doubt regarding whether fentanyl is more demand-led or supply-led (11).

However, it is important to note that currently, the synthetic opioid crisis is centered in North America. In 2022, in the United States (US), there were 71,238 deaths linked to use of synthetic opioids such as fentanyl (12), and in 2021, there were at least 10,000 fentanyl seizures in the US which weighed over 10,000 kg in total (13). In Europe, however, among 12 countries providing seizure data to the European Union Early Warning System, in 2021, there were only 187 recorded fentanyl seizures (weighting 5.5 kg in total) (14). Heroin has largely been replaced by fentanyls in the US, but heroin is still the most common illicit opioid in Europe (14). In Italy, an average of 0.74 tons of heroin have been seized annually between 2011 and 2021 (14).

Despite differences in the opioid and other drug landscapes between the US and Europe, new and effective approaches are needed to monitor shifting drug-related phenomena. For example, in the US, in 2021, there was an increase to 24,538 deaths linked to cocaine use and 32,856 deaths linked to other psychostimulant (mainly methamphetamine) use (12), but the vast majority of such cases involve co-use of opioids (15). A greater understanding of co-use (or co-exposure) of illicit fentanyl,

1
2
3 opioids, and other psychoactive drugs can help adapt and improve existing interventions aimed to
4 reduce overdose mortality, together with broad integrated public health strategies based on overdose
5 education and prevention and to support the drugs debate (16-18).
6
7
8
9

10 In this context, hair analysis has proved to be an easy and effective tool to investigate the
11 prevalence of use of psychoactive substances, since the keratin matrix allows for the detection of past
12 drug exposure and for the investigation of the chronological profile of the exposure to one or multiple
13 compounds. Furthermore, hair analysis is now based on multiclass methods for both well-known and
14 emerging compounds, allowing for the investigation of different consumption patterns, including co-
15 use of common drugs (including prescription opioids), as well as occasional vs. frequent NSO use or
16 exposure (19-21). While several papers have described multianalyte screening methods capable of
17 detecting NSOs (22), few have presented results from real samples (23-25). Polydrug use has generally
18 been shown to be common based on the aforementioned studies, involving several psychoactive
19 substances and not only heroin (26, 27). In general, fentanyl has been the most frequently detected
20 compound among the class of fentologs (8, 19), suggesting that it is the most prevalent molecule
21 while the less common analogs tend to be co-used with other drugs and are thus not consumed in
22 isolation. [Other typical matters of current discussion \(in order to provide a definitive interpretation
23 of positive vs. negative results\) are: \(i\) the meaning of quantitative results, in terms of occasional and
24 frequent use or exposure\), and \(ii\) the identification of proper metabolites to discriminate direct
25 exposure from potential external contamination.](#)
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46

47 In this paper, we present [our](#) ultra-high performance liquid chromatography tandem mass
48 spectrometry (UHPLC-MS/MS) analysis of a subset of hair samples collected in the US and in Italy
49 [based on purposive sampling methods](#). All samples were screened for fentologs, prescription opioids,
50 nitazens, bupropion, and other common controlled psychoactive drugs.
51
52
53
54
55
56
57

58 **2. Materials and methods**

59 *2.1. Reagents and standards*

60

All chemicals, including methanol, formic acid and acetonitrile, were purchased from Sigma-Aldrich (Milan, Italy) while ultra-pure water was obtained using a Milli-Q® UF-Plus apparatus (Millipore, Bedford, MA, USA). The analytical standards of the target analytes and deuterated internal standards (norfentanyl-D5, fentanyl-D5 and oxycodone-D6) were purchased from LGC Promochem (Milan, Italy) and Sigma-Aldrich (Milan, Italy) (purity >99%, concentration between 0.1 mg/mL and 1 mg/mL), or kindly provided by the Italian National Early Warning System (provided at a concentration of 0.02 mg/mL). [The list of target analytes is presented in Table 1.](#) All stock standard solutions were prepared in methanol at 1 mg/mL and stored at -20°C until used. Working solution of 42 analytes (identified among the most common synthetic opioids and those recently observed by the warning systems) and internal standard solution were prepared at the final concentration of 1 µg/mL by dilution with methanol.

2.2. Sample collection and preparation

In this study, we focus on two purposive samples of adults—from the US and from Italy. Hair samples were collected in 2022 in the US (60 samples, from an ongoing rapid street reporting surveillance study being conducted throughout various US cities by the National Drug Early Warning System) (28) and in Italy (60 samples, from harm reduction services in Northern Italy), according to international guidelines (29). With regard to the 60 US samples, we focused on samples provided by participants who reported past-12 month use of heroin and/or fentanyl (n=18), 21 participants reporting past-12-month use of at least one novel psychoactive substance (NPS; who did not report heroin or fentanyl use; n=21), and a random sample of 21 participants who did not report heroin, fentanyl, or NPS use (n=21). In order to nullify any further risk related to data sharing and to safeguard the privacy of sample donors, in the US, all samples were collected anonymously. Italian samples were made anonymous by alphanumeric codes and used only in our laboratory. The risk of re-identification was also nullified. Furthermore, subjects provided informed consent to be tested for drug exposure. The study protocol for hair sample collection and testing for US samples was reviewed

1
2
3 and approved by the Institutional Review Board (IRB) of the University of Florida. The study
4
5 protocol for hair sample collection in Italy was approved by the Bioethical Committee of the
6
7 University of Turin.
8
9

10 All samples were analyzed up to a maximum of the proximal 12 cm, since the study aimed to
11
12 explore the intake of drugs in the 12 months prior to collection (assuming a normal hair growth rate
13
14 of 1 cm per month). When hair was ≤ 6 cm, it was analyzed in its entire length. When hair was longer
15
16 than 6 cm (54 samples), two segments were prepared for analysis (with one representing roughly the
17
18 past six months and the other representing roughly the previous 6-12 months). Therefore, a total of
19
20 174 separate segments was analyzed. The targeted screening for common drugs was performed using
21
22 previously published and fully validated methods (30, 31). Existing procedures for novel opioids (32,
23
24 33) were adjusted to expand the panel of screened molecules. A partial validation was performed,
25
26 aimed to verify sensitivity and specificity. The limits of detection are presented in Table S1, while
27
28 data for trueness and precision at three different concentration levels are presented in Table S2.
29
30
31
32

33 All samples were treated with a procedure developed on-purpose for the keratin matrix. About
34
35 50 mg of hair was decontaminated by an initial wash with 1-mL dichloromethane followed by a
36
37 second wash with 1-mL methanol, each one performed under 3 min stirring. The dried hair was
38
39 pulverized using six steel balls stirring in a Precellys[®] homogenizer. The pulverized samples were
40
41 extracted by keeping them immersed in 0.5-mL methanol added with 2.5 μ L of an internal standards
42
43 mixture (final concentration of 0.01 ng/mg) at $+55 \pm 5$ C for 15 h. Lastly, the organic phase was
44
45 collected and an aliquot of 3 μ L was directly injected into the UHPLC–MS-MS system. A calibration
46
47 curve in the range 10-250 pg/mg was also prepared by spiking the proper quantities of analytical
48
49 standards into a blank hair sample.
50
51
52
53
54
55

56 2.3 Instrumentation

57
58 UHPLC separation was performed with a Phenomenex Kinetex C18 column (100 \times 2.1 mm,
59
60 1.7 μ m) maintained at 45 °C on the SCIEX ExionLC[™] AC system. The mobile phases consisted of

1
2
3 water (A) and acetonitrile (B), both with formic acid 5 mM. The LC flow rate was set at 0.5 mL/min
4
5 and the mobile phase eluted under the following linear gradient conditions: (A:B, v:v) isocratic
6
7 elution at 95:5 for 0.5 min, from 95:5 to 5:95 in 7.5 min, isocratic elution at 5:95 for 0.5 min and final
8
9 re-equilibration for 1.5 min to the initial condition. The total run time was 10 min. All analyses were
10
11 performed using a mass spectrometer equipped with a quadrupole trap SCIEX triple Quad™ 7500
12
13 mass spectrometer (Sciex, Darmstadt, Germany) system equipped with an OptiFlow Pro ion source
14
15 with an analytical probe and E Lens. The ionization source was operated with electrospray ionization
16
17 (ESI) in the positive mode. For each transition, compound-specific parameters such as collision
18
19 energy (CE) were also optimized after infusion of the standard solution. A single acquisition method
20
21 was created using the Scheduled MRM algorithm in SCIEX OS software 2.0. Three MRM transitions
22
23 were monitored for each targeted analyte. The full list of the target analytes, the monitored transitions,
24
25 and their instrumental parameter are reported in Table 1.
26
27
28
29
30

31 32 33 2.4 Statistical Analysis

34
35 We used descriptive statistics to describe the number of segments testing positive for various
36
37 drugs in each country, and among positive cases we also described the range of levels of molecules
38
39 detected. Within the Italian sample, we also computed Spearman correlations to determine the extent
40
41 to which level of detection of each drug was correlated.
42
43
44
45

46 47 3. Results and discussion

48 49 50 51 Testing for NSO

52
53 In the 60 samples collected in the US (comprising of 83 segments), at least one opioid was
54
55 detected in 16 segments (19.3%). Fentanyl was detected above the LOD (estimated at 5 pg/mg) in 14
56
57 segments (16.9%). The range of measured concentrations of fentanyl was extremely wide, ranging
58
59 from 13 pg/mg through 7300 pg/mg, with a mean value of 1377 pg/mg and a median of 382 pg/mg.
60

1
2
3 Only 4 segments measured below 100 pg/mg (0.1 ng/mg). In eight segments, external contamination
4 was excluded because the metabolite norfentanyl was also detected in the range 32 pg/mg – 2300
5 pg/mg, with a mean value of 809 pg/mg and a median of 209 pg/mg. Another promising marker of
6
7
8 active fentanyl use, beta-hydroxyfentanyl (34), was detected in six cases, in the range 17 pg/mg -
9
10
11
12 1400 pg/mg. However, when beta-hydroxyfentanyl was present, norfentanyl was as well. Overall, the
13
14 main metabolites were detected in the majority of hair samples testing positive for fentanyl supporting
15
16 the possibility to ascertain active use. Acetylfentanyl, which is suggestive of clandestine production,
17
18 was detected in 3 cases (range: 129 pg/mg – 265 pg/mg) and 4-ANPP, which is a precursor of
19
20 fentanyl, in 7 cases (range: 23 pg/mg-2200 pg/mg), confirming that these two molecules are often
21
22 present in hair samples from people exposed to fentanyl, as a by-product of either metabolism or
23
24 synthesis of fentanyl. One further fentalog, despropionyl-p-fluorofentanyl, was detected in one
25
26 sample at the concentration of 25 pg/mg, together with fentanyl at 2900 pg/mg. The sporadic
27
28 occurrence of the other fentalogs has different possible explanations: (i) low prevalence within the
29
30 populations assessed at the time of the sample collection, (ii) poor incorporation or low stability in
31
32 the keratin matrix, and/or (iii) insufficient sensitivity of the analytical method in relation to the low
33
34 effective dosage. A summary of results is presented in Table 2.

35
36
37
38
39
40 Three prescription opioids were detected in seven segments (8.4%), usually together with
41
42 fentanyl. Only two segments (collected from the same subject) followed a different trend, with
43
44 fentanyl below the LOD and hydrocodone measured at 37 pg/mg and 46 pg/mg, respectively. The
45
46 trace level of fentanyl detected might indicate unintended exposure as an adulterant or contaminant
47
48 if the drugs ~~was~~ were obtained illegally. The nitazene compounds and bupropion, which appeared to
49
50 have a significant presence in the NPS opioid market in 2019 and 2020 (3), were not detected.
51
52
53 Although the number of samples analyzed in this study was relatively small, the non-detection of
54
55 emerging opioids is coherent with the modern drug scenario, in which the typical life cycle of a new
56
57 substance is generally short. Most new drugs appear to remain in circulation less than six months and
58
59
60

1
2
3 up to one year, but then rapidly decline, disappear, and then are replaced by other newly emerging
4
5 synthetic substances (3).
6

7
8 Novel and prescription synthetic opioid identification was much less common in the 60
9
10 samples collected in Italy. In particular, fentalogs were never detected, while only five segments
11
12 (5.5%) were positive for at least one compound such as hydrocodone, oxycodone, or tramadol. Two
13
14 subjects were positive for all three prescription opioids. In one case, two segments were obtained
15
16 from the same sample, showing the same trend of consumption (hydrocodone at 19 and 22 pg/mg,
17
18 respectively).
19
20

21 22 23 24 Testing for common drugs

25
26 Samples were considered positive in accordance with international cut-offs for parent drugs
27
28 and metabolites (35). In the group of samples from the US, cocaine was the most prevalent substance
29
30 found in the samples, with 19 segments (22.9%) resulting above the cut-off for either cocaine or its
31
32 metabolite benzoylecgonine (BZE). Cocaethylene was detected above 0.05 ng/mg only in five cases.
33
34 The 6-acetylmorphine (6-MAM) as marker of heroin use was identified in five segments (6.0%), and
35
36 all samples positive for 6-MAM also tested positive for BZE. While it is not possible to discriminate
37
38 whether cocaine and heroin were taken simultaneously or in rapid sequence, the fact that the two
39
40 substances were used in the same six months is remarkable. Use of cannabis-derived products was
41
42 verified by the presence of Δ^9 -tetrahydrocannabinol (THC) in only eight segments (9.6%). It is
43
44 noteworthy that the frequent use of amphetamine/methamphetamine/MDMA as a whole was
45
46 observed in 26 segments (31.3%), of which 14 tested positive also to cocaine. Ketamine was detected
47
48 in only one segment, in contradiction with the increasing trend recently reported, especially in the
49
50 New York City area (36, 37).
51
52
53

54
55
56 An exhaustive comparison of patterns of drug use between the US and Italy based on the
57
58 group of results hereby presented is not possible, nor is this the goal of our research. However, a
59
60 striking difference emerges from the results obtained from the samples collected in Italy. Among 91

1
2
3 segments, a total of 41 (45.1%) tested positive for exposure ketamine in the range 245-8500 pg/mg
4
5 (mean value 2324 pg/mg, median 1496 pg/mg). The large majority of samples positive for ketamine
6
7 also tested positive for MDMA and/or cocaine, showing a trend of potential co-use of stimulating and
8
9 dissociative substances. Overall, THC was still the most prevalent parent drug, with 73 positive
10
11 segments (80.2%). The use of heroin, proven by the presence of 6-MAM, was identified in eight
12
13 segments (8.8%).
14
15

16
17 Correlation matrices for the measured levels of common drugs in the Italian population of hair
18
19 samples are presented in Figure 1. High correlation coefficients suggest that subjects who were more
20
21 exposed to ketamine were more exposed to amphetamine, possibly due to a co-use of the substances
22
23 in certain recreational contexts.
24
25

26 27 28 4. Conclusions

29
30 Hair analysis can help to retrospectively explore trends in drug use, and incorporating hair
31
32 testing into epidemiology studies or surveillance studies can provide opportunity for relatively rapid
33
34 dissemination of results (including public alerts) to both the scientific community and populations at
35
36 risk.
37
38

39
40 _____ In this analysis focusing on hair samples collected in the US and in Italy, we tested for use or
41
42 exposure to fentalogs, prescription opioids, and more common controlled drugs including ketamine.
43
44 Results suggest that currently fentalogs continue to be a US (or North American) phenomenon, with
45
46 no detected cases in Italy despite high prevalence of detection of other drugs within this country.
47
48 However, we checkedinvestigated samples from two different populations within these two countries,
49
50 and results should not be directly used to indicate prevalence of drug use, as we used purposive
51
52 sampling. As such, results are not generalizable to US or Italian populations, but rather present a
53
54 snapshot of drug use within select populations in each country. Indeed, all cases of synthetic opioid
55
56 detection were in the US, but we focused on a sample in which many participants reported recent
57
58 synthetic opioid use. Most use of common party drugs such as ketamine and MDMA were detected
59
60

1
2
3 in the Italian sample, but we must keep in mind that these were individuals receiving harm reduction
4 services associated with nightlife. People who attend nightclubs in particular tend to report higher
5 prevalence of use of such drugs than the general population (38-40). Prevalence of past-year ketamine
6 use among young adults is estimated to be <1% in both the US and in Europe (14, 41), although
7 seizures of the drug appear to be increasing at a similar rate (14, 42). While prevalence of past-year
8 heroin use is estimated to be <1% in both the US and Europe (14, 43), in the US, synthetic opioids
9 such as fentanyl analogs indeed are more available and have been involved in hundreds of thousands
10 of deaths in recent years (13, 44). As such, it is important to note both where biological specimens
11 are collected but also the populations from which they are obtained. This is because results will vary
12 in particular across high-risk populations (e.g., nightclub attendees, people who utilize drug checking
13 services) and the general population.
14
15
16
17
18
19
20
21
22
23
24
25
26
27

28
29 Thanks to the longer detection window of hair (in comparison to much shorter detection
30 windows provided by urine, saliva, and blood), drug exposures occurring 1-2 weeks up through a
31 year before hair collection can provide retrospective results to inform scientists and public health
32 practitioners about the diffusion of drugs in their countries. Hair analysis results based on real hair
33 samples can provide information regarding both intentional and unintentional exposure to NPS/NSO,
34 both with and without use of common controlled drugs. As such, hair testing can serve as an addition
35 to epidemiology studies that seek to incorporate biological testing with survey research. The
36 combination of surveys and hair testing can thus be used to monitor drug exposure in a more effective
37 manner than using surveys or biological testing alone.
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Acknowledgments

Research reported in this publication was supported by the National Institute on Drug Abuse of the National Institutes of Health under Award Numbers U01DA051126 and R01DA044207. The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Institutes of Health. Prof. Alberto Salomone acknowledges support from the Project CH4.0 under the MUR program "Dipartimenti di Eccellenza 2023-2027" (CUP: D13C22003520001) funding from "Implementazione dell'identificazione e studio degli effetti delle NPS: Sviluppo di una multicentrica di ricerca per potenziare la base dati dell'Osservatorio Nazionale Tossicodipendenze e del Sistema di Allerta Precoce" (CUP I55E22000320001) We acknowledge the NDEWS team for the data collection methods and interviewing efforts.

Funding

J. Palamar is funded by the National Institutes of Health (NIH) (R01DA044207 and U01DA051126). A. Salomone received funding from "Implementazione dell'identificazione e studio degli effetti delle NPS: Sviluppo di una multicentrica di ricerca per potenziare la base dati dell'Osservatorio Nazionale Tossicodipendenze e del Sistema di Allerta Precoce" (CUP I55E22000320001) and from Fondazione C.R.T. "Approcci innovativi per comprendere la diffusione di nuove sostanze stupefacenti nella popolazione"

References

1. Ciccarone D. The triple wave epidemic: Supply and demand drivers of the US opioid overdose crisis. *Int J Drug Policy*. 2019;71:183-8. <https://doi.org/10.1016/j.drugpo.2019.01.010>
2. Zoorob M. Fentanyl shock: The changing geography of overdose in the United States. *Int J Drug Policy*. 2019;70:40-6. <https://doi.org/10.1016/j.drugpo.2019.04.010>
3. Vandeputte MM, Krotulski AJ, Papsun DM, Logan BK, Stove CP. The rise and fall of isotonitazene and bupropion: Two recent stars in the synthetic opioid firmament. *J Anal Toxicol*. 2022;46:115-21+. <https://doi.org/10.1093/jat/bkab082>
4. Laing MK, Ti L, Marmel A, Tobias S, Shapiro AM, Laing R, et al. An outbreak of novel psychoactive substance benzodiazepines in the unregulated drug supply: Preliminary results from a community drug checking program using point-of-care and confirmatory methods. *Int J Drug Policy*. 2021;93:103169. <https://doi.org/10.1016/j.drugpo.2021.103169>
5. Tupper W, McCrae K, Garber I, Lysyshyn M, Wood E. Initial results of a drug checking pilot program to detect fentanyl adulteration in a Canadian setting. *Drug Alcohol Depend*. 2018;190:242-5. <https://doi.org/10.1016/j.drugalcdep.2018.06.020>
6. Tobias S, Shapiro AM, Grant CJ, Patel P, Lysyshyn M, Ti L. Drug checking identifies counterfeit alprazolam tablets. *Drug Alcohol Depend*. 2021;218:108300. <https://doi.org/10.1016/j.drugalcdep.2020.108300>
7. May C, Downey C, Power JD, Kavanagh PV. An unusual detection of tert-butyl-4-anilinopiperidine-1-carboxylate in seizures of falsified 'Xanax' tablets and in items in a suspected heroin seizure submitted by Irish law enforcement. *Drug Test Anal*. 2020;12(9):1387-92. <https://doi.org/10.1002/dta.2884>
8. Palamar JJ, Salomone A, Bigiarini R, Vincenti M, Acosta P, Tofighi B. Testing hair for fentanyl exposure: A method to inform harm reduction behavior among individuals who use heroin. *Am J Drug Alcohol Abuse*. 2019;45(1):90-6. <https://doi.org/10.1080/00952990.2018.1550652>
9. DiSalvo P, Cooper G, Tsao J, Romeo M, Laskowski LK, Chesney G, et al. Fentanyl-contaminated cocaine outbreak with laboratory confirmation in New York City in 2019. *Am J Emerg Med*. 2020;40:103-5. <https://doi.org/10.1016/j.ajem.2020.12.002>
10. Park JN, Rashidi E, Foti K, Zoorob M, Sherman S, Alexander GC. Fentanyl and fentanyl analogs in the illicit stimulant supply: Results from U.S. drug seizure data, 2011-2016. *Drug Alcohol Depend*. 2021;218:108416. <https://doi.org/10.1016/j.drugalcdep.2020.108416>
11. Mars SG, Rosenblum D, Ciccarone D. Illicit fentanyls in the opioid street market: Desired or imposed? *Addiction*. 2019;114(5):774-80. <https://doi.org/10.1111/add.14474>
12. Centers for Disease Control and Prevention. U.S. overdose deaths in 2021 increased half as much as in 2020 – but are still up 15%. 2022. Available from: https://www.cdc.gov/nchs/pressroom/nchs_press_releases/2022/202205.htm. Accessed July 12, 2023.
13. Palamar JJ, Ciccarone D, Rutherford C, Keyes KM, Carr TH, Cottler LB. Trends in seizures of powders and pills containing illicit fentanyl in the United States, 2018 through 2021. *Drug Alcohol Depend*. 2022:109398. <https://doi.org/10.1016/j.drugalcdep.2022.109398>
14. European Monitoring Centre for Drugs and Drug Addiction. European Drug Report 2023: Trends and Developments; 2023. Available from:

https://www.emcdda.europa.eu/publications/european-drug-report/2023_en. Accessed July 12, 2023.

15. Kariisa M, Scholl L, Wilson N, Seth P, Hoots B. Drug overdose deaths involving cocaine and psychostimulants with abuse potential - United States, 2003-2017. *MMWR Morb Mortal Wkly Rep.* 2019;68(17):388-95. <https://doi.org/10.15585/mmwr.mm6817a3>
16. Martinez S, Jones JD, Brandt L, Campbell ANC, Abbott R, Comer SD. The increasing prevalence of fentanyl: A urinalysis-based study among individuals with opioid use disorder in New York City. *Am J Addict.* 2021;30(1):65-71. <https://doi.org/10.1111/ajad.13092>
17. Palamar JJ, Salomone A, Barratt MJ. Drug checking to detect fentanyl and new psychoactive substances. *Curr Opin Psychiatry.* 2020;33(4):301-5. <https://doi.org/10.1097/yco.0000000000000607>
18. Bach H, Jenkins V, Aledhaim A, Moayedi S, Schenkel SM, Kim HK. Prevalence of fentanyl exposure and knowledge regarding the risk of its use among emergency department patients with active opioid use history at an urban medical center in Baltimore, Maryland. *Clin Toxicol.* 2020;58(6):460-5. <https://doi.org/10.1080/15563650.2019.1657583>
19. Salomone A, Bigiarini R, Palamar JJ, McKnight C, Vinsick L, Amante E, et al. Toward the interpretation of positive testing for fentanyl and its analogs in real hair samples: Preliminary considerations. *J Anal Toxicol.* 2020;44(4):362-9. <https://doi.org/10.1093/jat/bkz102>
20. Larabi IA, Martin M, Etting I, Pfau G, Edel Y, Alvarez JC. Development and validation of liquid chromatography-tandem mass spectrometry targeted screening of 16 fentanyl analogs and U-47700 in hair: Application to 137 authentic samples. *Drug Test Anal.* 2020;12(9):1298-308. <https://doi.org/10.1002/dta.2868>
21. Chas J, Bauer R, Larabi IA, Peytavin G, Roux P, Cua E, et al. Evaluation of drug abuse by hair analysis and self-reported use among MSM under PrEP: Results from a French substudy of the ANRS-IPERGAY Trial. *J Acquir Immune Defic Syndr.* 2021;86(5):552-61. <https://doi.org/10.1097/qai.0000000000002610>
22. Palmquist KB, Truver MT, Shoff EN, Krotulski AJ, Swortwood MJ. Review of analytical methods for screening and quantification of fentanyl analogs and novel synthetic opioids in biological specimens. *J Forensic Sci.* 2023; online ahead of print. <https://doi.org/10.1111/1556-4029.15282>
23. Platosz NA, Binz TM, Baumgartner MR, Lendoiro E, de Castro A, Concheiro M. Quantification of Classic, Prescription and synthetic opioids in hair by LC-MS-MS. *J Anal Toxicol.* 2021;45(9):943-9. <https://doi.org/10.1093/jat/bkaa152>
24. Freni F, Moretti M, Radaelli D, Carelli C, Osculati AMM, Tronconi L, et al. Determination of fentanyl and 19 derivatives in hair: Application to an Italian population. *J Pharm Biomed Anal.* 2020;189:113476. <https://doi.org/10.1016/j.jpba.2020.113476>
25. Ramírez Fernández MDM, Wille SMR, Jankowski D, Hill V, Samyn N. Development of an UPLC-MS/MS method for the analysis of 16 synthetic opioids in segmented hair, and evaluation of the polydrug history in fentanyl analogue users. *Forensic Sci Int.* 2020;307:110137. <https://doi.org/10.1016/j.forsciint.2019.110137>
26. Rammohan I, Jain S, Sun S, Marks C, Milloy MJ, Hayashi K, et al. Latent polydrug use patterns and the provision of injection initiation assistance among people who inject drugs in three North American settings. *Drug Alcohol Rev.* 2023. <https://doi.org/10.1111/dar.13687>
27. van Amsterdam J, Pierce M, van den Brink W. Predictors and motives of polydrug use in opioid users. A narrative review. *Curr Opin Psychiatry.* 2023;36(4):301-7. <https://doi.org/10.1097/yco.0000000000000875>
28. Cottler LB, Goldberger BA, Nixon SJ, Striley CW, Barenholtz E, Fitzgerald ND, et al. Introducing NIDA's new National Drug Early Warning System. *Drug Alcohol Depend.* 2020;217:108286. <https://doi.org/10.1016/j.drugalcdep.2020.108286>

- 1
2
3 29. Salomone A, Tsanaclis L, Agius R, Kintz P, Baumgartner MR. European guidelines for
4 workplace drug and alcohol testing in hair. *Drug Test Anal.* 2016;8(10):996-1004.
5 <https://doi.org/10.1002/dta.1999>
- 6 30. Di Corcia D, Salomone A, Gerace E. Analysis of drugs of abuse in hair samples by
7 ultrahigh-performance liquid chromatography-tandem mass spectrometry (UHPLC-MS/MS).
8 *Methods Mol Biol.* 2018;1810:107-14. https://doi.org/10.1007/978-1-4939-8579-1_10
- 9 31. Di Corcia D, D'Urso F, Gerace E, Salomone A, Vincenti M. Simultaneous determination in
10 hair of multiclass drugs of abuse (including THC) by ultra-high performance liquid
11 chromatography-tandem mass spectrometry. *J Chromatogr B Analyt Technol Biomed Life Sci.*
12 2012;899:154-9. <https://doi.org/10.1016/j.jchromb.2012.05.003>
- 13 32. Salomone A, Palamar JJ, Gerace E, Di Corcia D, Vincenti M. Hair testing for drugs of abuse
14 and new psychoactive substances in a high-risk population. *J Anal Toxicol.* 2017;41(5):376-81.
15 <https://doi.org/10.1093/jat/bkx020>
- 16 33. Salomone A, Palamar JJ, Bigiarini R, Gerace E, Di Corcia D, Vincenti M. Detection of
17 fentanyl analogs and synthetic opioids in real hair samples. *J Anal Toxicol.* 2019;43(4):259-65.
18 <https://doi.org/10.1093/jat/bky093>
- 19 34. Salomone A, Di Corcia D, Negri P, Kolia M, Amante E, Gerace E, et al. Targeted and
20 untargeted detection of fentanyl analogues and their metabolites in hair by means of UHPLC-
21 QTOF-HRMS. *Anal Bioanal Chem.* 2021;413(1):225-33. <https://doi.org/10.1007/s00216-020-02994-x>
- 22 35. Favretto D, Cooper G, Andraus M, Sporkert F, Agius R, Appenzeller B, et al. The Society
23 of Hair Testing consensus on general recommendations for hair testing and drugs of abuse testing in
24 hair. *Drug Test Anal.* 2023; online ahead of print. <https://doi.org/10.1002/dta.3526>
- 25 36. Arango E, Toriello A, Rosario Z, Cooper G. Increasing prevalence of ketamine in drivers in
26 New York City including the identification of 2-fluoro-deschloroketamine. *J Anal Toxicol.*
27 2021;45(8):799-806. <https://doi.org/10.1093/jat/bkab057>
- 28 37. Palamar JJ, Keyes KM. Trends in drug use among electronic dance music party attendees in
29 New York City, 2016-2019. *Drug Alcohol Depend.* 2020;209:107889.
30 <https://doi.org/10.1016/j.drugalcdep.2020.107889>
- 31 38. Palamar JJ, Le A, Rutherford C, Keyes KM. Exploring potential bellwethers for drug-related
32 mortality in the general population: A case for sentinel surveillance of trends in drug use among
33 nightclub/festival attendees. *Subst Use Misuse.* 2022;58(2):188-97.
34 <https://doi.org/10.1080/10826084.2022.2151315>
- 35 39. Palamar JJ, Rutherford C, Cleland CM, Keyes KM. Concerts, bars, parties, and raves:
36 Differential risk for drug use among high school seniors according to venue attendance. *Subst*
37 *Abus.* 2022;43(1):785-91. <https://doi.org/10.1080/08897077.2021.2010253>
- 38 40. Palamar JJ, Griffin-Tomas M, Ompad DC. Illicit drug use among rave attendees in a
39 nationally representative sample of US high school seniors. *Drug Alcohol Depend.* 2015;152:24-31.
40 <https://doi.org/10.1016/j.drugalcdep.2015.05.002>
- 41 41. Palamar JJ, Rutherford C, Keyes KM. Trends in ketamine use, exposures, and seizures in
42 the United States. *Am J Public Health.* 2021;111(11):2046-9.
43 <https://doi.org/10.2105/ajph.2021.306486>
- 44 42. Palamar JJ, Wilkinson ST, Carr TH, Rutherford C, Cottler LB. Trends in illicit ketamine
45 seizures in the US from 2017-2022. *JAMA Psychiatry.* 2023;80(7):750-751.
- 46 43. Center for Behavioral Health Statistics and Quality. Results from the 2021 National Survey
47 on Drug Use and Health: Detailed tables. Rockville, MD: Substance Abuse and Mental Health
48 Services Administration; 2022. Available from [https://www.samhsa.gov/data/report/2021-nsduh-](https://www.samhsa.gov/data/report/2021-nsduh-detailed-tables)
49 [detailed-tables](https://www.samhsa.gov/data/report/2021-nsduh-detailed-tables). Accessed July 12, 2023.
- 50 44. Hedegaard H, Miniño AM, Warner M. Drug overdose deaths in the United States, 1999-
51 2019. *NCHS Data Brief.* 2020;394:1-8. Available from
52 <https://www.cdc.gov/nchs/data/databriefs/db394-H.pdf>. Accessed July 12, 2023.
- 53
54
55
56
57
58
59
60

Table 1: List (ordered by RT) of the monitored transitions, their instrumental parameters and the related internal standard for the screened compounds.

<u>Compound</u>	<u>Retention time, min</u>	<u>Precursor mass Q1 m/z</u>	<u>Fragments mass Q3 m/z</u>	<u>CE (V)</u>	<u>EP (V)</u>	<u>Internal Standard</u>
<u>Acetyl norfentanyl</u>	<u>2.10</u>	<u>219.1</u>	<u>84.1</u> <u>55.1</u> <u>56.1</u>	<u>23</u> <u>48</u> <u>42</u>	<u>8</u>	<u>Fentanyl-D5</u>
<u>Methoxyacetyl norfentanyl</u>	<u>2.10</u>	<u>249.1</u>	<u>84.1</u> <u>55.1</u> <u>56</u>	<u>22</u> <u>55</u> <u>40</u>	<u>8</u>	<u>Fentanyl-D5</u>
<u>Oxycodone</u>	<u>2.10</u>	<u>316.0</u>	<u>298.1</u> <u>241.1</u> <u>256.1</u>	<u>25</u> <u>38</u> <u>34</u>	<u>8</u>	<u>Oxycodone-D6</u>
<u>Hydrocodone</u>	<u>2.20</u>	<u>300.0</u>	<u>199.1</u> <u>171.1</u> <u>128.0</u>	<u>42</u> <u>51</u> <u>74</u>	<u>8</u>	<u>Fentanyl-D5</u>
<u>Norfentanyl</u>	<u>2.60</u>	<u>233.0</u>	<u>84.1</u> <u>150.1</u> <u>55.0</u>	<u>24</u> <u>22</u> <u>50</u>	<u>8</u>	<u>Norfentanyl-D5</u>
<u>Metodesnitazene</u>	<u>2.60</u>	<u>338.1</u>	<u>100.0</u> <u>72.0</u> <u>121.0</u>	<u>23</u> <u>53</u> <u>50</u>	<u>8</u>	<u>Fentanyl-D5</u>
<u>3-methyl Norfentanyl</u>	<u>2.70</u>	<u>247.1</u>	<u>98.1</u> <u>150.1</u> <u>69.0</u>	<u>23</u> <u>26</u> <u>42</u>	<u>8</u>	<u>Fentanyl-D5</u>
<u>Furanyl Norfentanyl</u>	<u>2.70</u>	<u>271.2</u>	<u>84</u> <u>56.1</u> <u>55</u>	<u>20</u> <u>41</u> <u>54</u>	<u>8</u>	<u>Fentanyl-D5</u>
<u>Tramadol</u>	<u>2.80</u>	<u>264.1</u>	<u>58.1</u> <u>246.1</u>	<u>46</u> <u>15</u>	<u>8</u>	<u>Fentanyl-D5</u>
<u>Butyryl Norfentanyl</u>	<u>3.00</u>	<u>247.2</u>	<u>84.0</u> <u>177.2</u> <u>55.0</u>	<u>24</u> <u>21</u> <u>55</u>	<u>8</u>	<u>Fentanyl-D5</u>
<u>Etodesnitazene</u>	<u>3.00</u>	<u>352.1</u>	<u>100.1</u> <u>71.9</u> <u>107.1</u>	<u>26</u> <u>57</u> <u>60</u>	<u>8</u>	<u>Fentanyl-D5</u>
<u>Remifentanyl</u>	<u>3.10</u>	<u>377.1</u>	<u>317.2</u> <u>228.0</u> <u>116.1</u>	<u>22</u> <u>27</u> <u>37</u>	<u>8</u>	<u>Fentanyl-D5</u>
<u>Butyrylfentanyl carboxy metabolite</u>	<u>3.10</u>	<u>381.1</u>	<u>188.2</u> <u>105.1</u> <u>260.1</u>	<u>34</u> <u>56</u> <u>34</u>	<u>8</u>	<u>Fentanyl-D5</u>

<u>OH-thioentanyl</u>	<u>3.10</u>	<u>359.1</u>	<u>192.1</u> <u>146.1</u> <u>111.0</u>	<u>32</u> <u>32</u> <u>50</u>	<u>8</u>	<u>Fentanyl-D5</u>
<u>Valeryl p-fluoro fentanyl</u>	<u>3.20</u>	<u>395.1</u>	<u>188.2</u> <u>105.0</u> <u>274.1</u>	<u>32</u> <u>57</u> <u>33</u>	<u>8</u>	<u>Fentanyl-D5</u>
<u>Acetylfentanyl</u>	<u>3.20</u>	<u>323.0</u>	<u>188.2</u> <u>105.0</u> <u>103.0</u>	<u>38</u> <u>38</u> <u>86</u>	<u>8</u>	<u>Fentanyl-D5</u>
<u>Ocfentanyl</u>	<u>3.20</u>	<u>371.1</u>	<u>105.1</u> <u>188.2</u> <u>134.0</u>	<u>50</u> <u>31</u> <u>38</u>	<u>8</u>	<u>Fentanyl-D5</u>
<u>beta-OH-fentanyl</u>	<u>3.30</u>	<u>353.2</u>	<u>186.1</u> <u>204.2</u> <u>335.2</u>	<u>32</u> <u>28</u> <u>26</u>	<u>8</u>	<u>Fentanyl-D5</u>
<u>4-ANPP</u>	<u>3.40</u>	<u>281.0</u>	<u>188.2</u> <u>105.0</u> <u>103.0</u>	<u>24</u> <u>41</u> <u>63</u>	<u>8</u>	<u>Fentanyl-D5</u>
<u>Alfentanyl</u>	<u>3.50</u>	<u>417.0</u>	<u>268.3</u> <u>197.2</u> <u>165.0</u>	<u>24</u> <u>35</u> <u>47</u>	<u>8</u>	<u>Fentanyl-D5</u>
<u>Acrylfentanyl</u>	<u>3.50</u>	<u>335.1</u>	<u>188.2</u> <u>105.0</u> <u>132.1</u>	<u>30</u> <u>50</u> <u>42</u>	<u>8</u>	<u>Fentanyl-D5</u>
<u>Despropionyl-p- fluorofentanyl</u>	<u>3.50</u>	<u>299.2</u>	<u>188.1</u> <u>105.0</u> <u>134.0</u>	<u>24</u> <u>39</u> <u>32</u>	<u>8</u>	<u>Fentanyl-D5</u>
<u>Flunitazene</u>	<u>3.50</u>	<u>371.1</u>	<u>100.1</u> <u>109.1</u> <u>72.1</u>	<u>33</u> <u>65</u> <u>58</u>	<u>8</u>	<u>Fentanyl-D5</u>
<u>Fentanyl</u>	<u>3.50</u>	<u>337.1</u>	<u>188.2</u> <u>105.0</u> <u>132.1</u>	<u>32</u> <u>49</u> <u>42</u>	<u>8</u>	<u>Fentanyl-D5</u>
<u>Metonitazene</u>	<u>3.50</u>	<u>383.0</u>	<u>100.0</u> <u>72.1</u> <u>121.0</u>	<u>26</u> <u>58</u> <u>38</u>	<u>8</u>	<u>Fentanyl-D5</u>
<u>U-47700</u>	<u>3.50</u>	<u>328.9</u>	<u>204.1</u> <u>286.1</u> <u>206.1</u>	<u>36</u> <u>24</u> <u>34</u>	<u>8</u>	<u>Fentanyl-D5</u>
<u>4-methylfentanyl</u>	<u>3.60</u>	<u>351.1</u>	<u>91</u> <u>202.1</u> <u>119.1</u>	<u>51</u> <u>30</u> <u>35</u>	<u>8</u>	<u>Fentanyl-D5</u>
<u>AH-7921</u>	<u>3.60</u>	<u>329.0</u>	<u>173.0</u> <u>284.1</u> <u>286.1</u>	<u>40</u> <u>23</u> <u>24</u>	<u>8</u>	<u>Fentanyl-D5</u>
<u>Furanilfentanyl</u>	<u>3.60</u>	<u>375.0</u>	<u>188.2</u> <u>105.0</u> <u>103.0</u>	<u>28</u> <u>52</u> <u>82</u>	<u>8</u>	<u>Fentanyl-D5</u>

<u>Brorphine</u>	<u>3.70</u>	<u>402.0</u>	<u>218.2</u> <u>104.1</u> <u>218.2</u>	<u>29</u> <u>63</u> <u>35</u>	<u>8</u>	<u>Fentanyl-D5</u>
<u>Carfentanyl</u>	<u>3.70</u>	<u>395.0</u>	<u>335.2</u> <u>246.1</u> <u>113.0</u>	<u>25</u> <u>34</u> <u>34</u>	<u>8</u>	<u>Fentanyl-D5</u>
<u>Cyclopropylfentanyl</u>	<u>3.70</u>	<u>349.1</u>	<u>188.1</u> <u>105.0</u> <u>132.0</u>	<u>32</u> <u>51</u> <u>40</u>	<u>8</u>	<u>Fentanyl-D5</u>
<u>N-pyrrolidino etonitazene</u>	<u>3.70</u>	<u>395.0</u>	<u>98.0</u> <u>107.0</u> <u>56.0</u>	<u>27</u> <u>70</u> <u>82</u>	<u>8</u>	<u>Fentanyl-D5</u>
<u>Isotonitazene</u>	<u>3.70</u>	<u>411.2</u>	<u>100.0</u> <u>106.9</u> <u>72.0</u>	<u>20</u> <u>52</u> <u>42</u>	<u>8</u>	<u>Fentanyl-D5</u>
<u>Butyrylfentanyl</u>	<u>3.80</u>	<u>351.2</u>	<u>188.2</u> <u>105.1</u> <u>230.2</u>	<u>31</u> <u>49</u> <u>31</u>	<u>8</u>	<u>Fentanyl-D5</u>
<u>Phenyl fentanyl</u>	<u>3.80</u>	<u>385.2</u>	<u>188.2</u> <u>105</u> <u>134.2</u>	<u>29</u> <u>51</u> <u>36</u>	<u>8</u>	<u>Fentanyl-D5</u>
<u>Sufentanyl</u>	<u>3.90</u>	<u>387.0</u>	<u>238.1</u> <u>355.1</u> <u>111.0</u>	<u>26</u> <u>26</u> <u>46</u>	<u>8</u>	<u>Fentanyl-D5</u>
<u>4-F-butylfentanyl</u>	<u>3.90</u>	<u>369.1</u>	<u>188.1</u> <u>105.0</u> <u>248.1</u>	<u>33</u> <u>55</u> <u>33</u>	<u>8</u>	<u>Fentanyl-D5</u>
<u>Phenylacetyl fentanyl</u>	<u>4.10</u>	<u>399.2</u>	<u>105</u> <u>188.2</u> <u>134.1</u>	<u>55</u> <u>32</u> <u>39</u>	<u>8</u>	<u>Fentanyl-D5</u>
<u>MT-45</u>	<u>4.20</u>	<u>349.1</u>	<u>181.1</u> <u>166.2</u> <u>169.2</u>	<u>36</u> <u>46</u> <u>25</u>	<u>8</u>	<u>Fentanyl-D5</u>
<u>beta-phenyl Fentanyl</u>	<u>4.30</u>	<u>413.2</u>	<u>188.2</u> <u>105.0</u> <u>292.1</u>	<u>35</u> <u>55</u> <u>37</u>	<u>8</u>	<u>Fentanyl-D5</u>
<u>Butonitazene</u>	<u>4.4</u>	<u>425.2</u>	<u>100.1</u> <u>72.0</u> <u>107.0</u>	<u>31</u> <u>67</u> <u>75</u>	<u>8</u>	<u>Fentanyl-D5</u>

Table 2 Summary of results obtained from 60 real hair samples collected in the U.S. All concentrations are in pg/mg. Two segments from the same sample are referred to as *a* and *b*

<u>Sample</u>	<u>Fentanyl</u>	<u>Norfentanyl</u>	<u>4-ANPP</u>	<u>β-OH-fentanyl</u>	<u>Other fentalogs</u>	<u>Prescription opioids</u>
<u>1</u>	<u>90</u>	=	=	=	=	=
<u>2</u>	<u>384</u>	<u>43</u>	<u>29</u>	<u>71</u>	=	=
<u>3</u>	<u>501</u>	<u>69</u>	=	=	=	=
<u>4a</u>	<u>2900</u>	<u>1700</u>	<u>747</u>	<u>1400</u>	<u>Acetyl fentanyl 157</u> <u>Despropionyl p-fluorofentanyl 25</u>	<u>Tramadol</u> <u>1400</u>
<u>4b</u>	<u>2900</u>	<u>2000</u>	<u>737</u>	<u>1300</u>	<u>Acetyl fentanyl 129</u>	<u>Tramadol</u> <u>1200</u>
<u>5a</u>	=	=	=	=	=	<u>Hydrocodone</u> <u>37</u>
<u>5b</u>	=	=	=	=	=	<u>Hydrocodone</u> <u>46</u>
<u>6</u>	<u>15</u>	=	=	=	=	<u>Hydrocodone</u> <u>26</u> <u>Tramadol 13</u> <u>Oxycodone</u> <u>128</u>
<u>7</u>	<u>13</u>	=	=	=	=	=
<u>8</u>	<u>103</u>	=	=	=	=	=
<u>9</u>	<u>331</u>	<u>32</u>	<u>23</u>	=	=	=
<u>10</u>	<u>1800</u>	<u>150</u>	<u>73</u>	<u>60</u>	=	=
<u>11</u>	<u>143</u>	=	=	=	=	=
<u>12a</u>	<u>2800</u>	<u>268</u>	<u>223</u>	<u>17</u>	=	<u>Hydrocodone</u> <u>78</u> <u>Tramadol 25</u>
<u>12b</u>	<u>7300</u>	<u>2300</u>	<u>2200</u>	<u>980</u>	<u>Acetyl fentanyl 265</u>	<u>Tramadol</u> <u>1000</u>
<u>13</u>	<u>61</u>	=	=	=	=	=

Table 1: List of the monitored transitions, their instrumental parameters and the related internal standard for the screened compounds.

Compound	Retention time, min	Precursor mass Q1 m/z	Fragments mass Q3 m/z	CE (V)	EP (V)	Internal Standard
4-ANPP	3.40	281.0	188.2 105.0 103.0	24 41 63	8	Fentanyl-D5
Acetylfentanyl	3.20	323.0	188.2 105.0 103.0	38 38 86	8	Fentanyl-D5
Carfentanyl	3.70	395.0	335.2 246.1 113.0	25 34 34	8	Fentanyl-D5
Fentanyl	3.50	337.1	188.2 105.0 132.1	32 49 42	8	Fentanyl-D5
Furanilfentanyl	3.60	375.0	188.2 105.0 103.0	28 52 82	8	Fentanyl-D5
Hydrocodone	2.20	300.0	199.1 171.1 128.0	42 51 74	8	Fentanyl-D5
Norfentanyl	2.60	233.0	84.1 150.1 55.0	24 22 50	8	Norfentanyl-D5
Remifentanyl	3.10	377.1	317.2 228.0 116.1	22 27 37	8	Fentanyl-D5
Tramadol	2.80	264.1	58.1 246.1	46 15	8	Fentanyl-D5
Alfentanyl	3.50	417.0	268.3 197.2 165.0	24 35 47	8	Fentanyl-D5
Oxycodone	2.10	316.0	298.1 241.1 256.1	25 38 34	8	Oxycodone-D6
Sufentanyl	3.90	387.0	238.1 355.1 111.0	26 26 46	8	Fentanyl-D5
U-47700	3.5	328.9	204.1 286.1 206.1	36 24 34	8	Fentanyl-D5
4-F-butylfentanyl	3.90	369.1	188.1 105.0	33 55	8	Fentanyl-D5

			248.1	33		
4-methylfentanyl	3.60	351.1	91 202.1 119.1	51 30 35	8	Fentanyl-D5
Acrylfentanyl	3.50	335.1	188.2 105.0 132.1	30 50 42	8	Fentanyl-D5
AH-7921	3.60	329.0	173.0 284.1 286.1	40 23 24	8	Fentanyl-D5
Butyrylfentanyl	3.80	351.2	188.2 105.1 230.2	31 49 31	8	Fentanyl-D5
Cyclopropylfentanyl	3.70	349.1	188.1 105.0 132.0	32 51 40	8	Fentanyl-D5
MT-45	4.20	349.1	181.1 166.2 169.2	36 46 25	8	Fentanyl-D5
Ocfentanyl	3.20	371.1	105.1 188.2 134.0	50 31 38	8	Fentanyl-D5
3-methyl Norfentanyl	2.70	247.1	98.1 150.1 69.0	23 26 42	8	Fentanyl-D5
Acetyl-norfentanyl	2.10	219.1	84.1 55.1 56.1	23 48 42	8	Fentanyl-D5
Butyrylfentanyl carboxy-metabolite	3.10	381.1	188.2 105.1 260.1	34 56 34	8	Fentanyl-D5
Despropionyl-p- fluorofentanyl	3.5	299.2	188.1 105.0 134.0	24 39 32	8	Fentanyl-D5
Valeryl p-fluoro fentanyl	3.20	395.1	188.2 105.0 274.1	32 57 33	8	Fentanyl-D5
beta-OH-fentanyl	3.30	353.2	186.1 204.2 335.2	32 28 26	8	Fentanyl-D5
beta-phenyl Fentanyl	4.30	413.2	188.2 105.0 292.1	35 55 37	8	Fentanyl-D5
Butyryl Norfentanyl	3.00	247.2	84.0 177.2 55.0	24 21 55	8	Fentanyl-D5
Furanyl Norfentanyl	2.70	271.2	84	20	8	Fentanyl-D5

			56.1 55	41 54		
Methoxyacetyl norfentanyl	2.10	249.1	84.1 55.1 56	22 55 40	8	Fentanyl-D5
OH-thioentanyl	3.10	359.1	192.1 146.1 111.0	32 32 50	8	Fentanyl-D5
Phenylacetyl fentanyl	4.10	399.2	105 188.2 134.1	55 32 39	8	Fentanyl-D5
Phenyl fentanyl	3.80	385.2	188.2 105 134.2	29 51 36	8	Fentanyl-D5
Brorphine	3.70	402.0	218.2 104.1 218.2	29 63 35	8	Fentanyl-D5
Butonitazene	4.4	425.2	100.1 72.0 107.0	31 67 75	8	Fentanyl-D5
Etodesnitazene	3.00	352.1	100.1 71.9 107.1	26 57 60	8	Fentanyl-D5
Flunitazene	3.50	371.1	100.1 109.1 72.1	33 65 58	8	Fentanyl-D5
Metodesnitazene	2.60	338.1	100.0 72.0 121.0	23 53 50	8	Fentanyl-D5
Metonitazene	3.50	383.0	100.0 72.1 121.0	26 58 38	8	Fentanyl-D5
N-pyrrolidino etonitazene	3.70	395.0	98.0 107.0 56.0	27 70 82	8	Fentanyl-D5
Isotonitazene	3.68	411.2	100.0 106.9 72.0	20 52 42	8	Fentanyl-D5

Table 2 Summary of results obtained from 60 real hair samples collected in the U.S. All concentrations are in pg/mg. Two segments from the same sample are referred to as *a* and *b*.

Sample	Fentanyl	Norfentanyl	4-ANPP	β -OH-fentanyl	Other fentalogs	Prescription opioids
1	90	-	-	-	-	-
2	384	43	29	71	-	-
3	501	69	-	-	-	-
4a	2900	1700	747	1400	Acetyl fentanyl 157 Despropionyl p-fluorofentanyl 25	Tramadol 1400
4b	2900	2000	737	1300	Acetyl fentanyl 129	Tramadol 1200
5a	-	-	-	-	-	Hydrocodone 37
5b	-	-	-	-	-	Hydrocodone 46
						Hydrocodone 26
6	15	-	-	-	-	Tramadol 13 Oxycodone 128
7	13	-	-	-	-	-
8	103	-	-	-	-	-
9	331	32	23	-	-	-
10	1800	150	73	60	-	-
11	143	-	-	-	-	-
12a	2800	268	223	17	-	Hydrocodone 78 Tramadol 25
12b	7300	2300	2200	980	Acetyl fentanyl 265	Tramadol 1000
13	61	-	-	-	-	-

Legend Figure 1

Correlation matrices for the measured levels of common drugs of abuse in the Italian population

For Peer Review

Table 1: List (ordered by RT) of the monitored transitions, their instrumental parameters and the related internal standard for the screened compounds.

Compound	Retention time, min	Precursor mass Q1 m/z	Fragments mass Q3 m/z	CE (V)	EP (V)	Internal Standard
Acetyl norfentanyl	2.10	219.1	84.1	23	8	Fentanyl-D5
			55.1	48		
			56.1	42		
Methoxyacetyl norfentanyl	2.10	249.1	84.1	22	8	Fentanyl-D5
			55.1	55		
			56	40		
Oxycodone	2.10	316.0	298.1	25	8	Oxycodone-D6
			241.1	38		
			256.1	34		
Hydrocodone	2.20	300.0	199.1	42	8	Fentanyl-D5
			171.1	51		
			128.0	74		
Norfentanyl	2.60	233.0	84.1	24	8	Norfentanyl-D5
			150.1	22		
			55.0	50		
Metodesnitazene	2.60	338.1	100.0	23	8	Fentanyl-D5
			72.0	53		
			121.0	50		
3-methyl Norfentanyl	2.70	247.1	98.1	23	8	Fentanyl-D5
			150.1	26		
			69.0	42		
Furanyl Norfentanyl	2.70	271.2	84	20	8	Fentanyl-D5
			56.1	41		
			55	54		
Tramadol	2.80	264.1	58.1	46	8	Fentanyl-D5
			246.1	15		
Butyryl Norfentanyl	3.00	247.2	84.0	24		Fentanyl-D5

			177.2	21	8	
			55.0	55		
Etodesnitazene	3.00	352.1	100.1	26		Fentanyl-D5
			71.9	57	8	
			107.1	60		
Remifentanyl	3.10	377.1	317.2	22		Fentanyl-D5
			228.0	27	8	
			116.1	37		
Butyrylfentanyl carboxy metabolite	3.10	381.1	188.2	34		Fentanyl-D5
			105.1	56	8	
			260.1	34		
OH-thiofentanyl	3.10	359.1	192.1	32		Fentanyl-D5
			146.1	32	8	
			111.0	50		
Valeryl p-fluoro fentanyl	3.20	395.1	188.2	32		Fentanyl-D5
			105.0	57	8	
			274.1	33		
Acetylfentanyl	3.20	323.0	188.2	38		Fentanyl-D5
			105.0	38	8	
			103.0	86		
Ocfentanyl	3.20	371.1	105.1	50		Fentanyl-D5
			188.2	31	8	
			134.0	38		
beta-OH-fentanyl	3.30	353.2	186.1	32		Fentanyl-D5
			204.2	28	8	
			335.2	26		
4-ANPP	3.40	281.0	188.2	24		Fentanyl-D5
			105.0	41	8	
			103.0	63		
Alfentanyl	3.50	417.0	268.3	24		Fentanyl-D5
			197.2	35	8	
			165.0	47		

Acrylfentanyl	3.50	335.1	188.2 105.0 132.1	30 50 42	8	Fentanyl-D5
Despropionyl-p-fluorofentanyl	3.50	299.2	188.1 105.0 134.0	24 39 32	8	Fentanyl-D5
Flunitazene	3.50	371.1	100.1 109.1 72.1	33 65 58	8	Fentanyl-D5
Fentanyl	3.50	337.1	188.2 105.0 132.1	32 49 42	8	Fentanyl-D5
Metonitazene	3.50	383.0	100.0 72.1 121.0	26 58 38	8	Fentanyl-D5
U-47700	3.50	328.9	204.1 286.1 206.1	36 24 34	8	Fentanyl-D5
4-methylfentanyl	3.60	351.1	91 202.1 119.1	51 30 35	8	Fentanyl-D5
AH-7921	3.60	329.0	173.0 284.1 286.1	40 23 24	8	Fentanyl-D5
Furanilfentanyl	3.60	375.0	188.2 105.0 103.0	28 52 82	8	Fentanyl-D5
Brorphine	3.70	402.0	218.2 104.1 218.2	29 63 35	8	Fentanyl-D5
Carfentanyl	3.70	395.0	335.2 246.1	25 34	8	Fentanyl-D5

			113.0	34		
Cyclopropylfentanyl	3.70	349.1	188.1 105.0 132.0	32 51 40	8	Fentanyl-D5
N-pyrrolidino etonitazene	3.70	395.0	98.0 107.0 56.0	27 70 82	8	Fentanyl-D5
Isotonitazene	3.70	411.2	100.0 106.9 72.0	20 52 42	8	Fentanyl-D5
Butyrylfentanyl	3.80	351.2	188.2 105.1 230.2	31 49 31	8	Fentanyl-D5
Phenyl fentanyl	3.80	385.2	188.2 105 134.2	29 51 36	8	Fentanyl-D5
Sufentanyl	3.90	387.0	238.1 355.1 111.0	26 26 46	8	Fentanyl-D5
4-F-butylfentanyl	3.90	369.1	188.1 105.0 248.1	33 55 33	8	Fentanyl-D5
Phenylacetyl fentanyl	4.10	399.2	105 188.2 134.1	55 32 39	8	Fentanyl-D5
MT-45	4.20	349.1	181.1 166.2 169.2	36 46 25	8	Fentanyl-D5
beta-phenyl Fentanyl	4.30	413.2	188.2 105.0 292.1	35 55 37	8	Fentanyl-D5

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Butonitazene	4.4	425.2	100.1 72.0 107.0	31 67 75	8	Fentanyl-D5
--------------	-----	-------	------------------------	----------------	---	-------------

For Peer Review

Table 2 Summary of results obtained from 60 real hair samples collected in the U.S. All concentrations are in pg/mg. Two segments from the same sample are referred to as *a* and *b*

Sample	Fentanyl	Norfentanyl	4-ANPP	β -OH-fentanyl	Other fentalogs	Prescription opioids
1	90	-	-	-	-	-
2	384	43	29	71	-	-
3	501	69	-	-	-	-
4a	2900	1700	747	1400	Acetyl fentanyl 157 Despropionyl p-fluorofentanyl 25	Tramadol 1400
4b	2900	2000	737	1300	Acetyl fentanyl 129	Tramadol 1200
5a	-	-	-	-	-	Hydrocodone 37
5b	-	-	-	-	-	Hydrocodone 46
						Hydrocodone 26
6	15	-	-	-	-	Tramadol 13 Oxycodone 128
7	13	-	-	-	-	-
8	103	-	-	-	-	-
9	331	32	23	-	-	-
10	1800	150	73	60	-	-
11	143	-	-	-	-	-
12a	2800	268	223	17	-	Hydrocodone 78 Tramadol 25
12b	7300	2300	2200	980	Acetyl fentanyl 265	Tramadol 1000
13	61	-	-	-	-	-

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60




Figure 1

131x114mm (300 x 300 DPI)

Table S1: Limits of Detection (LOD) for the screened compounds. LOD were calculated from five injections of the lowest point of calibration curve.

Compound	LOD (pg/mg)
4-ANPP	1.7
Acetylfentanyl	2.8
Carfentanyl	2.2
Fentanyl	3.7
Furanilfentanyl	1.6
Hydrocodone	6.2
Norfentanyl	3.3
Remifentanyl	0.6
Tramadol	4.3
Alfentanyl	0.8
Oxycodone	1.7
Sufentanyl	3.5
U-47700	3.6
4-F-butylfentanyl	2.2
4-methylfentanyl	1.0
Acrylfentanyl	5.1

AH-7921	3.5
Butyrylfentanyl	1.8
Ciclopropylfentanyl	2.8
MT-45	1.5
Ocfentanyl	1.6
3-methyl Norfentanyl	1.2
Acetyl norfentanyl	1.7
Butyrylfentanyl carboxy metabolite	1.9
Despropionyl-p-fluorofentanyl	3.1
Valeryl p-fluoro fentanyl	3.5
Beta-OH-fentanyl	0.8
Beta-phenyl Fentanyl	1.2
Butiryl Norfentanyl	1.7
Furanyl Norfentanyl	1.2
Methoxyacetyl norfentanyl	5.5
OH-ThioFentanyl	1.0
Phenylacetyl fentanyl	5.0

Phenyl fentanyl	1.8
Brorphine	0.7
Butonitazene	3.0
Etodesnitazene	5.4
Flunitazene	1.6
Metodesnitazene	8.0
Metonitazene	3.1
N-pyrrolidino etonitazene	4.1
Isotonitazene	5.6

Table S2: Precision and trueness values at three different concentrations. A partial validation was performed for the compounds which were detected in real samples

Compound	Concentration (pg/mg)	Precision (CV%)	Trueness (BIAS%)
Fentanyl	10	14%	11%
	50	15%	-4%
	200	18%	-1%
Norfentanyl	10	5%	-20%
	50	6%	8%
	200	7%	0.1%
4-ANPP	10	18%	14%
	50	17%	2%
	200	12%	5%
β -OH-fentanyl	10	19%	15%
	50	20%	-8%
	200	20%	-1%
Acetyl fentanyl	10	5%	-14%
	50	8%	14%
	200	12%	-4%
Despropionyl p-fluorofentanyl	10	17%	18%
	50	19%	-17%
	200	17%	-0.2%
Tramadol	10	12%	-18%
	50	15%	9%
	200	10%	-1%
Hydrocodone	10	20%	-20%
	50	18%	16%
	200	19%	-2%
Oxycodone	10	14%	-12%
	50	7%	-2%
	200	15%	-1%

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

For Peer Review