

AperTO - Archivio Istituzionale Open Access dell'Università di Torino

Spectral Analysis for Weed Identification in Rice Fields

This is the author's manuscript

Original Citation:

Availability:

This version is available <http://hdl.handle.net/2318/1985610> since 2024-06-18T12:22:33Z

Published version:

DOI:10.13140/RG.2.2.10326.46400

Terms of use:

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)

Spectral Analysis for Weed Identification in Rice Fields

Bennani Z., Papandrea G., Beltramo A., Di Furia I., Sopegno A., Biglia A., Fogliatto S., Vidotto F.
University of Turin, Department of Agricultural, Forest and Food Sciences (DISAFA), Grugliasco Italy

International Temperate Rice Conference
New Orleans, USA June 5th – 8th

INTRODUCTION

Site Specific Weed Management (SSWM) has been recognized as a method to reduce herbicide use for a more sustainable weed management. Local application of herbicides aided by weed mapping allows the reduction of herbicides without impact on yield production.

In this study, we assess weed recognition in rice fields through hyperspectral data and calculated vegetations indices.

Specifically, we focus on weed species that are present in rice fields such as *Echinochloa crus-galli*, and *Persicaria maculosa* or species present in the levees such as *Digitaria sanguinalis* and *Potentilla reptans*.

METHODS

Field study

3 locations
(NW Italy)

Palestro
Olcenengo
Livorno Ferraris



Hyperspectral readings

- Spectral Evolution® RS-5400 spectroradiometer
- Readings on topmost leaf of at least 12 plants/species

- Spectral signature
- Vegetation indices
 - Normalized difference vegetation index (NDVI 705) (Gitelson and Merzlyak, 1994)
 - Carotenoids Index (CARI) (Zhou et al., 2017)
 - Enhanced Vegetation Index (EVI) (Huete et al., 1997)

Species discrimination based on leaf reflectance (software: PAST)



Species of interest



RESULTS




Figure 1. Light reflectance of rice and weed species in the range 400 – 1400 nm.




Figure 2. Vegetative indices calculated for the different species. For each index, the letters indicate homogeneous statistical groups.

- The Linear Discriminant Analysis (LDA) resulted in a linear model that distinguishes between ORYSA, POLLA, POLPE, and ECHCG with an **accuracy rate of 92%**.
- The model considers 4 coefficients based on the reflectance data between 400-1400 nm represented in their corresponding axes (**Figures 3 and 4**).
- Axis 1 describes 64% of the variability** between species, and when combined with axes 2 or 3 describes 85% or 74% of the observed variability, respectively.
- Axis 1 allows the distinction between monocots and dicots** with negative scores for the first and positive scores for the latter (**Figure 3**).
- A clear distinction can be observed between POLSS and PTLRE through axis 2 (**Figure 3**).
- Even though the combination of axes 2 and 3 describes only 30% of the variability, it shows a better discrimination between the grasses DIGSA, ECHCG, and ORYSA (**Figure 4**).





Figure 4. Rice and weeds discrimination with leaf reflectance: axes 2 – 3 at 95% confidence interval

- The spectral signatures (**Figure 1**) showed variation especially in the range 400-1400 nm.
- POLSS shows a **higher reflectance in the range 700-1400 nm** as typically occurs in dicotyledonous species compared to monocots (Jacquemoud and Ustin, 2019).
- The analysis of the vegetative indices allowed the **distinction between rice plants and weeds** by creating statistically homogenous groups (**Figure 2**).
- The **carotenoid index (CARI) resulted in the most categories (3)** by discriminating between the dicots POLSS and PTLRE, the monocots ECHCG and DIGSA, and ORYSA.
- The **Enhanced Vegetation Index (EVI)** was significantly higher in **POLSS** compared to the other species.
- NDVI 705** allowed a **distinction among the grasses** (DIGSA, ECHCG, and ORYSA).

CONCLUSION

This study substantiates the potential of spectral analysis for the distinction between rice and weed species, especially through the development of distinctive indices after discriminant analysis. Further studies are necessary to include other species and to upscale the methodology to hyperspectral scans acquired via drone flights.

REFERENCES

- Gitelson A. and Mark N. M. 1994. "Spectral Reflectance Changes Associated with Autumn Senescence of Aesculus Hippocastanum L. and Acer Platanoides L. Leaves. Spectral Features and Relation to Chlorophyll Estimation." *Journal of Plant Physiology* 143 (3): 286–92. [https://doi.org/10.1016/S0176-1617\(11\)81633-0](https://doi.org/10.1016/S0176-1617(11)81633-0).
- Huete A. R., Liu H. Q., Batchily K., and van Leeuwen W. 1997. "A Comparison of Vegetation Indices over a Global Set of TM Images for EOS-MODIS." *Remote Sensing of Environment* 59 (3): 440–51. [https://doi.org/10.1016/S0034-4257\(96\)00112-5](https://doi.org/10.1016/S0034-4257(96)00112-5).
- Jacquemoud S. and Ustin S. 2019. "Leaf Optical Properties." <https://doi.org/10.1017/9781108686457>.
- Zhou X., Huang W., Kong W., Ye H., Dong Y., and Casa R. 2017. "Assessment of Leaf Carotenoids Content with a New Carotenoid Index: Development and Validation on Experimental and Model Data." *International Journal of Applied Earth Observation and Geoinformation* 57 (May):24–35. <https://doi.org/10.1016/j.jag.2016.12.005>.

ACKNOWLEDGEMENTS

This work was supported by the grant Agritech National Research Center funded by European Union Next-Generation EU (Piano Nazionale di Ripresa e Resilienza (PNRR) - Missione 4 Componente 2, Investimento 1.4—D.D. n. 3175 18/12/2021 CN00000022).