

AperTO - Archivio Istituzionale Open Access dell'Università di Torino

**Medial anterior prefrontal cortex stimulation down-regulates implicit reactions to threats and prevents the return of fear**

**This is the author's manuscript**

*Original Citation:*

*Availability:*

This version is available <http://hdl.handle.net/2318/1985790> since 2024-10-11T08:45:10Z

*Published version:*

DOI:10.7554/eLife.85951

*Terms of use:*

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)

## eLife's Review Process

eLife works to improve the process of peer review so that it more effectively conveys the assessment of expert reviewers to authors, readers and other interested parties. In the future we envision a system in which research is first published as a preprint and the outputs of peer review are the primary way research is assessed, rather than journal title.

Our editorial process produces two outputs: i) an assessment by peers designed to be posted alongside a preprint for the benefit of the readers; ii) detailed feedback on the manuscript for the authors, including requests for revisions and suggestions for improvement.

Therefore we want to change how we construct and write peer reviews to make them useful to both authors and readers in a way that better reflects the work you put into reading and thinking about a paper.

eLife reviews now have three parts:

- An **evaluation summary** (in two or three sentences) that captures the major conclusions of the review in a concise manner, accessible to a wide audience.
- A **public review** that details the strengths and weaknesses of the manuscript before you, and discusses whether the authors' claims and conclusions are justified by their data.
- A set of private **recommendations for the authors** that outline how you think the science and its presentation could be strengthened.

All three sections will be used as the basis for an eLife publishing decision, which will, as always, be made after a consultation among the reviewers and editor. Each of the **public reviews** will be published (anonymously) alongside the preprint, together with a response from the authors if they choose. In the case of papers we reject after review, the authors can choose to delay posting until their paper has been published elsewhere.

If this is your first time going through this new process, we ask that you take some time to read our [Reviewer Guide](#), which discusses how we see each section will be used, what it should contain, and what we hope it accomplishes. And we remind you that, with the shift of reviews from private correspondence to public discourse, it is more important than ever that reviews are written in a **clear and constructive manner** appropriate for a public audience and mindful of the impact language choices might have on the authors.

## Information about the manuscript

### Medial anterior prefrontal cortex stimulation down-regulates implicit reactions to threats and prevents the return of fear

Tracking no: 04-01-2023-RA-eLife-85951R2

**Impact statement:** Implicit defensive reactions to a learned threat-predictive cue and novel stimuli may be significantly down-regulated through a repetitive Transcranial Magnetic Stimulation procedure over the medial anterior prefrontal cortex in humans.

**Competing interests:** No competing interests declared

#### Author contributions:

Eugenio Manassero: Conceptualization; Formal analysis; Investigation; Methodology; Writing - original draft; Writing - review and editing  
Giulia Concina: Formal analysis; Investigation  
Maria Clarissa Chantal Caraig: Formal analysis; Investigation  
Pietro Sarasso: Investigation  
Adriana Salatino: Investigation  
Raffaella Ricci: Supervision; Methodology; Project administration; Writing - review and editing  
Benedetto Sacchetti: Conceptualization; Supervision; Funding acquisition; Methodology; Writing - original draft; Project administration; Writing - review and editing

#### Funding:

Compagnia di San Paolo (CSP): Benedetto Sacchetti, CSTO167503; Fondazione Giovanni Gorla (FGG): Eugenio Manassero; Fondazione CRT (CRT Foundation): Benedetto Sacchetti; Banca d'Italia (Bank of Italy): Benedetto Sacchetti; Fondazione Cassa di Risparmio di Verona Vicenza Belluno e Ancona (Fondazione Cariverona): Benedetto Sacchetti; Italian Ministry of University and Research (MIUR): Benedetto Sacchetti  
The funders had no role in study design, data collection and interpretation, or the decision to submit the work for publication.

**Data Availability:**

All data generated or analysed during this study are included in the manuscript and supporting file.

N/A

**Ethics:**

Human Subjects: Yes Ethics Statement: Each participant provided written informed consent after receiving a complete description of the experimental procedures. All experimental procedures were approved by the Bioethics Committee of the University of Turin (protocols N. 19961 and N. 161427).

Clinical Trial: No Animal Subjects: No

1 **Title: Medial anterior prefrontal cortex stimulation down-regulates implicit reactions to**  
2 **threats and prevents the return of fear**

3

4 **Authors:** Eugenio Manassero<sup>1§</sup>, Giulia Concina<sup>1§</sup>, Maria Clarissa Chantal Caraig<sup>1</sup>, Pietro Sarasso<sup>2</sup>,  
5 Adriana Salatino<sup>2</sup>, Raffaella Ricci<sup>2</sup> and Benedetto Sacchetti<sup>1\*</sup>

6

7 **Affiliations:**

8 <sup>1</sup> Rita Levi-Montalcini Department of Neurosciences, University of Turin, Corso Raffaello 30,  
9 10125 Turin, Italy

10 <sup>2</sup> Department of Psychology, University of Turin, Via Giuseppe Verdi 10, 10124 Turin, Italy

11

12 <sup>§</sup>These authors contributed equally to the work

13 <sup>\*</sup>Correspondence to: [benedetto.sacchetti@unito.it](mailto:benedetto.sacchetti@unito.it)

14 **Abstract**

15 Down-regulating emotional overreactions toward threats is fundamental for developing treatments  
16 for anxiety and post-traumatic disorders. The prefrontal cortex (PFC) is critical for top-down  
17 modulatory processes, and despite previous studies adopting repetitive Transcranial Magnetic  
18 Stimulation (rTMS) over this region provided encouraging results in enhancing extinction, no  
19 studies have hitherto explored the effects of stimulating the medial anterior PFC (aPFC,  
20 encompassing the Brodmann area 10) on threat memory and generalization. Here we showed that  
21 rTMS over the aPFC applied before threat memory retrieval immediately decreases implicit  
22 reactions to learned and novel stimuli in humans. These effects enduringly persisted one week later  
23 in the absence of rTMS. No effects were detected on explicit recognition. Critically, rTMS over the  
24 aPFC resulted in a more pronounced reduction of defensive responses compared to rTMS targeting  
25 the dorsolateral PFC. These findings reveal a previously unexplored prefrontal region, the  
26 modulation of which can efficiently and durably inhibit implicit reactions to learned threats. This  
27 represents a significant advancement towards the long-term deactivation of exaggerated responses  
28 to threats.

## 29 **Introduction**

30 Emotional memories related to past threat experiences allow humans to predict future dangers and  
31 trigger adaptive defensive reactions when encountering learned threat-signaling cues<sup>1</sup>. However,  
32 extremely dangerous situations may lead to psychological disorders<sup>2</sup>. Furthermore, the ability to  
33 generalize defensive reactions to new stimuli enables organisms to anticipate potential threats and  
34 respond to them based on similar perilous experiences lived in the past. On the other hand,  
35 evaluation mechanisms excessively biased toward threat generalization (i.e. overgeneralization)  
36 may underlie anxiety disorders and trauma<sup>3</sup>. At the base of these processes, in a previous work<sup>4</sup> we  
37 observed that autonomic-implicit and cognitive-explicit tunings may diverge when humans are  
38 exposed to the same new stimuli, where cognitive generalization may enable a flexible evaluation  
39 of incoming cues to develop adaptive predictions of potential dangers. **The crosswise presence of**  
40 **overgeneralization in anxiety diseases, and the dissociation between autonomic and cognitive**  
41 **defensive response patterns, highlight the importance of including both implicit and explicit**  
42 **generalization tasks to characterize fear-related processes in humans.**

43         Attempting to down-regulate the emotional overreactions toward threat-predictive and new  
44 stimuli is one of the main routes for developing effective treatments for anxiety and post-traumatic  
45 disorders. Common approaches such as pharmacological treatments and cognitive-behavioral  
46 therapy (CBT) have demonstrated partial efficacy<sup>5</sup>, and recent evidence suggests that the functional  
47 outcome of behavioral methods may depend on the extent to which the prefrontal cortex is recruited  
48 during these processes<sup>6</sup>. Hence, new intervention strategies influencing the prefrontal dynamics  
49 would represent an important advance in the field<sup>7</sup>.

50         Previous studies adopted transcranial direct current stimulation (tDCS) or transcranial  
51 electrical stimulation (tES) to disrupt the consolidation of these memories<sup>8-10</sup>, potentiate extinction  
52 processes<sup>11,12</sup>, and narrow threat generalization patterns<sup>13</sup>, leading to contradictory results.  
53 According to one work<sup>8</sup>, cathodal stimulation over the dorsolateral prefrontal cortex (dlPFC)  
54 disrupted threat memory consolidation, with no enhancing effect of anodal stimulation. In contrast,

55 other studies found an increase in implicit responses with anodal stimulation<sup>9</sup> and no effect of  
56 cathodal stimulation<sup>10</sup> over the same site. Moreover, one study employing anodal stimulation over  
57 the dlPFC<sup>12</sup> revealed an improvement in extinction learning but no delayed effects on the recall of  
58 the extinction memory. A further investigation<sup>11</sup> reported that low-frequency alternating-current  
59 (AC) stimulation of the medial prefrontal cortex (mPFC) augmented the defensive responses,  
60 whereas direct-current (DC) stimulation widened threat generalization profiles.

61 An alternative neurostimulation approach is repetitive transcranial magnetic stimulation  
62 (rTMS), which ensures greater focality<sup>14,15</sup>. Some rTMS studies targeted the mPFC<sup>16</sup> and the  
63 posterior PFC<sup>17</sup> to obtain a successful enhancement of extinction learning, while others<sup>18,19</sup> targeted  
64 the dlPFC to disrupt threat-memory reconsolidation. Indeed, most rTMS-based research targeting  
65 the PFC has pursued an improvement of fear extinction, which may be followed by a return of fear  
66 with a change of context (i.e. renewal)<sup>20</sup> where prevention of relapse over time is the main  
67 challenge for therapies dedicated to post-traumatic and anxiety disorders. No previous studies  
68 reported significant effects in down-modulating the defensive responses triggered by a learned  
69 threatening stimulus without adopting fear extinction.

70 So far, human brain stimulation studies have been mainly focused on the dorsolateral region  
71 of the PFC<sup>7</sup>, partly because other prefrontal areas involved in the top-down regulation of subcortical  
72 threat-detection systems –such as the ventromedial PFC (vmPFC), are too deep to be reached with  
73 TMS<sup>17</sup>. However, within the PFC, a brain structure that is emerging to be engaged in downstream  
74 emotional regulation is the anterior prefrontal cortex (aPFC), also known as the frontopolar cortex  
75 or rostral frontal cortex. The aPFC encompasses the most anterior portion of the prefrontal cortex  
76 (Brodmann area 10, BA 10)<sup>21</sup> and extends over a wider cortical space in humans than in other  
77 species<sup>22</sup>. Even if it has not been included in fear network models so far, many studies<sup>23–25</sup>  
78 highlighted its role in emotional down-regulation. Anatomical projections have been found between  
79 the lateral<sup>25,26</sup> and the medial aPFC<sup>27</sup> and the amygdala, and functional connectivity has been  
80 detected between the aPFC and the vmPFC during fear down-regulation<sup>28</sup>. Notably, hypoactivation,

81 reduced connectivity, and altered thickness of aPFC were reported in PTSD patients<sup>29–32</sup>, whereas a  
82 longitudinal study<sup>33</sup> showed that strong activation of the aPFC resulted in a higher resilience against  
83 PTSD onset. Accordingly, enhanced aPFC activity and potentiated aPFC-vmPFC connectivity were  
84 detected after an effective therapy in PTSD patients<sup>34</sup>. Crucially, the aPFC is a surface area easily  
85 accessible with rTMS. However, to our knowledge, no study has been so far conducted to explore  
86 the effects of aPFC stimulation on the expression of a threat memory without extinction learning in  
87 humans.

88 In our current study, we posited that applying rTMS to the aPFC could influence implicit  
89 defensive responses to a learned threat-predictive stimulus and/or the conscious recognition of it.  
90 Subsequently, we explored additional hypotheses. The second hypothesis centered on the potential  
91 extension of rTMS dampening effects to new stimuli, thereby reducing threat generalization. The  
92 third hypothesis focused on the enduring persistence of rTMS effects on defensive responses over  
93 time. The final hypothesis proposed that the dampening effects achieved by stimulating the aPFC  
94 might surpass those observed when targeting the dorsolateral PFC.



95 **Results**

96 **aPFC-focused rTMS effects on implicit defensive reactions toward threat-predictive and new**  
97 **cues**

98 To explore the effects of an aPFC-centered rTMS on the implicit responses to a learned threat, we  
99 designed a three-session experiment starting with a threat learning session followed by an implicit  
100 retention test and a follow-up implicit re-test (Figure 1).

101 During the learning session, participants learned to associate an auditory cue (conditioned  
102 stimulus, CS, 800Hz) with a mild electric stimulation (unconditioned stimulus, US, individually  
103 calibrated intensity) in a given environment (context A). We adopted a single-cue learning  
104 paradigm because it more ecologically reflects real-life traumatic experiences<sup>35-39</sup>. To validate the  
105 between-groups homogeneity in the painful stimuli perception, we compared the post-conditioning  
106 US ratings, and we observed no significant differences between groups (Student's unpaired  $t$  test,  
107  $t_{(58)} = 0.799$ ,  $P = 0.428$ ,  $\eta_p^2 = 0.011$ ) (Table 1). We also did not observe significant differences  
108 between groups in SCRs to the CS during the preconditioning phase ( $t_{(58)} = 0.418$ ,  $P = 0.677$ ,  $\eta_p^2 =$   
109  $0.003$ ), to the CS during the conditioning phase ( $2 \times 15$  mixed ANOVA; main effect of group:  $F_{(1,52)}$   
110  $= 2.367$ ,  $P = 0.130$ ,  $\eta_p^2 = 0.044$ ; main effect of trial:  $F_{(8,762,455,600)} = 13.366$ ,  $P < 0.001$ ,  $\eta_p^2 = 0.204$ ;  
111 group  $\times$  trial interaction:  $F_{(8,762,455,600)} = 1.619$ ,  $P = 0.109$ ,  $\eta_p^2 = 0.030$ ; Student's unpaired  $t$  test on  
112 the averaged response,  $t_{(58)} = 1.290$ ,  $P = 0.202$ ,  $\eta_p^2 = 0.028$ ), nor to the US during the conditioning  
113 phase ( $t_{(58)} = 1.011$ ,  $P = 0.316$ ,  $\eta_p^2 = 0.017$ ) (Figure 2-figure supplement 1).

114 One week later, we tested the implicit memory of the learned association in control sham-  
115 stimulated subjects and in those who received rTMS over the aPFC shortly before the memory test.  
116 To locate this brain region, which corresponds to the BA 10<sup>40</sup>, we positioned the coil over the  
117 frontopolar midline electrode (Fpz) adopting the international 10–20 electroencephalogram (EEG)  
118 coordinate system<sup>41</sup> since previous rTMS studies<sup>16,42,43</sup> ensured this placement reached the aPFC.  
119 An offline 10-min session of 1Hz-rTMS targeting this neural site (aPFC,  $n = 30$ ) was applied

120 immediately before memory retrieval (Figure 2A). Control subjects underwent a 10-min sham  
121 stimulation procedure over the same cortical area (sham,  $n = 30$ ).

122 Memory retention was tested in a different environment from that where the learning had  
123 occurred (context B) to avoid any contextual influence on retrieval<sup>4,44–47</sup>. Indeed, the context shift  
124 for this session mirrors a real-life treatment setting –which unlikely takes place in the threatening  
125 location. To test implicit threat memory, we performed an implicit recognition task in which  
126 subjects were exposed to the CS while being recorded in their evoked autonomic reactions (i.e.,  
127 electrodermal skin conductance responses, SCRs). No US shocks were delivered during this phase.  
128 Besides the CS, participants were presented with two novel but perceptually similar tones (NS<sub>1</sub>,  
129 1000Hz; NS<sub>2</sub>, 600Hz) to study threat generalization. Auditory frequencies of NSs were selected to  
130 obtain a slowly decaying gradient of defensive tunings<sup>4,48,49</sup>. To test the effects of rTMS on memory  
131 retention, we compared the between-group differences as well as the within-group differences from  
132 the acquisition phase to the testing phase through a  $2 \times 2$  mixed ANOVA. This analysis yielded a  
133 not significant main effect of group ( $F_{(1,58)} = 2.015$ ,  $P = 0.161$ ,  $\eta_p^2 = 0.034$ ), a not significant main  
134 effect of phase ( $F_{(1,58)} = 0.053$ ,  $P = 0.818$ ,  $\eta_p^2 = 0.001$ ) and a significant group  $\times$  phase interaction  
135 ( $F_{(1,58)} = 13.445$ ,  $P = 0.001$ ,  $\eta_p^2 = 0.188$ ). Simple main effects analysis revealed no significantly  
136 different mean CS-evoked SCRs between groups during the conditioning phase ( $P = 0.506$ ;  
137 Bonferroni corrected). On the contrary, during the test phase subjects that received rTMS over the  
138 aPFC exhibited weakened CS-related SCRs than those observed in the sham group ( $P = 0.006$ ;  
139 Bonferroni corrected). Moreover, the aPFC group showed reduced autonomic responses to the CS  
140 from conditioning to test ( $P = 0.008$ ; Bonferroni corrected) whereas the sham group displayed  
141 increased mean SCRs to the CS from conditioning to test ( $P = 0.018$ ; Bonferroni corrected) (Figure  
142 2B,C). This data indicates that the rTMS procedure affected SCRs triggered by memory retrieval  
143 performed shortly after rTMS. To the best of our knowledge, this is the first evidence that brain  
144 stimulation may promptly attenuate implicit defensive reactions during memory retrieval.

145 In the test session, we also analyzed threat generalization to the NSs through a  $2 \times 3$  mixed  
146 ANOVA, which showed a significant main effect of group ( $F_{(1,58)} = 5.310$ ,  $P = 0.025$ ,  $\eta_p^2 = 0.084$ ),  
147 a not significant main effect of tone ( $F_{(2,116)} = 0.690$ ,  $P = 0.504$ ,  $\eta_p^2 = 0.012$ ) and a not significant  
148 group  $\times$  tone interaction ( $F_{(2,116)} = 1.301$ ,  $P = 0.276$ ,  $\eta_p^2 = 0.022$ ), revealing that the aPFC group  
149 displayed overall attenuated responses to tones relative to the sham condition (Figure 2D,E).

150 We next sought to disambiguate whether the rTMS effects were due to a general down-  
151 regulation of electrodermal responsivity, or whether they specifically targeted the threat memory.  
152 To this end, subjects were presented with an unconditioned threatening stimulus consisting of a  
153 female scream sample (unconditioned stimulus 2, US<sub>2</sub>) while being recorded in their SCRs. No  
154 significant differences emerged between conditions ( $t_{(58)} = 0.334$ ,  $P = 0.739$ ,  $\eta_p^2 = 0.002$ ), indicating  
155 that the rTMS did not cause an overall inhibition of electrodermal reactivity (Figure 2F).

156 To test whether and to what extent rTMS-related outcomes endured beyond the after-effect  
157 window and persisted over a long-term period, we planned a follow-up session. One week after the  
158 threat memory retrieval test, all participants returned to the conditioning room (context A) and  
159 underwent a re-testing phase, identical to the testing one except for the absence of rTMS  
160 administration. This phase also allowed us to test a possible renewal effect<sup>20</sup> since subjects were re-  
161 exposed to the original threatening environment.

162 Concerning the implicit responses to the CS, a  $2 \times 2$  mixed ANOVA showed a not  
163 significant main effect of group ( $F_{(1,58)} = 1.952$ ,  $P = 0.168$ ,  $\eta_p^2 = 0.033$ ), a significant main effect of  
164 phase ( $F_{(1,58)} = 7.690$ ,  $P = 0.007$ ,  $\eta_p^2 = 0.117$ ) and a significant group  $\times$  phase interaction ( $F_{(1,58)} =$   
165  $9.966$ ,  $P = 0.003$ ,  $\eta_p^2 = 0.147$ ). Simple main effects analysis revealed that participants of the aPFC  
166 group persisted in displaying weaker SCRs than those observed in the sham group ( $P = 0.006$ ;  
167 **Bonferroni corrected**). Moreover, the aPFC group persisted in showing a decrease of defensive  
168 reactions to the CS from conditioning to follow-up ( $P < 0.001$ ; **Bonferroni corrected**), while the  
169 sham group did not display significantly different SCRs in the two phases ( $P = 0.787$ ; **Bonferroni**  
170 **corrected**) (Figure 2G,H).

171           These findings support an enduring effect of the aPFC-rTMS in attenuating the long-term  
172 implicit defensive responses to the learned threat-predictive cue, even with the re-exposition to the  
173 environment where threat learning had occurred. We next analyzed the autonomous response  
174 patterns to the female scream sample (unconditioned stimulus 2, US<sub>2</sub>) and again we found that  
175 reactions did not differ between groups ( $t_{(58)} = 0.057$ ,  $P = 0.955$ ,  $\eta_p^2 < 0.001$ ) (Figure 2I). Thus, the  
176 persistent effect was expressed notwithstanding an unaffected electrodermal overall reactivity.

177           An important aspect to consider is that rTMS application over the forehead can be  
178 subjectively perceived as unpleasant. We, therefore, investigated whether an rTMS-related  
179 discomfort before memory retrieval might have provoked habituation to unpleasant stimulations,  
180 leading to a reduction in SCR levels during CS presentations. We repeated the entire experiment in  
181 one further group (ctrl discomfort,  $n = 10$ ) by replacing the rTMS procedure with a 10-min  
182 discomfort-inducing procedure over the same site of the forehead to mimic the rTMS-evoked  
183 unpleasant sensations in the absence of neural stimulation effects. This group showed no  
184 significantly different CS-evoked SCR levels to those of the sham group during the test session as  
185 well as during the follow-up session (Figure 2-figure supplement 2). Thus, the discomfort  
186 experienced during the rTMS procedure did not contribute to the reduction of electrodermal  
187 responses observed in the aPFC-stimulated group.

188

### 189 **aPFC-focused rTMS effects on the explicit memory recognition and perceptual discrimination**

190 We then investigated the effect of rTMS over the aPFC on the retention of explicit-declarative  
191 threat memories. A further group of subjects that received the identical 1Hz-rTMS procedure over  
192 the aPFC (aPFC-E,  $n = 21$ ) and a further control group (sham-E,  $n = 21$ ) underwent an explicit two-  
193 alternative forced-choice (2AFC) recognition task, in which they were presented with a random  
194 sequence of tone pairs, each composed of the CS and one of the two NSs. Subjects were asked to  
195 consciously identify which stimulus of each pair was the one previously paired with the US (i.e., the  
196 CS), and to provide a subjective confidence level for each choice using a scale ranging from 0

197 (completely unsure) to 10 (completely sure)<sup>4,50</sup>. Both groups reported not significantly different  
198 post-conditioning US ratings ( $t_{(40)} = 0.339$ ,  $P = 0.737$ ,  $\eta_p^2 = 0.003$ ) and successfully identified the  
199 CS amongst the NSs with an accuracy level above the 50% chance level (aPFC-E:  $t_{(20)} = 9.226$ ,  $P <$   
200  $0.001$ ,  $\eta_p^2 = 0.810$ ; sham-E:  $t_{(20)} = 14.240$ ,  $P < 0.001$ ,  $\eta_p^2 = 0.910$ ). A between-groups comparison  
201 ( $t_{(40)} = 1.114$ ,  $P = 0.272$ ,  $\eta_p^2 = 0.030$ ) showed no differences in the explicit recognition accuracy  
202 (Figure 3A). The two groups were not differently confident when making their choices ( $t_{(40)} =$   
203  $0.842$ ,  $P = 0.405$ ,  $\eta_p^2 = 0.017$ ) (Figure 3B), thereby supporting the lack of rTMS-related effects.

204 Next, since a previous study<sup>13</sup> targeting the vmPFC modulated perceptual discrimination  
205 processes, we implemented a 2AFC perceptual task in which we investigated the ability of  
206 participants to sensory discriminate between the CS and the two NSs by collecting binary ‘same or  
207 different’ judgments as well as confidence ratings. The perceptual discrimination test yielded no  
208 significant between-groups differences in accuracy ( $t_{(40)} = 1.362$ ,  $P = 0.181$ ,  $\eta_p^2 = 0.044$ ) as well as  
209 confidence levels ( $t_{(40)} = 0.917$ ,  $P = 0.365$ ,  $\eta_p^2 = 0.021$ ). Indeed, both groups discriminated the CS  
210 from the NSs with high precision (aPFC-E:  $0.980 \pm 0.015$  SEM; sham-E:  $1.000 \pm 0.000$  SEM) and  
211 with no different confidence levels (aPFC-E:  $9.409 \pm 0.153$  SEM; sham-E:  $9.586 \pm 0.117$  SEM),  
212 thereby showing no rTMS effects on sensory abilities.

213 These data suggest that the pre-retrieval rTMS procedure over the aPFC did not affect the  
214 explicit recognition nor the perceptual discrimination of a learned threat.

215 During the follow-up session, explicit recognition patterns demonstrated an over-chance  
216 accuracy level for each group (aPFC-E:  $t_{(20)} = 13.780$ ,  $P < 0.001$ ,  $\eta_p^2 = 0.905$ ; sham-E:  $t_{(20)} = 7.162$ ,  
217  $P < 0.001$ ,  $\eta_p^2 = 0.720$ ). Again, here there were no between-group differences ( $t_{(40)} = 1.024$ ,  $P =$   
218  $0.312$ ,  $\eta_p^2 = 0.026$ ) since both groups achieved a high recognition accuracy (Figure 3C). Groups did  
219 also not report different confidence levels ( $t_{(40)} = 0.084$ ,  $P = 0.934$ ,  $\eta_p^2 < 0.001$ ) (Figure 3D).

220 As in the case of the previous session, we did not observe significant between-group  
221 differences in the perceptual discrimination ( $t_{(40)} = 1.000$ ,  $P = 0.323$ ,  $\eta_p^2 = 0.024$ ) and the respective  
222 confidence ratings ( $t_{(40)} = 0.149$ ,  $P = 0.882$ ,  $\eta_p^2 < 0.001$ ). Indeed, the discrimination accuracy

223 (aPFC-E:  $1.000 \pm 0.000$  SEM; sham-E:  $0.993 \pm 0.007$  SEM) and the self-assessed confidence  
224 (aPFC-E:  $9.598 \pm 0.147$  SEM; sham-E:  $9.633 \pm 0.182$  SEM) were high in each condition.

225

## 226 **Topographical selectivity of rTMS effects on implicit defensive responses to threat-predictive** 227 **and new cues**

228 To ascertain the topographical selectivity, in one further condition (OC,  $n = 30$ ) we applied the  
229 rTMS over the left occipital cortex as an active control site (Figure 4A) and we contrasted its  
230 implicit reactions with those of the group stimulated over the aPFC.

231 No differences emerged between the two conditions in terms of post-conditioning US  
232 ratings ( $t_{(58)} = 0.000$ ,  $P = 1.000$ ,  $\eta_p^2 = 0.000$ ) (Table 1), SCR responses to the CS during the  
233 preconditioning phase ( $t_{(58)} = 1.037$ ,  $P = 0.304$ ,  $\eta_p^2 = 0.018$ ), to the CS during the conditioning phase  
234 ( $2 \times 15$  mixed ANOVA; main effect of group:  $F_{(1,54)} = 0.124$ ,  $P = 0.726$ ,  $\eta_p^2 = 0.002$ ; main effect of  
235 trial:  $F_{(9,368,505,856)} = 13.341$ ,  $P < 0.001$ ,  $\eta_p^2 = 0.198$ ; group  $\times$  trial interaction:  $F_{(9,368,505,856)} = 0.994$ ,  $P$   
236  $= 0.445$ ,  $\eta_p^2 = 0.018$ ; Student's unpaired  $t$  test on the averaged response,  $t_{(58)} = 0.162$ ,  $P = 0.872$ ,  $\eta_p^2$   
237  $< 0.001$ ), and to the US during the conditioning phase ( $t_{(58)} = 1.210$ ,  $P = 0.231$ ,  $\eta_p^2 = 0.025$ ) (Figure  
238 4-figure supplement 1).

239 Next, we analyzed implicit reactions toward the CS in both conditioning and test sessions. A  
240  $2 \times 2$  mixed ANOVA revealed a not significant group main effect ( $F_{(1,58)} = 2.952$ ,  $P = 0.091$ ,  $\eta_p^2 =$   
241  $0.048$ ), a not significant phase main effect ( $F_{(1,58)} = 2.027$ ,  $P = 0.160$ ,  $\eta_p^2 = 0.034$ ), and a significant  
242 group  $\times$  phase interaction ( $F_{(1,58)} = 4.705$ ,  $P = 0.034$ ,  $\eta_p^2 = 0.075$ ). CS-related SCRs did not differ  
243 between groups during conditioning ( $P = 0.798$ ; Bonferroni corrected) but, during the test, the aPFC  
244 group exhibited weaker defensive responses than the OC group ( $P = 0.019$ ; Bonferroni corrected).  
245 Unlike the aPFC group, whose implicit reactions to the CS diminished from conditioning to test ( $P$   
246  $= 0.014$ ; Bonferroni corrected), the OC group's responses did not differ in the two phases ( $P =$   
247  $0.600$ ; Bonferroni corrected) (Figure 4B,C).

248 No significant between-group differences were observed in implicit responses to new tones  
249 ( $2 \times 3$  mixed ANOVA; main effect of group:  $F_{(1,58)} = 2.775$ ,  $P = 0.101$ ,  $\eta_p^2 = 0.046$ ; main effect of  
250 tone:  $F_{(2,116)} = 5.857$ ,  $P = 0.004$ ,  $\eta_p^2 = 0.092$ ; group  $\times$  tone interaction:  $F_{(2,116)} = 3.739$ ,  $P = 0.027$ ,  $\eta_p^2$   
251  $= 0.061$ ). While the CS triggered weaker reactions in the aPFC group ( $P = 0.019$ ; Bonferroni  
252 corrected), both the NS<sub>1</sub> ( $P = 0.203$ ; Bonferroni corrected) and the NS<sub>2</sub> ( $P = 0.323$ ; Bonferroni  
253 corrected) elicited not significantly different responses in the two conditions. These findings  
254 underscored the selectivity of divergent rTMS effects in the aPFC and OC groups specifically for  
255 the CS. Fear tuning analysis of the aPFC group's implicit reactions unveiled no differences in SCR  
256 amplitudes elicited by the CS and the NS<sub>1</sub> ( $P = 0.378$ ; Bonferroni corrected), by the CS and the NS<sub>2</sub>  
257 ( $P = 1.000$ ; Bonferroni corrected), and by the NSs ( $P = 0.552$ ; Bonferroni corrected). In the case of  
258 the OC group, implicit reactions were not different for the CS and the NS<sub>1</sub> ( $P = 0.876$ ; Bonferroni  
259 corrected) but the NS<sub>2</sub> evoked lower SCRs than the CS ( $P < 0.001$ ; Bonferroni corrected) and the  
260 NS<sub>1</sub> ( $P = 0.041$ ; Bonferroni corrected) (Figure 4D,E). Furthermore, no significant group differences  
261 were detected in SCRs elicited by US<sub>2</sub> during the test session ( $t_{(58)} = 0.175$ ,  $P = 0.862$ ,  $\eta_p^2 < 0.001$ )  
262 (Figure 4F).

263 The distinctive pattern toward the learned threatening cue persisted during the follow-up  
264 session ( $2 \times 2$  mixed ANOVA; main effect of group:  $F_{(1,58)} = 2.141$ ,  $P = 0.149$ ,  $\eta_p^2 = 0.036$ ; main  
265 effect of phase:  $F_{(1,58)} = 26.023$ ,  $P < 0.001$ ,  $\eta_p^2 = 0.310$ ; group  $\times$  phase interaction:  $F_{(1,58)} = 3.167$ ,  $P$   
266  $= 0.080$ ,  $\eta_p^2 = 0.052$ ). The aPFC group continued to react more dimly to the CS compared to the OC  
267 group ( $P = 0.026$ ; Bonferroni corrected). Both the aPFC ( $P < 0.001$ ; Bonferroni corrected) and the  
268 OC ( $P = 0.022$ ; Bonferroni corrected) groups showed decreased responses relative to conditioning  
269 (Figure 4G,H). Conversely, no significant differences were observed in SCRs evoked by US<sub>2</sub>  
270 during the follow-up session ( $t_{(58)} = 0.574$ ,  $P = 0.568$ ,  $\eta_p^2 = 0.006$ ) (Figure 4I).

271

272 **Comparison between the effects of rTMS administered over the anterior versus the**  
273 **dorsolateral prefrontal cortex**

274 Next, we asked whether the findings we obtained by targeting the aPFC were finely specific for this  
275 site or, alternatively, they overlapped with those observed by targeting other prefrontal sub-regions.  
276 For this purpose, in one further group (dlPFC,  $n = 30$ ) we applied the same rTMS procedure over  
277 the left dorsolateral PFC (Figure 5A) and we then compared the implicit patterns of this group with  
278 those displayed by the aPFC condition. We selected the left dlPFC since previous studies<sup>e.g. 17</sup>  
279 targeted the left hemisphere for testing the rTMS effects on the PFC, and some evidence<sup>see 7</sup>  
280 suggested that inhibitory tDCS and rTMS over the left dlPFC may disrupt threat memory  
281 consolidation.

282 We found no significant differences between the two conditions in the post-conditioning US  
283 ratings ( $t_{(58)} = 0.908$ ,  $P = 0.368$ ,  $\eta_p^2 = 0.014$ ) (Table 1), in SCRs to the CS during the  
284 preconditioning phase ( $t_{(58)} = 0.967$ ,  $P = 0.337$ ,  $\eta_p^2 = 0.016$ ), to the CS during the conditioning phase  
285 ( $2 \times 15$  mixed ANOVA; main effect of group:  $F_{(1,51)} = 0.026$ ,  $P = 0.873$ ,  $\eta_p^2 = 0.001$ ; main effect of  
286 trial:  $F_{(8,026,409,333)} = 12.135$ ,  $P < 0.001$ ,  $\eta_p^2 = 0.192$ ; group  $\times$  trial interaction:  $F_{(8,026,409,333)} = 1.042$ ,  $P$   
287  $= 0.403$ ,  $\eta_p^2 = 0.020$ ; Student's unpaired  $t$  test on the averaged response,  $t_{(58)} = 0.378$ ,  $P = 0.707$ ,  $\eta_p^2$   
288  $= 0.002$ ), and to the US during the conditioning phase ( $t_{(58)} = 1.752$ ,  $P = 0.085$ ,  $\eta_p^2 = 0.050$ ) (Figure  
289 5-figure supplement 1).

290 Then we compared the implicit reactions toward the CS during conditioning and test  
291 sessions. A  $2 \times 2$  mixed ANOVA indicated a not significant main effect of group ( $F_{(1,58)} = 1.874$ ,  $P$   
292  $= 0.176$ ,  $\eta_p^2 = 0.031$ ), a not significant main effect of phase ( $F_{(1,58)} = 0.122$ ,  $P = 0.729$ ,  $\eta_p^2 = 0.002$ ),  
293 and a significant group  $\times$  phase interaction ( $F_{(1,58)} = 10.810$ ,  $P = 0.002$ ,  $\eta_p^2 = 0.157$ ). CS-evoked  
294 SCRs did not differ between the two groups during conditioning ( $P = 0.647$ ; Bonferroni corrected)  
295 while during the test we found weaker defensive responses in the aPFC group relative to the dlPFC  
296 group ( $P = 0.009$ ; Bonferroni corrected). At odds with the aPFC group whose implicit reactions to  
297 the CS were diminished from conditioning to test ( $P = 0.013$ ; Bonferroni corrected), the dlPFC  
298 group increasingly responded during the test relative to conditioning ( $P = 0.042$ ; Bonferroni



299 corrected) (Figure 5B,C). This incremental trend is in line with a previous study that delivered a  
300 1Hz-rTMS protocol over the left dlPFC<sup>18</sup>.

301 We found no between-groups differences in the implicit responses to the new tones ( $2 \times 3$   
302 mixed ANOVA; main effect of group:  $F_{(1,58)} = 3.967$ ,  $P = 0.051$ ,  $\eta_p^2 = 0.064$ ; main effect of tone:  
303  $F_{(2,116)} = 2.819$ ,  $P = 0.064$ ,  $\eta_p^2 = 0.046$ ; group  $\times$  tone interaction:  $F_{(2,116)} = 3.286$ ,  $P = 0.041$ ,  $\eta_p^2 =$   
304  $0.054$ ) since to both the NS<sub>1</sub> ( $P = 0.188$ ; Bonferroni corrected) and the NS<sub>2</sub> ( $P = 0.110$ ; Bonferroni  
305 corrected) were not significantly different. These data showed that the divergent rTMS effects in the  
306 aPFC and the dlPFC groups were selective for the CS. Fear tuning analysis of the dlPFC group's  
307 implicit reactions revealed no different SCR amplitudes elicited by the CS and the NS<sub>1</sub> ( $P = 0.158$ ;  
308 Bonferroni corrected) and by the NSs ( $P = 0.721$ ; Bonferroni corrected), but the NS<sub>2</sub> evoked lower  
309 SCRs than the CS ( $P = 0.014$ ; Bonferroni corrected) (Figure 5D,E). We also detected no significant  
310 differences between groups in the SCRs elicited by the US<sub>2</sub> during the test session ( $t_{(58)} = 1.762$ ,  $P =$   
311  $0.083$ ,  $\eta_p^2 = 0.051$ ) (Figure 5F).

312 The different pattern toward the learned threatening cue was replicated during the follow-up  
313 session ( $2 \times 2$  mixed ANOVA; main effect of group:  $F_{(1,58)} = 3.751$ ,  $P = 0.058$ ,  $\eta_p^2 = 0.061$ ; main  
314 effect of phase:  $F_{(1,58)} = 3.114$ ,  $P = 0.083$ ,  $\eta_p^2 = 0.051$ ; group  $\times$  phase interaction:  $F_{(1,58)} = 15.248$ ,  $P$   
315  $< 0.001$ ,  $\eta_p^2 = 0.208$ ) since the aPFC group persisted in more dimly reacting to the CS with respect  
316 to the dlPFC group ( $P = 0.001$ ; Bonferroni corrected), and the aPFC group endured in displaying  
317 attenuated responses relative to conditioning ( $P < 0.001$ ; Bonferroni corrected) while the dlPFC  
318 group did not ( $P = 0.136$ ; Bonferroni corrected) (Figure 5G,H). No significant differences were  
319 instead observed in SCRs evoked by the US<sub>2</sub> during the follow-up session ( $t_{(58)} = 1.927$ ,  $P = 0.059$ ,  
320  $\eta_p^2 = 0.060$ ) (Figure 5I).

321 These findings demonstrated that rTMS over the left dorsolateral PFC did not diminish  
322 implicit defensive reactions in the absence of an extinction paradigm, as in other previous studies<sup>16–</sup>  
323 <sup>18</sup>. Meanwhile, rTMS targeting the aPFC proved to be effective in achieving this outcome.

324 **Discussion**

325 In this study, we found that implicit reactions to both learned and novel stimuli were significantly  
326 down-regulated following a 1Hz-rTMS procedure over the aPFC.

327 So far, most rTMS studies targeting the prefrontal cortex have been conducted to enhance  
328 fear extinction processes. A study<sup>16</sup> administering one session of 10Hz-rTMS over the mPFC  
329 observed enhancement of extinction learning. These behavioral results were mirrored by the  
330 functional near-infrared spectroscopy (fNIRS) findings, which revealed increased mPFC activity in  
331 the stimulated group relative to the sham group<sup>16</sup>. Subsequently, Raij *et al.* (2018) delivered brief  
332 20Hz-rTMS trains over the left posterior PFC –a region that showed robust functional connectivity  
333 with the vmPFC– during extinction learning and found a reduction of defensive responses during  
334 extinction recall.

335 Our study differs from the previous ones because we tested rTMS effects over the medial  
336 anterior prefrontal cortex (medial BA 10), and we did not include extinction training before  
337 retrieval. We observed a significant decrease in defensive reactions shortly after rTMS, and this  
338 effect was maintained until the follow-up session. Thus, we identified a previously unexplored  
339 prefrontal region, the modulation of which can efficiently and durably inhibit implicit reactions to  
340 learned threats. These dampening effects may be due to the fact that rTMS over the aPFC have  
341 directly modulated the defensive responses activated by the implicit threat memory trace.  
342 Alternatively, the rTMS procedure over the aPFC may have inhibited the recall of the CS-US  
343 association, preventing the defensive responses from being activated by the CS. This possibility  
344 would be in line with a large body of literature on humans<sup>see 51</sup> which demonstrates the importance  
345 of the medial PFC for value-based processing.

346 **Autonomic reactions to the new tones in the aPFC group relative to the sham control group**  
347 **did not support the conclusion that rTMS targeted threat generalization, leaving open the question**  
348 **of the specificity of rTMS effects. However, the lack of between-group differences in the autonomic**  
349 **responses to the US<sub>2</sub> seems to suggest that the observed effect may be memory-related and not due**

350 to a general dampening of autonomic reactivity. Interestingly, defensive responses toward the NSs  
351 were decreased following the stimulation of the left occipital cortex (OC group, BA18/19). This  
352 effect might be explained by the fact that anatomical and functional reciprocal projections between  
353 the medial BA10 and visual association cortices (including BA17/18/19) have been traced via the  
354 fronto-occipital fasciculus (FOF) of the human brain<sup>52–54</sup> but see 27.

355       Regarding the persistence of inhibitory effects during the follow-up session, different factors  
356 may have contributed to this result. Firstly, the inhibition of SCR responses induced by rTMS  
357 during the mnemonic retention test could have persistently reduced such conditioned responses  
358 even at a distance from the treatment. Moreover, the inhibition of these responses during the test  
359 might have boosted the extinction of these responses, contributing to keeping them low over time.  
360 On this possibility, it should be pointed out that one core knowledge about extinction is that under  
361 certain circumstances –such as a simple passage of time (i.e., spontaneous recovery) or a change in  
362 surrounding context (i.e., renewal)– extinguished reactions triggered by the CS may reoccur, giving  
363 rise to the phenomenon known as return of fear<sup>20,55,56</sup>. To test potential renewal phenomena, which  
364 have not been investigated in the aforementioned studies<sup>16,17</sup>, we opted for a context-shift amongst  
365 the learning (context A), the test (context B), and the follow-up phase (context A), and we found  
366 down-regulated defensive reactions in both the test and the follow-up phases. These data  
367 demonstrated that the aPFC-rTMS protocol long-term reduced threat memory expression in a  
368 different context as well as in the context in which the threatening experience had occurred, thus  
369 preventing the return of fear. Finally, we cannot exclude that the rTMS applied immediately before  
370 the mnemonic retention test interfered with the reconsolidation process that is known to occur after  
371 this test<sup>19</sup>, resulting in a persistent impairment in the retention of this mnemonic trace.

372       To potentiate the neural activity of the PFC, both the aforementioned studies<sup>16,17</sup> adopted  
373 high-frequency rTMS protocols –which are conventionally considered excitatory of proximal brain  
374 activity<sup>57</sup>. In our study, we adopted a low-frequency rTMS protocol –which is conventionally  
375 considered inhibitory<sup>57</sup>. Recent evidence, however, challenged this common frequency-dependent

376 rule<sup>58</sup>. Resting-state functional magnetic resonance imaging (fMRI) studies demonstrated that 1Hz-  
377 rTMS protocols may also induce downstream distal effects and enhance functional connectivity  
378 amongst the brain regions located underneath the coil and remote brain areas of the stimulated  
379 neural network<sup>59</sup>. Additionally, some studies<sup>60,61</sup> reported that 1Hz-rTMS procedures delivered over  
380 the PFC may paradoxically increase regional cerebral blood flow (rCBF).

381 The dorsolateral PFC is another prefrontal region that is assumed to be critically involved in  
382 threat learning<sup>62,63</sup> and the down-regulation of the cortico-meso-limbic network<sup>64</sup>. One  
383 investigation<sup>18</sup> probed the effects of a 1Hz-rTMS over the dlPFC after memory reactivation to  
384 disrupt threat-memory reconsolidation. Stimulated groups failed to discriminate between  
385 threatening and safe stimuli, with an increase in autonomic responses to these last ones. A more  
386 recent study<sup>19</sup> adopted the continuous theta-burst stimulation (cTBS) over the right dlPFC during  
387 the reconsolidation window and successfully decreased the defensive responses for threat  
388 memories. In our study, we found an immediate and long-term reduction of defensive responses to  
389 the CS only in subjects that were stimulated over the aPFC, while reactions to the NSs were  
390 decreased in both conditions. This evidence suggests that targeting the aPFC might represent a more  
391 promising approach for therapeutic applications. The lack of any down-regulation of CS-evoked  
392 reactions that we found in the dlPFC group, at odds with previous studies targeting the same  
393 cortical area<sup>18,19</sup>, might be due either to the fact that we did not adopt an extinction paradigm, or to  
394 the different brain stimulation approach (rTMS vs cTBS).

395 The neural mechanisms by which rTMS over the aPFC decreases threat-conditioned  
396 responses can be manifold. Fear memories are formed and retrieved by an intricate neural network  
397 encompassing the amygdala<sup>65</sup>, the cerebellum<sup>66-68</sup>, and sensory cortices<sup>46,69-77</sup>. Indeed, previous  
398 evidence showed both structural connections between the aPFC and the amygdala<sup>25-27</sup> and a  
399 connectivity pathway of downstream modulation from the aPFC to the vmPFC<sup>34</sup>. This projection is  
400 activated during fear regulation<sup>28</sup>, possibly supporting the vmPFC in top-down modulating the  
401 amygdala<sup>78</sup>. Through the direct or indirect connections of the aPFC with these areas, it might be that

402 the effects of focal manipulations of aPFC activity reflect complex and dynamic changes in the  
403 overall neural network state and/or influence the activity of some of these areas.

404 Although previous studies enlightened the role of the medial BA10 and BA10–posterior  
405 hippocampus functional connectivity in episodic memory retrieval<sup>see 79</sup>, we did not detect any  
406 rTMS-driven effect on explicit recognition memory. The observed divergence between autonomous  
407 and declarative patterns might have been due to a selective rTMS action upon the neural system  
408 supporting implicit threat processing, which has been widely dissociated from the neural system  
409 underlying explicit memory processes<sup>80–82</sup>. Critically, an rTMS procedure that shapes implicit  
410 overreactions to learned threats without affecting conscious knowledge of danger might represent a  
411 strategic advantage for therapeutic applications.

412 Since prevention of relapse is the main challenge for therapies dedicated to post-traumatic  
413 and anxiety disorders, our findings may represent an advance in this direction by providing a  
414 potential strategy to deactivate emotional overreactions and, most of all, to prevent the return of  
415 fear. Future research perspectives might consist of exploring this rTMS application over the aPFC  
416 in clinical populations displaying high levels of anxiety or suffering from anxiety disorders and  
417 PTSDs.

## 418 **Materials and Methods**

### 419 **Participants**

420 All participants ( $n = 183$ ) were healthy volunteers (mean age:  $23.86 \pm 2.90$ , 74 males and 109  
421 females) with no history of psychiatric disorders, neurological illnesses, cardiovascular diseases,  
422 illegal drug use, musical training, or any other exclusion criteria for rTMS administration<sup>83</sup>. During  
423 the pre-experimental screening phase, each volunteer was also administered the *State-Trait Anxiety*  
424 *Inventory Form Y*<sup>84,85</sup>, and those who showed a score  $>80$  in the sum of the two subscales (State +  
425 Trait anxiety) were not included in the sample (see Table 1 for all groups' mean State-Trait Anxiety  
426 Inventory scores). Participants were then randomly assigned to each experimental condition, based  
427 on sex and age (see Table 1 for all groups' mean age and sex distribution). We discarded eleven  
428 participants because of a complete absence of skin conductance responses (SCRs) during the test  
429 session, leaving a total of 172 participants. Each participant provided written informed consent after  
430 receiving a complete description of the experimental procedures. All experimental procedures were  
431 performed in accordance with the ethical standards of the Declaration of Helsinki and were  
432 approved by the Bioethics Committee of the University of Turin (protocols N. 19961 and N.  
433 161427).

434

### 435 **Auditory stimuli**

436 Auditory stimuli were pure sine wave tones with oscillation frequencies of 800Hz (CS), 1000Hz  
437 (NS<sub>1</sub>), and 600Hz (NS<sub>2</sub>), lasting 6s with onset/offset ramps of 5ms. Tones were digitally generated  
438 using Audacity 2.1.2 software (Audacity® freeware). The unconditioned threatening stimulus (US<sub>2</sub>)  
439 consisted of a woman scream sample lasting 4s. All auditory stimuli were binaurally delivered  
440 through headphone speakers (Direct Sound EX29) at 50 dB intensity. All experimental scenarios  
441 were controlled by Presentation® 21.1 software (NeuroBehavioral Systems, Berkeley, CA).

442

443

444

### 445 **Preconditioning**

446 This phase consisted of the presentation of 4 trials of the CS (800Hz) with an inter-trial-interval  
447 (ITI) randomly ranging between 21s and 27s. SCRs were recorded during this phase to provide a  
448 baseline response pattern to the 800Hz tone for each participant. At the end of this phase,  
449 participants were asked to confirm whether the tones were easily audible but not too loud or  
450 annoying.

451

### 452 **Unconditioned stimulus calibration procedure**

453 Before starting with the calibration procedure, systolic and diastolic blood pressure was measured to  
454 prevent possible hypo-arousal reactions caused by basal hypotension. The unconditioned stimulus  
455 (US) consisted of a mild electrical shock (train pulse at 50Hz lasting 200ms, with a single pulse  
456 duration of 1000 $\mu$ s) generated with a direct current stimulator (DS7A Constant Current Stimulator,  
457 Digitimer). Impulses were delivered through a bar stimulating electrode connected by a Velcro strap  
458 on the upper surface of the dominant hand's index finger. The electrical stimulation intensity was  
459 individually calibrated through a staircase procedure<sup>4,50,86</sup>, starting with a low current near the  
460 perceptible tactile threshold (~0.5 mA). Participants were asked to rate the painfulness of each train  
461 pulse on a scale ranging from 0 (not painful at all), 1 (pain threshold) to 10 (highly painful if  
462 protracted in time). At the end of the procedure, the US amplitude was then set at the current level  
463 (mA) corresponding to the mean rating of '7' on the subjective analog scale.

464

### 465 **Conditioning**

466 After a 1-min resting period, participants underwent a single-cue auditory threat conditioning,  
467 which consisted of the presentation of 15 trials of the conditioned stimulus (CS, 800Hz), with an  
468 ITI randomly ranging between 21s and 27s. The CS co-terminated with the US 12 times (80%  
469 reinforcement rate). Subjects were not informed about any possible CS-US contingency. To validate

470 the threat learning experience, immediately following this phase subjects rated the painfulness of  
471 the US using the same analog scale as in the preconditioning calibration procedure (see Table 1 for  
472 all groups' US current intensity and US analog ratings).

473

#### 474 **Transcranial Magnetic Stimulation**

475 Transcranial Magnetic Stimulation was performed with a Magstim Rapid<sup>2</sup> Stimulator (Magstim  
476 Co., Whitland, Dyfed, UK). A 70-mm figure-of-eight coil was positioned over the subject's M1  
477 cortical area at the optimum scalp position to elicit a contraction of the contralateral abductor  
478 *pollicis brevis* muscle (APB). Resting motor threshold (rMT) was defined as the minimum  
479 stimulation intensity that induced a visible finger movement in at least 5 out of 10 single pulses  
480 over the right-hand area of the left primary motor cortex<sup>16,87</sup>. After having determined each  
481 individual's rMT, we applied a single train of 1Hz-rTMS<sup>88,89</sup> for a total duration of 10 min (600  
482 pulses) to the target area. The rTMS intensity was set at 80% of the rMT for subjects whose rMT  
483 was  $\leq 50\%$  of the machine's maximum deliverable power (e.g., the intensity corresponded to 40%  
484 of the maximum power when the rMT was equal to 50% of the same parameter). For subjects with  
485 an rMT  $> 50\%$ , the stimulation intensity was always set to a ceiling corresponding to 40% of the  
486 machine's maximum deliverable power (see Table 1 for each group's mean rMT and mean  
487 stimulation intensity). During the rTMS procedure participants were seated in a comfortable  
488 recliner that we adjusted to allow their upper body to be in a sloped position, thus ensuring an  
489 optimal positioning of the coil.

490 To target the medial anterior portion of the prefrontal cortex (BA 10; aPFC and aPFC-E groups),  
491 the coil was centered over Fpz (10% of nasion-inion distance) according to the international 10–20  
492 electroencephalogram (EEG) system<sup>41</sup> (Figure 1). This placement should –with an rTMS reach of  
493 1.5 to 2 cm beneath the scalp<sup>90,91</sup>– ensure the targeting of the medial aPFC as in previous  
494 studies<sup>16,42,43</sup> and avoid the targeting of the dorsomedial prefrontal cortex (dmPFC), which would  
495 have been localizable with a scalp-based heuristic approach of 25.84% nasion-inion distance<sup>92</sup>. In



496 the case of left occipital cortex stimulation (OC group), the coil was positioned over O1 using the  
497 10–20 EEG system (BA 18/19), which functionally corresponds to associative visual cortices V3,  
498 V4, and V5<sup>93,94</sup> (Figure 1). For the stimulation of the left dorsolateral prefrontal cortex (dlPFC  
499 group), the coil was placed over F3 using the 10–20 EEG system (BA 8/9)<sup>18,95</sup> (Figure 1). For sham  
500 stimulation (sham and sham-E groups), the coil was centered over Fpz and positioned perpendicular  
501 to the scalp surface, so that no effective stimulation reached the brain during the procedure but  
502 allowed subjects to feel a comparable coil-scalp contact and hear the same noise as in real  
503 stimulation (Figure 1).

504 All participants were blinded to their experimental condition (i.e., active or sham), and were not  
505 informed about the potential cognitive or emotional effects of the stimulation.

506

#### 507 **Discomfort-inducing procedure**

508 The discomfort-inducing procedure mirrored the rTMS protocol and consisted of the delivery of  
509 mild electrical shocks (single 1-Hz train of 600 pulses lasting 10min, with a single pulse duration of  
510 500 $\mu$ s to mimic the duration of a single TMS pulse) generated with a direct current stimulator  
511 (DS7A Constant Current Stimulator, Digitimer). Impulses were delivered through two cup-  
512 stimulating electrodes attached to the surface of the subject's forehead in correspondence with Fpz  
513 according to the 10-20 EEG system. As in the case of the US calibration, the electrical stimulation  
514 intensity was individually calculated through a staircase procedure<sup>86</sup>, starting with a low current  
515 near the perceptible tactile threshold (~0.5 mA). Participants were asked to evaluate the perceived  
516 discomfort of each pulse on a scale from 0 (no discomfort) to 10 (high discomfort). At the end of  
517 the procedure, the shock amplitude was set at the current level (mA) corresponding to the mean  
518 rating of '4' on the subjective analog scale. To quantify the habituation to the uncomfortable  
519 stimulations, at the end of every minute of the 10-min procedure (i.e. every 60 pulses), subjects  
520 were requested to rate the level of the present discomfort on the same scale adopted during the  
521 calibration procedure.

522

### 523 **Implicit recognition test**

524 After a 1-min resting period, participants underwent this task, which consisted of the presentation of  
525 12 auditory stimuli in a completely random sequence:  $4 \times \text{CS}$ ,  $4 \times \text{NS}_1$ ,  $4 \times \text{NS}_2$ , with an ITI whose  
526 duration randomly ranged between 21s and 27s. SCRs were recorded throughout this phase, and the  
527 stimulating electrode was kept attached to create the expectation of receiving the US<sup>44</sup>. Differently  
528 from other paradigms<sup>48,96–98</sup>, here no shocks were delivered to avoid any reacquisition effect<sup>4,50</sup>.

529

### 530 **Implicit unconditioned threatening test**

531 This task was designed to elicit an unconditioned electrodermal response and consisted of the  
532 presentation of 4 trials of a woman scream sample lasting 4s, with an ITI randomly ranging between  
533 21s and 27s. SCRs were recorded throughout this phase, and the stimulating electrode was kept  
534 attached.

535

### 536 **Two-alternative forced-choice (2AFC) explicit recognition test**

537 This procedure involves the presentation of two stimuli on each trial and the subject chooses the one  
538 that was previously encoded (i.e. the first or the second one). As in our previous works<sup>4,50</sup>, a 2AFC  
539 design was preferred over a new-old paradigm, which involves one single stimulus on each trial,  
540 and the subject judges whether the stimulus has been previously encoded (old), or whether it is new.  
541 Our choice was motivated by the evidence that a 2AFC task improves recognition performance and  
542 discourages response biases such as the familiarity-based decision bias, namely the heuristic to  
543 endorse novel cues as ‘old’ when their familiarity is high<sup>99</sup>.

544 The task consisted of the presentation of 16 tone-pairs, each composed of the CS (800Hz) and one  
545 of the two NSs ( $\text{NS}_1$ , 1000Hz or  $\text{NS}_2$ , 600Hz) in a completely random sequence:  $4 \times \text{CS vs NS}_1$ ,  $4 \times$   
546  $\text{NS}_1 \text{ vs CS}$ ,  $4 \times \text{CS vs NS}_2$ ,  $4 \times \text{NS}_2 \text{ vs CS}$ . On each trial, the two stimuli were presented with an  
547 intra-trial-interval of 1000ms. After each pair offset, an ITI randomly ranging between 21s and 27s

548 occurred. Participants were explained that in each couple of sounds, there was a tone that they had  
549 heard on the first session (one week before or, in the case of the follow-up session, two weeks  
550 before) and a new tone. Participants were then instructed to recognize and verbally report which one  
551 (the first or the second) was the tone heard in the first session, paired with the US-shock (CS).  
552 Participants were further asked to verbally provide a confidence rating about each response, on a  
553 scale from 0 (completely unsure) to 10 (completely sure). No feedback was supplied. As in the  
554 implicit task, the stimulating electrode was kept attached, but no shock was delivered.

### 555 556 **Two-alternative forced-choice (2AFC) perceptual discrimination test**

557 The task consisted of the presentation of 7 pairs of auditory stimuli (i.e. CS vs NS<sub>1</sub>, NS<sub>1</sub> vs CS, CS  
558 vs NS<sub>2</sub>, NS<sub>2</sub> vs CS, CS vs CS, NS<sub>1</sub> vs NS<sub>1</sub>, NS<sub>2</sub> vs NS<sub>2</sub>) with a 1000-ms intra-pair-interval in a  
559 completely random sequence (ITI randomly ranging between 21s and 27s). For each pair, subjects  
560 were asked to report whether the two tones were “the same tone or different tones”, and to provide a  
561 confidence rating on an analog scale from 0 (completely unsure) to 10 (completely sure). No  
562 feedback was supplied, and the stimulating electrode was kept attached.

### 563 564 **Psychophysiological recording and analysis**

565 Event-related skin conductance responses (SCRs) were used as an implicit index of defensive  
566 responses. To record the autonomic signal, two Ag-AgCl non-polarizable electrodes filled with  
567 isotonic paste were attached to the index and middle fingers of the non-dominant hand by Velcro  
568 straps. The transducers were connected to the GSR100C module of the BIOPAC MP-150 system  
569 (BIOPAC Systems, Goleta, CA) and signals were recorded at a channel sampling rate of 1000 Hz.  
570 SCR waveforms were analyzed offline using AcqKnowledge 4.1 software (BIOPAC Systems,  
571 Goleta, CA), and were performed blindly to the subject’s experimental condition and the  
572 randomized sequence of stimuli. Each SCR was evaluated as event-related if the trough-to-peak  
573 deflection occurred 1–6 s (for the CS and the NSs) or 1–4 s (for the US<sub>2</sub>) after the stimulus onset,

574 the duration was comprised between 0.5 and 5.0 s, and the amplitude was greater than 0.02 micro  
575 siemens ( $\mu\text{S}$ ). Responses that did not fit these criteria were scored zero. To account for inter-  
576 individual variability, these raw values were then scaled according to each participant's average  
577 unconditioned response by dividing each response by the mean unconditioned stimulus (US)  
578 response during the conditioning phase<sup>100,101</sup>. Scaled SCR data were square-root transformed to  
579 normalize the distributions<sup>102</sup>.

580

### 581 **Statistical analyses**

582 We computed the appropriate sample size based on a power analysis performed through G\*Power  
583 3.1.9.2. For the main statistics, i.e. mixed ANOVA (within-between interaction) with two groups  
584 and two measurements, with the following input parameters:  $\alpha$  equal to 0.05, power (1- $\beta$ ) equal to  
585 0.95, and a hypothesized effect size ( $f$ ) equal to 0.25, the estimated sample size resulted in  $n = 30$   
586 per experimental group.

587 Since most variables passed the D'Agostino-Pearson omnibus normality test, parametric statistics  
588 were adopted in each experiment.

589 To test the between-group differences in post-conditioning US ratings, preconditioning **mean** SCRs  
590 levels, **mean** SCRs to the **CS and the** US during conditioning, and **mean** SCRs to the US<sub>2</sub> during the  
591 test and the follow-up sessions, we performed Student's unpaired  $t$  tests. **Potential differences in**  
592 **CS-related SCRs over the 15 trials of the conditioning phase were tested through  $2 \times 15$  mixed**  
593 **ANOVAs with Group (aPFC vs sham, aPFC vs OC, aPFC vs dlPFC) as between-subject variable**  
594 **and Trial (1–15) as within-subject variable.**

595 To test the potential between-group differences in the implicit reactions to the CS during the  
596 conditioning session, the test session, and the follow-up session, as well as the within-group  
597 differences from conditioning to test/follow-up phases, we computed  $2 \times 2$  mixed ANOVAs with  
598 Group (aPFC vs sham, aPFC vs OC, aPFC vs dlPFC, sham vs ctrl discomfort) as between-subject  
599 variable and Phase (conditioning vs test, conditioning vs follow-up) as within-subject variable.

600 Bonferroni adjustment was applied for simple main effects analyses. To compare between-group  
601 and within-group responses to the CS and the NSs during the test session, we performed  $2 \times 3$   
602 mixed ANOVAs with Group (aPFC *vs* sham, aPFC *vs* OC, aPFC *vs* dlPFC) as between-subject  
603 variable and Tone (NS<sub>1</sub>, CS, and NS<sub>2</sub>) as within-subject variable. Bonferroni adjustment was  
604 applied for simple main effects analyses.

605 To test the between-group differences in the explicit recognition and respective confidence ratings,  
606 as well as in the perceptual discrimination and respective confidence ratings during the test and the  
607 follow-up sessions (aPFC-E *vs* sham-E), we performed Student's unpaired *t* tests. To test whether  
608 explicit recognition levels were significantly higher than the 50% chance level for each condition  
609 during the test and the follow-up sessions, we calculated Student's one sample *t* tests against 0.50.

610 For each ANOVA we assessed the Sphericity assumption through Mauchly's Test. Where it was  
611 violated, we applied the Greenhouse-Geisser correction accordingly.

612 The null hypothesis was rejected at  $P < 0.05$  significance level. All statistical analyses were  
613 performed using SPSS Statistics 22 (IBM) and Prism 9 (GraphPad).

614 **Acknowledgements**

615 We thank all the subjects for their participation in this study. We also thank Melania Lattuada,  
616 Veronica Cintori, Ester Fusaro, Alessia Di Blasi, and Adriana Monno for their help in the  
617 experimental procedures and for their continuous support and advice. This work was supported by  
618 the “Compagnia di San Paolo, Progetto d’Ateneo”, University of Turin 2017 (CSTO167503), by the  
619 Fondazione Giovanni Gorla and Fondazione CRT (Talenti della Società Civile, ed. 2018), by the  
620 Grant “Progetti di ricerca di Rilevante Interesse Nazionale (PRIN)” 2017 (Project n.  
621 20178NNRCR\_002) from the Italian Ministry of University and Research (MIUR), Fondazione  
622 Cariverona 2020, Fondazione CRT 2021, and Banca D’Italia 2023.

623

624 **Competing interests**

625 The authors declare no financial interests or potential conflicts of interest.

626 **References**

- 627 1. DiFazio, L. E., Fanselow, M. & Sharpe, M. J. The effect of stress and reward on encoding  
628 future fear memories. *Behav. Brain. Res.* **417**, 113587 (2022).
- 629 2. Wilker, S., Elbert, T. & Kolassa, I. T. The downside of strong emotional memories: how  
630 human memory-related genes influence the risk for posttraumatic stress disorder—a selective  
631 review. *Neurobiol. Learn. Mem.* **112**, 75-86 (2014).
- 632 3. Dunsmoor, J. E. & Paz, R. Fear generalization and anxiety: behavioral and neural mechanisms.  
633 *Biol. Psychiatry* **78**, 336-343 (2015).
- 634 4. Manassero, E., Mana, L., Concina, G., Renna, A. & Sacchetti, B. Implicit and explicit systems  
635 differently predict possible dangers. *Sci. Rep.* **9**, 1-12 (2019).
- 636 5. Taylor, S., Abramowitz, J. S. & McKay, D. Non-adherence and nonresponse in the treatment of  
637 anxiety disorders. *J. Anxiety Disord.* **26**, 583-589 (2012).
- 638 6. Fonzo, G. A. et al. PTSD psychotherapy outcome predicted by brain activation during  
639 emotional reactivity and regulation. *Am. J. Psychiatry* **174**, 1163-1174 (2017).
- 640 7. Marković, V., Vicario, C. M., Yavari, F., Salehinejad, M. A. & Nitsche, M. A. A systematic  
641 review on the effect of transcranial direct current and magnetic stimulation on fear memory and  
642 extinction. *Front. Hum. Neurosci.* **15**, 1-26 (2021).
- 643 8. Asthana, M. et al. Effects of transcranial direct current stimulation on consolidation of fear  
644 memory. *Front. Psychiatry* **4**, 107 (2013).
- 645 9. Mungee, A., Kazzer, P., Feeser, M., Nitsche, M. A., Schiller, D. & Bajbouj, M. Transcranial  
646 direct current stimulation of the prefrontal cortex: a means to modulate fear memories.  
647 *Neuroreport* **25**, 480-484 (2014).
- 648 10. Mungee, A., Burger, M. & Bajbouj, M. No effect of cathodal transcranial direct current  
649 stimulation on fear memory in healthy human subjects. *Brain. Sci.* **6**, 55 (2016).
- 650 11. Abend, R. et al. Modulation of fear extinction processes using transcranial electrical  
651 stimulation. *Transl. Psychiatry* **6**, e913-e913 (2016).

- 652 12. Van't Wout, M., Mariano, T. Y., Garnaat, S. L., Reddy, M. K., Rasmussen, S. A. & Greenberg,  
653 B. D. Can transcranial direct current stimulation augment extinction of conditioned fear? *Brain*  
654 *Stimul.* **9**, 529-536 (2016).
- 655 13. Roesmann, K. et al. Transcranial direct current stimulation of the ventromedial prefrontal  
656 cortex modulates perceptual and neural patterns of fear generalization. *Biol. Psychiatry Cogn.*  
657 *Neurosci. Neuroimaging* **7**, 210-220 (2022).
- 658 14. Miniussi, C. et al. Efficacy of repetitive transcranial magnetic stimulation/transcranial direct  
659 current stimulation in cognitive neurorehabilitation. *Brain stimul.* **1**, 326-336 (2008).
- 660 15. Elder, G. J. & Taylor, J. P. Transcranial magnetic stimulation and transcranial direct current  
661 stimulation: treatments for cognitive and neuropsychiatric symptoms in the neurodegenerative  
662 dementias? *Alzheimer's Res. Ther.* **6**, 1-11 (2014).
- 663 16. Guhn, A. et al. Medial prefrontal cortex stimulation modulates the processing of conditioned  
664 fear. *Front. Behav. Neurosci.* **8**, 44 (2014).
- 665 17. Rajj, T., Nummenmaa, A., Marin, M. F., Porter, D., Furtak, S., Setsompop, K. & Milad, M. R.  
666 Prefrontal cortex stimulation enhances fear extinction memory in humans. *Biol. Psychiatry* **84**,  
667 129-137 (2018).
- 668 18. Borgomaneri, S., Battaglia, S., Garofalo, S., Tortora, F., Avenanti, A. & Di Pellegrino, G.  
669 State-dependent TMS over prefrontal cortex disrupts fear-memory reconsolidation and prevents  
670 the return of fear. *Curr. Biol.* **30**, 3672-3679 (2020).
- 671 19. Su, S. et al. Continuous theta-burst stimulation over the right dorsolateral prefrontal cortex  
672 disrupts fear memory reconsolidation in humans. *iScience* **25**, 103614 (2022).
- 673 20. Vervliet, B., Craske, M. G. & Hermans, D. Fear extinction and relapse: state of the art. *Annu.*  
674 *Rev. Clin. Psychol.* **9**, 215-248 (2013).
- 675 21. Ramnani, N. & Owen, A. M. Anterior prefrontal cortex: insights into function from anatomy  
676 and neuroimaging. *Nat. Rev. Neurosci.* **5**, 184-194 (2004).



- 677 22. Semendeferi, K., Armstrong, E., Schleicher, A., Zilles, K. & van Hoesen, G. W. Prefrontal  
678 cortex in humans and apes: a comparative study of area 10. *Am. J. Phys. Anthropol.* **114**, 224-  
679 241 (2001).
- 680 23. Volman, I. et al. Reduced serotonin transporter availability decreases prefrontal control of the  
681 amygdala. *J. Neurosci.* **33**, 8974-8979 (2013).
- 682 24. Koch, S. B. J., Mars, R. B., Toni, I. & Roelofs, K. Emotional control, reappraised. *Neurosci.*  
683 *Biobehav. Rev.* **95**, 528-534 (2018).
- 684 25. Bramson, B. et al. Human lateral frontal pole contributes to control over emotional approach-  
685 avoidance actions. *J. Neurosci.* **40**, 2925-2934 (2020).
- 686 26. Folloni, D. et al. Dichotomous organization of amygdala/temporal-prefrontal bundles in both  
687 humans and monkeys. *Elife* **8**, 1-23 (2019).
- 688 27. Peng, K., Steele, S. C., Becerra, L. & Borsook, D. Brodmann area 10: collating, integrating and  
689 high level processing of nociception and pain. *Prog. Neurobiol.* **161**, 1-22 (2018).
- 690 28. Klumpers, F. et al. Prefrontal mechanisms of fear reduction after threat offset. *Biol. Psychiatry*  
691 **68**, 1031-1038 (2010).
- 692 29. Lanius, R. A. et al. Functional connectivity of dissociative responses in posttraumatic stress  
693 disorder: a functional magnetic resonance imaging investigation. *Biol. Psychiatry* **57**, 873-884  
694 (2005).
- 695 30. Morey, R. A., Petty, C. M., Cooper, D. A., LaBar, K. S. & McCarthy, G. Neural systems for  
696 executive and emotional processing are modulated by symptoms of posttraumatic stress  
697 disorder in Iraq War veterans. *Psychiatry Res. Neuroimaging* **162**, 59-72 (2008).
- 698 31. Sadeh, N. et al. Neurobiological indicators of disinhibition in posttraumatic stress disorder.  
699 *Hum. Brain Mapp.* **36**, 3076-3086 (2015).
- 700 32. Sadeh, N. et al. SKA2 methylation is associated with decreased prefrontal cortical thickness  
701 and greater PTSD severity among trauma-exposed veterans. *Mol. Psychiatry* **21**, 357-363  
702 (2016).

- 703 33. Kaldewaij, R., Koch, S. B., Hashemi, M. M., Zhang, W., Klumpers, F. & Roelofs, K. Anterior  
704 prefrontal brain activity during emotion control predicts resilience to post-traumatic stress  
705 symptoms. *Nat. Hum. Behav.* **5**, 1055-1064 (2021).
- 706 34. Fonzo, G. A. et al. Selective effects of psychotherapy on frontopolar cortical function in PTSD.  
707 *Am. J. Psychiatry* **174**, 1175-1184 (2017).
- 708 35. Resnik, J. & Paz, R. Fear generalization in the primate amygdala. *Nat. Neurosci.* **18**, 188-190  
709 (2015).
- 710 36. Wong, A. H. & Lovibond, P. F. Rule-based generalisation in single-cue and differential fear  
711 conditioning in humans. *Biol. Psychol.* **129**, 111-120 (2017).
- 712 37. Grosso, A., Santoni, G., Manassero, E., Renna, A. & Sacchetti, B. A neuronal basis for fear  
713 discrimination in the lateral amygdala. *Nat. Commun.* **9**, 1-12 (2018).
- 714 38. Concina, G., Cambiaghi, M., Renna, A. & Sacchetti, B. Coherent activity between the  
715 prelimbic and auditory cortex in the slow-gamma band underlies fear discrimination. *J.*  
716 *Neurosci.* **38**, 8313-8328 (2018).
- 717 39. Grosso, A., Cambiaghi, M., Milano, L., Renna, A., Sacco, T. & Sacchetti, B. Region-and layer-  
718 specific activation of the higher order auditory cortex Te2 after remote retrieval of fear or  
719 appetitive memories. *Cereb. Cortex* **27**, 3140-3151 (2017).
- 720 40. Hanlon, C. A., Dowdle, L. T., Gibson, N. B., Li, X., Hamilton, S., Canterbury, M. & Hoffman,  
721 M. Cortical substrates of cue-reactivity in multiple substance dependent populations:  
722 transdiagnostic relevance of the medial prefrontal cortex. *Transl. Psychiatry* **8**, 1-8 (2018).
- 723 41. Jasper, H. H. The ten-twenty electrode system of the International Federation.  
724 *Electroencephalogr. Clin. Neurophysiol.* **10**, 370-375 (1958).
- 725 42. Herrmann, M. J., Katzorke, A., Busch, Y., Gromer, D., Polak, T., Pauli, P. & Deckert, J.  
726 Medial prefrontal cortex stimulation accelerates therapy response of exposure therapy in  
727 acrophobia. *Brain Stimul.* **10**, 291-297 (2017).

- 728 43. Karmann, A. J., Maihöfner, C., Lautenbacher, S., Sperling, W., Kornhuber, J. & Kunz, M. The  
729 role of prefrontal inhibition in regulating facial expressions of pain: a repetitive transcranial  
730 magnetic stimulation study. *J. Pain.* **17**, 383-391 (2016).
- 731 44. Ameli, R., Ip, C. & Grillon, C. Contextual fear-potentiated startle conditioning in humans:  
732 Replication and extension. *Psychophysiology* **38**, 383-390 (2001).
- 733 45. Maren, S., Phan, K. L. & Liberzon, I. The contextual brain: implications for fear conditioning,  
734 extinction and psychopathology. *Nat. Rev. Neurosci.* **14**, 417-428 (2013).
- 735 46. Sacco, T. & Sacchetti, B. Role of secondary sensory cortices in emotional memory storage and  
736 retrieval in rats. *Science* **329**, 649-656 (2010).
- 737 47. Sacchetti, B., Baldi, E., Tassoni, G. & Bucherelli, C. Memorization of contextual and CS  
738 conditioned fear response (freezing) in a one-trial acquisition paradigm. *Arch. Ital. Biol.* **137**,  
739 235-248 (1999).
- 740 48. Onat, S. & Büchel, C. The neuronal basis of fear generalization in humans. *Nat. Neurosci.* **18**,  
741 1811-1818 (2015).
- 742 49. Laufer, O., Israeli, D. & Paz, R. Behavioral and neural mechanisms of overgeneralization in  
743 anxiety. *Curr. Biol.* **26**, 713-722 (2016).
- 744 50. Manassero, E., Giordano, A., Raimondo, E., Cicolin, A. & Sacchetti, B. Sleep deprivation  
745 during memory consolidation, but not before memory retrieval, widens threat generalization to  
746 new stimuli. *Front. Neurosci.* **781**, 1-12 (2022).
- 747 51. Hiser, J. & Koenigs, M. The multifaceted role of the ventromedial prefrontal cortex in emotion,  
748 decision making, social cognition, and psychopathology. *Biol. Psychiatry* **83**, 638-647 (2018).
- 749 52. Catani, M. et al. Short frontal lobe connections of the human brain. *Cortex* **48**, 273-291 (2012).
- 750 53. Orr, J. M., Smolker, H. R. & Banich, M. T. Organization of the human frontal pole revealed by  
751 large-scale DTI-based connectivity: implications for control of behavior. *PloS one* **10**, 1-23  
752 (2015).

- 753 54. Wu, Y., Sun, D., Wang, Y. & Wang, Y. Subcomponents and connectivity of the inferior fronto-  
754 occipital fasciculus revealed by diffusion spectrum imaging fiber tracking. *Front. Neuroanat.*  
755 **10**, 1-13 (2016).
- 756 55. Pavlov, I. P. *Conditioned Reflexes*. (London: Oxford Univ. Press., 1927).
- 757 56. Rachman, S. Studies in desensitization III: speed of generalization. *Behav. Res. Ther.* **4**, 7-15  
758 (1966).
- 759 57. Pascual-Leone, A., Tormos, J. M., Keenan, J., Tarazona, F., Cañete, C. & Catalá, M. D. Study  
760 and modulation of human cortical excitability with transcranial magnetic stimulation. *Clin.*  
761 *Neurophysiol.* **15**, 333-343 (1998).
- 762 58. Luber, B. & Deng, Z. D. Application of non-invasive brain stimulation in psychophysiology.  
763 In: Berntson, G. G., Cacioppo, J. T. & Tassinary, L. G. (Eds.). *Handbook of Psychophysiology*,  
764 *4th ed.* (Cambridge University Press, Cambridge, pp. 116-150, 2016).
- 765 59. Beynel, L., Powers, J. P. & Appelbaum, L. G. Effects of repetitive transcranial magnetic  
766 stimulation on resting-state connectivity: A systematic review. *NeuroImage* **211**, 116596  
767 (2020).
- 768 60. Eisenegger, C., Treyer, V., Fehr, E. & Knoch, D. Time-course of “off-line” prefrontal rTMS  
769 effects—a PET study. *NeuroImage* **42**, 379-384 (2008).
- 770 61. Nahas, Z. et al. Unilateral left prefrontal transcranial magnetic stimulation (TMS) produces  
771 intensity-dependent bilateral effects as measured by interleaved BOLD fMRI. *Biol. Psychiatry*  
772 **50**, 712-720 (2001).
- 773 62. Dunsmoor, J. E., Bandettini, P. A. & Knight, D. C. Impact of continuous versus intermittent  
774 CS-UCS pairing on human brain activation during Pavlovian fear conditioning. *Behav.*  
775 *Neurosci.* **121**, 635-642 (2007).
- 776 63. Dunsmoor, J. E., Bandettini, P. A. & Knight, D. C. Neural correlates of unconditioned response  
777 diminution during Pavlovian conditioning. *NeuroImage* **40**, 811-817 (2008).

- 778 64. Vicario, C. M., Salehinejad, M. A., Felmingham, K., Martino, G. & Nitsche, M. A. A  
779 systematic review on the therapeutic effectiveness of non-invasive brain stimulation for the  
780 treatment of anxiety disorders. *Neurosci. Biobehav. Rev.* **96**, 219-231 (2019).
- 781 65. Pape, H. C. & Pare, D. Plastic synaptic networks of the amygdala for the acquisition,  
782 expression, and extinction of conditioned fear. *Physiol. Rev.* **90**, 419-463 (2010).
- 783 66. Sacchetti, B., Baldi, E., Lorenzini, C. A. & Bucherelli, C. Cerebellar role in fear-conditioning  
784 consolidation. *PNAS* **99**, 8406-8411 (2002).
- 785 67. Sacchetti, B., Scelfo, B., Tempia, F. & Strata, P. Long-term synaptic changes induced in the  
786 cerebellar cortex by fear conditioning. *Neuron* **42**, 973-982 (2004).
- 787 68. Zhu, L., Sacco, T., Strata, P. & Sacchetti, B. Basolateral amygdala inactivation impairs  
788 learning-induced long-term potentiation in the cerebellar cortex. *PloS one* **6**, e16673 (2011).
- 789 69. Grosso, A. et al. The higher order auditory cortex is involved in the assignment of affective  
790 value to sensory stimuli. *Nat. Comm.* **6**, 1-14 (2015).
- 791 70. Manassero, E., Renna, A., Milano, L. & Sacchetti, B. Lateral and basal amygdala account for  
792 opposite behavioral responses during the long-term expression of fearful memories. *Sci. Rep.* **8**,  
793 1-12 (2018).
- 794 71. Cambiaghi, M. et al. Higher-order sensory cortex drives basolateral amygdala activity during  
795 the recall of remote, but not recently learned fearful memories. *J. Neurosci.* **36**, 1647-1659  
796 (2016a).
- 797 72. Cambiaghi M, Grosso A, Renna A. & Sacchetti B. Differential recruitment of auditory cortices  
798 in the consolidation of recent auditory fearful memories. *J. Neurosci.* **36**, 8586-8597 (2016b).
- 799 73. Concina, G., Renna, A., Grosso, A. & Sacchetti, B. The auditory cortex and the emotional  
800 valence of sounds. *Neurosci. Biobehav. Rev.* **98**, 256-264 (2019).
- 801 74. You, Y., Brown, J. & Li, W. Human sensory cortex contributes to the long-term storage of  
802 aversive conditioning. *J. Neurosci.* **41**, 3222-3233 (2021).

- 803 75. You, Y., Novak, L. R., Clancy, K. J. & Li, W. Pattern differentiation and tuning shift in human  
804 sensory cortex underlie long-term threat memory. *Curr. Biol.* **32**, 2067-2075 (2022).
- 805 76. Ojala, K. E., Staib, M., Gerster, S., Ruff, C. C. & Bach, D. R. Inhibiting human aversive  
806 memory by transcranial theta-burst stimulation to the primary sensory cortex. *Biol. psychiatry*  
807 **92**, 149-157 (2022).
- 808 77. Monfils, M. H. The high road to inhibiting fear memories. *Biol. Psychiatry* **92**, 102-103 (2022).
- 809 78. Motzkin, J. C., Philippi, C. L., Wolf, R. C., Baskaya, M. K. & Koenigs, M. Ventromedial  
810 prefrontal cortex is critical for the regulation of amygdala activity in humans. *Biol. Psychiatry*  
811 **77**, 276-284 (2015).
- 812 79. Faran, Y. A comment on the connection between BA10 and episodic memory. *Front. Behav.*  
813 *Neurosci.* **17**, 1-3 (2023).
- 814 80. Bechara, A., Tranel, D., Damasio, H., Adolphs, R., Rockland, C. & Damasio, A. R. Double  
815 dissociation of conditioning and declarative knowledge relative to the amygdala and  
816 hippocampus in humans. *Science* **269**, 1115-1118 (1995).
- 817 81. LaBar, K. S. & Cabeza, R. Cognitive neuroscience of emotional memory. *Nat. Rev. Neurosci.*  
818 **7**, 54-64 (2006).
- 819 82. Knight, D. C., Waters, N. S. & Bandettini, P. A. Neural substrates of explicit and implicit fear  
820 memory. *NeuroImage* **45**, 208-214 (2009).
- 821 83. Rossi, S. et al. Safety and recommendations for TMS use in healthy subjects and patient  
822 populations, with updates on training, ethical and regulatory issues: Expert Guidelines. *Clin.*  
823 *Neurophysiol.* **132**, 269-306 (2021).
- 824 84. Spielberger, C. D., Gorsuch, R. L., Lushene, R., Vagg, P. R. & Jacobs, G. A. *Manual for the*  
825 *State-Trait Anxiety Inventory*. (Palo Alto: Consulting Psychologists Press, 1983).
- 826 85. Pedrabissi, L. & Santinello, M. *Nuova versione italiana dello STAI forma Y [New Italian*  
827 *version of the STAI form Y]*. (Firenze: Organizzazioni Speciali, 1989).
- 828 86. Cornsweet, T. N. The staircase-method in psychophysics. *Am. J. Psychol.* **75**, 485-491 (1962).

- 829 87. Westin, G. G., Bassi, B. D., Lisanby, S. H. & Luber, B. New York State Psychiatric Institute,  
830 N. Y. Determination of motor threshold using visual observation overestimates transcranial  
831 magnetic stimulation dosage: safety implications. *Clin. Neurophysiol.* **125**, 142-147 (2014).
- 832 88. Ando, A. et al. Embodied simulation and ambiguous stimuli: The role of the mirror neuron  
833 system. *Brain Res.* **1629**, 135-142 (2015).
- 834 89. Salatino, A. et al. Transcranial magnetic stimulation of posterior parietal cortex modulates line-  
835 length estimation but not illusory depth perception. *Front. Psychol.* **10**, 1169 (2019).
- 836 90. Epstein, C. M., Schwartzberg, D. G., Davey, K. R. & Sudderth, D. B. Localizing the site of  
837 magnetic brain stimulation in humans. *Neurology* **40**, 666-670 (1990).
- 838 91. Rudiak, D. & Marg, E. Finding the depth of magnetic brain stimulation: a re-evaluation.  
839 *Electroencephalogr.* **93**, 358-371 (1994).
- 840 92. Mir-Moghtadaei, A., Giacobbe, P., Daskalakis, Z. J., Blumberger, D. M. & Downar, J.  
841 Validation of a 25% nasion–inion heuristic for locating the dorsomedial prefrontal cortex for  
842 repetitive transcranial magnetic stimulation. *Brain. Stimul.* **9**, 793-795 (2016).
- 843 93. Rojas, G. M., Alvarez, C., Montoya, C. E., De la Iglesia-Vaya, M., Cisternas, J. E. & Gálvez,  
844 M. Study of resting-state functional connectivity networks using EEG electrodes position as  
845 seed. *Front. Neurosci.* **12**, 235 (2018).
- 846 94. Brighina, F., Ricci, R., Piazza, A., Scalia, S., Giglia, G. & Fierro, B. Illusory contours and  
847 specific regions of human extrastriate cortex: evidence from rTMS. *Eur. J. Neurosci.* **17**, 2469-  
848 2480 (2003).
- 849 95. Mir-Moghtadaei, A. et al. Concordance between BeamF3 and MRI-neuronavigated target sites  
850 for repetitive transcranial magnetic stimulation of the left dorsolateral prefrontal cortex. *Brain.*  
851 *Stimul.* **8**, 965-973 (2015).
- 852 96. Lissek, S., Bradford, D. E., Alvarez, R. P., Burton, P., Espensen-Sturges, T., Reynolds, R. C. &  
853 Grillon, C. Neural substrates of classically conditioned fear-generalization in humans: a  
854 parametric fMRI study. *Soc. Cogn. Affect. Neurosci.* **9**, 1134-1142 (2014).

- 855 97. Dunsmoor, J. E., Kroes, M. C., Braren, S. H. & Phelps, E. A. Threat intensity widens fear  
856 generalization gradients. *Behav. Neurosci.* **131**, 168-175 (2017).
- 857 98. Holt, D. J., Boeke, E. A., Wolthusen, R. P., Nasr, S., Milad, M. R. & Tootell, R. B. A  
858 parametric study of fear generalization to faces and non-face objects: relationship to  
859 discrimination thresholds. *Front. Hum. Neurosci.* **8**, 1-12 (2014).
- 860 99. Macmillan, N. A. & Creelman, C. D. Detection theory: A user's guide. (New York: Psychology  
861 press, 2004).
- 862 100. Schiller, D., Monfils, M. H., Raio, C. M., Johnson, D. C., LeDoux, J. E. & Phelps, E. A.  
863 Preventing the return of fear in humans using reconsolidation update mechanisms. *Nature* **463**,  
864 49-53 (2010).
- 865 101. Battaglia, S., Garofalo, S. & Di Pellegrino, G. Context-dependent extinction of threat  
866 memories: Influences of healthy aging. *Sci. Rep.* **8**, 1-13 (2018).
- 867 102. Lykken, D. T. & Venables, P. H. Direct measurement of skin conductance: A proposal for  
868 standardization. *Psychophysiology* **8**, 656-672 (1971).



869 **Table 1. Experimental groups' descriptive, experimental, and clinical data.** The table reports,  
870 for each experimental condition: sample size (*N*), sex distribution (F = Female, M = Male), mean  
871 age, State-Trait Anxiety Inventory Form Y (STAI-Y) State subscale score during session 1 (S1),  
872 session 2 (S2), and session 3 (S3), and Trait subscale score, US current intensity (mA), post-  
873 conditioning US rating, rTMS resting motor threshold (rMT), rTMS power, and discomfort  
874 stimulation (DS) current intensity (mA). All data are mean  $\pm$  standard deviation.

875

Group	<i>N</i>	Sex	Age	STAI-Y State (S1)	STAI-Y State (S2)	STAI-Y State (S3)	STAI-Y Trait	US (mA)	US rating	rTMS rMT	rTMS power	DS (mA)
aPFC	30	18F 12M	24.45 $\pm$ 3.78	30.97 $\pm$ 4.07	32.47 $\pm$ 7.16	30.60 $\pm$ 6.04	39.27 $\pm$ 6.18	4.92 $\pm$ 2.06	5.28 $\pm$ 0.90	58.20 $\pm$ 6.40	39.73 $\pm$ 1.11	-
sham	30	18F 12M	23.35 $\pm$ 2.35	33.23 $\pm$ 5.86	32.70 $\pm$ 7.74	31.87 $\pm$ 6.51	38.77 $\pm$ 4.02	4.88 $\pm$ 2.45	5.47 $\pm$ 0.88	-	-	-
OC	30	18F 12M	24.14 $\pm$ 2.62	32.33 $\pm$ 5.51	31.53 $\pm$ 7.57	30.60 $\pm$ 6.75	39.03 $\pm$ 5.12	4.99 $\pm$ 3.17	5.28 $\pm$ 1.06	60.90 $\pm$ 6.67	39.70 $\pm$ 1.47	-
dIPFC	30	18F 12M	23.91 $\pm$ 3.15	31.70 $\pm$ 5.40	30.83 $\pm$ 7.04	30.13 $\pm$ 5.88	39.17 $\pm$ 5.85	5.16 $\pm$ 2.43	5.57 $\pm$ 1.45	58.77 $\pm$ 5.89	39.90 $\pm$ 0.40	-
aPFC-E	21	13F 8M	24.39 $\pm$ 2.43	31.71 $\pm$ 4.89	30.90 $\pm$ 5.66	30.48 $\pm$ 4.96	38.29 $\pm$ 6.21	5.13 $\pm$ 1.86	5.43 $\pm$ 0.94	58.67 $\pm$ 7.16	39.52 $\pm$ 1.54	-
sham-E	21	13F 8M	23.83 $\pm$ 2.73	33.10 $\pm$ 5.59	31.48 $\pm$ 5.54	30.38 $\pm$ 7.73	38.29 $\pm$ 5.22	5.27 $\pm$ 3.19	5.31 $\pm$ 1.31	-	-	-
ctrl discomfort	10	5F 5M	22.34 $\pm$ 3.67	34.40 $\pm$ 4.20	36.50 $\pm$ 6.47	34.20 $\pm$ 5.98	39.70 $\pm$ 4.03	6.97 $\pm$ 4.14	5.65 $\pm$ 1.11	-	-	6.65 $\pm$ 2.25

876

877 **Figure legends**

878

879 **Figure 1. Schematic diagram depicting the experimental outline and rTMS conditions.** In the  
880 first session (day 1, context A), participants underwent a single-cue threat conditioning in which a  
881 tone (CS) was paired with a mild electrical shock (US). In the second session (day 8, context B), a  
882 1Hz-rTMS procedure was actively applied over the medial anterior prefrontal cortex (aPFC,  $n = 30$ ;  
883 aPFC-E,  $n = 21$ ), sham-applied over the same site (sham,  $n = 30$ ; sham-E,  $n = 21$ ), actively applied  
884 over the left occipital cortex (OC,  $n = 30$ ) and over the left dorsolateral prefrontal cortex (dlPFC,  $n$   
885  $= 30$ ). In the implicit conditions (aPFC, sham, OC, dlPFC), subjects underwent an implicit test  
886 during which they were presented with the CS and two new stimuli (NS<sub>1</sub> and NS<sub>2</sub>) and then an  
887 unconditioned threat test while being recorded in their SCRs. In the explicit conditions (aPFC-E,  
888 sham-E), participants underwent an explicit 2AFC recognition task during which they were  
889 presented with tone pairs each composed of the CS and one of the two NSs, and they were asked to  
890 recognize the CS providing a confidence level for each choice. Last, participants underwent a 2AFC  
891 perceptual discrimination test, in which they had to judge whether the two tones in each pair (CS  
892 and/or NSs) were “the same tone” or “different tones”. The third session (day 15, context A) was  
893 identical to the second one except for the absence of the rTMS.

894 **Figure 2. Effects of rTMS over the aPFC on immediate and remote implicit threat memory,**  
895 **threat generalization to new stimuli, and overall electrodermal responsivity. (A)** Simulation of  
896 rTMS effects on the neural tissue of the medial aPFC (medial BA 10), performed with SimNIBS  
897 4.0 software. The magnitude of the electric field is expressed in V/m. **(B, C)** Dot plot and line chart  
898 representing the mean SCRs elicited by the CS during the conditioning session and test session in  
899 the two different conditions. Groups' reactions were not different during the conditioning phase,  
900 whereas during the test phase the group stimulated over the aPFC showed attenuated implicit  
901 reactions relative to the sham condition. The aPFC group displayed reduced autonomic reactions to  
902 the CS from conditioning to test, while the sham group showed an increase in defensive responses.  
903 **(D, E)** Implicit reactions to all the tones (NS<sub>1</sub>, CS, and NS<sub>2</sub>) during the test session were decreased  
904 in the aPFC group relative to the sham group. Although we found a significant main effect of Group  
905 and no Group × Tone interaction effect, we reported the statistical significance marks of simple  
906 main effects. **(F)** Implicit reactions to the US<sub>2</sub> during the test session were not different between  
907 conditions, showing no rTMS effects on the overall electrodermal responsivity. **(G, H)** In the  
908 follow-up session, the aPFC group enduringly demonstrated reduced implicit reactions to the CS  
909 relative to the sham group and to the conditioning phase. **(I)** Implicit reactions to the US<sub>2</sub> during the  
910 follow-up session were not different between groups. \*  $P < 0.05$ , \*\*  $P < 0.01$ , \*\*\*  $P < 0.001$ . All  
911 data are mean and SEM. 2×2 mixed ANOVA followed by Bonferroni-adjusted *post hoc*  
912 comparisons (B, C, G, H); 2×3 mixed ANOVA followed by Bonferroni-adjusted *post hoc*  
913 comparisons (D, E); Student's unpaired *t* test (F, I).

914 **Figure 3. Effects of rTMS over the aPFC on immediate and remote explicit threat memory.**  
915 (A) During the test session, explicit recognition patterns were not different between the group  
916 stimulated over the aPFC and the sham group. (B) During the test session, confidence ratings did  
917 not differ between the two conditions. (C) During the follow-up session, aPFC-E and sham-E  
918 groups identified the CS between the NSs in a not different manner. (D) During the follow-up  
919 session, aPFC-E and sham-E groups were not differently confident about their explicit choices. All  
920 data are mean and SEM. Student's unpaired *t* test (A, B, C, D).

921 **Figure 4. Selective effects of rTMS over the aPFC and the left OC on the defensive responses**  
922 **to threat-predictive cues. (A)** Simulation of rTMS effects on the neural tissue of the left OC (BA  
923 18/19), performed with SimNIBS 4.0 software. The magnitude of the electric field is expressed in  
924 V/m. **(B, C)** Dot plot and line chart representing the mean SCRs elicited by the CS during the  
925 conditioning session and test session in the OC group, compared with the same aPFC group of Fig.  
926 2. The two groups did not differently respond during the conditioning phase, but during the test  
927 phase the group stimulated over the aPFC showed weaker reactions than the OC group. While the  
928 defensive reactions of the aPFC group decreased from conditioning to test, those of the OC group  
929 remained not differently high. **(D, E)** Implicit reactions to NSs during the test session did not differ  
930 between groups. In the OC group, the responses elicited by the NS<sub>2</sub> were lower than those evoked  
931 by the CS and the NS<sub>1</sub>. **(F)** Implicit reactions to the US<sub>2</sub> during the test session were not different  
932 between groups. **(G, H)** In the follow-up session, the aPFC group persisted in showing reduced  
933 implicit reactions to the CS relative to the OC group. Defensive reactions of both groups decreased  
934 from the conditioning phase. Although we found a significant main effect of Phase and no Group ×  
935 Phase interaction effect, we reported the statistical significance marks of simple main effects. **(I)**  
936 Implicit reactions to the US<sub>2</sub> during the follow-up session were not different between groups. \*  $P <$   
937 0.05, \*\*\*  $P <$  0.001. All data are mean and SEM. 2×2 mixed ANOVA followed by Bonferroni-  
938 adjusted *post hoc* comparisons (B, C, G, H); 2×3 mixed ANOVA followed by Bonferroni-adjusted  
939 *post hoc* comparisons (D, E); Student's unpaired *t* test (F, I).

940 **Figure 5. Different effects of rTMS over the aPFC and the left dlPFC on immediate and**  
941 **remote implicit threat memory.** (A) Simulation of rTMS effects on the neural tissue of the left  
942 dlPFC (BA 8/9), performed with SimNIBS 4.0 software. The magnitude of the electric field is  
943 expressed in V/m. (B, C) Dot plot and line chart representing the mean SCRs elicited by the CS  
944 during the conditioning session and test session in the dlPFC group, compared with the same aPFC  
945 group of Fig. 2. The two conditions did not differently react during the conditioning phase, whereas  
946 during the test phase the group stimulated over the aPFC displayed lower reactions than the dlPFC  
947 group. Implicit reactions of the aPFC group decreased from conditioning to test, while those of the  
948 dlPFC group increased. (D, E) Implicit reactions to NSs during the test session did not differ  
949 between groups. In the dlPFC group, the responses elicited by the NS<sub>2</sub> were lower than those  
950 evoked by the CS. (F) The two groups did not differently react to the US<sub>2</sub> during the test session.  
951 (G, H) In the follow-up session, the aPFC group persisted in more dimly reacting to the CS relative  
952 to the dlPFC group and to the conditioning phase. (I) Implicit reactions to the US<sub>2</sub> during the  
953 follow-up session were not different between groups. \*  $P < 0.05$ , \*\*  $P < 0.01$ , \*\*\*  $P < 0.001$ . All  
954 data are mean and SEM. 2×2 mixed ANOVA followed by Bonferroni-adjusted *post hoc*  
955 comparisons (B, C, G, H); 2×3 mixed ANOVA followed by Bonferroni-adjusted *post hoc*  
956 comparisons (D, E); Student's unpaired *t* test (F, I).

957 **Figure 2-figure supplement 1. Implicit reactions during preconditioning (CS) and**  
958 **conditioning (CS, US) in the aPFC and sham groups.** (A) Dot plot representing the mean SCRs  
959 elicited by the CS during the preconditioning phase in the aPFC group ( $n = 30$ ) compared with the  
960 sham group ( $n = 30$ ). Implicit reactions were not significantly different. (B) Mean SCRs elicited by  
961 the US during the conditioning phase in the aPFC group compared with the sham group. Responses  
962 were not significantly different. (C) Mean SCRs evoked by the CS over the 15 trials of the  
963 conditioning phase, and averaged SCRs (trials 1-15) in the aPFC and sham groups. Autonomic  
964 reactions were not significantly different. All data are mean and SEM. Student's unpaired  $t$  test (A,  
965 B, C);  $2 \times 15$  mixed ANOVA (C).


966 **Figure 2-figure supplement 2. Effects of a discomfort-inducing procedure on immediate and**  
967 **remote implicit threat memory. (A)** Mean discomfort ratings provided by the subjects of the ctrl  
968 discomfort group ( $n = 10$ ) during the 10-min discomfort-inducing procedure. **(B)** Dot plot  
969 representing the mean SCRs elicited by the CS during the conditioning phase and the test phase in  
970 the ctrl discomfort group and the sham group ( $n = 30$ ). The groups' reactions were not significantly  
971 different during the conditioning phase as well as during the test phase ( $2 \times 2$  mixed ANOVA; main  
972 effect of group:  $F_{(1,38)} = 0.712$ ,  $P = 0.404$ ; main effect of phase:  $F_{(1,38)} = 1.713$ ,  $P = 0.198$ ; group  $\times$   
973 phase interaction:  $F_{(1,38)} = 0.956$ ,  $P = 0.335$ ). **(C)** In the follow-up session, SCRs to the CS did not  
974 differ between groups ( $2 \times 2$  mixed ANOVA; main effect of group:  $F_{(1,38)} = 1.335$ ,  $P = 0.255$ ; main  
975 effect of phase:  $F_{(1,38)} = 0.042$ ,  $P = 0.838$ ; group  $\times$  phase interaction:  $F_{(1,38)} = 0.175$ ,  $P = 0.678$ ). All  
976 data are mean and SEM.




977 **Figure 4-figure supplement 1. Implicit reactions during preconditioning (CS) and**  
978 **conditioning (CS, US) in the aPFC and OC groups. (A)** Dot plot representing the mean SCRs  
979 elicited by the CS during the preconditioning phase in the aPFC group ( $n = 30$ ) compared with the  
980 OC group ( $n = 30$ ). Implicit reactions were not significantly different. **(B)** Mean SCRs elicited by  
981 the US during the conditioning phase in the aPFC group compared with the OC group. Responses  
982 were not significantly different. **(C) Mean SCRs evoked by the CS over the 15 trials of the**  
983 **conditioning phase, and averaged SCRs (trials 1-15) in the aPFC and OC groups. Autonomic**  
984 **reactions were not significantly different.** All data are mean and SEM. Student's unpaired  $t$  test (A,  
985 B, C);  $2 \times 15$  mixed ANOVA (C).










986 **Figure 5-figure supplement 1. Implicit reactions during preconditioning (CS) and**  
987 **conditioning (CS, US) in the aPFC and dlPFC groups.** (A) Dot plot representing the mean SCRs  
988 elicited by the CS during the preconditioning phase in the aPFC group ( $n = 30$ ) compared with the  
989 dlPFC group ( $n = 30$ ). Implicit reactions were not significantly different. (B) Mean SCRs elicited by  
990 the US during the conditioning phase in the aPFC group compared with the dlPFC group.  
991 Responses were not significantly different. (C) Mean SCRs evoked by the CS over the 15 trials of  
992 the conditioning phase, and averaged SCRs (trials 1-15) in the aPFC and dlPFC groups. Autonomic  
993 reactions were not significantly different. All data are mean and SEM. Student's unpaired  $t$  test (A,  
994 B, C);  $2 \times 15$  mixed ANOVA (C).





# Experimental outline





















## rTMS conditions




**A** medial anterior prefrontal cortex**B** conditioning CS vs test CS**C** conditioning CS vs test CS**D** test CS vs NSs**E** test CS vs NSs**F** test US<sub>2</sub>**G** conditioning CS vs follow-up CS**H** conditioning CS vs follow-up CS**I** follow-up US<sub>2</sub>




**A** left occipital cortex**B** conditioning CS vs test CS**C** conditioning CS vs test CS**D** test CS vs NSs**E** test CS vs NSs**F** test US<sub>2</sub>**G** conditioning CS vs follow-up CS**H** conditioning CS vs follow-up CS**I** follow-up US<sub>2</sub>

**A** left dorsolateral prefrontal cortex**B** conditioning CS vs test CS**C** conditioning CS vs test CS**D** test CS vs NSs**E** test CS vs NSs**F** test US<sub>2</sub>**G** conditioning CS vs follow-up CS**H** conditioning CS vs follow-up CS**I** follow-up US<sub>2</sub>


### A preconditioning CS



### B conditioning US




### C conditioning CS







**A** discomfort-inducing procedure




**B** conditioning CS vs test CS




**C** conditioning CS vs follow-up CS




### A preconditioning CS




### B conditioning US




### C conditioning CS



### A preconditioning CS



### B conditioning US



### C conditioning CS

