

AperTO - Archivio Istituzionale Open Access dell'Università di Torino

**Determination of trace antibiotics in water and milk via preconcentration and cleanup using activated carbons**

**This is the author's manuscript**

*Original Citation:*

*Availability:*

This version is available <http://hdl.handle.net/2318/1876180> since 2025-01-24T15:52:41Z

*Published version:*

DOI:10.1016/j.foodchem.2022.132695

*Terms of use:*

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)

# Determination of trace antibiotics in water and milk via preconcentration and cleanup using activated carbon

Pengyun Liu<sup>1</sup>, Zhilin Wu<sup>1</sup>, Alessandro Barge<sup>1</sup>, Luisa Boffa<sup>1</sup>, Katia Martina,<sup>1</sup>

Giancarlo Cravotto<sup>1,2\*</sup>

<sup>1</sup> Department of Drug Science and Technology, University of Turin, via P. Giuria 9,  
Turin, 10125, Italy.

<sup>2</sup> World-Class Research Center "Digital biodesign and personalized healthcare",  
Sechenov First Moscow State Medical University, Moscow, 119991 Russia.

\*Correspondence: giancarlo.cravotto@unito.it (G. Cravotto), Tel: +39.011.670.7183,

Fax: +39.011.670.7162.

Abbreviations: ABX, antibiotics; SMM, sulfamonomethoxine sodium; OTC, oxytetracycline; CEF, ceftiofur hydrochloride; MAR, marbofloxacin; MRLs, maximum residue limits; HPLC, high pressure liquid chromatography; SPE, solid phase extraction; RE, recovery efficiency; ACs, activated carbons; CPAC, coconut powdered AC;  $S_{BET}$ , Brunauer-Emmett-Teller surface area;  $V_{Meso}$ , mesopore volume;  $V_{Micro}$ , micropore volume; PGAC, peat granular AC; WPAC, wood powder AC; MeCN, acetonitrile; TFA, trifluoroacetic acid; MeOH, methanol; EtOH, ethanol;  $NH_4OH$ , ammonia solution; EDTA, ethylenediaminetetraacetic acid disodium salt hydrate; t-BuOH, t-butanol; n-PrOH, n-propanol; MW, molecular weight; MV, molar volume;  $LogK_{OW}$ , octanol-water partition coefficient;  $S_{Water}$ , solubility of ABX in water;  $S_{EtOH}$ , saturated mole fraction solubility of ABX in EtOH;  $pK_a$ , dissociation constant; NHA, number of H-bond acceptors; NHD, number of H-bond donors; Refs., references;  $Bp$ , boiling point;  $\epsilon$ , dielectric constant; EDA,  $\pi$ - $\pi$  electron donor-acceptor; TC, tetracycline; LODs, limit of detections; LOQs, limit of quantitations; LRs, linear ranges; RSD, relative standard deviation; FA, formic acid; DCM, methylene chloride; MSPD, magnetic solid phase dispersion; MSPE, magnetic solid phase extraction.

36 **Abstract**

37 CPAC-SPE-HPLC (coconut powdered activated carbon -SPE- HPLC) has been developed  
38 for the determination of antibiotic (ABX), sulfamonomethoxine sodium (SMM),  
39 oxytetracycline (OTC), ceftiofur hydrochloride (CEF) and marbofloxacin (MAR), in water and  
40 milk. Over 99.0% SMM and OTC were recovered from 20 mL of 0.5 µg/mL ABX solution  
41 using 10 mg-CPAC for adsorption and 2 mL of 30% NH<sub>4</sub>OH/EtOH (1/19 v/v) for elution.  
42 Similarly, over 99.0% CEF and MAR were recovered using 15 mg-CPAC and 2 mL of 30%  
43 NH<sub>4</sub>OH/n-PrOH (1/19 v/v). Moreover, the *REs* of various ABX from 5-80 mL of 0.02-2.00  
44 µg/mL medicated milk containing 10 mM EDTA are ordered as follows: OTC (99.3%), SMM  
45 (99.1%) > CEF (68.9%) > MAR (61.4%). No interference towards HPLC analysis were  
46 observed with elution using 2 mL of 30% NH<sub>4</sub>OH/EtOH (1/19 v/v). Furthermore, much lower  
47 limit of detections (0.02 µg/mL) than the maximum residual limits from European Commission  
48 (0.075-0.100 µg/mL) were obtained.

49

50 **Keywords:** *Antibiotic residues in milk; Preconcentration; Cleanup; Activated carbon; Solid phase*  
51 *extraction.*

52

53

## 54 **1. Introduction**

55 Antibiotics (ABX) sulfamonomethoxine sodium (SMM), oxytetracycline (OTC),  
56 ceftiofur hydrochloride (CEF) and marbofloxacin (MAR), etc. are currently frequently  
57 used in European dairy farms, and their residues in milk are cause for serious concern  
58 in Europe. The European Commission has adopted maximum residue limits (MRLs) of  
59 SMM (100 µg/L), OTC (100 µg/L), CEF (100 µg/L), and MAR (75 µg/L) in milk to  
60 guarantee the safety of dairy products and the health of human beings [1]. Thus, the  
61 development of QuEChERS (quick, easy, cheap, effective, rugged and safe) methods  
62 for determining the levels of trace ABX in milk is important for dairy farmers, milk  
63 processors, regulatory authorities and researchers [2,3].

64 The methods for determining trace ABX in milk include chromatographic methods,  
65 microbiological approaches, immunochemical techniques and biosensors, etc. [2–5],  
66 with high pressure liquid chromatography (HPLC) seeing extensive use. As milk  
67 contains complex ingredients, such as protein, lactose, fat and inorganic ions, the  
68 determination of ABX can be disturbed by milk ingredients during HPLC analysis [6].  
69 This means that sample pretreatment is the most critical procedure as it can eliminate  
70 interference from milk ingredients (cleanup) and enrich ABX in magnitude  
71 (preconcentration) in order to achieve accurate ABX determination [2,7,8]. Several  
72 pretreatment methods, including solvent extraction, solid phase extraction (SPE) and  
73 centrifugal ultrafiltration, etc., have been developed [2,9,10]. SPE is an updated  
74 pretreatment method that is based on low-pressure liquid chromatography and liquid-  
75 solid phase equilibrium, and has been widely used to separate target analytes from

76 samples, preconcentrate ABX and clean samples for HPLC analyses [2,11]. Compared  
77 to solvent extraction, SPE is more convenient and efficient. However, deproteinization  
78 and the follow-up of centrifugation, sample loading, evaporation, reconstitution and  
79 filtration are still necessary [10]. If the deproteinization procedure of milk is skipped,  
80 high throughput of milk in the SPE column can cause blockage, an unstable flowrate  
81 and inconsistent adsorption times. In addition, either MeOH or H<sub>2</sub>O are usually used to  
82 clean up SPE columns after sample loading to further eliminate the interference of  
83 impurities [5,6,11].

84 Adsorption efficiency is directly associated with the recovery efficiency (*RE*) of  
85 ABX when using SPE. Various adsorbents, such as molecularly imprinted polymers,  
86 metal-organic frameworks and carbon materials, etc. have been developed for milk  
87 pretreatment [2,8,12]. Of these, activated carbons (ACs) besides the low cost, high  
88 adsorption capacity and recyclability, it shows the widest availability. ACs are ideal  
89 adsorbents as they also possess distinctive physicochemical properties, such as rich  
90 mesopores, high specific surface areas and a variety of functional groups [13,14]. The  
91 ABX in milk can be adsorbed onto ACs via physical interactions, such as electrostatic  
92 interactions,  $\pi$ - $\pi$  bonds, H-bonds, etc., and the loaded ABX on the ACs can then be  
93 eluted by different eluents [15]. Interestingly, mesoporous ACs belong to the group of  
94 “restricted access materials”. The narrow pore diameter of ACs only allows low  
95 molecular weight ABX to enter, while high molecular weight proteins or fats cannot  
96 enter mesopores [16]. The procedure of chemical deproteinization can therefore  
97 probably be skipped when ACs are used for the recovery of ABX from milk.

108 Furthermore, chelates that are formed between ABX, metal ions ( $\text{Ca}^{2+}$  and  $\text{Mg}^{2+}$ )  
109 and protein in milk can diminish the adsorption of ABX [2,5,6]. Chelating agents,  
110 including EDTA, oxalic acid and citric acid, have been added to milk to promote the  
111 recovery of ABX [2,5,6,9].

112 This study aims to develop an AC-SPE-HPLC approach for the determination of  
113 trace SMM, OTC, CEF and MAR in milk. Based on the physicochemical properties of  
114 ABX and eluents, such as acidity and alkalinity, dissociation, polarity, H-bonds, etc.,  
115 the adsorption and elution conditions of the SPE for the recovery of the model ABX  
116 from water were optimized. Furthermore, the *REs*, using different ACs, of ABX at  
117 various spiked concentrations in milk were evaluated in the absence and presence of  
118 EDTA. Finally, the proposed AC-SPE-HPLC method was verified and compared with  
119 methods from other published works.

## 110 **2. Materials and Methods**

### 111 **2.1. Materials and chemicals**

112 Pasteurized commercial skimmed milk was purchased from a local branch of a  
113 major supermarket (Turin, Italy) and stored at room temperature. Preliminary analyses  
114 demonstrated that the purchased milk was ABX-free. The main ingredients in 100 mL  
115 of milk were carbohydrate (4.8 g), sugar (4.8 g), protein (3.4 g), saturated fat (0.5 g),  
116 fatty acid (0.3 g) and salt (0.13 g). The medicated milk samples were prepared daily by  
117 adding the stock ABX solutions into the milk. The initial pH of raw, 10 mg/L CEF-, 10  
118 mg/L SMM-, 10 mg/L MAR-, and 10 mg/L OTC-spiked milk samples were determined  
119 to be 6.46, 6.56, 6.50, 6.56, and 6.58, respectively.

120 Coconut powdered AC (CPAC) was proven to act as a mesoporous adsorbent in  
121 our previous studies [13,15], and was used as the core adsorbent for the pretreatment  
122 of milk via SPE in this study. The Brunauer-Emmett-Teller (BET) surface area ( $S_{\text{BET}}$ ),  
123 mesopore volume ( $V_{\text{Meso}}$ ), micropore volume ( $V_{\text{Micro}}$ ) and average pore size of CPAC  
124 were measured to be 1952 m<sup>2</sup>/g, 1.57 cm<sup>3</sup>/g, 1.76 cm<sup>3</sup>/g and 3.95 nm, respectively.  
125 CPAC was purchased from ACEF S.P.A. Piacenza (Italy). Peat granular AC (PGAC)  
126 and wood powder AC (WPAC) were provided by Merck-Sigma-Aldrich, Milan (Italy).  
127 The textural properties of PGAC, WPAC and CPAC have been presented in the  
128 literature [13].

129 SMM (400 mg/mL, Daimeton 40, IZO Srl), OTC (92.7 mg/mL, Oextra MV 10,  
130 Huvepharma), CEF (50 mg/mL, Ceva Santé Animale), and MAR (100 mg/mL,  
131 Vetoquinol) were used as the model ABX in this work. The initial pH of 10 mg/L CEF-,  
132 10 mg/L SMM-, 10 mg/L MAR-, and 10 mg/L OTC aqueous solution were determined  
133 to be 5.51, 6.67, 5.63, and 6.67, respectively. Their chemical structures, major  
134 physicochemical properties and toxicological parameters are shown in Table S1 and in  
135 Scheme 1 (in Section 3.1.3), respectively. The information of solvents and water used  
136 was listed in the supplementary material (Section S1.0).

## 137 2.2. Apparatus and instrument

138 The analysis of water and milk samples was performed using a range of apparatus,  
139 including a high-speed refrigerated centrifuge (Allegra® 64R, Beckman Coulter, US),  
140 a UV-Vis spectrometer (Cary 60 UV-vis Spectrophotometer, USA), a HPLC system  
141 (Waters Corp., Milford, MA, USA) coupled with a diode array detector (UV/DAD,

142 Waters Corp., Milford, MA, USA), an automatic sampler (Waters Corp., Milford, MA,  
143 USA) and a binary HPLC pump (Waters Corp., Milford, MA, USA). An Xterra RP 18  
144 separation column (5  $\mu\text{m}$ , 150 mm $\times$ 4.6 mm; Phenomenex, Torrance, CA, USA) was  
145 used with gradient elution and UV-DAD acquisition.

146 The maximum absorption wavelengths for the UV-Vis analysis of SMM, OTC,  
147 CEF and MAR in water were found to be 273, 267, 288 and 293 nm, respectively. The  
148 HPLC-DAD analysis of the model ABX was performed according to the methods  
149 described in [Table S2](#). Briefly, the Xterra RP 18 separation column was pre-equilibrated  
150 for 21 min with 0.1% TFA in H<sub>2</sub>O (Phase A) and 0.1% TFA in MeCN (Phase B). A needle  
151 installed onto an automatic sampler, used for injections, was auto-washed with  
152 MeOH/H<sub>2</sub>O (1/1 v/v) before each sample injection. The injection volume was set as 20  
153  $\mu\text{L}$ . The samples were loaded and analyzed using HPLC-DAD at room temperature,  
154 and the flowrate of the mobile phase was set at 0.8 mL/min. The total running time for  
155 the analysis of the model ABX ranged from 26 to 32 min. After the analysis of a batch  
156 of samples, the separation column was washed with H<sub>2</sub>O (phase A) and MeCN (phase  
157 B) for 50 min. The identification of ABX was performed by comparing the UV  
158 absorption spectra and retention times of the milk samples and standard solutions.

### 159 **2.3. Preconcentration and cleanup of milk samples**

160 The pretreatment procedures, namely, the preconcentration and cleanup of trace  
161 ABX in milk for HPLC analysis, are shown in [Fig. 1](#).






Fig. 1. Procedures of pre-concentration and cleanup for the HPLC analysis of trace ABX in milk.

It is necessary to note that different amounts of EDTA had previously been added to the 5-80 mL milk samples so that 10 mM EDTA was contained in each. These were then shaken for 5 min to ensure that the EDTA adequately interacted with the milk matrix and that the milk flowed smoothly in the SPE column [2,5,6,9].

The follow-up procedures were conducted as follows. Firstly, either 10 mg powder AC or 50 mg granular AC were placed into the barrel of a glass syringe (i.d. 1.5 cm), and glass wool was fitted above and below the AC layer to prevent the AC washing out. Secondly, the 5-80 mL medicated-milk samples were passed through the AC column continuously. As a result, trace ABX were absorbed onto the AC and the white milk flowed out from the AC column. The AC column was then flushed, using 1 mL of distilled water, to remove the small amount of milk that adhered to the AC, and the

175 small amount of water in the AC-column was extruded out. Afterwards, 2 mL of eluent  
176 was passed through the ABX-loaded AC column to elute the adsorbed ABX and the  
177 elutant was collected in a glass vial. Furthermore, 2 mL of the elutant were concentrated,  
178 via mild N<sub>2</sub> purge, to 0.5 mL at room temperature. Some insoluble matter was observed  
179 and was separated by centrifugation at 26000 rpm for 1 min. Finally, the supernatant  
180 was collected and stored at 4 °C for HPLC analysis.

#### 181 **2.4. Evaluation of recovery efficiency**

182 The medicated milk samples containing 0.02, 0.10, 0.50 and 1.00 µg/mL ABX  
183 were used for the evaluation of the *REs*. The method for preconcentration and cleanup  
184 using AC-SPE is described in [Section 2.3](#). *RE* depended on adsorption/desorption  
185 efficiency and was calculated in accordance with [Eq. \(1\)](#):

$$186 \quad RE (\%) = \frac{C_d V_d}{C_0 V_0} \times 100\% \quad (1)$$

187 where,  $C_d$  (µg/mL) is the determined concentration of ABX in the concentrated elutant,  
188  $V_d$  (mL) is the volume of the concentrated elutant,  $C_0$  (µg/mL) is the spiked  
189 concentration of ABX in the milk sample, and  $V_0$  (mL) is the volume of the milk sample  
190 that passed through the AC-SPE column.

191 All experiments were repeated at least twice and errors are shown as the difference  
192 between the highest measured values and the average value in parallel experiments.  
193 When errors are not visible in the figures, they are smaller than the symbols  
194 representing the average values.


### 195 **3. Results and discussion**

#### 196 **3.1. Optimization of AC-SPE for ABX recovery from water**


197 The feasibility of using ACs as an adsorbent in SPE is the primary concern here.  
198 Although ACs are undoubtedly superior adsorbents, the desorption of organic  
199 adsorbates from them is generally considered to be a considerable challenge [17,18]. It  
200 was therefore necessary to demonstrate the high *RE* of ABX using ACs in water. CPAC  
201 has been proven to provide superior adsorption/desorption performance in our previous  
202 studies [13,15]. In the preliminary study, using water samples instead of milk samples  
203 made the whole AC-SPE operation easier, more economical and more environmentally  
204 friendly. The effects of CPAC amount, the flowrate of the water sample, the type,  
205 volume and flowrate of the eluent and the use of various ABX were evaluated in the  
206 search of optimal conditions on the *REs*.

##### 207 **3.1.1. Optimization of adsorption conditions**

208 Both the amount of adsorbent and the flowrate of the water sample through the  
209 SPE column are critical factors that influence the *RE* [8,19]. Firstly, the *REs* of the  
210 model ABX were evaluated with various doses of CPAC (5-15 mg). 20 mL of 0.5  
211  $\mu\text{g/mL}$  aqueous solutions of ABX were passed through the CPAC-columns at a flowrate  
212 of 1.00 mL/min (Adsorption) and 2 mL of 30%  $\text{NH}_4\text{OH}/\text{EtOH}$  (1/19 v/v) were passed  
213 through the CPAC-columns at a flowrate of 0.07 mL/min (Elution) at room temperature,  
214 and the results are presented in Fig. 2a.



215



216


217 **Fig. 2.** Effect of CPAC amount on *REs* of ABX in water. (a) *REs* using various amount of CPAC and 30%  
 218 NH<sub>4</sub>OH/EtOH (1/19 v/v) at room temperature (Adsorption conditions: 20 mL of 0.5 μg/mL ABX were passed  
 219 through CPAC columns at 1.00 mL/min. Elution conditions: 2 mL of 30% NH<sub>4</sub>OH/EtOH (1/19 v/v) were passed  
 220 through the ABX-loaded columns at 0.07 mL/min); (b) *REs* of MAR and CEF using various amount of CPAC and  
 221 30% NH<sub>4</sub>OH/n-PrOH (1/19 v/v) at room temperature (Adsorption conditions: 20 mL of 0.5 μg/mL ABX were passed  
 222 through CPAC columns at 1.00 mL/min. Elution conditions: 2 mL of 30% NH<sub>4</sub>OH/n-PrOH (1/19 v/v) were passed  
 223 through the ABX-loaded columns at 0.07 mL/min).

224 As shown in [Fig. 2a](#), the *REs* of SMM and OTC are much higher than those of  
 225 CEF and MAR. The reason will be discussed in [Section 3.1.3](#). In addition, increasing  
 226 CPAC amount initially increases the *REs*, the peak *REs*, 99.4% (SMM), 99.2% (OTC),

227 62.1% (CEF) and 60.4% (MAR), are achieved with 10 mg of CPAC, and then the *REs*  
228 decrease slowly with increasing the amount of CPAC. Generally, the increased amount  
229 of adsorbent provided more adsorption sites, which ensured that ABX was efficiently  
230 adsorbed. However, it is difficult to elute ABX at the higher adsorbent amount (> 10  
231 mg of CPAC). The effects of eluent type and amount are discussed in [Section 3.1.2](#).

232 To further increase the *REs* of CEF and MAR, the effect of CPAC amount on the  
233 *REs* of CEF and MAR with elution using 30% NH<sub>4</sub>OH/n-PrOH (1/19 v/v) was further  
234 investigated. 20 mL of 0.5 µg/mL aqueous solutions of CEF and MAR were passed  
235 through the 5-15 mg CPAC-columns at a flowrate of 1.00 mL/min at room temperature,  
236 and 2 mL of 30% NH<sub>4</sub>OH/n-PrOH (1/19 v/v) were used for the elution. The results are  
237 shown in [Fig. 2b](#). Obviously, the *REs* of CEF and MAR increase with increasing of  
238 CPAC amount and the highest *REs* reach 102.9% for CEF and 99.1% for MAR with 15  
239 mg of CPAC. As compared with elution using 30% NH<sub>4</sub>OH/EtOH (1/19 v/v), the *REs*  
240 of CEF and MAR increase by 66% and 81.2% using 30% NH<sub>4</sub>OH/n-PrOH (1/19 v/v)  
241 as eluent, respectively. Importantly, higher amount of the adsorbent (15 mg of CPAC)  
242 are required to fully catch CEF and MAR from water.

243 In addition, the adsorption efficiency of ABX is also dependent on the flowrate of  
244 the water sample. For example, 20 mL of 0.5 µg/mL MAR aqueous solutions were  
245 passed through the 10 mg CPAC-columns at various water-sample flowrates (0.30-2.00  
246 mL/min) at room temperature, and elution with 30% NH<sub>4</sub>OH/EtOH (1/19 v/v) or 30%  
247 NH<sub>4</sub>OH/n-PrOH (1/19 v/v), and the results are shown in [Fig. 3](#).



248

249 Fig. 3. Effect of water-sample flowrates on the *REs* of MAR at room temperature (Adsorption conditions: 20 mL of  
 250 0.5 µg/mL MAR were passed through 10 mg-CPAC columns. Elution conditions: 2 mL of eluents were passed  
 251 through the MAR-loaded columns at 0.07 mL/min).

252

As shown in Fig. 3, using 10 mg of CPAC for adsorption and 2 mL of 30%  
 253 NH<sub>4</sub>OH/EtOH (1/19 v/v) for elution, no obvious effects of water-sample flowrates  
 254 between 0.30-1.00 mL/min were observed, but the *REs* of MAR drops from 60.4% to  
 255 51.9% as the flowrate increases from 1.00 to 2.00 mL/min. Therefore, the higher sample  
 256 flowrate results in an insufficient adsorption time and lower *REs* [19]. The similar  
 257 effects of water-sample flowrates were observed with 10 mg of CPAC for adsorption  
 258 and 2 mL of 30% NH<sub>4</sub>OH/n-PrOH (1/19 v/v) for elution.

259

The effect of water-sample flowrates on *REs* of MAR with a higher CPAC amount  
 260 was further studied. With 15 mg of CPAC for adsorption and 2 mL of 30% NH<sub>4</sub>OH/n-  
 261 PrOH (1/19 v/v) for elution, as expected, the *REs* reached 99.1%, 98.1%, 97.7% and  
 262 98.2% at the sample flowrates of 0.30, 0.70, 1.00 and 2.00 mL/min, respectively. It  
 263 indicates that no obvious effects of water-sample flowrate on the *REs* of MAR were  
 264 observed, 15 mg of CPAC is sufficient for catching trace MAR and 30% NH<sub>4</sub>OH/n-

265 PrOH (1/19 v/v) is an ideal eluent.

### 266 3.1.2. Optimization of elution conditions

267 The elution conditions, e.g. the type, volume and flowrate of the eluent, determine  
268 the selectivity and efficiency of elution [5,8,20]. Acidity and alkalinity, polarity  
269 (hydrophobicity, dielectric constant and LogK<sub>ow</sub> value) and the H-bond interaction (H-  
270 donor/acceptor count) of eluents are the critical factors for the elution of organics from  
271 ACs [13]. The major physicochemical properties of the eluents are listed in Table 1.


272 Table 1. Major physicochemical properties of eluents (Data were adopted from Pubchem and Drugbank databases).

<i>Eluent</i>	<i>MW</i> ( <i>g/mol</i> )	<i>Bp</i> ( <i>°C</i> )	<i>LogK<sub>ow</sub></i>	<i>pK<sub>a</sub></i>	<i>NHA</i>	<i>NHD</i>	<i>ε</i>
H <sub>2</sub> O	18	100.0	-	15.7	1	1	80.4
HOAc	60	117.9	-0.17	4.8	2	1	6.2
NH <sub>4</sub> OH	35	38.0	-	9.2	1	2	2.5
MeCN	41	81.6	-0.34	8.8	1	0	38.8
MeOH	32	64.7	-0.77	13.1	1	1	32.7
EtOH	46	78.2	-0.31	15.9	1	1	24.6
n-PrOH	60	97.2	0.25	16.1	1	1	20.3
t-BuOH	74	84.2	0.35	19.2	1	1	10.9
n-BuOH	74	117.8	0.88	16.1	1	1	17.5

273 **Note:** MW, molecular weight; *Bp*, boiling point; LogK<sub>ow</sub>, octanol-water partition coefficient; pK<sub>a</sub>, dissociation  
274 constant; NHA, number of H-bond acceptors; NHD, number of H-bond donors; ε, dielectric constant.  
275

276 Referring to our previous work and according to pre-experiments, 30%  
277 NH<sub>4</sub>OH/EtOH (1/19 v/v) was proved to be an ideal eluent for the desorption of organics  
278 from CPAC. Based on Fig. 2a, the superior REs of SMM and OTC (> 99%) were  
279 achieved with elution using 30% NH<sub>4</sub>OH/EtOH (1/19 v/v), but the REs of CEF and  
280 MAR were around 60%. In contrast, more than 99% of CEF and MAR were recovered  
281 with 15 mg of CPAC for adsorption and 2 mL of 30% NH<sub>4</sub>OH/n-PrOH (1/19 v/v) for  
282 elution (Fig. 2b). Therefore, CEF and MAR were selected as the model ABX to evaluate

283 the role of various eluents on their *REs* [8,20]. Typically, 20 mL of 0.5 µg/mL CEF or  
 284 MAR aqueous solutions were passed through the 10 mg CPAC-columns at a water-  
 285 sample flowrate of 1.00 mL/min at room temperature. The *REs* of CEF and MAR using  
 286 various eluents are shown in Fig. 4.



287  
 288 Fig. 4. *REs* of CEF and MAR in water using CPAC columns and various eluents at room temperature (Adsorption  
 289 conditions: 20 mL of 0.5 µg/mL CEF or MAR were passed through 10 mg-CPAC columns at 1.00 mL/min. Elution  
 290 conditions: 2 mL of various eluents were passed through the ABX-loaded columns at 0.07 mL/min).

291 As shown in Fig. 4, remarkable different *REs* of CEF and MAR were observed  
 292 with elution using various eluents. CEF is a hydrophobic acidic compound with a lower  
 293  $S_{\text{water}}$  (100 mg/L), a higher  $\text{Log}K_{\text{OW}}$  value (1.2), and a lower  $\text{p}K_{\text{a}}$  value (2.83). In contrast,  
 294 MAR is a relatively hydrophilic acidic compound with higher  $S_{\text{water}}$  (2600 mg/L), a  
 295 lower  $\text{Log}K_{\text{OW}}$  value (-0.53), and a higher  $\text{p}K_{\text{a}}$  value (5.38), and MAR can be dissolved  
 296 in EtOH, DMSO, DMF, etc.[21].

297 Considering the different physicochemical properties of CEF and MAR and the



298 like-dissolves-like rule, MeOH, EtOH, n-PrOH, t-BuOH or n-BuOH with various  
299 polarities were used to elute CEF or MAR from CPAC, respectively (Table 1). The  
300 results show that the *REs* of CEF were 57.1%, 44.3% and 28.8% with elution using n-  
301 PrOH, n-BuOH and t-BuOH, respectively, while the *REs* of MAR obtained with elution  
302 using MeOH, EtOH and n-PrOH reached 15.2%, 29.8% and 22.0%, respectively. In  
303 addition, the very polar aprotic or protic solvents, water and MeCN are ineffective for  
304 eluting both of CEF and MAR.

305 The acidic and alkali eluents, HOAc and 30% NH<sub>4</sub>OH, were evaluated next. HOAc  
306 is known as a common carboxylic acid with high polarity and potential for H-bonding  
307 with the –COOH, –F or –NH<sub>2</sub> groups that exist in CEF or MAR molecules. NH<sub>4</sub>OH is  
308 a common alkali reagent with high polarity and potential for H-bonding, which favors  
309 the desorption of acidic CEF and MAR. The results show that the *REs* obtained only  
310 reached 13.2% (CEF) and 12.7% (MAR) with elution using HOAc, as well as 25.1%  
311 (CEF) and 23.8% (MAR) with elution using 30% NH<sub>4</sub>OH, respectively. The  
312 effectiveness of 30% NH<sub>4</sub>OH is higher than that of HOAc due to the dissociation of –  
313 COOH in the CEF and MAR molecules under alkali conditions [22].


314 To further improve the *REs* of CEF and MAR, acidic and alkali solvents were  
315 mixed with alcohols, such as MeOH, EtOH, n-PrOH, t-BuOH or n-BuOH to form co-  
316 solvents, so that the acidity and alkalinity, polarity and H-bond interaction of the eluents  
317 can synergistically enhance the desorption. As expected, the *REs* of CEF obtained by  
318 adding 30% NH<sub>4</sub>OH to n-PrOH, t-BuOH, and n-BuOH were significantly increased to  
319 75.8%, 46.0% and 54.9%, respectively. Meanwhile, the *REs* of MAR obtained by

320 adding 30% NH<sub>4</sub>OH to MeOH, EtOH and n-PrOH were extremely increased to 52.0%,  
321 60.4% and 71%, respectively. The more effective elution of CEF and MAR by adding  
322 30% NH<sub>4</sub>OH to n-PrOH than other alcohols could be attributed the appropriate polarity  
323 of the 30% NH<sub>4</sub>OH/n-PrOH [8,23,24]. In contrast, the *REs* of MAR obtained using  
324 HOAc/MeOH (1/19 v/v) and HOAc/EtOH (1/19 v/v) were 20.5% and 24.5%,  
325 respectively, due to the variation of the H-bond interactions and the –COOH  
326 dissociation of MAR [8,23,24]. In summary, Lewis bases solvents as eluents exhibit  
327 higher desorption efficiency than neutral and Lewis acid solvents [23,24]. The acidity  
328 or alkalinity, the polarity and the H-bond interactions of an eluent jointly determined  
329 the desorption efficiency and the co-solvent 30% NH<sub>4</sub>OH/n-PrOH (1/19 v/v) was an  
330 optimal eluent.


331 To further improve the desorption and to minimize the consumption of eluents, the  
332 effect of eluent volume on the *REs* of CEF and MAR was studied. In the preliminary  
333 study, 20 mL of 0.5 µg/mL MAR were passed through 10 mg-CPAC columns at 1.00  
334 mL/min, and then 1-10 mL of 30% NH<sub>4</sub>OH/EtOH (1/19 v/v) were passed through the  
335 MAR-loaded columns at 0.07 mL/min. As a result, the *REs* of MAR reached 36.6%,  
336 60.4%, 60.75, 61.0%, 61.8 and 64.3% with elution using 1, 2, 4, 6, 8 and 10 mL of the  
337 eluent, respectively. The *REs* of MAR increased remarkably as the eluent volume  
338 increased from 1 to 2 mL, but only a slight increase in *REs* occurred after 2 mL.

339 To further understand the effect of eluent volume on the *REs*, 30% NH<sub>4</sub>OH/n-PrOH  
340 (1/19 v/v) was used as eluent. 20 mL of 0.5 µg/mL MAR or CEF were passed through  
341 15 mg-CPAC columns at 1.00 mL/min, and then 1-4 mL of the eluent were passed

342 through the ABX loaded columns at 0.07 mL/min. The results are reported in Fig. 5a.



343



344

345 **Fig. 5.** Optimizing the elution processes of CEF and MAR from CPAC at room temperature. (a) Effect of eluent  
346 volumes on the *REs*-of CEF and MAR (Adsorption conditions: 20 mL of 0.5 µg/mL ABX were passed through 15  
347 mg-CPAC columns at 1.00 mL/min. Elution conditions: 1-4 mL of 30% NH<sub>4</sub>OH/n-PrOH (1/19 v/v) were passed  
348 through the ABX-loaded columns at 0.07 mL/min); (b) Effect of eluent flowrates on the *REs* of MAR (Adsorption  
349 conditions: 20 mL of 0.5 µg/mL MAR were passed through CPAC columns at 1.00 mL/min. Elution conditions: 2  
350 mL of 30% NH<sub>4</sub>OH/n-PrOH (1/19 v/v) were passed through the MAR-loaded columns at various flowrates).

351 As shown in Fig. 5a, the *REs* of MAR and CEF increased from 56.6% to 99.1%,  
352 and from 76.7% to 102.9% as the volume of 30% NH<sub>4</sub>OH/n-PrOH (1/19 v/v) increased  
353 from 1 to 2 mL, respectively, but no changes in the *REs* were observed after 2 mL. It is  
354 similar to the effects of the 30% NH<sub>4</sub>OH/EtOH (1/19 v/v) volume on the *REs* of MAR

355 and previous study's result [25]. The results reveal that 15 mg CPAC is sufficient for  
356 full adsorption of CEF and MAR from water, and 2 mL of 30% NH<sub>4</sub>OH/n-PrOH (1/19  
357 v/v) is a better eluent for eluting CEF and MAR from CPAC.

358 Furthermore, the desorption efficiency of ABX is also dependent on the flowrate  
359 of eluents [26]. The *REs* of MAR from loaded CPAC columns were evaluated at various  
360 flowrates of 2 mL of 30% NH<sub>4</sub>OH/n-PrOH (1/19 v/v) at room temperature, and the  
361 results are shown in Fig. 5b. Using 10 or 15 mg of CPAC for adsorption and 2 mL of  
362 30% NH<sub>4</sub>OH/n-PrOH (1/19 v/v) for elution, no obvious effects of eluent flowrates  
363 between 0.07-0.15 mL/min were observed, but *REs* of MAR significantly drops as the  
364 flowrate increases from 0.15 to 0.50 mL/min (Fig. 5b). Obviously, a higher eluent  
365 flowrate results in an insufficient desorption time and lower *REs* [26]. Consequently,  
366 0.15 mL/min was proven to be the optimal flowrate for eluting MAR from CPAC.


### 367 3.1.3. Recovery mechanism of ABX from water

368 The *REs* of ABX are different despite the same analysis method being used as they  
369 all have different chemicals structures and physicochemical properties [2,5]. As shown  
370 in Fig. 2a, the *REs* of SMM and OTC are much higher than those of CEF and MAR,  
371 suggesting that 10 mg-CPAC as an absorbent and 2 mL of 30% NH<sub>4</sub>OH/EtOH (1/19  
372 v/v) as an eluent are highly efficient for the recovery of trace SMM and OTC. In contrast,  
373 the *REs* of CEF and MAR reached 102.9% and 99.1%, respectively, as 15 mg CPAC  
374 and 2 mL of 30% NH<sub>4</sub>OH/n-PrOH (1/19 v/v) were used (Fig. 2b).

375 Generally, the  $S_{BET}$ ,  $V_{Meso}$  and  $V_{Meso}$  dominate the adsorption performance of ACs  
376 [17,27,28]. As mentioned in Section 2.1, CPAC possesses a very large  $S_{BET}$  and rich

377 micro- and mesopores with a reasonable fraction of micro- and mesopores (1.12) [13].  
 378 Therefore, CPAC is a superior adsorbent toward ABX in water and milk [15].  
 379 Adsorption of ABX onto ACs is a complex process as it is closely associated with the  
 380 physicochemical property of ACs, the molecule structure, property and geometric  
 381 barrier of ABX, and the adsorption conditions, etc. The critical physicochemical  
 382 properties of ABX, such as electrostatic interaction and Lewis acid-base interaction  
 383 ( $pK_a$ ), hydrophobicity ( $S_{water}$  and  $LogK_{OW}$ ), electronic coupling (the  $\pi$ - $\pi$  electron donor-  
 384 acceptor (EDA) interactions and H-bonding), pore filling (molar volume and molecular  
 385 size), etc., influence the adsorption affinity [27–31].

386 The major physicochemical properties, chemical structures and the molecular sizes  
 387 of model ABX are listed in Table S1 and Scheme 1.



Scheme 1 Chemical structures and molecular sizes of the model ABX (The molecular sizes were calculated from the Chem3D Program).

394 As shown in [Scheme 1](#), SMM contains a sulfonamide group, which connects with  
395 an aniline ring and an aromatic heterocyclic group. The unionized form of SMM has  
396 the smallest molecule size among the four model ABX, while CPAC has a high  $V_{\text{Meso}}$   
397 and  $S_{\text{BET}}$ , which favor the adsorption of low-sized sulfonamide ABX due to the pore-  
398 filling effect [\[27,31,32\]](#). In addition, the strong electron-withdrawing sulfonamide  
399 group causes the decrease of the  $\pi$ -electron density in aromatic rings and the  
400 heterocyclic group, and hence act as  $\pi$ -electron acceptors. In contrast, the  $\pi$ -electron-  
401 rich regions on the graphene surface of CPAC serve as  $\pi$ -electron donors. Such  $\pi$ - $\pi$   
402 EDA interactions enhance the adsorption of SMM on CPAC [\[27,29\]](#). Furthermore, the  
403 rich O-containing groups ( $4.04\pm 0.98\%$ ) of CPAC favor the adsorption of relatively  
404 hydrophilic SMM [\[29,30\]](#). Finally, electrostatic interactions and H-bonding may play  
405 additional roles [\[30,33\]](#).

406 OTC belongs to tetracycline ABX, possessing one tetracycline (TC) ring and  
407 multiple hydroxyl (-OH), N,N, dimethyl (-N(CH<sub>3</sub>)<sub>2</sub>) and amide (-C(O)NH<sub>2</sub>) groups on  
408 the TC ring. The molecular size of OTC is the second smaller among the four model  
409 ABX. Similar to the above, the higher adsorption affinity of OTC onto CPAC may be  
410 mainly attributed to the high  $V_{\text{Meso}}$ ,  $S_{\text{BET}}$  of CPAC, and the pore-filling effect [\[18,34\]](#).  
411 In addition, the electrostatic interaction and the  $\pi$ - $\pi$  interactions between OTC and  
412 CPAC should be involved in the adsorption of OTC [\[35\]](#). The multiple -OH, -N(CH<sub>3</sub>)<sub>2</sub>  
413 and -C(O)NH<sub>2</sub> groups in the OTC molecule are electron donors and induce strong  
414 conjugation with the  $\pi$ -electrons on aromatic rings and interact electrostatically with  
415 the positively charged regions on CPAC. The H-bonding between -OH group and O-

416 containing functional groups on CPAC is also important due to the appropriate contents  
417 of O-containing groups on CPAC. Also, O-containing groups facilitate the adsorption  
418 of relatively hydrophilic OTC [27,36].

419 Compared with the higher adsorption affinity of SMM and OTC on CPAC, the  
420 adsorption efficiencies of CEF and MAR onto CPAC are relatively lower (Fig. 2). CEF  
421 is the hydrochloride salt of ceftiofur, containing a  $\beta$ -lactam core structure (a cephem)  
422 coupling with a furan-carboxylic-thioester, an amide, an iminomethoxy, and a 2-amino-  
423 4-thiazoyl group. CEF has the largest molecular size and the highest hydrophobicity  
424 ( $\text{Log}K_{\text{ow}}$ : 1.2) among the four model ABX. The large molecular size of CEF causes a  
425 significant size-exclusion effect or geometric barrier, reducing the affinity with CPAC  
426 [27,31,32,34,37]. Meanwhile, its high hydrophobicity is unfavorable to the complex  
427 interactions of CEF with the rich O-containing groups on the surface of CPAC. Finally,  
428 acidic groups ( $-\text{COOH}$ ) in hydrophobic compounds negatively influence the adsorption  
429 on ACs due to the formation of large and dense water clusters [28,38].

430 MAR is one of fluoroquinolones, containing a bicyclic core structure related to the  
431 substance 4-quinolone, a fluorine atom in the chemical structure,  $-\text{COOH}$ , 3,5-  
432 dimethylpiperazine, carbonyl, etc. CEF has the second larger molecular size and the  
433 second higher hydrophobicity ( $\text{Log}K_{\text{ow}}$ : -0.5) among the four model ABX. As  
434 mentioned above, both larger molecular size and higher hydrophobicity are unfavorable  
435 to the adsorption of MAR on CPAC. It is similar to CEF,  $-\text{COOH}$  group in the MAR  
436 may reduce its affinity with CPAC [28,38]. Additionally, F-containing organics may  
437 exhibit a weaker affinity with ACs [39].

438 The effects of eluents on the elution of CEF and MAR from CPAC have been  
439 discussed in detail in [Section 3.1.2](#). It can be concluded that the acidity or alkalinity,  
440 polarity and H-bond interactions of eluent jointly dominate the elution performance of  
441 ABX from CPAC and the co-solvent 30% NH<sub>4</sub>OH/n-PrOH (1/19 v/v) is an appropriate  
442 eluent to elute CEF and MAR from CPAC [\[8,23,24\]](#). In contrast, 30% NH<sub>4</sub>OH/EtOH  
443 (1/19 v/v) is appropriate to effectively elute SMM and OTC. The prominent difference  
444 between these two eluents is the difference in polarities of EtOH and n-PrOH. The  
445 ionized SMM at the presence of NH<sub>4</sub>OH facilitates its elution by the polar eluent of 30%  
446 NH<sub>4</sub>OH/EtOH (1/19 v/v) based on the like-dissolves-like rule [\[29\]](#). Similarly, OTC with  
447 the strongest polarity among the four model ABX can be effectively eluted by 30%  
448 NH<sub>4</sub>OH/EtOH (1/19 v/v). Meanwhile, the strong H-bond interactions of 30%  
449 NH<sub>4</sub>OH/EtOH enhance the elution of SMM and OTC.

### 450 **3.2. Optimization of AC-SPE for ABX recovery from milk**

451 The major challenge encountered in the analysis of ABX in milk is separating the  
452 impurities, e.g. fat, protein, etc., and increasing the concentration of trace ABX [\[2,5\]](#).  
453 Using the optimization of AC-SPE for ABX recovery from water as a base, the  
454 technique was further optimized to validate the feasibility of recovering ABX from milk,  
455 and this included determining the *REs* of ABX using various adsorbents at different  
456 concentrations, determining the limit of detections (LODs), the limit of quantitations  
457 (LOQs), the linear ranges (LRs) and errors. Finally, the AC-SPE process was compared  
458 to other methods reported for the analysis of model ABX in milk.



459 **3.2.1. Recovery of ABX using various ACs at different concentrations**

460 To find an appropriate adsorbent, the *REs* of ABX were obtained at various  
 461 concentrations in different milk volumes using 50 mg PGAC, 10 mg WPAC or CPAC.  
 462 2 mL of 30% NH<sub>4</sub>OH/EtOH (1/19 v/v) were used to elute ABX from AC columns. As  
 463 a result, 0.5 mL HPLC samples were prepared from 10-80 mL of 0.02-1.00 µg/mL  
 464 medicated milk using AC-SPE; the analytes in milk were concentrated 20-160 fold.

465 It is difficult for large-volume milk to pass through the powder AC-SPE columns  
 466 due to the higher viscosity. Therefore, EDTA had previously been added to the milk  
 467 samples (containing 10 mM EDTA) to ensure the smooth flow of the large-volume milk  
 468 samples through the CPAC columns [2,5], which are designated as CPAC'. The results  
 469 of *REs* are summarized in Table 2.

470 Table 2. *REs* of ABX at various concentrations from milk using different ACs.

<i>ABX</i>	<i>C<sub>0</sub></i> (µg/mL)	<i>V<sub>s</sub></i> (mL)	<i>RE</i> (%) (mean, n=3)			
			PGAC	WPAC	CPAC	CPAC'
SMM	0.02	80	-	-	-	99.48±1.28
	0.10	40	4.89±0.06	47.82±0.17	98.18±1.03	99.53±0.51
	0.50	20	4.69±0.01	45.04±2.55	94.89±1.11	98.89±0.90
	1.00	10	4.24±0.00	40.57±2.26	94.79±0.68	98.59±0.75
OTC	0.02	80	-	-	-	99.67±0.42
	0.10	40	5.81±0.17	49.25±0.54	81.33±3.52	99.33±0.17
	0.50	20	4.49±0.03	45.70±0.75	79.78±1.7	99.30±0.06
	1.00	10	4.48±0.11	42.10±0.34	78.47±2.36	99.07±1.34
CEF	0.02	80	-	-	-	69.13±1.40
	0.10	40	1.16±0.02	11.81±0.30	38.41±1.26	69.01±0.16
	0.50	20	1.02±0.03	8.11±0.29	36.17±0.33	68.62±0.32
	1.00	10	0.89±0.00	8.30±0.02	38.36±3.33	68.79±0.32
MAR	0.02	80	-	-	-	61.93±1.05
	0.10	40	1.00±0.01	22.80±0.69	41.83±2.54	61.89±1.19
	0.50	20	0.90±0.09	21.47±0.19	38.65±0.02	60.98±0.37
	1.00	10	0.90±0.10	19.25±0.81	37.08±1.20	60.81±0.19

471 Adsorption conditions: 10-80 mL of medicated milk were passed through 10 mg-WPAC, CPAC and 50 mg-PGAC  
 472 columns at 1.00 mL/min at room temperature. Elution conditions: 2 mL of 30% NH<sub>4</sub>OH/EtOH (1/19 v/v) were

473 passed through the ABX-loaded AC columns at 0.07 mL/min at room temperature.

474 **Note:**  $C_0$ , initial concentration of ABX;  $V_s$ , volume of the sample;  $RE$ , recovery efficiency.

475


476 As shown in [Table 2](#), the  $RE$  ranges of various ABX using CPAC, WPAC and  
477 PGAC are 37.6%-96.0%, 9.4%-45.6% and 0.93%-4.93%, respectively. The  $REs$   
478 positively correlate with the  $S_{BET}$  of the ACs; the increasing  $S_{BET}$  provides more  
479 adsorption sites. The mean  $REs$  of ABX with CPAC at different concentrations in milk  
480 are ordered as follows: SMM (96.0%) > OTC (79.9%) > MAR (39.2%) > CEF (37.6%).  
481 The  $REs$  obtained from the milk samples are somewhat lower than those from the water  
482 samples, and this may be the result of the possible chelation of ABX with milk  
483 components. As for the differences in the  $REs$  of various ABX in milk, the order of  $REs$   
484 is similar to that obtained in water samples and reasons are discussed in [Section 3.1.3](#).

485 Surprisingly, adding EDTA to the milk samples not only made the milk flow more  
486 smoothly, but also significantly improved the  $REs$  of the model ABX. As listed on  
487 CPAC' in [Table 2](#), the mean  $REs$  with EDTA of SMM, OTC, CEF and MAR at different  
488 concentrations are 99.1%, 99.3%, 68.9% and 61.4%, respectively. These are 3.3%,  
489 24.4%, 56.7%, and 83.0% higher, respectively, than those obtained in the absence of  
490 EDTA. This remarkable improvement in  $REs$  may be attributed to the effective  
491 separation of ABX/proteins/metal ion (e.g.  $Ca^{2+}$ ) chelates by EDTA [9]. Thus, higher  
492  $REs$  can be obtained by CPAC-SPE for the analysis of ABX in milk.

493 Furthermore, 15 mg of CPAC and 30%  $NH_4OH/n-PrOH$  (1/19 v/v) have been  
494 proven to be effective adsorbent dose and eluent for the recovery of CEF and MAR in  
495 water samples. Aiming to increase the  $REs$  of CEF and MAR, 15 mg of CPAC and 30%

496  $\text{NH}_4\text{OH}/n\text{-PrOH}$  (1/19 v/v) were used as adsorbent and eluent. Unfortunately, higher  
497 *REs* cannot be achieved after unremitting efforts due to the interference of milk  
498 components. Hence it is necessary to promote the recovery of CEF and MAR in milk  
499 by using AC-SPE method in the future.

500 Besides the preconcentration of ABX from milk, the separation or removal  
501 efficiency of major components in milk is another concern. Fig. 6 shows the appearance  
502 of the samples during SPE process.



503  
504 Fig. 6 Appearance of samples collected during the pretreatment procedure for HPLC analysis.

505 As shown in Fig. 6, most of the white milk components that remained in the AC  
506 columns were washed out using 1 mL of distilled water, and very few residual white  
507 components were further removed from the elutant via centrifugation.

508 Fig. S1 illustrates a HPLC comparison of samples from blank milk, samples that

509 were recovered from medicated milk by using 10 mg of CPAC for adsorption and 2 mL  
510 of 30% NH<sub>4</sub>OH/EtOH (1/19 v/v) for elution, as well as standard ABX aqueous  
511 solutions. As shown in Fig. S1, only one peak appears for the standard CEF aqueous  
512 solution and this is at 12.6 min, while no peak was observed around 12.6 min for the  
513 sample from blank milk under the same HPLC conditions. This indicates that milk  
514 components did not interference with the CEF analysis. In contrast, the peak for CEF  
515 appears at around 12.6 min in the sample recovered from the CEF-spiked milk using  
516 CPAC-SPE. In addition, two tiny peaks, at around 7.8 min and 11.8 min, are found, and  
517 these probably arise from the impurities in the CEF stock solution. Similarly, the milk  
518 components did not interference in the OTC analysis, although three tiny peaks are  
519 found in the sample recovered from the OTC-spiked milk. Fortunately, the peaks do not  
520 disturb the OTC analysis. The HPLC analyses of MAR and SMM are better still as no  
521 problematic peaks are observed in the samples recovered from the MAR- and SMM-  
522 spiked milk. Moreover, EDTA did not appear in any HPLCs, which is consistent with a  
523 previous study [7]. Overall, the interference of milk components with the HPLC  
524 analysis of ABX by using CPAC-SPE can be neglected.

### 525 3.2.2. Validation of the CPAC-SPE-HPLC method

526 The European Commission Decision (2002/657/EC) on the performance of  
527 analytical methods and the interpretation of results was published in 2002 [40]. The  
528 CPAC-SPE-HPLC method for detecting trace ABX in milk was validated, with  
529 reference to the above Decision, by investigating LOD, LOQ, LR/R<sup>2</sup> (correlation  
530 coefficient), reproducibility and intraday/interday precision. Based on the MRLs of

531 SMM, OTC, CEF and MAR regulated by European Commission [1], LOD and LOQ  
 532 value for the model ABX were set as 0.020 µg/mL. Therefore, 80 mL of 0.020 µg/mL  
 533 medicated milk containing 10 mM EDTA were preconcentrated and cleaned up via  
 534 adsorption with 10 mg-CPAC columns, elution with 2 mL of 30% NH<sub>4</sub>OH/EtOH (1/19  
 535 v/v)), and purging with N<sub>2</sub> gas to prepare 0.5 mL of HPLC samples. Hence, the trace  
 536 ABX was concentrated 160-fold in the HPLC samples, and the *REs* of SMM, OTC,  
 537 CEF and MAR reached 99.48%, 99.67%, 69.13% and 61.93%, respectively.

538 In this range of 0.02-2.00 µg/mL of ABX, the linear correlation of ABX  
 539 concentrations in medicated milk and HPLC samples was investigated using CPAC-  
 540 SPE. The high *R*<sup>2</sup> values (0.993-0.998) for the four model ABX indicate that the linear  
 541 correlation between ABX concentrations in medicated milk and HPLC samples is  
 542 superior. The reproducibility of CPAC-SPE-HPLC was investigated in six repeated  
 543 tests. Reproducibility is expressed as the relative standard deviation (RSD, %) and  
 544 shown in italics in parentheses in Table 3.

545 Table 3. Mean *REs* (%) (n = 6) of ABX in milk using CPAC-SPE-HPLC, and reproducibility (RSD, %).

<i>C</i> <sub>ABX</sub> (µg/mL)	<i>SMM</i>	<i>OTC</i>	<i>CEF</i>	<i>MAR</i>
0.02	99.70 (6.3)	99.93 (6.5)	69.69 (4.0)	61.69 (2.4)
0.10	99.61 (2.4)	99.61 (6.4)	69.28 (3.2)	61.72 (2.0)
0.50	99.48 (4.0)	99.76 (3.1)	68.84 (2.2)	60.58 (2.2)
1.00	99.66 (5.1)	99.23 (3.2)	68.16 (3.7)	61.02 (1.2)

546 Note: *C*<sub>ABX</sub>, initial concentration of ABX in milk.

547

548 As listed in Table 3, the RSDs are in the range of 2.4%-6.3%, 3.1%-6.5%, 2.2%-  
 549 4.0%, and 1.2%-2.4% for SMM, OTC, CEF and MAR, respectively, demonstrating that  
 550 CPAC-SPE-HPLC has good determination reproducibility.

551 Furthermore, the precision of this method was investigated in terms of intraday and  
552 interday assays. The intraday/interday precision of CPAC-SPE-HPLC was evaluated  
553 using 0.1  $\mu\text{g/mL}$  and 1.0  $\mu\text{g/mL}$  medicated milk, which was consecutively tested five  
554 times in the same day ( $n = 5$ ) and also over three different days ( $n = 15$ ). For 0.1  $\mu\text{g/mL}$   
555 ABX in milk, the intraday RSD values obtained ranged from 1.2% to 6.4%, whereas  
556 the interday RSD values were in the range of 5.5–8.3%. For 1.0  $\mu\text{g/mL}$  ABX in milk,  
557 the intraday RSD values obtained ranged from 0.5% to 5.1%, while the interday RSDs  
558 were slightly higher, ranging from 4.5% to 7.8%.

### 559 3.2.3. Comparison with previous analytical methods

560 The developed CPAC-SPE-HPLC method was compared with previous SPE  
561 methods for the determination of trace SMM, OTC, CEF and MAR in milk. The  
562 sample-preparation procedure, amount of adsorbent, the volume of eluent, analytical  
563 instrument, *REs*, LOD and LOQ of each method are summarized in [Table S3](#) and  
564 discussed in [Section S2.0](#).

565 As seen in [Table S3](#), the higher *RE* of trace SMM and OTC in milk was achieved  
566 in this work over the previous SPE methods. The LOD and LOQ values for OTC (0.02  
567  $\mu\text{g/mL}$ ) are comparable with the previous data, but the previous methods possessed  
568 lower LOD and LOQ for SMM. More importantly, the LOD and LOQ of SMM and  
569 OTC (0.02  $\mu\text{g/mL}$ ) obtained in this work were lower than their MRLs (0.10  $\mu\text{g/mL}$ ). In  
570 comparison, the *REs* of CEF and MAR obtained in this work are lower than those  
571 reported in previous studies. Due to using higher amounts of adsorbents (200-500 mg),  
572 the previous LOD or/and LOQ of CEF and MAR are significantly lower. In contrast,

573 the LOD and LOQ of CEF and MAR (0.02  $\mu\text{g/mL}$ ) obtained with 10-15 mg CPAC in  
574 this work are still lower than the MRL (0.10 or 0.075  $\mu\text{g/mL}$ ).

575 Overall, the simplified sample-preparation procedure for the analysis of trace ABX  
576 in milk not only favors increasing *REs*, but also reduces the consumption of solvents  
577 and other materials, time and energy. AC-SPE-HPLC possesses inherent advantages,  
578 including the wide availability of powder AC sources, low consumption and costs,  
579 convenient operation, reliability and superior reproducibility, etc. Besides, the AC-  
580 SPE-HPLC method can be widely applied and easily verified and any commercial SPE  
581 cartridges are not required.

#### 582 **4. Conclusions**

583 A simple, accurate, reliable, and miniaturized CPAC-SPE-HPLC method has been  
584 successfully developed for the effective preconcentration, cleanup and determination  
585 of SMM, OTC, CEF and MAR in milk. Superior *REs* (>99%) of the model ABX in  
586 water were obtained in preliminary study. Subsequently, the conditions were further  
587 improved to obtain higher *REs* of the four model ABX in milk. 10 mg of CPAC and 2  
588 mL of 30%  $\text{NH}_4\text{OH}/\text{EtOH}$  (1/19 v/v) were scanned as the appropriate adsorbent and  
589 eluent. The mean *REs* with EDTA of SMM, OTC, CEF and MAR at different  
590 concentrations are 99.1%, 99.3%, 68.9% and 61.4%, respectively. The AC-SPE-HPLC  
591 method exhibits a good inter/intraday precision. To further improve AC-SPE-HPLC  
592 analysis of ABX in milk and expand the range of applications (water and food), in  
593 future, more species of ACs, such as ACs derived from different precursors, modified  
594 ACs, magnetic ACs and carbon composites, should be applied and evaluated as

595 adsorbents for SPE pretreatment.

596

597



## 598 **Acknowledgement**

599 Liu P. acknowledges the support of the China Scholarship Council (grant No.  
600 201909505008). Authors are grateful to Dr. Federica Calsolaro, Dr. Fabio Buccioli and  
601 Dr. Giorgio Grillo for technical assistance.

602

## 603 **Author Information**

### 604 **Corresponding Author**

605 **Giancarlo Cravotto** - Department of Drug Science and Technology, University of  
606 Turin, via Pietro Giuria 9, Turin, 10125, Italy.

607 Email: [giancarlo.cravotto@unito.it](mailto:giancarlo.cravotto@unito.it) Tel: +39.011.670.7183, Fax: +39.011.670.7162.

608

### 609 **Authors**

610 **Pengyun Liu, Zhilin Wu, Alessandro Barge, Luisa Boffa and Katia Martina** -

611 Department of Drug Science and Technology, University of Turin, via Pietro Giuria 9,  
612 Turin, 10125, Italy.

## 613 **Funding**

614 This research was supported by the Fondazione CRT “*Sviluppo di tecnologie*  
615 *integrate per l'eliminazione dei residui di antibiotici dal latte vaccino*”, by the Ministry  
616 of Science and Higher Education of the Russian Federation: World-Class Research  
617 Center, Sechenov First Moscow State Medical University and also by the China  
618 Scholarship Council.

## 619 **Declaration of interest**

620 The authors declare that they have no known competing financial interests or  
621 personal relationships that could have appeared to influence the work reported in this  
622 paper.

623

## 624 References

- 625 [1] Commission, E. (2010). Commission Regulation (EU) No 37/2010 of 22  
626 December 2009 on pharmacologically active substances and their classification  
627 regarding maximum residue limits in foodstuffs of animal origin. *OffJ Eur Union*,  
628 *15*, 1–72.
- 629 [2] Khatibi, S.A., Hamidi, S., & Siahi-Shadbad, M.R. (2021). Current trends in  
630 sample preparation by solid-phase extraction techniques for the determination of  
631 antibiotic residues in foodstuffs: a review. *Critical Reviews in Food Science and*  
632 *Nutrition*, *61*, 3361-3382. <https://doi.org/10.1080/10408398.2020.1798349>.
- 633 [3] Kantiani, L., Farré, M., & Barceló, D. (2009). Analytical methodologies for the  
634 detection of  $\beta$ -lactam antibiotics in milk and feed samples. *TrAC Trends in*  
635 *Analytical Chemistry*, *28*, 729–744. <https://doi.org/10.1016/j.trac.2009.04.005>.
- 636 [4] Greño, M., Castro-Puyana, M., Garcia, M.Á., & Marina, M.L. (2018). Analysis  
637 of antibiotics by CE and CEC and their use as chiral selectors: An update.  
638 *Electrophoresis*, *39*, 235–259. <https://doi.org/10.1002/elps.201700306>.
- 639 [5] Rossi, R., Saluti, G., Moretti, S., Diamanti, I., Giusepponi, D., & Galarini, R.  
640 (2018). Multiclass methods for the analysis of antibiotic residues in milk by liquid  
641 chromatography coupled to mass spectrometry: a review. *Food Additives &*  
642 *Contaminants Part A*, *35*, 241–257.  
643 <https://doi.org/10.1080/19440049.2017.1393107>.
- 644 [6] Koesukwiwat, U., Jayanta, S., & Leepipatpiboon, N. (2007). Solid-phase  
645 extraction for multiresidue determination of sulfonamides, tetracyclines, and  
646 pyrimethamine in Bovine's milk. *Journal of Chromatography A*, *1149*, 102–111.  
647 <https://doi.org/10.1016/j.chroma.2007.02.075>.
- 648 [7] Narola, B., Singh, A., Mitra, M., Santhakumar, P., & Chandrashekhar, T. (2011).  
649 A validated reverse phase HPLC method for the determination of disodium EDTA  
650 in meropenem drug substance with UV-detection using precolumn derivatization  
651 technique. *Analytical chemistry Insights*, *6*, ACI-S5953.  
652 <https://doi.org/10.4137%2FACI.S5953>.
- 653 [8] Xu, J.-J., An, M., Yang, R., Tan, Z., Hao, J., Cao, J., Peng, L.-Q., & Cao, W.  
654 (2016). Determination of tetracycline antibiotic residues in honey and milk by  
655 miniaturized solid phase extraction using chitosan-modified graphitized  
656 multiwalled carbon nanotubes. *Journal of agricultural and Food chemistry*, *64*,  
657 2647–2654. <https://doi.org/10.1021/acs.jafc.6b00748>.
- 658 [9] Kishida, K. (2011). Simplified extraction of tetracycline antibiotics from milk  
659 using a centrifugal ultrafiltration device. *Food chemistry*, *126*, 687–690.  
660 <https://doi.org/10.1016/j.foodchem.2010.11.021>.
- 661 [10] Horton, R., Randall, L., Bailey-Horne, V., Heinrich, K., Sharman, M., Brunton,  
662 L., La Ragione, R., & Jones, J. (2015). Degradation of cefquinome in spiked milk  
663 as a model for bioremediation of dairy farm waste milk containing cephalosporin  
664 residues. *Journal of applied microbiology*, *118*, 901–910.  
665 <https://doi.org/10.1111/jam.12765>.

- 666 [11] Lv, Y.-K., Wang, L.-M., Yang, L., Zhao, C.-X., & Sun, H.-W. (2012). Synthesis  
667 and application of molecularly imprinted poly (methacrylic acid)–silica hybrid  
668 composite material for selective solid-phase extraction and high-performance  
669 liquid chromatography determination of oxytetracycline residues in milk. *Journal*  
670 *of Chromatography A*, 1227, 48–53.  
671 <https://doi.org/10.1016/j.chroma.2011.12.108>.
- 672 [12] Bitas, D., & Samanidou, V. (2018). Molecularly imprinted polymers as extracting  
673 media for the chromatographic determination of antibiotics in milk. *Molecules*,  
674 23, 316. <https://doi.org/10.3390/molecules23020316>.
- 675 [13] Ge, X., Wu, Z., M., Manzoli, Wu, Z., & Cravotto, G. (2020). Feasibility and the  
676 mechanism of desorption of phenolic compounds from activated carbons.  
677 *Industrial & Engineering Chemistry Research*, 59, 12223–12231.  
678 <https://doi.org/10.1021/acs.iecr.0c01402>.
- 679 [14] Wu, Z., Liu, P., Wu, Z., & Cravotto, G. (2021). In situ Modification of Activated  
680 Carbons by Oleic Acid under Microwave Heating to Improve Adsorptive  
681 Removal of Naphthalene in Aqueous Solutions. *Processes*, 9, 391.  
682 <https://doi.org/10.3390/pr9020391>.
- 683 [15] Ge, X., Wu, Z., Manzoli, M., Bonelli, B., Mantegna, S., Kunz, W., & Cravotto,  
684 G. (2021). Adsorptive decontamination of antibiotic-spiked water and milk using  
685 commercial and modified activated carbons. *Journal of Environmental Chemical*  
686 *Engineering*, 9, 105544. <https://doi.org/10.1016/j.jece.2021.105544>.
- 687 [16] Dmitrienko, S.G., Kochuk, E.V., Tolmacheva, V.V., Apyari, V.A., & Zolotov, Y.A.  
688 (2015). Determination of the total content of some sulfonamides in milk using  
689 solid-phase extraction coupled with off-line derivatization and  
690 spectrophotometric detection. *Food chemistry*, 188, 51–56.  
691 <https://doi.org/10.1016/j.foodchem.2015.04.123>.
- 692 [17] Li, L., Quinlivan, P.A., & Knappe, D.R. (2002). Effects of activated carbon surface  
693 chemistry and pore structure on the adsorption of organic contaminants from  
694 aqueous solution. *Carbon*, 40, 2085–2100. [https://doi.org/10.1016/S0008-](https://doi.org/10.1016/S0008-6223(02)00069-6)  
695 [6223\(02\)00069-6](https://doi.org/10.1016/S0008-6223(02)00069-6).
- 696 [18] Hubetska, T., Kobylinska, N., & Garcia, J.R. (2020). Efficient adsorption of  
697 pharmaceutical drugs from aqueous solution using a mesoporous activated carbon.  
698 *Adsorption*, 26, 251–266. <https://doi.org/10.1007/s10450-019-00143-0>.
- 699 [19] Bruno, F., Curini, R., Corcia, A.D., Nazzari, M., Pallagrosi, M. (2002). An  
700 original approach to determining traces of tetracycline antibiotics in milk and  
701 eggs by solid-phase extraction and liquid chromatography/mass spectrometry.  
702 *Rapid Communications Mass Spectrometry*, 16, 1365–1376.  
703 <https://doi.org/10.1002/rcm.724>.
- 704 [20] Chemat, F., Vian, M.A., & Cravotto, G. (2012). Green extraction of natural  
705 products: concept and principles. *International journal of molecular sciences*, 13,  
706 8615–8627. <https://doi.org/10.3390/ijms13078615>.
- 707 [21] Cayam chemical company. (2018). Product information. Retrieved from  
708 <https://www.caymanchem.com/pdfs/24174.pdf>. Accessed 2021, December 21.
- 709 [22] Gibbons, P.M. (2014). TOPICS IN MEDICINE AND SURGERY. *Journal of*

- 710 *Exotic Pet Medicine*, 23, 21–38. <http://dx.doi.org/10.1053/j.jepm.2013.11.007>.
- 711 [23] Zamora, F., Sabio, E., Román, S., González-García, C.M., & Ledesma, B. (2010).  
712 Modelling the adsorption of p-Nitrophenol by the Boyd method in conjunction  
713 with the finite element method. *Adsorption Science & Technology*, 28, 671–687.  
714 <https://doi.org/10.1260/2F0263-6174.28.8-9.671>.
- 715 [24] Tamon, H., Saito, T., Kishimura, M., Okazaki, M., & Toei, R. (1990). Solvent  
716 regeneration of spent activated carbon in wastewater treatment. *Journal of*  
717 *chemical engineering of Japan*, 23, 426–432. <https://doi.org/10.1252/jcej.23.426>.
- 718 [25] Seyhan Bozkurt, S., Erdogan, D., Antep, M., Tuzmen, N., & Merdivan, M. (2016).  
719 Use of ionic liquid based chitosan as sorbent for preconcentration of  
720 fluoroquinolones in milk, egg, fish, bovine, and chicken meat samples by solid  
721 phase extraction prior to HPLC determination. *Journal of Liquid Chromatography*  
722 *& Related Technologies*, 39, 21–29.  
723 <https://doi.org/10.1080/10826076.2015.1116010>.
- 724 [26] El-Shahat, M., Burham, N., & Azeem, S.A. (2010). Flow injection analysis–solid  
725 phase extraction (FIA–SPE) method for preconcentration and determination of  
726 trace amounts of penicillins using methylene blue grafted polyurethane foam.  
727 *Journal of hazardous materials*, 177, 1054–1060.  
728 <https://doi.org/10.1016/j.jhazmat.2010.01.027>.
- 729 [27] Ji, L., Liu, F., Xu, Z., Zheng, S., & Zhu, D. (2010). Adsorption of pharmaceutical  
730 antibiotics on template-synthesized ordered micro-and mesoporous carbons.  
731 *Environmental science & technology*, 44, 3116–3122.  
732 <https://doi.org/10.1021/es903716s>.
- 733 [28] Min, P. (2020). Biochar adsorption of antibiotics and its implications to  
734 remediation of contaminated soil. *Water, Air, & Soil Pollution*, 23, 11–15.  
735 <https://doi.org/10.1007/s11270-020-04551-9>.
- 736 [29] Ji, L., Chen, W., Zheng, S., Xu, Z., & Zhu, D. (2009). Adsorption of sulfonamide  
737 antibiotics to multiwalled carbon nanotubes. *Langmuir*, 25, 11608–11613.  
738 <https://doi.org/10.1021/la9015838>.
- 739 [30] Ahmed, M.B., Zhou, J.L., Ngo, H.H., & Guo, W. (2015). Adsorptive removal of  
740 antibiotics from water and wastewater: progress and challenges. *Science of the*  
741 *Total Environment*, 532, 112–126. <https://doi.org/10.1016/j.scitotenv.2015.05.130>.
- 742 [31] Li, C., Zhu, X., He, H., Fang, Y., Dong, H., Lü, J., Li, J., & Li, Y. (2019).  
743 Adsorption of two antibiotics on biochar prepared in air-containing atmosphere:  
744 influence of biochar porosity and molecular size of antibiotics. *Journal of*  
745 *Molecular Liquids*, 274, 353–361. <https://doi.org/10.1016/j.molliq.2018.10.142>.
- 746 [32] Ji, L., Wan, Y., Zheng, S., & Zhu, D. (2011). Adsorption of tetracycline and  
747 sulfamethoxazole on crop residue-derived ashes: implication for the relative  
748 importance of black carbon to soil sorption. *Environmental Science & Technology*,  
749 45, 5580–5586. <https://doi.org/10.1021/es200483b>.
- 750 [33] Xiang, Y., Xu, Z., Wei, Y., Zhou, Y., Yang, X., Yang, Y., Yang, J., Zhang, J., Luo,  
751 L., & Zhou, Z. (2019). Carbon-based materials as adsorbent for antibiotics  
752 removal: mechanisms and influencing factors. *Journal of environmental*  
753 *management*, 237, 128–138. <https://doi.org/10.1016/j.jenvman.2019.02.068>.

- 754 [34] Li, M., Zhao, Z., Wu, X., Zhou, W., & Zhu, L. (2017). Impact of mineral  
755 components in cow manure biochars on the adsorption and competitive adsorption  
756 of oxytetracycline and carbaryl. *RSC advances*, 7, 2127–2136.  
757 <https://doi.org/10.1039/C6RA26534K>.
- 758 [35] Gao, M., Zhang, Y., Gong, X., Song, Z., & Guo, Z. (2018). Removal mechanism  
759 of di-n-butyl phthalate and oxytetracycline from aqueous solutions by nano-  
760 manganese dioxide modified biochar. *Environmental Science and Pollution*  
761 *Research*, 25, 7796–7807. <https://doi.org/10.1007/s11356-017-1089-5>.
- 762 [36] Ma, X., & Agarwal, S. (2016). Adsorption of emerging ionizable contaminants on  
763 carbon nanotubes: advancements and challenges. *Molecules*, 21, 628.  
764 <https://doi.org/10.3390/molecules21050628>.
- 765 [37] Huang, S.-C., Chung, T.-W., & Wu, H.-T. (2021). Effects of Molecular Properties  
766 on Adsorption of Six-Carbon VOCs by Activated Carbon in a Fixed Adsorber.  
767 *ACS omega*, 6, 5825–5835. <https://doi.org/10.1021/acsomega.0c06260>.
- 768 [38] Franz, M., Arafat, H.A., & Pinto, N.G. (2000). Effect of chemical surface  
769 heterogeneity on the adsorption mechanism of dissolved aromatics on activated  
770 carbon. *Carbon*, 38, 1807–1819. [https://doi.org/10.1016/S0008-6223\(00\)00012-9](https://doi.org/10.1016/S0008-6223(00)00012-9).
- 771 [39] Mezzari, I.A. (2006). Predicting the adsorption capacity of activated carbon for  
772 organic contaminants from fundamental adsorbent and adsorbate properties.  
773 (Master thesis) North Carolina State University, Raleigh.
- 774 [40] Commission, E. (2002). Commission Decision 2002/657/EC of 12 August 2002  
775 implementing Council Directive 96/23/EC concerning the performance of  
776 analytical methods and the interpretation of results. *Official Journal of European*  
777 *Communities*, 50, 8–36.

778

# Determination of trace antibiotics in water and milk via preconcentration and cleanup using activated carbon

Pengyun Liu<sup>1</sup>, Zhilin Wu<sup>1</sup>, Alessandro Barge<sup>1</sup>, Luisa Boffa<sup>1</sup>, Katia Martina<sup>1</sup>,  
Giancarlo Cravotto<sup>1,2\*</sup>

<sup>1</sup> Department of Drug Science and Technology, University of Turin, via P. Giuria 9,  
Turin, 10125, Italy.

<sup>2</sup> World-Class Research Center "Digital biodesign and personalized healthcare",  
Sechenov First Moscow State Medical University, Moscow, 119991 Russia.

\*Correspondence: giancarlo.cravotto@unito.it (G. Cravotto), Tel: +39.011.670.7183,  
Fax: +39.011.670.7162.

**Number of pages: 10**

**Number of figure: 1**

**Number of tables: 3**

---

Abbreviations: ABX, antibiotics; SMM, sulfamonomethoxine sodium; OTC, oxytetracycline; CEF, ceftiofur hydrochloride; MAR, marbofloxacin; MRLs, maximum residue limits; HPLC, high pressure liquid chromatography; SPE, solid phase extraction; RE, recovery efficiency; ACs, activated carbons; CPAC, coconut powdered AC;  $S_{\text{BET}}$ , Brunauer-Emmett-Teller surface area;  $V_{\text{Meso}}$ , mesopore volume;  $V_{\text{Micro}}$ , micropore volume; PGAC, peat granular AC; WPAC, wood powder AC; MeCN, acetonitrile; TFA, trifluoroacetic acid; MeOH, methanol; EtOH, ethanol;  $\text{NH}_4\text{OH}$ , ammonia solution; EDTA, ethylenediaminetetraacetic acid disodium salt hydrate; t-BuOH, t-butanol; n-PrOH, n-propanol; MW, molecular weight; MV, molar volume;  $\text{Log}K_{\text{OW}}$ , octanol-water partition coefficient;  $S_{\text{Water}}$ , solubility of ABX in water;  $S_{\text{EtOH}}$ , saturated mole fraction solubility of ABX in EtOH;  $\text{p}K_{\text{a}}$ , dissociation constant; NHA, number of H-bond acceptors; NHD, number of H-bond donors; Refs., references; *Bp*, boiling point;  $\epsilon$ , dielectric constant; EDA,  $\pi$ - $\pi$  electron donor-acceptor; TC, tetracycline; LODs, limit of detections; LOQs, limit of quantitations; LRs, linear ranges; RSD, relative standard deviation; FA, formic acid; DCM, methylene chloride; MSPD, magnetic solid phase dispersion; MSPE, magnetic solid phase extraction.

## **S1.0 The information of solvents and water**

Acetonitrile (MeCN,  $\geq 99.9\%$ ), trifluoroacetic acid (TFA,  $\geq 99\%$ ) and methanol (MeOH, 99.9%) were obtained from Sigma-Aldrich (France). Ethanol (EtOH,  $\geq 99.8\%$ ) was provided by Sigma-Aldrich (UK). Acetic acid (HOAc, 100%) was purchased from VWR International (European Commission). The ammonia solution (30%  $\text{NH}_4\text{OH}$ ) was provided by CARLO ERBA Reagents S.r.l. (Italy). Ethylenediaminetetraacetic acid disodium salt hydrate (EDTA,  $>99\%$ ), n-butanol (n-BuOH,  $>98\%$ ), t-butanol (t-BuOH, 99%), and n-propanol (n-PrOH, 99.8%) were purchased from Alfa Aesar (Thermo Fisher Scientific, Germany).

Milli-Q water was obtained from a Milli-Q Reference A+System (Merck Millipore, Darmstadt, Germany) and used for the preparation of HPLC mobile phases. Deionized water (conductivity  $\leq 2 \mu\text{S}/\text{cm}$ ) was used to prepare standard solutions of ABX, which were refrigerated at 4 °C.

## S2.0 Comparison with previous analytical methods


All of the samples prepared using SPE have undergone analysis with HPLC, LC-MS or -MS/MS, or UPLC-MS in the past. The present protocol exhibited higher *RE* than the previous SPE methods for the analysis of trace SMM in milk [45,46], as can be seen in Table S3. The previous methods possessed lower LOD and LOQ since high amounts of adsorbents (80-500 mg) were used. In this work, only 10 mg of CPAC was used as the adsorbent and the LOD and LOQ values of SMM (0.02 µg/mL) are also lower than the MRL (0.10 µg/mL). In addition, toxic solvents, such as DCM, MeCN and MeOH, and chemical deproteinization were not required in this study. Similarly, we achieved a higher *RE* for OTC here than the two previous SPE methods [11,47]. The LOD and LOQ values for OTC (0.02 µg/mL) are comparable with the previous data and are also lower than the MRL (0.10 µg/mL), although less CPAC (10 mg) was used as an adsorbent here, as 50 mg of adsorbent was used in the previous method. In addition, chemical deproteinization and redissolution were not required in this study. In terms of toxicity, the use of EtOH as an eluent in this study is better than the MeOH that was used in other studies.

The *RE* of CEF (68.9%) that was obtained using a 10 mg-CPAC column in this work is comparable with that (69.0%) obtained using a 500 mg-Carbograph 4 cartridge-SPE in a previous study [48]. However, the previously reported LOQ is significantly lower than the one in that study since very different amounts of adsorbents were used. Compared with the *RE* (97.0%) obtained in another study using 200 mg-Oasis HLB-MSPD, the *RE* of CEF in this work is significantly lower, but the LOD and LOQ of



CEF (0.02  $\mu\text{g/mL}$ ) are comparable with the previous data, and are also lower than the MRL (0.10  $\mu\text{g/mL}$ ), although less CPAC (10 mg) was used as the adsorbent [49]. A further benefit is that toxic solvents, such as DCM, MeCN and MeOH, and the redissolution of analytes are not required in this study. The *RE* of MAR (61.4%) that was obtained using a 10 mg-CPAC column in this work is considerably lower than those (82.5%-106.5%) obtained in three previous studies. Moreover, the previous methods had lower LOD and LOQ, but 0.020  $\mu\text{g/mL}$  of LOD or LOQ in this work is still sufficient to meet the requirements for the MRL of MAR (0.075  $\mu\text{g/mL}$ ). However, a remarkably lower amount of adsorbent (10 mg of CPAC) was used in this study than the 200-500 mg of adsorbents used in the other studies [25,50,51]. Furthermore, the toxic solvent MeOH and chemical deproteinization were not required in this study.

**Figure S1**



**Fig. S1.** Comparison of HPLCs from various samples. (a) sample from blank milk, (b) sample recovered from medicated milk by using 10 mg of CPAC for adsorption and 2 mL of 30%  $\text{NH}_4\text{OH}/\text{EtOH}$  (1/19 v/v) for elution, (c) standard ABX aqueous solutions (50  $\mu\text{g}/\text{mL}$  MAR, 100  $\mu\text{g}/\text{mL}$  CEF, OTC or SMM).

## Table S1

Table S1. Major physicochemical properties and toxicological parameters of the model ABX (Part of the data was adopted from the Pubchem and Drugbank databases).

<i>ABX</i>	<i>CAS</i>	<i>MW</i> (g/mol)	<i>MV</i> (cm <sup>3</sup> /mol)	<i>LogK<sub>ow</sub></i>	<i>S<sub>water</sub></i> (mg/L)	<i>S<sub>EtOH</sub></i> (10 <sup>3</sup> )	<i>pK<sub>a</sub></i>	<i>NHA</i>	<i>NHD</i>	<i>MRLs</i> (μg/L)	<i>LD<sub>50</sub></i> (mg/kg)	<i>Refs.</i>
SMM	38006-08-5	302	213.3	-0.8	10000	-	6.33	8	2	100	5620	[15,41]
											Rat oral	
OTC	2058-46-0	460	270.3	-0.9	300	<sup>a</sup> 12.0	3.27	11	8	100	>2000	[15,42,43]
											Colinus virginianus oral	
CEF	103980-44-5	560	290.9	1.2	100	23.3	2.83	12	4	100	-	[15]
MAR	115550-35-1	362	226.4	-0.5	2600	116.6	5.38	8	1	75	>0.002	[15,44]
											Mouse oral	

<sup>a</sup>The unit is mg/L; <sup>b</sup> the unit is mg/L/4 hr.

**Note:** *MW*, molecular weight; *MV*, molar volume; *LogK<sub>ow</sub>*, Octanol-water partition coefficient; *S<sub>water</sub>*, solubility in water; *S<sub>EtOH</sub>*, saturated mole fraction solubility of ABX in EtOH, calculated according to the literature [44]; *pK<sub>a</sub>*, dissociation constant; *NHA*, number of H-bond acceptors; *NHD*, number of H-bond donors; *MRLs*, maximum residue limits; *LD<sub>50</sub>*, amount of a material, given all at once, which causes the death of 50% (one half) of a group of test animals; *Refs.*, references.

## Table S2

Table S2. The HPLC conditions for the determination of the model ABX.

<i>ABX</i>	<i>Wavelength</i> ( <i>nm</i> )	<i>Retention time</i> ( <i>min</i> )	<i>Running time</i> ( <i>min</i> )	<i>Mobile phase</i>	
				<i>Phase A (%)</i>	<i>Phase B (%)</i>
				<i>0.1%TFA in H<sub>2</sub>O</i>	<i>0.1%TFA in MeCN</i>
SMM	273	25.483	0	99	1
			25	0	100
			32	0	100
OTC	267	13.313	0	98	2
			25	0	100
			26	0	100
CEF	265	12.510	0	98	2
			5	70	30
			20	0	100
			26	0	100
MAR	298	14.018	0	98	2
			25	0	100
			26	0	100

**Table S3**

**Table S3.** Comparison of the proposed CPAC-SPE-HPLC method with previous SPE methods for the determination of the model ABX in milk.

<i>ABX</i>	<i>Sample-preparation procedures</i>	<i>Eluent</i>	<i>Analytical instrument</i>	<i>RE (%)</i>	<i>LOD (µg/mL)</i>	<i>LOQ (µg/mL)</i>	<i>Refs.</i>
SMM	Agitation, 500 mg-carbograph 4 cartridge-SPE, redissolution, filtration.	1.5 mL MeOH- 6 mL DCM/ MeOH (80/20 v/v) acidified with 10 mM TFA continuously	LC-ES -MS	97.3	<sup>a</sup> 0.001	<sup>a</sup> 0.003	[45]
	Deproteinization, 80 mg-magnetic graphene oxide nanocomposite -MSPE, evaporation, redissolution.	1.0 mL MeCN containing 5 % NH <sub>4</sub> OH (v/v)	HPLC	86.8	0.00008	0.00025	[46]
	10 mg-CPAC-MSPE, concentration by N <sub>2</sub> , centrifugation.	2 mL 30% NH <sub>4</sub> OH/EtOH (1/19 v/v)	HPLC	<sup>b</sup> 99.1	0.02	0.02	This study
OTC	Deproteinization, desolvation, redissolution, filtration, 50 mg-molecularly imprinted poly (methacrylic acid) –silica hybrid composite -SPE.	2 mL MeOH/ HOAc (60/40 v/v)	LC-MS/MS	<sup>b</sup> 99.2	0.014	-	[11]
	Deproteinization, C <sub>18</sub> -SPE, constant volume.	0.75 mL MeOH	UPLC- MS/MS	<sup>b</sup> 98.8	-	-	[47]
	10 mg CPAC-SPE, concentration by N <sub>2</sub> , centrifugation.	2 mL 30% NH <sub>4</sub> OH/EtOH (1/19 v/v)	HPLC	<sup>b</sup> 99.3	0.02	0.02	This study
CEF	500 mg-carbograph 4 cartridge-SPE, desolvation by N <sub>2</sub> , redissolution, filtration.	6 mL DCM/ MeOH (80/20 v/v) acidified with 50 mM FA	HPLC	69.0	-	0.001	[48]
	200 mg-Oasis HLB- MSPD, evaporation to dryness, redissolution.	1 mL MeOH –2 mL MeCN continuously	HPLC	97.0	<sup>a</sup> 0.0118	<sup>a</sup> 0.0357	[49]
	10 mg-CPAC-SPE, concentration by N <sub>2</sub> , centrifugation.	2 mL 30% NH <sub>4</sub> OH/EtOH (1/19 v/v)	HPLC	<sup>b</sup> 68.9	0.02	0.02	This study
MAR	Centrifugation, PLRP-cartridges-SPE.	9 mL H <sub>2</sub> O/MeOH ( 80/20 v/v) plus 0.5% FA (1.5 mL)	HPLC	106.5	0.0001	-	[50]
	Deproteinization, filtration, redissolution, centrifugation, 500 mg-Oasis HLB cartridges-SPE, evaporation to dryness, redissolution.	10 mL MeOH containing 1.5% (w/v) HOAc	LC-MS	<sup>b</sup> 100.5	0.00009	0.00031	[51]
	Deproteinization, 200 mg-ionic liquid based chitosan-SPE, desolvation by N <sub>2</sub> , constant volume.	5 mL 20% NH <sub>3</sub> (v/v MeOH)	LC-MS/MS	82.5	0.00423	-	[25]
	10 mg-CPAC-SPE, concentration by N <sub>2</sub> , centrifugation.	2 mL 30% NH <sub>4</sub> OH/EtOH (1/19 v/v)	HPLC	<sup>b</sup> 61.4	0.02	0.02	This study

<sup>a</sup> The unit: ppm; <sup>b</sup> with EDTA in medicated milk.


**Note:** *RE*, recovery efficiency; *LODs*, limit of detections; *LOQs*, limit of quantitations; *Refs*, references; *FA*, Formic acid; *DCM*, Methylene chloride; *MSPD*, Magnetic solid phase dispersion; *MSPE*, Magnetic solid phase extraction.

## References

- [11] Lv, Y.-K., Wang, L.-M., Yang, L., Zhao, C.-X., & Sun, H.-W. (2012). Synthesis and application of molecularly imprinted poly (methacrylic acid)–silica hybrid composite material for selective solid-phase extraction and high-performance liquid chromatography determination of oxytetracycline residues in milk. *Journal of Chromatography A*, *1227*, 48–53. <https://doi.org/10.1016/j.chroma.2011.12.108>.
- [15] Ge, X., Wu, Z., Manzoli, M., Bonelli, B., Mantegna, S., Kunz, W., & Cravotto, G. (2021). Adsorptive decontamination of antibiotic-spiked water and milk using commercial and modified activated carbons. *Journal of Environmental Chemical Engineering*, *9*, 105544. <https://doi.org/10.1016/j.jece.2021.105544>.
- [25] Seyhan Bozkurt, S., Erdogan, D., Antep, M., Tuzmen, N., & Merdivan, M. (2016). Use of ionic liquid based chitosan as sorbent for preconcentration of fluoroquinolones in milk, egg, fish, bovine, and chicken meat samples by solid phase extraction prior to HPLC determination. *Journal of Liquid Chromatography & Related Technologies*, *39*, 21–29. <https://doi.org/10.1080/10826076.2015.1116010>.
- [41] Fu, L., Huang, T., Wang, S., Wang, X., Su, L., Li, C., & Zhao, Y. (2017). Toxicity of 13 different antibiotics towards freshwater green algae *Pseudokirchneriella subcapitata* and their modes of action. *Chemosphere*, *168*, 217–222. <https://doi.org/10.1016/j.chemosphere.2016.10.043>.
- [42] O’Neil, M.J., Smith, A., Heckelman, P.E., & Budavari, S. (2001). The merck index-An encyclopedia of chemicals, drugs, and biologicals. whitehouse station. NJ: Merck and Co, Inc., 767, 4342.
- [43] Yalkowsky, S., & He, Y. (2003). An extensive compilation of aqueous solubility data for organic compounds extracted from the AQUASOL database. *Handbook Aqueous Solubility Data* (p 377). 377.
- [44] Wu, Z., Li, W., Yu, P., Fan, X., Sun, H., Zhao, H., & Zhang, Y. (2019). Measurement and Correlation of Solubility of Marbofloxacin in 12 Pure Solvents from 283.15 to 328.15 K. *Journal of Chemical & Engineering Data*, *64*, 5275–5281. <https://doi.org/10.1021/acs.jced.9b00490>.
- [45] Cavaliere, C., Curini, R., Di Corcia, A., Nazzari, M., & Samperi, R. (2003). A simple and sensitive liquid chromatography- mass spectrometry confirmatory method for analyzing sulfonamide antibacterials in milk and egg. *Journal of agricultural Food chemistry*, *51*, 558–566. <https://doi.org/10.1021/jf020834w>.
- [46] Wang, Y., Liu, L., Xiao, C., Chen, L., Yang, P., Liu, Q., Wang, J., & Liu, X. (2016). Rapid determination of trace sulfonamides in milk by graphene oxide-based magnetic solid phase extraction coupled with HPLC–MS/MS. *Food Analytical Methods*, *9*, 2521–2530. <https://doi.org/10.1007/s12161-016-0433-6>.
- [47] Mei-Ratliff, Y. (2012). Determination of the antibiotic oxytetracycline in commercial milk by solid-phase extraction: a high-performance liquid chromatography (HPLC) experiment for quantitative instrumental analysis. *Journal of Chemical Education*, *89*, 656–659. <https://doi.org/10.1021/ed900065y>.
- [48] Bruno, F., Curini, R., Corcia, A.D., Nazzari, M., & Samperi, R. (2001). Solid-

- phase extraction followed by liquid chromatography- mass spectrometry for trace determination of  $\beta$ -lactam antibiotics in bovine milk. *Journal of agricultural and Food chemistry*, 49, 3463–3470. <https://doi.org/10.1021/jf010046r>.
- [49] Karageorgou, E.G., Samanidou, V.F., & Papadoyannis, I.N. (2012). Ultrasound-assisted matrix solid phase dispersive extraction for the simultaneous analysis of  $\beta$ -lactams (four penicillins and eight cephalosporins) in milk by high performance liquid chromatography with photodiode array detection. *Journal of Separation Science*, 35, 2599–2607. <https://doi.org/10.1002/jssc.201200514>.
- [50] Kantiani, L., Farré, M., & Barceló, D. (2011). Rapid residue analysis of fluoroquinolones in raw bovine milk by online solid phase extraction followed by liquid chromatography coupled to tandem mass spectrometry. *Journal of Chromatography A*, 1218, 9019–9027. <https://doi.org/10.1016/j.chroma.2011.09.079>.
- [51] Herrera-Herrera, A.V., Hernández-Borges, J., Rodríguez-Delgado, M.A., Herrero, M., & Cifuentes, A. (2011). Determination of quinolone residues in infant and young children powdered milk combining solid-phase extraction and ultra-performance liquid chromatography–tandem mass spectrometry. *Journal of Chromatography A*, 1218, 7608–7614. <https://doi.org/10.1016/j.chroma.2011.05.066>.

### Graphical abstract



Simple preconcentration and cleanup of trace antibiotics in milk using activated carbon (AC)-based solid phase extraction for HPLC analysis.