



AperTO - Archivio Istituzionale Open Access dell'Università di Torino

# **The Lane-Emden System on Cartan-Hadamard Manifolds: Asymptotics and Rigidity of Radial Solutions**

Original Citation: Published version: DOI:10.1093/imrn/rnae079 Terms of use: Open Access Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law. Availability: **This is a pre print version of the following article:** This version is available http://hdl.handle.net/2318/2010950 since 2024-11-13T17:07:18Z

(Article begins on next page)

# THE LANE-EMDEN SYSTEM ON CARTAN-HADAMARD MANIFOLDS: ASYMPTOTICS AND RIGIDITY OF RADIAL SOLUTIONS

MATTEO MURATORI AND NICOLA SOAVE

Abstract. We investigate existence and qualitative properties of globally defined and positive radial solutions of the Lane-Emden system, posed on a Cartan-Hadamard model manifold  $\mathbb{M}^n$ . We prove that, for critical or supercritical exponents, there exists at least a one-parameter family of such solutions. Depending on the stochastic completeness or incompleteness of  $\mathbb{M}^n$ , we show that the existence region stays one dimensional in the former case, whereas it becomes two dimensional in the latter. Then, we study the asymptotics at infinity of solutions, which again exhibit a dichotomous behavior between the stochastically complete (where both components are forced to vanish) and incomplete cases. Finally, we prove a rigidity result for finite-energy solutions, showing that they exist if and only if  $\mathbb{M}^n$  is isometric to  $\mathbb{R}^n$ .

### 1. INTRODUCTION

The Lane-Emden equation

$$
-\Delta u = u^p, \quad u > 0,
$$
\n<sup>(1.1)</sup>

with  $p > 0$ , is the prototype of semilinear elliptic equations, and has played a central role in the development of several tools in the analysis of nonlinear PDEs. Its system counterpart, known in the literature as Lane-Emden system, that is

$$
\begin{cases}\n-\Delta u = v^q \\
-\Delta v = u^p \\
u, v > 0,\n\end{cases}
$$
\n(1.2)

with  $p, q > 0$ , has also received a lot of attention in the recent years, but is far less understood. The purpose of this paper is to study existence and qualitative properties of radial solutions to (1.2) posed on a *Cartan-Hadamard manifold*  $\mathbb{M}^n$ , in the *critical or supercritical regime* of the exponents (see below). Recall that a Cartan-Hadamard manifold is a complete and simply connected Riemannian manifold with nonpositive sectional curvature. An important feature of this type of manifolds consists in the possibility of writing global polar coordinates centered at any reference point  $o \in \mathbb{M}^n$ , as the well-known Cartan-Hadamard theorem entails that the exponential map at  $o$  is a diffeomorphism between  $\mathbb{R}^n$  and  $\mathbb{M}^n$ .

Thus, in order to understand existence and properties of solutions to (1.2) on Cartan-Hadamard manifolds, and having in mind the Euclidean case, it appears natural to focus on the radial problem at first, upon requiring in addition that the ambient space itself is spherically symmetric. In this framework, we establish the existence of positive radial solutions on any Cartan-Hadamard model manifold (we refer to Subsection 1.2 for definitions and details on the geometric background). As a second step, focusing on uniqueness or multiplicity, and on the limit behavior of such solutions, we discover an interesting dichotomy: if the underlying manifold  $\mathbb{M}^n$  is *stochastically complete*, then the scenario is Euclidean like, namely we prove a uniqueness result and show that all solutions are such that both  $u$  and  $v$  vanish at infinity (see Theorem 1.1); if instead  $\mathbb{M}^n$  is *stochastically incomplete*, then we show a new phenomenon of multiple existence of positive solutions with strictly positive limits at infinity (see Theorem 1.2). Finally,

<sup>2020</sup> Mathematics Subject Classification. Primary: 58J05; 35B53; 35J47; 35J91. Secondary: 58J70; 34A12; 34A34; 34C40; 34D05.

Key words and phrases. Cartan-Hadamard manifolds; Lane-Emden system; Radial solutions; Stochastic completeness.

we establish a strong rigidity result in terms of the natural energies involved when dealing with (1.2): either they are all finite, and in this case the underlying manifold is necessarily  $\mathbb{R}^n$ , the problem is critical, and  $(u, v)$  belongs to a 1-parameter family, or they are all infinite (see Theorem 1.3). As a consequence of our methods of proof, we can actually extend all of our main results to a suitable class of Riemannian models that are not necessarily Cartan-Hadamard (see Corollary 1.5).

1.1. Motivation and the state of the art. Existence and qualitative properties of solutions to (1.1), posed in the Euclidean space  $\mathbb{R}^n$ ,  $n \geq 2$ , are by now well understood: in the *subcritical* regime, namely  $1 < p+1 < 2^* := \frac{2n}{n-2}$  (with  $2^* = \infty$  if  $n = 2$ ), the problem has no classical solutions, regardless of radial symmetry. In the *critical* case  $p + 1 = 2^*$  such solutions exist, are radially symmetric, and correspond to the extremals of the Sobolev inequality; in particular, they belong to the Sobolev space  $D^{1,2}(\mathbb{R}^n)$ , defined as the completion of  $C_c^{\infty}(\mathbb{R}^n)$  with respect to the norm

$$
||u||_{D^{1,2}(\mathbb{R}^n)}^2 := \int_{\mathbb{R}^n} |\nabla u|^2 dx.
$$

In the *supercritical* regime  $p + 1 > 2^*$ , radial classical solutions still exist, decay to 0 at infinity, but do not belong to the *energy space*  $D^{1,2}(\mathbb{R}^n)$ . We refer the interested reader to the excellent monograph [22], and to the corresponding bibliography, for further details.

The situation can be considerably different if  $(1.1)$  is posed on a Cartan-Hadamard manifold  $\mathbb{M}^n$ . A first remarkable difference is that positive radial solutions may exist even in the subcritical regime, both with finite and infinite  $D^{1,2}(\mathbb{M}^n)$  norm; for instance, this is the case if  $\mathbb{M}^n \equiv \mathbb{H}^n$  is the *n*-dimensional hyperbolic space [3, 18], or if  $\mathbb{M}^n$  is a more general model manifold satisfying suitable assumptions [2, Theorems 2.5 and 2.7]. Concerning the critical or supercritical regimes, the situation is somehow more rigid, in the following sense. If u is a radial solution to  $(1.1)$  on a Cartan-Hadamard model manifold  $\mathbb{M}^n$ , with  $p+1 \geq 2^*$  and  $||u||_{D^{1,2}(\mathbb{R}^n)} < +\infty$ , then  $\mathbb{M}^n$  is necessarily isometric to  $\mathbb{R}^n$  and u is therefore an Aubin-Talenti function [20, Theorem 1.3] (see also [2, Theorem 2.4] for a related result obtained under additional assumptions on  $\mathbb{M}^n$ , and [16] for a previous rigidity result concerning solutions that minimize the Sobolev quotient). Furthermore, the asymptotic behavior of radial solutions is strongly affected by the global geometric properties of the underlying manifold: if  $\mathbb{M}^n$  is *stochastically complete*, then all radial solutions tend to 0 at infinity; otherwise, if it is *stochastically incomplete*, each solution converges to a strictly positive constant at infinity [20, Theorem 1.4]. Additional asymptotic estimates can be found in [20, Theorem 1.5] and [2, Theorem 2.4].

We finally refer to  $[1, 5, 7, 8]$  and references therein for results regarding  $(1.1)$  and related inequalities posed on manifolds with nonnegative Ricci curvature, namely the case complementary to ours.

Concerning system  $(1.2)$ , the problem in  $\mathbb{R}^n$  presents several analogies with the corresponding scalar case. In particular, one can naturally introduce a subcritical regime

 $\frac{1}{p+1} + \frac{1}{q+1}$ 

 $p, q > 0, \frac{1}{1}$ 

a critical regime

$$
p, q > 0,
$$
 
$$
\frac{1}{p+1} + \frac{1}{q+1} = \frac{n-2}{n},
$$
 (1.3)

 $\frac{1}{n}$ ,

 $\frac{1}{q+1} > \frac{n-2}{n}$ 

and a supercritical regime

$$
p, q > 0 \,, \qquad \frac{1}{p+1} + \frac{1}{q+1} < \frac{n-2}{n} \,. \tag{1.4}
$$

For subcritical exponents, radial positive solutions do not exist [19], and it is conjectured that positive classical solutions do not exist at all. This has been rigorously proved only up to dimension  $n = 4$  [27] (see also [4, 21, 23, 24]). On the other hand, for critical or supercritical exponents radial positive solutions do exist [17,25,26]. In the critical case, they correspond to extremals for higher-order Sobolev inequalities [17] (see also [28]), while in the supercritical regime it is not expected that they belong to any natural Sobolev space.

As far as we know, problem (1.2) on Cartan-Hadamard (model) manifolds is untouched. The purpose of this paper is thus to investigate existence and qualitative properties of radial positive solutions, in the critical or supercritical regimes. This implies working in spatial dimension  $n \geq 3$ , otherwise conditions (1.3) and (1.4) are empty. As concerns the subcritical regime, that we do not address here, it is natural to wonder whether or not, on suitable Cartan-Hadamard manifolds that significantly differ from  $\mathbb{R}^n$ , globally positive solutions can exist. If we think of what happens in the Euclidean case (non-existence, as recalled above), and the results in  $[18]$  and  $[2]$  (existence for  $(1.1)$  on the hyperbolic space and on many other model manifolds), the answer still depends on the specific analytic-geometric properties of  $\mathbb{M}^n$ .

In order to state our main results in a precise form, it is necessary to recall first some standard definitions and introduce notations accordingly.

1.2. Main results and basic notions. We say that a noncompact Riemannian manifold  $\mathbb{M}^n$  is a model if there exists a pole  $o \in \mathbb{M}^n$  such that its metric g is given, in polar (or spherical) global coordinates about  $o$ , by

$$
g \equiv dr \otimes dr + \psi^2(r) g_{\mathbb{S}^{n-1}},
$$

where r is the Riemannian distance of a point  $x \equiv (r, \theta) \in \mathbb{R}^+ \times \mathbb{S}^{n-1}$  from o,  $g_{\mathbb{S}^{n-1}}$  stands for the usual round metric on the unit sphere  $\mathbb{S}^{n-1}$  and  $\psi$  is a  $C^1([0,+\infty)) \cap C^{\infty}((0,+\infty))$  positive function with  $\psi(0) = 0$  and  $\psi'(0) = 1$  (for a more complete introduction to model manifolds we refer the reader e.g. to [10, 11]). The Cartan-Hadamard assumption in this case is equivalent to the fact that  $\psi$  is in addition convex, due to the explicit expression of the sectional curvatures in terms of  $\psi$ , see for instance [14, Section 2]. A prototypical example is represented by the choice  $\psi(r) = \sinh r$ , which gives rise to a well-known realization of the *hyperbolic space*  $\mathbb{H}^n$ , whose sectional curvature is identically equal to  $-1$ .

For future convenience, we define

$$
\Theta(r) := \frac{1}{\psi^{n-1}(r)} \int_0^r \psi^{n-1} \, ds \qquad \forall r > 0 \, ; \tag{1.5}
$$

namely, Θ is the function accounting for the volume-surface ratio of geodesic balls centered at the pole. Its importance, for our purposes, is due to the fact that a model manifold  $\mathbb{M}^n$  (not necessarily of Cartan-Hadamard type) is stochastically complete if  $\Theta \notin L^1(\mathbb{R}^+)$ , whereas it is stochastically incomplete if  $\Theta \in L^1(\mathbb{R}^+)$ . This dichotomy will have a key role in our results. We refer to [11, Sections 3 and 6 for a deeper discussion, and we also point out the recent papers  $[12, 13]$  for new nonlinear analytic characterizations of stochastic (in)completeness for general manifolds.

By writing (1.2) in polar coordinates, it is not difficult to check (see for instance [2] or [20] for the details) that a radial solution is (represented by) a regular enough function  $(u, v) : [0, +\infty) \to \mathbb{R}^2$  solving the Cauchy problem

$$
\begin{cases}\n(\psi^{n-1}u')' + \psi^{n-1} |v|^{q-1}v = 0 & \text{for } r > 0 \\
(\psi^{n-1}v')' + \psi^{n-1} |u|^{p-1}u = 0 & \text{for } r > 0\n\end{cases}
$$
\n
$$
u'(0) = 0 = v'(0)
$$
\n
$$
u(0) = \xi, \quad v(0) = \eta,
$$
\n(1.6)

for some initial data  $(\xi, \eta) \in (0, +\infty)^2$ . Note that, although we are only interested in positive solutions, for technical reasons it is necessary to be able to deal with sign-changing solutions as well, whence the replacement of  $v^q$  and  $u^p$  with  $|v|^{q-1}v$  and  $|u|^{p-1}u$ , respectively. The fact that  $(1.6)$  gives rise to an everywhere positive solution is a highly nontrivial issue, which is actually false in general, and will be thoroughly addressed in Section 3. In what follows, we will say that  $(u, v)$  is a (radial) globally positive solution if it solves (1.6) for every  $r > 0$  and  $u, v > 0$  on  $[0, +\infty)$ . Clearly, any such a solution solves the Lane-Emden system (1.2).

At this point it is worth recalling that, in the Euclidean setting, from well-known results due to Serrin and Zou [25, 26] (see also [17]) the system (1.6) for  $\mathbb{M}^n \equiv \mathbb{R}^n$  and in the critical-supercritical regime

$$
\frac{1}{p+1} + \frac{1}{q+1} \le \frac{n-2}{n} \tag{1.7}
$$

admits a globally positive solution if and only if the initial data  $(\xi, \eta)$  satisfy the explicit relation

$$
\eta \equiv \eta(\xi) = c \,\xi^{\frac{p+1}{q+1}}\,,\tag{1.8}
$$

where c is a positive constant depending only on p, q, n. In particular, we observe that  $\xi \mapsto \eta(\xi)$  is a strictly increasing and continuous bijection of  $(0, +\infty)$  into itself. We stress that the specific form (1.8) of the function  $\eta(\xi)$  is crucially related to the natural scaling properties of (1.2) in  $\mathbb{R}^n$ , which however fail on model manifolds. Indeed, as we will see in a moment, such function is not explicit and its very definition, *i.e.* the fact that for every  $\xi > 0$  there exists a *unique* value of  $\eta$  ensuring global positivity, strongly depends on the stochastic completeness or incompleteness of  $\mathbb{M}^n$ .

Before stating our main results, for notational convenience, for any globally positive solution of (1.6) we set

$$
\ell_u := \lim_{r \to +\infty} u(r) \,, \qquad \ell_v := \lim_{r \to +\infty} v(r) \,,
$$

the existence of such limits being guaranteed by the monotonicity of both components, which readily follows from the differential equations in (1.6) (see Section 2).

**Theorem 1.1** (Globally positive solutions for stochastically complete manifolds). Let  $\mathbb{M}^n$  (n > 3) be a Cartan-Hadamard model manifold associated to a function  $\psi$  with  $\Theta \notin L^1(\mathbb{R}^+)$ . Let  $p, q > 0$  fulfill (1.7). Then, for each  $\xi > 0$  there exists one and only one value  $\eta \equiv \eta(\xi) > 0$  such that  $(\xi, \eta)$  gives rise to a globally positive solution  $(u, v)$  to  $(1.6)$ , which satisfies

$$
\ell_u=\ell_v=0\,.
$$

Moreover, the function  $\xi \mapsto \eta(\xi)$  is a continuous and strictly increasing bijection of  $(0, +\infty)$  into itself.

Hence, in the stochastically complete case the situation is Euclidean like, since there exists a specific continuous curve of initial data that give rise to globally positive solutions, except that it has no more the explicit expression (1.8). Furthermore, both components of the solution vanish as  $r \to +\infty$ . This agrees with the results of Serrin and Zou in [25, 26]. Instead, in the stochastically incomplete case the scenario is more complicated and marks a striking difference with respect to the Euclidean framework.

**Theorem 1.2** (Globally positive solutions for stochastically incomplete manifolds). Let  $\mathbb{M}^n$   $(n \geq 3)$  be a Cartan-Hadamard model manifold associated to a function  $\psi$  with  $\Theta \in L^1(\mathbb{R}^+)$ . Let  $p, q > 0$  fulfill (1.7). Then there exist two functions  $\eta_m$ ,  $\eta_M$  which are continuous and strictly increasing bijections of  $(0, +\infty)$ into itself, satisfying

$$
\eta_m(\xi) < \eta_M(\xi) \qquad \forall \xi > 0,\tag{1.9}
$$

$$
\limsup_{\xi \to +\infty} \left[ \eta_M(\xi) - \eta_m(\xi) \right] < +\infty \,, \tag{1.10}
$$

such that for each  $\xi > 0$  problem (1.6) admits a globally positive solution  $(u, v)$  if and only if

$$
\eta_m(\xi) \le \eta \le \eta_M(\xi). \tag{1.11}
$$

In addition, the following behavior at infinity holds:

$$
\int \ell_u > 0, \quad \ell_v = 0 \qquad \text{if } \eta = \eta_m(\xi), \tag{1.12a}
$$

$$
\begin{cases}\n\ell_u > 0, & \ell_v > 0 \\
\ell_u > 0, & \ell_v > 0\n\end{cases} \quad \text{if } \eta_m(\xi) < \eta < \eta_M(\xi),
$$
\n(1.12b)

$$
\left(\ell_u = 0, \ \ell_v > 0 \qquad \text{if } \eta = \eta_M(\xi). \right) \tag{1.12c}
$$



FIGURE 1. The regions of existence of a globally positive solution in terms of the initial data  $(\xi, \eta)$ , in the case of a stochastically complete (A) and incomplete (B) manifold, with the corresponding behavior of the limits at infinity  $(\ell_u, \ell_v)$  of the components.

Two symbolic instances of the global positivity region in the space of the parameters  $(\xi, \eta)$ , associated with a stochastically complete and a stochastically incomplete Cartan-Hadamard model manifold, respectively, are depicted in Figure 1 below.

We now focus on the possible existence of radial *finite-energy solutions* to (1.2) in the critical or supercritical cases. We recall that for the scalar problem (1.1) such solutions cannot exist unless  $\mathbb{M}^n \equiv \mathbb{R}^n$ (*i.e.* the manifold is isometric to the Euclidean space), as shown in  $[20]$ . Here we are able to reproduce the natural counterpart of this rigidity result for the Lane-Emden system.

**Theorem 1.3** (Energy rigidity). Let  $\mathbb{M}^n$  ( $n \geq 3$ ) be a Cartan-Hadamard model manifold associated to a function  $\psi$ , and let  $p, q > 0$  fulfill (1.7). Suppose that, for some  $(\xi, \eta) \in (0, +\infty)^2$ , there exists a radial solution  $(u, v)$  to  $(1.2)$  such that

$$
\int_0^{+\infty} u'v' \psi^{n-1} ds < +\infty \quad or \quad \int_0^{+\infty} u^{p+1} \psi^{n-1} ds < +\infty \quad or \quad \int_0^{+\infty} v^{q+1} \psi^{n-1} ds < +\infty.
$$
  
\nThen  $\mathbb{M}^n \equiv \mathbb{R}^n$ ,  
\nand  
\n
$$
\frac{1}{p+1} + \frac{1}{q+1} = \frac{n-2}{n}
$$

 $\epsilon$ 

$$
\int_0^{+\infty} u'v' \,\psi^{n-1} \, ds + \int_0^{+\infty} u^{p+1} \,\psi^{n-1} \, ds + \int_0^{+\infty} v^{q+1} \,\psi^{n-1} \, ds < +\infty. \tag{1.13}
$$

**Remark 1.1.** We point out that on  $\mathbb{M}^n \equiv \mathbb{R}^n$  radial solutions do comply with (1.13) for every critical pair  $(p, q)$ , see [17]. They can be obtained through a variational argument combined with a scaling property, and are characterized as extremals of higher-order Sobolev inequalities. As already observed, up to translations, radial solutions form a 1-parameter family (the parameter being the value of one component, say u, at  $r = 0$ ), and their asymptotic behavior is understood [15]. Differently from what happens in the scalar case  $(1.1)$ , their explicit expression is however unknown in general.

It is also worth to mention that any solution to (1.2) on  $\mathbb{R}^n$  with  $(u, v) \in L^{p+1}(\mathbb{R}^n) \times L^{q+1}(\mathbb{R}^n)$  is radially symmetric [6]. This is proved by using an integral version of the moving planes method, and seems hardly adaptable on general manifolds.

We now state a key preliminary proposition regarding the "first zeros" of a solution to  $(1.6)$ , which may be of independent interest and will play an important role in the proof of our main results.

**Proposition 1.4.** Let  $\mathbb{M}^n$  (n > 3) be a Cartan-Hadamard model manifold associated to a function  $\psi$ , and let  $p, q > 0$  fulfill (1.7). Then, given any  $(\xi, \eta) \in (0, +\infty)^2$ , problem (1.6) coupled with

 $u(R) = v(R) = 0$  for some  $R > 0$ 

has no positive solution on  $(0, R)$ .

Note that this can be read as a nonexistence result for the radial homogeneous Dirichlet problem in Riemannian balls.

We conclude with a result in the spirit of [2, Theorem 2.2], which is essentially a consequence of our methods of proof, showing that the Cartan-Hadamard assumption can be slightly relaxed.

Corollary 1.5. Let  $\mathbb{M}^n$   $(n \geq 3)$  be a noncompact model manifold associated to a function  $\psi$ , and let  $p, q > 0$  fulfill (1.7). Then Theorems 1.1, 1.2, 1.3 and Proposition 1.4 still hold provided the function

$$
\mathcal{V}(r) := \left(\int_0^r \psi^{n-1} \, ds\right)^{\frac{pq-1}{2(p+1)(q+1)}} \qquad \forall r \in [0, +\infty) \tag{1.14}
$$

is convex.

Remark 1.2. In the critical case, the exponent in formula (1.14) attains its minimum value (within the critical-supercritical regime), which is precisely  $\frac{1}{n-1}$ . After some routine calculations (see *e.g.* the proof of Proposition 2.5 below), one can check that the convexity of  $V$  is equivalent to

$$
\int_0^r \frac{\psi^n \psi''}{(\psi')^2} ds \ge 0 \quad \forall r \in (0, +\infty).
$$

In particular, we emphasize the fact that there is room for  $\psi''$  to be negative somewhere, which means that the (radial) curvatures of  $\mathbb{M}^n$  are allowed to be positive is some small region. Clearly, this is a fortiori admissible in the supercritical regime.

1.3. Paper organization. We devote Section 2 to the proof of some useful inequalities and local existence results for the solutions to  $(1.6)$  (including Proposition 1.4). In Section 3 we show that, for a given  $\xi > 0$ , there exists at least one  $\eta > 0$  for which (1.6) yields a globally positive solution. This will require a number of preliminary technical tools. In Section 4 we prove our main results regarding the complete structure of the region of global positivity in the initial-data space  $(\xi, \eta)$  and the asymptotics of solutions as  $r \to +\infty$ , that is Theorems 1.1 and 1.2. Finally, Section 5 contains the proof of the rigidity Theorem 1.3, and in Section 6 we establish the generalization of our main results stated in Corollary 1.5.

For notational convenience, from here on we set  $a \vee b := \max\{a, b\}$  and  $a \wedge b := \min\{a, b\}.$ 

### 2. Preliminary properties of radial solutions

In this section, we first establish some basic local existence results for problem (1.6), and then we focus on a key Pohozaev-type inequality and its consequences.

From here on, unless otherwise specified, we take for granted that  $\mathbb{M}^n$  is a Cartan-Hadamard model manifold of dimension  $n \geq 3$ . In particular, we observe that the differential equations appearing in the system  $(1.6)$  can be rewritten as

$$
u'' + (n-1)\frac{\psi'}{\psi}u' + |v|^{q-1}v = 0, \qquad v'' + (n-1)\frac{\psi'}{\psi}v' + |u|^{p-1}u = 0,
$$
\n(2.1)

or, by integrating,

$$
\psi^{n-1}(r) u'(r) = -\int_0^r |v|^{q-1} v \psi^{n-1} ds, \qquad \psi^{n-1}(r) v'(r) = -\int_0^r |u|^{p-1} u \psi^{n-1} ds. \tag{2.2}
$$

In the sequel, both  $(2.1)$  and  $(2.2)$  will be very useful to our purposes.

2.1. Local existence and continuation lemmas. The existence and uniqueness of a local positive solution to (1.6) is rather classical, but for the sake of completeness we provide a full proof.

**Lemma 2.1.** Let  $p, q > 0$ . Then for every  $(\xi, \eta) \in (0, +\infty)^2$  there exists  $\rho \equiv \rho(\xi, \eta) > 0$ , depending continuously on  $(\xi, \eta)$ , such that problem (1.6) has a unique positive solution for  $r \in (0, \rho)$ .

Proof. Let

$$
X := \left\{ (u, v) \in C([0, \rho]; \mathbb{R}^2) : ||(u, v) - (\xi, \eta)||_{\infty} \leq \frac{1}{2} \xi \wedge \eta \right\},\
$$

where  $\rho$  has to be chosen later, and  $\|(\cdot, \cdot)\|_{\infty}$  denotes the norm obtained as the maximum between the usual  $L^{\infty}$  norms of the two components on  $[0, \rho]$ . Note that, by construction, since  $\xi, \eta > 0$  the set X is formed by positive functions. Let also  $F \equiv (F_1, F_2) : X \to C([0, \rho]; \mathbb{R}^2)$  be defined by

$$
F_1(u, v)(r) := \xi - \int_0^r \frac{1}{\psi^{n-1}(s)} \left( \int_0^s v^q \psi^{n-1} dt \right) ds,
$$
  

$$
F_2(u, v)(r) := \eta - \int_0^r \frac{1}{\psi^{n-1}(s)} \left( \int_0^s u^p \psi^{n-1} dt \right) ds.
$$

It is not difficult to check that

$$
|F_1(u,v)(r) - \xi| \le \left(\frac{3}{2}\eta\right)^q \int_0^r \Theta ds,
$$
  

$$
|F_2(u,v)(r) - \eta| \le \left(\frac{3}{2}\xi\right)^p \int_0^r \Theta ds,
$$
\n(2.3)

and

$$
|F_1(u_1, v_1)(r) - F_1(u_2, v_2)(r)| \leq \left(\sup_{\tau \in \left[\frac{1}{2}\eta, \frac{3}{2}\eta\right]} q \tau^{q-1}\right) \|v_1 - v_2\|_{\infty} \int_0^r \Theta ds,
$$
  

$$
|F_2(u_1, v_1)(r) - F_2(u_2, v_2)(r)| \leq \left(\sup_{\tau \in \left[\frac{1}{2}\xi, \frac{3}{2}\xi\right]} p \tau^{p-1}\right) \|u_1 - u_2\|_{\infty} \int_0^r \Theta ds,
$$
\n(2.4)

for every  $r \in [0, \rho]$  and every  $(u_1, v_1), (u_2, v_2) \in X$ . Since  $\Theta$  is locally bounded (it is smooth on  $(0, +\infty)$ ) and behaves like  $r^2$  near  $r = 0$ , from (2.3) and (2.4) it follows easily that there exists  $\rho > 0$ , depending on  $\psi$  and n, and in a continuous fashion on  $(\xi, \eta)$ , such that F is a contraction mapping from X into itself. Thus, by the Banach fixed-point theorem,  $F$  has a unique fixed point in  $X$ , which is clearly a positive solution of  $(1.6)$  in  $(0, \rho)$ .

The local solution constructed above is always positive for small  $r$ , and by classical ODE theory can be extended to a maximal interval  $[0, T)$ , possibly changing sign. Note that, if  $p \wedge q \geq 1$ , the extension and its lifetime  $T > 0$  only depend on the initial data  $(\xi, \eta)$ . However, if  $p \wedge q < 1$ , the uniqueness of the solution may fail past any point  $\bar{r} > 0$  where either  $u(\bar{r}) = 0$  or  $v(\bar{r}) = 0$ ; thus, the value of T can in this case also depend on the specific chosen extension, and not only on  $(\xi, \eta)$ . Nevertheless, we will see that it is still possible to obtain a quantitative lower bound on T.

Let us now set

$$
R_{\xi,\eta} := \sup \{ r \in [0,T) : \ u(t) \wedge v(t) > 0 \quad \forall t \in (0,r) \}, \tag{2.5}
$$

namely the size of the maximal positivity interval of the solution. Note that, by definition, if  $R_{\xi,\eta} < +\infty$ either  $u(R_{\xi,\eta}) = 0$  or  $v(R_{\xi,\eta}) = 0$ , and  $R_{\xi,\eta}$  is uniquely determined by  $\xi$  and  $\eta$  (as opposed to T), since the solution is unique as long as it stays positive. Moreover, still in the case where  $R_{\xi,\eta}$  is finite, the continuation theorem for ODE ensures that  $T > R_{\xi,\eta}$ , and one between u and v must necessarily change sign past  $r = R_{\xi,\eta}$ , due to the strong maximum principle.

As a straightforward consequence of the definition of  $R_{\xi,\eta}$  and (2.2), we have the following fundamental monotonicity result, along with the characterization of the limits at infinity in the stochastically complete case.

**Lemma 2.2.** Let  $p, q > 0$  and  $(\xi, \eta) \in (0, +\infty)^2$ . Let  $(u, v)$  solve  $(1.6)$ . Then

$$
u'(r) < 0 \quad and \quad v'(r) < 0 \qquad \forall r \in (0, R_{\xi, \eta}).
$$

In particular, if  $R_{\xi,n} = +\infty$  (namely  $(u, v)$  is a globally positive solution) there exist finite the limits

$$
\ell_u:=\lim_{r\to+\infty}u(r)\geq 0\,,\qquad \ell_v:=\lim_{r\to+\infty}v(r)\geq 0\,.
$$

Corollary 2.3. Let  $\Theta \notin L^1(\mathbb{R}^+)$ . Let  $p, q > 0$  and  $(\xi, \eta) \in (0, +\infty)^2$ . Then, if  $(u, v)$  is a globally positive solution to (1.6), it satisfies

$$
\ell_u=\ell_v=0\,.
$$

*Proof.* Suppose by contradiction that one of these limits, say  $\ell_u$ , is strictly positive (if  $\ell_u = 0$  and  $\ell_v > 0$ the argument can be repeated in the same way). Upon integrating  $(2.2)$  on  $(r, +\infty)$ , we deduce that

$$
v(r) - \ell_v = \int_r^{+\infty} \frac{1}{\psi^{n-1}(s)} \left( \int_0^s u^p \, \psi^{n-1} \, dt \right) ds \qquad \forall r > 0 \, .
$$

The above integral can be easily estimated by recalling that  $u$  is monotone decreasing due to Lemma 2.2, hence

which is clearly not possible. 
$$
v(r) - \ell_v \ge \ell_u^p \int_r^{+\infty} \Theta ds = +\infty \qquad \forall r > 0,
$$

In the proof of Proposition 3.4, that is the crucial fact that under (1.7) for every  $\xi > 0$  there exists at least one  $\eta \equiv \eta(\xi) > 0$  that gives rise to a globally positive solution, we will need the continuous dependence of (suitable extensions of) the solutions to (1.6) with respect to  $(\xi, \eta)$  also beyond the positivity radius  $R_{\xi,\eta}$ . Again, this follows from standard ODE theory if  $p \land q \geq 1$ , but since (1.7) allows one exponent to be strictly smaller than 1 (at least in dimension  $n \geq 5$ ), we need an argument which covers all these cases.

**Lemma 2.4.** Let  $p, q > 0$  and  $(\xi, \eta) \in (0, +\infty)^2$ . Let  $(\bar{u}, \bar{v})$  be the solution to (1.6) provided by Lemma 2.1. Let us fix any  $\sigma \in (\rho, R_{\xi,\eta})$ . Then there exists an extension of  $(\bar{u}, \bar{v})$  whose maximal existence interval contains  $[0, \sigma + \beta]$ , where

$$
\beta := \min\left\{\frac{\sigma}{2^q}\left(-1 + \sqrt{1 + \frac{C_1}{\sigma^2}}\right), \frac{\sigma}{2^p}\left(-1 + \sqrt{1 + \frac{C_2}{\sigma^2}}\right)\right\} > 0,
$$
\n(2.6)

for some positive constants  $C_1, C_2 > 0$  depending continuously on  $(\xi, \eta)$  and independent of  $\sigma$ .

*Proof.* We let Y denote the subset of  $C([\sigma, \sigma + \beta]; \mathbb{R}^2)$  consisting of all functions  $(u, v)$  which satisfy

$$
||(u,v) - (\bar{u},\bar{v})(\sigma)||_{\infty} \leq \xi \wedge \eta,
$$
\n(2.7)

with  $\sigma$  as in the statement and  $\beta > 0$  to be chosen later. By what we recalled above, we can assume that  $(\bar{u}, \bar{v})$  exists up to  $r = \sigma$ . Let also  $F \equiv (F_1, F_2) : Y \to C([\sigma, \sigma + \beta]; \mathbb{R}^2)$  be defined by

$$
F_1(u,v)(r) := \bar{u}(\sigma) + \psi^{n-1}(\sigma) \bar{u}'(\sigma) \int_{\sigma}^r \frac{1}{\psi^{n-1}} ds - \int_{\sigma}^r \frac{1}{\psi^{n-1}(s)} \left( \int_{\sigma}^s |v|^{q-1} v \, \psi^{n-1} \, dt \right) ds,
$$
  
\n
$$
F_2(u,v)(r) := \bar{v}(\sigma) + \psi^{n-1}(\sigma) \, \bar{v}'(\sigma) \int_{\sigma}^r \frac{1}{\psi^{n-1}} ds - \int_{\sigma}^r \frac{1}{\psi^{n-1}(s)} \left( \int_{\sigma}^s |u|^{p-1} u \, \psi^{n-1} \, dt \right) ds.
$$

The set Y is clearly closed and convex, and it is easy to check that  $F$  is a continuous mapping from Y to  $C([\sigma, \sigma + \beta]; \mathbb{R}^2)$ , since  $p, q > 0$ . We aim at showing that F has a precompact image, and that for  $\beta > 0$  conveniently chosen it maps Y into itself. As a result, the Schauder fixed-point theorem (see

e.g. [9, Corollary 11.2]) will ensure the existence of a fixed point  $(\hat{u}, \hat{v})$  for F in Y, which is nothing but a solution of the Cauchy problem

$$
\begin{cases}\n\left(\psi^{n-1}\,\hat{u}'\right)' + \psi^{n-1}\left|\hat{v}\right|^{q-1}\hat{v} = 0 & \text{for } r \in (\sigma, \sigma + \beta) \\
\left(\psi^{n-1}\,\hat{v}'\right)' + \psi^{n-1}\left|\hat{u}\right|^{p-1}\hat{u} = 0 & \text{for } r \in (\sigma, \sigma + \beta) \\
\hat{u}'(\sigma) = \bar{u}'(\sigma), \quad \hat{v}'(\sigma) = \bar{v}'(\sigma), \\
\hat{u}(\sigma) = \bar{u}(\sigma), \quad \hat{v}(\sigma) = \bar{v}(\sigma).\n\end{cases}
$$

Therefore, because  $\bar{u}(\sigma), \bar{v}(\sigma) > 0$  and the solution is unique as long as it is positive, we can assert that  $(\hat{u}, \hat{v})$  is the desired extension. Let us then prove that the assumptions of the Schauder fixed-point theorem are met.

We start by showing that F has a precompact image. Since  $\sigma < R_{\xi,\eta}$ , by Lemma 2.2 we have that  $0 < \bar{u}(\sigma) < \xi$  and  $0 < \bar{v}(\sigma) < \eta$ . Hence, if  $(u, v) \in Y$ , it follows that  $||u||_{\infty} \leq 2\xi$  and  $||v||_{\infty} \leq 2\eta$ . Now, by the definition of F, for every  $\sigma \leq r_0 < r \leq \sigma + \beta$  it holds

$$
|F_1(u,v)(r) - F_1(u,v)(r_0)| \le \psi^{n-1}(\sigma) |\bar{u}'(\sigma)| \int_{r_0}^r \frac{1}{\psi^{n-1}} ds + \int_{r_0}^r \frac{1}{\psi^{n-1}(s)} \left( \int_{\sigma}^s |v|^q \psi^{n-1} dt \right) ds
$$
  

$$
\le C \int_{r_0}^r \frac{1}{\psi^{n-1}} ds + (2\eta)^q \int_{r_0}^r \Theta ds,
$$

where C is a positive constant depending on  $\sigma, \xi, \eta, n$  through  $\psi$  and  $\bar{u}$ . A similar expression can be derived for  $|F_2(u, v)(r) - F_2(u, v)(r_0)|$  and, recalling the regularity and positivity properties of  $\psi$  (along with the fact that  $\sigma > \rho > 0$ , we readily deduce the equicontinuity of  $F(Y)$ . Concerning the (quantitative) uniform boundedness, note that by  $(2.2)$  we have

$$
-\psi^{n-1}(\sigma)\,\bar{u}'(\sigma) = \int_0^\sigma \bar{v}^q \,\psi^{n-1}\,ds\,.
$$

Therefore, thanks to the monotonicity of  $\psi$ , it follows that

$$
|F_1(u,v)(r) - \bar{u}(\sigma)| \leq \psi^{n-1}(\sigma) |\bar{u}'(\sigma)| \int_{\sigma}^r \frac{1}{\psi^{n-1}} ds + \int_{\sigma}^r \frac{1}{\psi^{n-1}(s)} \left( \int_{\sigma}^s |v|^q \psi^{n-1} dt \right) ds
$$
  
\n
$$
\leq \left( \eta^q \int_0^{\sigma} \psi^{n-1} ds \right) \int_{\sigma}^{\sigma+\beta} \frac{1}{\psi^{n-1}} ds + (2\eta)^q \int_{\sigma}^{\sigma+\beta} \frac{1}{\psi^{n-1}(s)} \left( \int_{\sigma}^s \psi^{n-1} dt \right) ds \quad (2.8)
$$
  
\n
$$
\leq \eta^q \sigma \beta + (2\eta)^q \frac{\beta^2}{2},
$$

for all  $r \in [\sigma, \sigma + \beta]$ . Similarly, we obtain

$$
|F_2(u,v)(r) - \bar{v}(\sigma)| \le \xi^p \sigma \beta + (2\xi)^p \frac{\beta^2}{2}.
$$
\n(2.9)

This proves the  $F(Y)$  is also bounded and hence, by the Ascoli-Arzelà theorem, it is indeed a precompact subset of  $C([\sigma, \sigma + \beta]; \mathbb{R}^2)$ .

It remains to show that  $F: Y \to Y$  for suitable choice of  $\beta > 0$ . By (2.8) and (2.9), this is the case if

$$
\left[\eta^q \sigma \beta + (2\eta)^q \frac{\beta^2}{2}\right] \vee \left[\xi^p \sigma \beta + (2\xi)^p \frac{\beta^2}{2}\right] \leq \xi \wedge \eta.
$$

It is straightforward to verify that such inequality holds for  $\beta$  as in (2.6), provided

$$
C_1 = \frac{2^{q+1}\xi \wedge \eta}{\eta^q}, \qquad C_2 = \frac{2^{p+1}\xi \wedge \eta}{\xi^p}.
$$

Note, in particular, that  $C_1$  and  $C_2$  depend continuously on  $(\xi, \eta)$  and are independent of  $\sigma$ . The proof is thus complete.  $\Box$ 

If  $R_{\xi,\eta} < +\infty$ , by taking limits as  $\sigma \to R_{\xi,\eta}^-$  in (2.6) we deduce that the constructed solution to (1.6) is defined at least on the interval

$$
\left[0, R_{\xi,\eta} + \min\left\{\frac{R_{\xi,\eta}}{2^q} \left(-1 + \sqrt{1 + \frac{C_1}{R_{\xi,\eta}^2}}\right), \frac{R_{\xi,\eta}}{2^p} \left(-1 + \sqrt{1 + \frac{C_2}{R_{\xi,\eta}^2}}\right)\right\}\right] \supseteq [0, R_{\xi,\eta}].
$$

As already mentioned, such solution must change sign past  $R_{\xi,\eta}$ , and it may not be unique beyond this threshold if  $p \wedge q < 1$ .

2.2. Some fundamental identities and inequalities. Given a solution to  $(1.6)$ , we introduce the associated energy function

$$
F_{(u,v)}(r) := u'(r)v'(r) + \frac{1}{p+1} |u(r)|^{p+1} + \frac{1}{q+1} |v(r)|^{q+1},
$$

along with the Pohozaev function

$$
P_{(u,v)}(r) := \left(\int_0^r \psi^{n-1} ds\right) F_{(u,v)}(r) + \psi^{n-1}(r) \left(\frac{u(r)v'(r)}{p+1} + \frac{u'(r)v(r)}{q+1}\right).
$$

**Proposition 2.5.** Let  $p, q > 0$  and  $(\xi, \eta) \in (0, +\infty)^2$ . Let  $(u, v)$  solve  $(1.6)$ , with maximal existence interval  $[0, T)$ . Then

$$
F'_{(u,v)}(r) = -2(n-1)\frac{\psi'(r)}{\psi(r)}u'(r)v'(r)
$$
\n(2.10)

and

$$
P'_{(u,v)}(r) = K(r)u'(r)v'(r)
$$
\n(2.11)

for all  $r \in (0,T)$ , where

$$
K(r) := \left(\frac{1}{p+1} + \frac{1}{q+1} - \frac{n-2}{n}\right)\psi^{n-1}(r) - \frac{2(n-1)}{n}\left(\int_0^r \frac{\psi^n \psi''}{(\psi')^2} ds\right) \frac{\psi'(r)}{\psi(r)}.
$$

Proof. Formula  $(2.10)$  follows by direct calculations, using  $(2.1)$ . Taking advantage of the latter and  $(2.2)$ , we can then compute the derivative of  $P_{(u,v)}$ :

$$
P'_{(u,v)}(r) = \psi^{n-1}(r) F_{(u,v)}(r) - 2(n-1) \left( \int_0^r \psi^{n-1} ds \right) \frac{\psi'(r)}{\psi(r)} u'(r) v'(r)
$$
  

$$
- \frac{\psi^{n-1}(r) |u(r)|^{p+1}}{p+1} - \frac{\psi^{n-1}(r) |v(r)|^{q+1}}{q+1} + \left( \frac{1}{p+1} + \frac{1}{q+1} \right) \psi^{n-1}(r) u'(r) v'(r)
$$
  

$$
= \left[ \left( 1 + \frac{1}{p+1} + \frac{1}{q+1} \right) \psi^{n-1}(r) - 2(n-1) \left( \int_0^r \psi^{n-1} ds \right) \frac{\psi'(r)}{\psi(r)} \right] u'(r) v'(r).
$$

Integrating by parts, recalling that  $\psi(0) = 0$  and  $\psi'(0) = 1$ , we have that

$$
\int_0^r \psi^{n-1} ds = \frac{1}{n} \int_0^r \frac{n \psi^{n-1} \psi'}{\psi'} ds = \frac{\psi^n(r)}{n \psi'(r)} + \frac{1}{n} \int_0^r \frac{\psi^n \psi''}{(\psi')^2} ds,
$$

and hence

$$
P'_{(u,v)}(r) = \left[ \left( 1 + \frac{1}{p+1} + \frac{1}{q+1} - 2\frac{n-1}{n} \right) \psi^{n-1}(r) - 2\frac{n-1}{n} \left( \int_0^r \frac{\psi^n \psi''}{(\psi')^2} ds \right) \frac{\psi'(r)}{\psi(r)} \right] u'(r) v'(r)
$$
  
=  $K(r)u'(r)v'(r)$ ,

that is  $(2.11)$ .

The above result yields a key monotonicity property for  $P_{(u,v)}$ , in the critical or supercritical case.

**Proposition 2.6.** Let  $p, q > 0$  fulfill  $(1.7)$ , and  $(\xi, \eta) \in (0, +\infty)^2$ . Let  $(u, v)$  solve  $(1.6)$ . Then  $K(r) \leq 0$ for every  $r > 0$ , with  $K(r) = 0$  if and only if equality holds in  $(1.7)$  and  $\psi''(s) = 0$  for every  $s \in (0, r)$ . In particular, we have that  $P_{(u,v)}(r) \leq 0$  for every  $r \in (0, R_{\xi,\eta})$ , with  $P_{(u,v)}(r) = 0$  if and only if equality holds in (1.7) and  $\psi''(s) = 0$  for every  $s \in (0, r)$ .

*Proof.* The first part of the thesis is a direct consequence of the definition of K, since  $\psi'' \geq 0$  everywhere on any Cartan-Hadamard model manifold. The second part follows from formula (2.11) and the fact that  $P_{(u,v)}(0) = 0$ , recalling also Lemma 2.2.

We are now in position to prove our nonexistence result for  $(1.6)$  on balls.

*Proof of Proposition 1.4.* Assume, by contradiction, that the Cauchy problem  $(1.6)$  admits a solution  $(u, v)$  for suitable initial data  $(\xi, \eta) \in (0, +\infty)^2$ , which is positive on  $(0, R)$  and satisfies  $u(R) = v(R) = 0$ for some  $R > 0$ . Then, by virtue of Lemma 2.2, we have that  $u'(R) < 0$  and  $v'(R) < 0$ . However, since  $(1.7)$  holds, the definition of  $P_{(u,v)}$  and Proposition 2.6 entail

$$
0 \ge P_{(u,v)}(R) = \left(\int_0^R \psi^{n-1} ds\right) u'(R)v'(R) > 0,
$$
 which is absurd.

As a consequence, we infer that in the critical or supercritical case either  $R_{\xi,\eta} = +\infty$  or  $R_{\xi,\eta} < +\infty$ and the two components of the solution do not vanish simultaneously at  $r = R_{\xi,n}$ .

Finally, we show a fundamental ordering property for positive solutions.

**Lemma 2.7.** Let  $p, q > 0$ . Let  $\xi_1 \ge \xi_2 > 0$  and  $\eta_2 > \eta_1 > 0$ . Then, if  $(u_1, v_1)$  and  $(u_2, v_2)$  are two positive solutions to (1.6) starting from  $(\xi_1, \eta_1)$  and  $(\xi_2, \eta_2)$ , respectively, in the common interval  $(0, b)$ for some  $b \in (0, +\infty]$ , the functions

$$
r \mapsto u_1(r) - u_2(r)
$$
 and  $r \mapsto v_2(r) - v_1(r)$ 

are strictly increasing in (0, b).

Proof. First of all, let us show that

$$
u_2(r) < u_1(r)
$$
 and  $v_2(r) > v_1(r)$   $\forall r \in (0, b).$  (2.12)

Note that it suffices to establish the right inequality only, as the validity of the latter implies (recall (2.2))

$$
u_2(r) = \xi_2 - \int_0^r \frac{1}{\psi^{n-1}(s)} \left( \int_0^s v_2^q \psi^{n-1} dt \right) ds < \xi_1 - \int_0^r \frac{1}{\psi^{n-1}(s)} \left( \int_0^s v_1^q \psi^{n-1} dt \right) ds = u_1(r) \tag{2.13}
$$
\n
$$
\forall r \in (0, b),
$$

namely the left inequality. To this aim, we observe that by continuity  $v_2 > v_1$  at least in a small neighborhood of 0. Hence, if  $v_2 > v_1$  failed to hold in the whole  $(0, b)$  there would exist some  $a \in (0, b)$ such that  $v_2 > v_1$  in  $(0, a)$  and  $v_2(a) = v_1(a)$ . However, by arguing exactly as in (2.13), this would yield  $u_2 < u_1$  in  $(0, a)$ , which in turn entails

$$
v_2(a) = \eta_2 - \int_0^a \frac{1}{\psi^{n-1}(s)} \left( \int_0^s u_2^p \psi^{n-1} dt \right) ds > \eta_1 - \int_0^a \frac{1}{\psi^{n-1}(s)} \left( \int_0^s u_1^p \psi^{n-1} dt \right) ds = v_1(a),
$$

a contradiction. Therefore,  $(2.12)$  holds, and since for all  $r \in (0, b)$  we have

$$
(u_1 - u_2)'(r) = \frac{1}{\psi^{n-1}(r)} \int_0^r (v_2^q - v_1^q) \psi^{n-1} ds \quad \text{and} \quad (v_2 - v_1)'(r) = \frac{1}{\psi^{n-1}(r)} \int_0^r (u_1^p - u_2^p) \psi^{n-1} ds,
$$

such derivatives are strictly positive.

## 3. Existence of (at least) one globally positive solution

Our goal here is to establish an existence result for globally positive solutions, that covers both the stochastically complete and incomplete cases. To this aim, we carefully adapt the strategy developed by Serrin and Zou in [25], and split the argument into some intermediate steps.

Using the same notation as in Section 2, let us introduce the sets

$$
A := \{ (\xi, \eta) \in (0, +\infty)^2 : R_{\xi, \eta} < +\infty \text{ and } v(R_{\xi, \eta}) > u(R_{\xi, \eta}) = 0 \},
$$
  

$$
B := \{ (\xi, \eta) \in (0, +\infty)^2 : R_{\xi, \eta} < +\infty \text{ and } u(R_{\xi, \eta}) > v(R_{\xi, \eta}) = 0 \}.
$$

Note that, in view of Proposition 1.4, if (1.7) holds then  $A \cup B$  accounts for the whole set of initial data that do not give rise to a globally positive solution of  $(1.6)$ . Since we will have to handle separately, in some parts of the proof, stochastically complete and incomplete manifolds, in the latter case we also define the quantity

$$
\theta := \int_0^{+\infty} \Theta \, dr \in (0, +\infty), \tag{3.1}
$$

where  $\Theta$  is the same function as in  $(1.5)$ .

In the next three lemmas we describe the main topological properties of A and B.

**Lemma 3.1.** Let  $p, q > 0$ . Then both  $A \neq \emptyset$  and  $B \neq \emptyset$ . More precisely: (i) If  $\mathbb{M}^n$  is stochastically complete, namely  $\Theta \notin L^1(\mathbb{R}^+),$  then

$$
s, t > 0, \quad t > s^{\frac{p+1}{q+1}} \qquad \Longrightarrow \qquad (\xi, \eta) \equiv (s, 2t) \in A,
$$

and

$$
s, t > 0, \quad t < s^{\frac{p+1}{q+1}} \qquad \Longrightarrow \qquad (\xi, \eta) \equiv (2s, t) \in B.
$$

(ii) If  $\mathbb{M}^n$  is stochastically incomplete, namely  $\Theta \in L^1(\mathbb{R}^+)$ , then

$$
s, t > 0
$$
,  $t > (\theta s^p) \vee \left(\frac{s}{\theta}\right)^{\frac{1}{q}}$   $\implies$   $(\xi, \eta) \equiv (s, 2t) \in A$ ,

and

$$
s, t > 0
$$
,  $s > (\theta t^q) \vee \left(\frac{t}{\theta}\right)^{\frac{1}{p}}$   $\implies$   $(\xi, \eta) \equiv (2s, t) \in B$ .

*Proof.* We prove only the statements regarding the set  $A$ , as those regarding the set  $B$  are completely analogous. Let  $s, t > 0$ . We consider the (local) solution to  $(1.6)$  with  $(\xi, \eta) \equiv (s, 2t)$ , and define

$$
I := (0, s) \times (t, 2t), \qquad R_I := \sup \{ r \in (0, R_{\xi, \eta}) : (u(r), v(r)) \in I \}.
$$

By integrating (2.2) and exploiting the monotonicity of the components, we obtain

$$
u(r) - s \le -t^q \int_0^r \Theta \, d\sigma \qquad \text{and} \qquad v(r) - 2t \ge -s^p \int_0^r \Theta \, d\sigma \tag{3.2}
$$

for all  $r \in (0, R_I)$ . If  $R_I < +\infty$ , then by continuity and again monotonicity either  $u(R_I) = 0$  and  $v(R_I) \geq t$  or  $u(R_I) > 0$  and  $v(R_I) = t$ . In the former case it is plain that  $(s, 2t) \in A$ , so the proof is complete. Therefore, in what follows we aim at ruling out, under the stated assumptions on  $(s, t)$ , both the possibilities  $R_I = +\infty$  and  $u(R_I) > 0$  with  $v(R_I) = t$ .

(*i*)  $\Theta \notin L^1(\mathbb{R}^+).$ 

If  $R_I = +\infty$ , then  $(u, v)$  is a globally positive solution of (1.6), so Lemma 2.2 implies that  $u(r) \to \ell_u \in$  $[0, s)$  and  $v(r) \to \ell_v \in [t, 2t)$  as  $r \to +\infty$ . However, this is in contradiction with Corollary 2.3, since

 $t > 0$ . Thus  $R_I < +\infty$ . Suppose now that  $u(R_I) > 0$  and  $v(R_I) = t$ . As  $\Theta \notin L^1(\mathbb{R}^+)$  and  $\Theta > 0$ , there exists  $r_0 \in (0, +\infty)$  such that

$$
\int_0^r \Theta \, d\sigma - \frac{t}{s^p} \begin{cases} < 0 & \text{if } r < r_0 \,, \\ = 0 & \text{if } r = r_0 \,, \\ > 0 & \text{if } r > r_0 \,. \end{cases}
$$

By (3.2), we have that

$$
t = v(R_I) \ge 2t - s^p \int_0^{R_I} \Theta dr,
$$

whence  $R_I \ge r_0$ , and in particular  $u(r_0) \ge u(R_I) > 0$ . On the other hand, still (3.2) entails

$$
u(r_0) \le s - t^q \int_0^{r_0} \Theta dr = s \left( 1 - \frac{t^{q+1}}{s^{p+1}} \right) < 0,
$$

where the last inequality follows from the assumptions on  $(s, t)$ . Therefore, we obtain a contradiction again.

(*ii*)  $\Theta \in L^1(\mathbb{R}^+).$ 

Recall that in this case our assumptions on  $(s, t)$  read  $t > \theta s^p$  and  $\theta t^q > s$ . If  $R_I = +\infty$ , then by taking the limit as  $r \to +\infty$  in (3.2) we obtain

$$
0\leq \ell_u\leq s-\theta t^q<0\,,
$$

a contradiction. Thus  $R_I < +\infty$ . Suppose now that  $u(R_I) > 0$  and  $v(R_I) = t$ . Still by (3.2) and the positivity of  $\Theta$ , it holds

$$
t = v(R_I) \ge 2t - s^p \int_0^{R_I} \Theta dr > 2t - \theta s^p
$$

,

whence it follows that  $\theta s^p > t$ , which is absurd.

**Lemma 3.2.** Let  $p, q > 0$ . Then both the sets A and B are open.

*Proof.* When  $p \wedge q \geq 1$ , the result is essentially a consequence of the continuous dependence of the solutions of (1.6) with respect to the initial data  $(\xi, \eta)$ . In order to deal with general exponents, so as to cover the non-Lipschitz case  $p \wedge q < 1$  as well, one can argue similarly to [25, Section 5], where the same issue was addressed in  $\mathbb{R}^n$ . Since the modifications required to adapt their proof to our Riemannian setting are minor, we only sketch the main points of the strategy.

First of all, we fix an arbitrary  $(\xi_0, \eta_0) \in A$  and let  $(u_0, v_0)$  denote the corresponding solution to (1.6) starting from  $(\xi_0, \eta_0)$ . From the definition of A, it follows that  $R_0 \equiv R_{\xi_0, \eta_0} < +\infty$ ,  $u_0(R_0) = 0$  and  $v_0(R_0) > 0$ , with  $(u_0, v_0)$  positive in  $[0, R_0)$ . The goal is to show that there exists  $\delta_0 > 0$  (small enough) such that if  $|\xi - \xi_0| \vee |\eta - \eta_0| < \delta_0$  then  $(\xi, \eta) \in A$ , *i.e.* the solution  $(u, v)$  to (1.6) starting from  $(\xi, \eta)$ satisfies  $R_{\xi,\eta} < +\infty$ ,  $u(R_{\xi,\eta}) = 0$  and  $v(R_{\xi,\eta}) > 0$ . To this aim, we proceed as follows.

i) There exist  $c > 0$  and  $\delta' \in (0,1)$  such that, if  $|\xi - \xi_0| \vee |\eta - \eta_0| < \delta'$ , then the maximal existence interval of  $(u, v)$  contains at least the common interval  $[0, R_0 + c]$ . Moreover, the uniform bounds

$$
|u(r)| \le 2\xi \le 2(\xi_0 + 1) \quad \text{and} \quad |v(r)| \le 2\eta \le 2(\eta_0 + 1) \qquad \forall r \in [0, R_0 + c]
$$
 (3.3)

hold. This can be achieved by means of Lemma 2.4, taking advantage of the fact that the constants  $C_1, C_2$  in (2.6) depend continuously on  $(\xi, \eta)$ , the functions belonging to the space Y comply with (2.7) and, in addition, continuous dependence of the solutions holds in every interval [0, S] for  $S < R_0$ arbitrarily close to  $R_0$ , since  $u_0, v_0 > 0$  in  $[0, S]$ . In particular, for  $\delta' \equiv \delta_S' > 0$  small enough, we have  $R_{\xi,\eta} > S.$ 

ii) There exist  $\varepsilon \in (0, c \wedge R_0)$  and  $\delta'' \in (0, \delta')$  such that, if  $|\xi - \xi_0| \vee |\eta - \eta_0| < \delta''$ , then

$$
|u'(r) - u'_0(R_0)| \le \frac{1}{2} |u'_0(R_0)| \quad \text{and} \quad |v'(r) - v'_0(R_0)| \le \frac{1}{2} |v'_0(R_0)| \qquad \forall r \in [R_0 - \varepsilon, R_0 + \varepsilon]. \tag{3.4}
$$

As concerns the left estimate in (3.4) (one argues analogously for the right one), upon using (2.2) and the triangle inequality it is not difficult to deduce that

$$
|u'(r) - u'_0(R_0)| \le |u'(R_0 - \varepsilon) - u'_0(R_0 - \varepsilon)| + |u'_0(R_0)| \left| \frac{\psi^{n-1}(R_0)}{\psi^{n-1}(r)} - 1 \right|
$$
  
+ 
$$
\frac{1}{\psi^{n-1}(r)} \int_{R_0 - \varepsilon}^r v^q \psi^{n-1} ds + \frac{1}{\psi^{n-1}(r)} \int_{R_0 - \varepsilon}^{R_0} v_0^q \psi^{n-1} ds,
$$
 (3.5)

for all  $r \in [R_0-\varepsilon, R_0+\varepsilon]$ . Thanks to (3.3), the last three terms on the right-hand side of (3.5) can be made arbitrarily small by choosing  $\varepsilon$  small enough depending only on  $R_0$ ,  $u'_0(R_0)$ ,  $\xi_0$ ,  $\eta_0$ ,  $q$ ,  $\psi$ ,  $n$ . On the other hand, the first term can also be made arbitrarily small upon requiring  $|\xi - \xi_0| \vee |\eta - \eta_0| < \delta''$ , for a suitable  $\delta'' \equiv \delta''_{\varepsilon} \in (0, \delta')$ , still due to continuous dependence (recall that  $(u_0, v_0)$  is positive in  $[0, R_0 - \varepsilon]$ ).

iii) We notice that, in the whole interval  $[R_0 - \varepsilon, R_0 + \varepsilon]$ , by virtue of (3.4) we have

$$
u'(r) \le \frac{1}{2}u'_0(R_0) < 0
$$
 and  $\frac{3}{2}v'_0(R_0) \le v'(r) \le \frac{1}{2}v'_0(R_0) < 0$ .

In particular, upon integration, for every  $\alpha \in (0, \varepsilon)$  we obtain

$$
v(R_0 + \varepsilon) \ge v(R_0 - \alpha) - \frac{3}{2} |v_0'(R_0)| (\alpha + \varepsilon) \quad \text{and} \quad u(R_0 + \varepsilon) \le u(R_0 - \alpha) - \frac{1}{2} |u_0'(R_0)| (\alpha + \varepsilon). \tag{3.6}
$$

With no loss of generality, we may further require  $\varepsilon$  and  $\alpha$  to be so small that

$$
\varepsilon \le \frac{v(R_0)}{6\,|v_0'(R_0)|} \qquad \text{and} \qquad 0 < u_0(R_0 - \alpha) \le \frac{1}{4} \, |u_0'(R_0)| \, \varepsilon \,, \tag{3.7}
$$

since  $v_0(R_0) > 0$  and  $u_0(R_0) = 0$ . In view of (3.6), (3.7) and continuous dependence on [0,  $R_0 - \alpha$ ], we can choose  $\delta_0 \in (0, \delta'')$  so small that, if  $|\xi - \xi_0| \vee |\eta - \eta_0| < \delta_0$ , then  $(u, v)$  is positive in  $[0, R_0 - \varepsilon]$ with

$$
v(R_0 + \varepsilon) > 0
$$
 and  $u(R_0 + \varepsilon) < 0$ .

Because both u and v are decreasing in  $[R_0 - \varepsilon, R_0 + \varepsilon]$ , this implies that  $R_{\xi,\eta} \in (R_0 - \varepsilon, R_0 + \varepsilon)$ and  $u(R_{\xi,\varepsilon})=0$ . Hence  $(\xi,\eta)\in A$  as desired (the proof for B is similar).

 $\Box$ 

**Lemma 3.3.** Let  $p, q > 0$ . Then there exist two functions  $\phi, \gamma$  which are continuous and strictly increasing bijections of  $(0, +\infty)$  into itself, such that

$$
\{(\xi, \eta) \in (0, +\infty)^2 : \xi < \phi(\eta)\} \subset A,
$$
  

$$
\{(\xi, \eta) \in (0, +\infty)^2 : \eta < \gamma(\xi)\} \subset B.
$$

*Proof.* Let  $s > 0$ , and define

$$
f(s) := \begin{cases} s^{\frac{p+1}{q+1}} & \text{if } \Theta \notin L^1(\mathbb{R}^+), \\ (\theta s^p) \vee (\frac{s}{\theta})^{\frac{1}{q}} & \text{if } \Theta \in L^1(\mathbb{R}^+). \end{cases}
$$

In both cases, it is clear that f is a continuous and strictly increasing bijection of  $(0, +\infty)$  into itself, with inverse function  $f^{-1}$ . Moreover, by Lemma 3.1, we have

$$
s, t > 0, \quad t > f(s) \qquad \Longrightarrow \qquad (\xi, \eta) \equiv (s, 2t) \in A.
$$

In particular, we deduce that

$$
\xi,\eta\in(0,+\infty)^2\,,\quad \xi
$$

This shows the part of the thesis regarding A, with

$$
\phi(\eta) := f^{-1}\left(\frac{\eta}{2}\right) \qquad \forall \eta > 0.
$$

The proof of the second part of the thesis, regarding the set  $B$ , is completely analogous and therefore we  $\Box$  omit it.

We are finally in position to prove the most important result of this section.

**Proposition 3.4.** Let  $p, q > 0$  fulfill (1.7). Then, for each  $\xi > 0$  there exists at least one value  $\eta \equiv$  $\eta(\xi) > 0$  such that  $(\xi, \eta)$  gives rise to a globally positive solution to (1.6). In particular, (1.2) on  $\mathbb{M}^n$  has at least a 1-parameter family of solutions.

*Proof.* Let  $\xi > 0$ . Thanks to Lemma 3.3, we have that  $(\xi, \overline{\eta}) \in B$  for every  $\overline{\eta} > 0$  such that  $\overline{\eta} < \gamma(\xi)$ . Moreover, still by Lemma 3.3, it follows that  $(\xi, \hat{\eta}) \in A$  for every  $\hat{\eta} > 0$  such that  $\hat{\eta} > \phi^{-1}(\xi)$ . Thus, since A and B are open (due to Lemma 3.2) and disjoint sets, there necessarily exists  $\eta > 0$  (depending on ξ) such that  $(\xi, \eta) \notin A \cup B$ . Let us consider the solution  $(u, v)$  to the Cauchy problem (1.6) with this initial datum. Then there are two possibilities: either  $R_{\xi,\eta} = +\infty$ , which means that  $(u, v)$  is a globally positive solution as desired, or  $R_{\xi,\eta} < +\infty$ , and in this case  $u(R_{\xi,\eta}) = v(R_{\xi,\eta}) = 0$ . However, the latter cannot occur in view of Proposition 1.4, so the proof is complete.  $\Box$ 

## 4. Structure of globally positive solutions: proof of Theorems 1.1 and 1.2

In this section we prove our main results concerning the structure, with respect to  $(\xi, \eta)$ , of the existence region of globally positive solutions, along with their asymptotic behavior. In the stochastically complete case, the tools introduced above are enough, thus we will start from the proof of Theorem 1.1. On the contrary, the situation is much more complicated in the stochastically incomplete case, as we will need several new preliminary results, hence we devote to it an entire subsection.

*Proof of Theorem 1.1.* The fact that  $\ell_u = \ell_v = 0$ , for any globally positive solution, has already been shown in Corollary 2.3. Hence, let us focus on the rest of the statement. From Proposition 3.4, we know that for each  $\xi > 0$  there exists at least one  $\eta > 0$  such that the solution to (1.6) is globally positive. Assume by contradiction that there exists another  $\bar{\eta} > 0$  such that also the solution to (1.6) with initial data  $(\xi, \bar{\eta})$ , which we denote by  $(\bar{u}, \bar{v})$ , is globally positive. With no loss of generality, we can suppose that  $\bar{\eta} > \eta$ . Then, upon applying Lemma 2.7 with  $\xi_1 = \xi_2 = \xi$ ,  $\eta_2 = \bar{\eta} > \eta = \eta_1$  and  $b = +\infty$ , we would infer that

$$
\ell_{\bar{v}} - \ell_v = \lim_{r \to +\infty} \left[ \bar{v}(r) - v(r) \right] > \bar{\eta} - \eta > 0,
$$

which is absurd still in view of Corollary 2.3. Therefore,  $\eta \equiv \eta(\xi)$  is uniquely identified and it is well defined a function  $F : (0, +\infty) \to (0, +\infty)$  that to every  $\xi > 0$  associates such value. In order to show that it is nondecreasing, let  $\xi_1 > \xi_2 > 0$ . If, by contradiction,  $F(\xi_1) < F(\xi_2)$  then by reasoning as above we would obtain  $\ell_{v_2} - \ell_{v_1} > 0$ , which is impossible. Now we observe that, by reversing the roles of p, q and  $\xi, \eta$  and repeating the above argument, one finds that it is well defined a function  $G : (0, +\infty) \to (0, +\infty)$ that to every  $\eta > 0$  associates the only value  $\xi \equiv \xi(\eta) > 0$  such that  $(\xi, \eta)$  gives rise to a globally positive solution to (1.6). By construction, it is plain that  $G(F(\xi)) = \xi$  and  $F(G(\eta)) = \eta$  for all  $\xi, \eta > 0$ , so F is a bijection of  $(0, +\infty)$  into itself with  $G = F^{-1}$ . On the other hand, since F is nondecreasing, the only possibility is that it is actually strictly increasing and continuous.  $\Box$ 

4.1. The stochastically incomplete case. First of all, note that if  $\mathbb{M}^n$  is stochastically incomplete, then of course it cannot be isometric to  $\mathbb{R}^n$ . In terms of the function  $\psi$ , this means that  $\psi''$  must be strictly positive somewhere. Therefore, in the light of Proposition 2.6, we deduce that in the critical or supercritical case  $P_{(u,v)}(r) \leq -C < 0$  for every r large enough. As a straightforward consequence, we have that

$$
\psi^{n-1}(r) \left[ u'(r)v(r) + u(r)v'(r) \right] \leq -C \qquad \forall r \geq r_0,
$$

which in turn implies that

$$
u(r)v(r) \ge C \int_{r}^{+\infty} \frac{1}{\psi^{n-1}} ds \qquad \forall r \ge r_0,
$$
\n(4.1)

for a suitable  $r_0 > 0$  large enough. Note that the integral on the right-hand side is finite as a trivial consequence of the fact that  $\Theta \in L^1(\mathbb{R}^+)$ . This basic estimate will be crucial in the proof of the next result, which is the cornerstone of this subsection.

**Proposition 4.1.** Let  $\Theta \in L^1(\mathbb{R}^+)$  and  $p, q > 0$  fulfill (1.7). Let  $(\xi, \eta) \in (0, +\infty)^2$ . Then, if  $(u, v)$  is a qlobally positive solution to  $(1.6)$ , it holds

$$
\lim_{r \to +\infty} u(r) \vee v(r) > 0.
$$

*Proof.* We argue by contradiction, assuming that both  $u(r)$  and  $v(r)$  vanish as  $r \to +\infty$ . In such case, integrating first the differential equations in  $(1.6)$  from r to s, and then integrating with respect to s from r to  $+\infty$ , we readily obtain the following identities:

$$
0 = 1 + \frac{u'(r)}{u(r)} \psi^{n-1}(r) \int_r^{+\infty} \frac{1}{\psi^{n-1}} ds - \frac{1}{u(r)} \int_r^{+\infty} \frac{1}{\psi^{n-1}(s)} \left( \int_r^s v^q \psi^{n-1} dt \right) ds,
$$
 (4.2)

and

$$
0 = 1 + \frac{v'(r)}{v(r)} \psi^{n-1}(r) \int_r^{+\infty} \frac{1}{\psi^{n-1}} ds - \frac{1}{v(r)} \int_r^{+\infty} \frac{1}{\psi^{n-1}(s)} \left( \int_r^s u^p \psi^{n-1} dt \right) ds,
$$
(4.3)

for all  $r > 0$ . Note that (1.7) implies  $pq > 1$ , hence we can and will assume with no loss of generality that  $p > 1$ . Now we consider three possibilities:

$$
\limsup_{r \to +\infty} \frac{v(r)}{u^p(r)} = +\infty \quad \text{and} \quad \liminf_{r \to +\infty} \frac{v(r)}{u^p(r)} = 0,
$$
\n(A)

$$
\liminf_{r \to +\infty} \frac{v(r)}{u^p(r)} > 0,
$$
\n(B)

$$
\limsup_{r \to +\infty} \frac{v(r)}{u^p(r)} < +\infty. \tag{C}
$$

Since  $(A)$ ,  $(B)$  and  $(C)$  cover all the scenarios, achieving a contradiction in each case will prove the thesis. Suppose that (A) holds; in particular, this entails the existence of a sequence  $r_m \to +\infty$  such that

$$
\lim_{m \to \infty} \frac{v(r_m)}{u^p(r_m)} = 0 \quad \text{and} \quad \left(\frac{v}{u^p}\right)'(r_m) = 0 \quad \forall m \in \mathbb{N} \,.
$$
 (4.4)

The rightmost identity is equivalent to

$$
v'(r_m) u^{-p}(r_m) - p v(r_m) u^{-p-1}(r_m) u'(r_m) = 0 \quad \forall m \in \mathbb{N},
$$

that is

$$
\frac{1}{p} \frac{v'(r_m)}{v(r_m)} = \frac{u'(r_m)}{u(r_m)} \qquad \forall m \in \mathbb{N},
$$

whence

$$
\frac{u'(r_m)}{u(r_m)}\,\psi^{n-1}(r_m)\int_{r_m}^{+\infty}\frac{1}{\psi^{n-1}}\,ds = \frac{1}{p}\frac{v'(r_m)}{v(r_m)}\,\psi^{n-1}(r_m)\int_{r_m}^{+\infty}\frac{1}{\psi^{n-1}}\,ds \ge -\frac{1}{p} \qquad \forall m \in \mathbb{N},\qquad(4.5)
$$

where the inequality follows from  $(4.3)$  evaluated at  $r = r_m$ . On the other hand, the leftmost identity in  $(4.4)$  implies the existence of some constant  $c > 0$  such that

$$
v(r_m) \leq c u^p(r_m) \qquad \forall m \in \mathbb{N} \,. \tag{4.6}
$$

Hence, using (4.2) with  $r = r_m$  we end up with

$$
0 = 1 + \frac{u'(r_m)}{u(r_m)} \psi^{n-1}(r_m) \int_{r_m}^{+\infty} \frac{1}{\psi^{n-1}} ds - \frac{1}{u(r_m)} \int_{r_m}^{+\infty} \frac{1}{\psi^{n-1}(s)} \left( \int_{r_m}^{s} v^q \psi^{n-1} dt \right) ds
$$
  
\n
$$
\geq \frac{p-1}{p} - \frac{v^q(r_m)}{u(r_m)} \int_{r_m}^{+\infty} \Theta ds \geq \frac{p-1}{p} - c^q u^{pq-1}(r_m) \int_{r_m}^{+\infty} \Theta ds
$$
\n(4.7)

for all  $m \in \mathbb{N}$ , where we took advantage of (4.5), (4.6) and the fact that v is decreasing. Since  $pq > 1$ ,  $p > 1$ , and  $\Theta \in L^1(\mathbb{R}^+)$ , letting  $m \to \infty$  in  $(4.7)$  we reach the contradiction  $0 \geq \frac{p-1}{p}$ .

Suppose instead that  $(B)$  holds. This means that there exists a constant  $c > 0$  such that

$$
u^p(r) \leq c v(r) \qquad \forall r \geq 0. \tag{4.8}
$$

Let us rule out the possibility that

$$
\limsup_{r \to +\infty} \frac{v'(r)}{v(r)} \psi^{n-1}(r) \int_r^{+\infty} \frac{1}{\psi^{n-1}} ds \le -1.
$$
\n(4.9)

Indeed, if (4.9) were satisfied, we could select  $0 < \epsilon < \frac{1}{p+1}$  and  $r_{\epsilon} > r_0$  such that

$$
\frac{v'(r)}{v(r)}\,\psi^{n-1}(r)\int_r^{+\infty}\frac{1}{\psi^{n-1}}\,ds\leq -(1-\epsilon)\qquad\forall r\geq r_\epsilon\,,
$$

and a simple integration of this differential inequality on  $(r_{\epsilon}, r)$  would yield

$$
v(r) \le C_{\epsilon} \left( \int_{r}^{+\infty} \frac{1}{\psi^{n-1}} ds \right)^{1-\epsilon} \qquad \forall r \ge r_{\epsilon}, \tag{4.10}
$$

for some constant  $C_{\epsilon} > 0$ . Hence, thanks to (4.1), we would infer that

$$
u(r) \ge \frac{C}{C_{\epsilon}} \left( \int_{r}^{+\infty} \frac{1}{\psi^{n-1}} ds \right)^{\epsilon} \qquad \forall r \ge r_{\epsilon}.
$$
\n(4.11)

But (4.10), (4.11) and (4.8) are inconsistent, as  $0 < \epsilon < \frac{1}{p+1}$ . Therefore, since (4.9) cannot hold, we deduce that there exist a sequence  $r_m \to +\infty$  and  $\alpha \in [0,1)$  such that

$$
\lim_{m \to \infty} \frac{v'(r_m)}{v(r_m)} \psi^{n-1}(r_m) \int_{r_m}^{+\infty} \frac{1}{\psi^{n-1}} ds = -\alpha.
$$
\n(4.12)

Taking  $r = r_m$  in (4.3), using the fact that u is decreasing along with (4.8), we obtain

$$
0 = 1 + \frac{v'(r_m)}{v(r_m)} \psi^{n-1}(r_m) \int_{r_m}^{+\infty} \frac{1}{\psi^{n-1}} ds - \frac{1}{v(r_m)} \int_{r_m}^{+\infty} \frac{1}{\psi^{n-1}(s)} \left( \int_{r_m}^s u^p \psi^{n-1} dt \right) ds
$$
  
\n
$$
\geq 1 + \frac{v'(r_m)}{v(r_m)} \psi^{n-1}(r_m) \int_{r_m}^{+\infty} \frac{1}{\psi^{n-1}} ds - \frac{u^p(r_m)}{v(r_m)} \int_{r_m}^{+\infty} \Theta ds
$$
  
\n
$$
\geq 1 + \frac{v'(r_m)}{v(r_m)} \psi^{n-1}(r_m) \int_{r_m}^{+\infty} \frac{1}{\psi^{n-1}} ds - c \int_{r_m}^{+\infty} \Theta ds
$$
 (4.13)

for all  $m \in \mathbb{N}$ . Hence, by passing to the limit in  $(4.13)$  as  $m \to \infty$ , due to  $(4.12)$  we reach the contradiction  $0 \geq 1 - \alpha$ .

Finally, suppose that  $(C)$  holds. This means that there exists a constant  $c > 0$  such that

$$
v(r) \leq c u^p(r) \qquad \forall r \geq 0. \tag{4.14}
$$

Similarly to  $(B)$ , we can rule out the possibility that

$$
\limsup_{r \to +\infty} \frac{u'(r)}{u(r)} \,\psi^{n-1}(r) \int_r^{+\infty} \frac{1}{\psi^{n-1}} \, ds \le -1 \,,
$$

otherwise for any  $0 < \epsilon < \frac{p}{p+1}$  and suitable constants  $r_{\epsilon} > r_0$  and  $C_{\epsilon} > 0$  we would deduce the inequalities

$$
u(r) \le C_{\epsilon} \left(\int_{r}^{+\infty} \frac{1}{\psi^{n-1}} ds\right)^{1-\epsilon} \quad \text{and} \quad v(r) \ge \frac{C}{C_{\epsilon}} \left(\int_{r}^{+\infty} \frac{1}{\psi^{n-1}} ds\right)^{\epsilon} \qquad \forall r \ge r_{\epsilon},
$$

which are inconsistent with (4.14). Hence, we can infer the existence of a sequence  $r_m \rightarrow +\infty$  and  $\beta \in [0, 1)$  such that

$$
\lim_{m \to \infty} \frac{u'(r_m)}{u(r_m)} \psi^{n-1}(r_m) \int_{r_m}^{+\infty} \frac{1}{\psi^{n-1}} ds = -\beta.
$$
\n(4.15)

Taking  $r = r_m$  in (4.2), by combining the decreasing monotonicity of v and (4.14) we reach

$$
0 = 1 + \frac{u'(r_m)}{u(r_m)} \psi^{n-1}(r_m) \int_{r_m}^{+\infty} \frac{1}{\psi^{n-1}} ds - \frac{1}{u(r_m)} \int_{r_m}^{+\infty} \frac{1}{\psi^{n-1}(s)} \left( \int_{r_m}^{s} v^q \psi^{n-1} dt \right) ds
$$
  
\n
$$
\geq 1 + \frac{u'(r_m)}{u(r_m)} \psi^{n-1}(r_m) \int_{r_m}^{+\infty} \frac{1}{\psi^{n-1}} ds - \frac{v^q(r_m)}{u(r_m)} \int_{r_m}^{+\infty} \Theta ds
$$
  
\n
$$
\geq 1 + \frac{u'(r_m)}{u(r_m)} \psi^{n-1}(r_m) \int_{r_m}^{+\infty} \frac{1}{\psi^{n-1}} ds - c^q u^{pq-1}(r_m) \int_{r_m}^{+\infty} \Theta ds
$$

for all  $m \in \mathbb{N}$ , which leads again to the contradiction  $0 \geq 1-\beta$  as  $m \to \infty$ , since (4.15) holds and  $pq > 1$ . The proof is thus complete.  $\Box$ 

The above proposition motivates a better understanding of the asymptotic behavior of globally positive solutions when  $\Theta \in L^1(\mathbb{R}^+)$ , as for the moment we only know that at least one of the two components has a strictly positive limit. This is the main purpose of the next intermediate results.

**Lemma 4.2.** Let  $p, q > 0$ . Let  $\xi > 0$  and  $\eta_2 > \eta_1 > 0$ . Then, if  $(u_1, v_1)$  and  $(u_2, v_2)$  are two globally positive solutions to (1.6) starting from  $(\xi, \eta_1)$  and  $(\xi, \eta_2)$ , respectively, for each  $\eta \in (\eta_1, \eta_2)$  there exists a globally positive solution to (1.6) starting from  $(\xi, \eta)$ .

*Proof.* A locally positive solution  $(u, v)$  to  $(1.6)$  always exists by shooting, and it continues to exist as long as it is positive (recall the results of Subsection 2.1). So, let  $b \equiv R_{\xi,\eta} \in (0, +\infty]$  be the largest number for which  $(u, v)$  is positive in the interval  $(0, b)$ . If, by contradiction,  $b < +\infty$  then its definition would imply that either  $u(b) = 0$  or  $v(b) = 0$ . However, due to Lemma 2.7, we can infer that

$$
u_2(r) < u(r) < u_1(r)
$$
 and  $v_1(r) < v(r) < v_2(r)$   $\forall r \in (0, b)$ ,

which are clearly inconsistent with both  $u(b) = 0$  and  $v(b) = 0$ , since  $(u_1, v_1)$  and  $(u_2, v_2)$  are globally positive solutions by assumption.  $\Box$ 

In the following, we let  $C_b([0, +\infty); \mathbb{R}^2)$  denote the space of globally bounded and continuous functions on  $[0, +\infty)$  with values in  $\mathbb{R}^2$ . For notational convenience, we set  $|(a_1, a_2)| := |a_1| \vee |a_2|$  for any  $a_1, a_2 \in \mathbb{R}$ .

**Lemma 4.3.** Let  $\Theta \in L^1(\mathbb{R}^+)$ . Given any  $C_1, C_2 > 0$ , the set

$$
Z := \left\{ (u, v) \in C_b([0, +\infty); \mathbb{R}^2) \middle| \quad \begin{array}{l} ||(u, v)||_{\infty} \leq C_1, \\ |(u(r), v(r)) - (u(s), v(s))| \leq C_2 \int_s^r \Theta \, dt & \forall r > s \geq 0 \end{array} \right\} \tag{4.16}
$$

is compact in  $C_b([0,+\infty);\mathbb{R}^2)$ .

Proof. Let  $\{(u_k, v_k)\}\subset Z$ . Since  $\Theta$  is locally bounded, from the definition of Z it follows that for every  $R > 0$  the sequence is uniformly bounded and uniformly Lipschitz in  $C([0, R]; \mathbb{R}^2)$ . Hence, by the Ascoli-Arzelà theorem and a standard diagonal procedure we can infer that there exist  $(\bar{u}, \bar{v}) \in C([0, +\infty); \mathbb{R}^2)$ and a subsequence  $\{(u_{k_j}, v_{k_j})\}$  such that

$$
(u_{k_j}, v_{k_j}) \underset{j \to \infty}{\longrightarrow} (\bar{u}, \bar{v}) \quad \text{in } C([0, R]; \mathbb{R}^2) \text{ for every } R > 0.
$$
 (4.17)

In particular, pointwise convergence takes place, which readily ensures that  $(\bar{u}, \bar{v}) \in Z$ . We are therefore left with proving that convergence is uniform in the whole half line  $[0, +\infty)$ . To this aim, for any arbitrary  $\varepsilon > 0$  we select  $R_{\varepsilon} > 0$  so large that

$$
\int_{R_{\varepsilon}}^{+\infty} \Theta \, dt < \frac{\varepsilon}{3C_2} \,. \tag{4.18}
$$

By virtue of (4.17), there exists  $j_{\varepsilon} \in \mathbb{N}$  such that

$$
\left| \left( u_{k_j}(r), v_{k_j}(r) \right) - (\bar{u}(r), \bar{v}(r)) \right| < \frac{\varepsilon}{3} \qquad \forall r \in [0, R_{\varepsilon}], \ \forall j > j_{\varepsilon} \,. \tag{4.19}
$$

On the other hand, for larger values of  $r$  we have

$$
\left| \left( u_{k_j}(r), v_{k_j}(r) \right) - \left( \bar{u}(r), \bar{v}(r) \right) \right|
$$
  
\n
$$
\leq \left| \left( u_{k_j}(r), v_{k_j}(r) \right) - \left( u_{k_j}(R_{\varepsilon}), v_{k_j}(R_{\varepsilon}) \right) \right| + \left| \left( u_{k_j}(R_{\varepsilon}), v_{k_j}(R_{\varepsilon}) \right) - \left( \bar{u}(R_{\varepsilon}), \bar{v}(R_{\varepsilon}) \right) \right|
$$
  
\n
$$
+ \left| \left( \bar{u}(R_{\varepsilon}), \bar{v}(R_{\varepsilon}) \right) - \left( \bar{u}(r), \bar{v}(r) \right) \right| \leq 2C_2 \int_{R_{\varepsilon}}^r \Theta \, dt + \frac{\varepsilon}{3} < \varepsilon \qquad \forall r > R_{\varepsilon}, \ \forall j > j_{\varepsilon},
$$
\n(4.20)

where we exploited the definition of Z along with (4.18). Taking the limit as  $j \to \infty$  in (4.19) and (4.20) we end up with

$$
\limsup_{j\to\infty}\sup_{r\in[0,+\infty)}\left|\left(u_{k_j}(r),v_{k_j}(r)\right)-\left(\bar{u}(r),\bar{v}(r)\right)\right|\leq\varepsilon,
$$

which completes the proof in view of the arbitrariness of  $\varepsilon$ .

**Lemma 4.4.** Let  $p, q > 0$ . Given  $(\xi, \eta) \in (0, +\infty)^2$ , suppose that  $(u, v)$  is a globally positive solution to (1.6). Let  $\{(\xi_k, \eta_k)\}\subset (0, +\infty)^2$  be a sequence that converges to  $(\xi, \eta)$ . Let  $(u_k, v_k)$  denote the local solution to (1.6) starting from  $(\xi_k, \eta_k)$  and  $[0, R_k)$  its maximal positivity interval according to (2.5), with  $R_k \equiv R_{\xi_k,\eta_k} \in (0,+\infty]$ , for each  $k \in \mathbb{N}$ . Then

$$
\lim_{k \to \infty} R_k = +\infty \quad \text{and} \quad (u_k, v_k) \underset{k \to \infty}{\longrightarrow} (u, v) \quad locally \text{ uniformly in } [0, +\infty).
$$

*Proof.* With no loss of generality, we may assume that  $R_k < +\infty$  for all  $k \in \mathbb{N}$ , so either  $u_k(R_k) = 0$  or  $v_k(R_k) = 0$ . For simplicity, and up to subsequences, we discuss the former case only (if instead  $v_k(R_k) = 0$ one argues similarly). Hence, from a further integration of  $(2.2)$  we obtain

$$
0 = \xi_k - \int_0^{R_k} \frac{1}{\psi^{n-1}(s)} \left( \int_0^s v_k^q \psi^{n-1} dt \right) ds \ge \xi_k - \eta_k^q \int_0^{R_k} \Theta ds,
$$
\n(4.21)

thus it is plain that  ${R_k}$  stays bounded away from zero. Still up to subsequences, we may therefore assume, in addition, that  $R_k \to R$  as  $k \to \infty$  for some  $R \in (0, +\infty]$ . Suppose by contradiction that  $R < +\infty$ . As a result, for every  $S \in (0, R)$  the sequence  $\{(u_k, v_k)\}\$ is eventually positive and lies in a set of the form  $(4.16)$ , up to replacing  $[0, +\infty)$  with  $[0, S]$ . A local version of Lemma 4.3 is therefore applicable and guarantees that, again up to subsequences, it converges in  $C([0, S]; \mathbb{R}^2)$  to a nonnegative function  $(\bar{u}, \bar{v})$ , so by passing to the limit in the integral formulations satisfied by  $(u_k, v_k)$  we find that

$$
\bar{u}(r) = \xi - \int_0^r \frac{1}{\psi^{n-1}(s)} \left( \int_0^s \bar{v}^q \, \psi^{n-1} \, dt \right) ds \qquad \forall r \in [0, S]
$$

and

$$
\bar{v}(r) = \eta - \int_0^r \frac{1}{\psi^{n-1}(s)} \left( \int_0^s \bar{u}^p \, \psi^{n-1} \, dt \right) ds \qquad \forall r \in [0, S].
$$

However, this means that  $(\bar{u}, \bar{v})$  solves the same problem as  $(u, v)$  in [0, S], and therefore it must coincide with the latter in such interval (recall that  $(u, v)$  is globally positive by assumption). In particular, we

can rewrite  $(4.21)$ , for large k, as

$$
0 = \xi_k - \int_0^{R_k} \frac{1}{\psi^{n-1}(s)} \left( \int_0^s v_k^q \psi^{n-1} dt \right) ds
$$
  
=  $\xi_k - \int_0^S \frac{1}{\psi^{n-1}(s)} \left( \int_0^s v_k^q \psi^{n-1} dt \right) ds - \int_S^{R_k} \frac{1}{\psi^{n-1}(s)} \left( \int_0^s v_k^q \psi^{n-1} dt \right) ds$   
 $\geq \xi_k - \int_0^S \frac{1}{\psi^{n-1}(s)} \left( \int_0^s v_k^q \psi^{n-1} dt \right) ds - \eta_k^q \int_S^{R_k} \Theta ds,$ 

whence, taking limits as  $k \to \infty$  and using the above established convergence, it follows that

$$
0 \ge \xi - \int_0^S \frac{1}{\psi^{n-1}(s)} \left( \int_0^s v^q \, \psi^{n-1} \, dt \right) ds - \eta^q \int_S^R \Theta \, ds = u(S) - \eta^q \int_S^R \Theta \, ds \, .
$$

Finally, letting  $S \uparrow R$ , we would end up with

 $0 \geq u(R)$ ,

which is in contradiction with the global positivity of u. As a result, the only possibility is that  $R_k \to +\infty$ as  $k \to \infty$ , and thus the previously shown uniform convergence holds (at least) locally in the whole  $[0, +\infty)$ .

**Lemma 4.5.** Let  $\Theta \in L^1(\mathbb{R}^+)$  and  $p, q > 0$ . Let  $\{(\xi_k, \eta_k)\} \subset (0, +\infty)^2$  be a sequence that converges to some  $(\xi, \eta) \in (0, +\infty)^2$ , such that  $(u_k, v_k)$  is a globally positive solution to (1.6) starting from  $(\xi_k, \eta_k)$ , for each  $k \in \mathbb{N}$ . Then

$$
(u_k, v_k) \longrightarrow_{k \to \infty} (u, v)
$$
 uniformly in  $[0, +\infty)$ ,

where  $(u, v)$  is a globally positive solution to  $(1.6)$  starting from  $(\xi, \eta)$ .

Proof. It is enough to notice that, due to the positivity and the monotonicity of the components, the inequalities

$$
0 \le u_k(r) \le \xi_k \quad \text{and} \quad 0 \le v_k(r) \le \eta_k \qquad \forall r \in [0, +\infty)
$$

hold for every  $k \in \mathbb{N}$ , thus it is readily seen that  $\{(u_k, v_k)\}\$ is contained in a set of the form  $(4.16)$ . Hence, by virtue of Lemma 4.3, it admits a uniformly convergent subsequence  $\{(u_{k_j}, v_{k_j})\}$  to some  $(u, v) \in C_b([0, +\infty); \mathbb{R}^2)$ , which is therefore also nonnegative. On the other hand, by passing to the limit in the integral identities

$$
u_k(r) = \xi_k - \int_0^r \frac{1}{\psi^{n-1}(s)} \left( \int_0^s v_k^q \psi^{n-1} dt \right) ds, \qquad v_k(r) = \eta_k - \int_0^r \frac{1}{\psi^{n-1}(s)} \left( \int_0^s u_k^p \psi^{n-1} dt \right) ds,
$$

we infer that  $(u, v)$  actually solves (1.6) with initial datum  $(\xi, \eta)$ , so it is identified as the globally positive solution starting from  $(\xi, \eta)$ . Since the argument can be repeated along every subsequence of  $\{(u_k, v_k)\}$ , the claimed result holds for the whole sequence the claimed result holds for the whole sequence.

**Lemma 4.6.** Let  $\Theta \in L^1(\mathbb{R}^+)$  and  $p, q > 0$ . Given  $(\xi, \eta) \in (0, +\infty)^2$ , suppose that  $(u, v)$  is a globally positive solution to  $(1.6)$ . Then:

- i) If  $\ell_u > 0$ , there exists  $\varepsilon > 0$  such that for every  $\overline{\eta} \in (\eta, \eta + \varepsilon)$  the solution to (1.6) starting from  $(\xi, \bar{\eta})$  is globally positive;
- ii) If  $\ell_v > 0$ , there exists  $\varepsilon \in (0, \eta)$  such that for every  $\hat{\eta} \in (\eta \varepsilon, \eta)$  the solution to (1.6) starting from  $(\xi, \hat{\eta})$  is globally positive.

Proof. We proceed in both cases with an argument by contradiction.

i) If the thesis were false, then there would exist a sequence  $\varepsilon_k \downarrow 0$  such that the local solution  $(u_k, v_k)$ to (1.6), starting from  $(\xi, \eta + \varepsilon_k)$ , has a maximal positivity interval  $[0, R_k)$  with  $R_k \in (0, +\infty)$ . Thanks to the comparison principle entailed by Lemma 2.7, it must necessarily be  $u_k$  that vanishes at  $r = R_k$ , so we can write

$$
0 = \xi - \int_0^R \frac{1}{\psi^{n-1}(s)} \left( \int_0^s v_k^q \psi^{n-1} dt \right) ds - \int_R^{R_k} \frac{1}{\psi^{n-1}(s)} \left( \int_0^s v_k^q \psi^{n-1} dt \right) ds
$$
  
 
$$
\geq \xi - \int_0^R \frac{1}{\psi^{n-1}(s)} \left( \int_0^s v_k^q \psi^{n-1} dt \right) ds - (\eta + \varepsilon_k)^q \int_R^{+\infty} \Theta ds,
$$
 (4.22)

where  $R \in (0, R_k)$  is arbitrary but fixed for the moment. On the other hand, Lemma 4.4 ensures that  $R_k \to +\infty$  and  $\{(u_k, v_k)\}\)$  converges locally uniformly to  $(u, v)$  as  $k \to \infty$ , thus we can pass to the limit in  $(4.22)$  to obtain

$$
0 \ge \xi - \int_0^R \frac{1}{\psi^{n-1}(s)} \left( \int_0^s v^q \, \psi^{n-1} \, dt \right) ds - \eta^q \int_R^{+\infty} \Theta \, ds = u(R) - \eta^q \int_R^{+\infty} \Theta \, ds \, .
$$

Hence, by finally letting  $R \to +\infty$ , we end up with the inequality

 $0 > \ell_u$ ,

which is absurd.

ii) Similarly, to case i), denying the thesis would imply the existence of a sequence  $\varepsilon_k \downarrow 0$  such that the local solution  $(u_k, v_k)$  to (1.6), starting from  $(\xi, \eta - \varepsilon_k)$ , has a maximal positivity interval  $[0, R_k)$ with  $R_k \in (0, +\infty)$ . Still by virtue of Lemma 2.7, we deduce that in this case the component that vanishes at  $r = R_k$  is necessarily v. Hence, as above we can write

$$
0 = \eta - \varepsilon_k - \int_0^R \frac{1}{\psi^{n-1}(s)} \left( \int_0^s u_k^p \psi^{n-1} dt \right) ds - \int_R^{R_k} \frac{1}{\psi^{n-1}(s)} \left( \int_0^s u_k^p \psi^{n-1} dt \right) ds
$$
  
 
$$
\geq \eta - \varepsilon_k - \int_0^R \frac{1}{\psi^{n-1}(s)} \left( \int_0^s u_k^p \psi^{n-1} dt \right) ds - \xi^p \int_R^{+\infty} \Theta ds
$$

for all  $R \in (0, R_k)$ . Therefore, by using again Lemma 4.4 and passing to the limit as  $k \to \infty$ , we obtain

$$
0 \ge \eta - \int_0^R \frac{1}{\psi^{n-1}(s)} \left( \int_0^s u^p \, \psi^{n-1} \, dt \right) ds - \xi^p \int_R^{+\infty} \Theta \, ds = v(R) - \xi^p \int_R^{+\infty} \Theta \, ds,
$$
  
s

that is

upon taking the limit as  $R \to +\infty$ , still a contradiction.

Before proving Theorem 1.2, we establish a useful quantitative bound on the values of the limits at infinity  $\ell_u$  and  $\ell_v$ .

**Proposition 4.7.** Let  $\Theta \in L^1(\mathbb{R}^+)$  and  $p, q > 0$  fulfill  $pq > 1$ . Let  $(\xi, \eta) \in (0, +\infty)^2$ . Then, if  $(u, v)$  is a globally positive solution to  $(1.6)$ , it holds

$$
\ell_u \le \frac{p^{\frac{q+1}{pq-1}}(q+1)^{\frac{q+2}{pq-1}}}{(p+1)^{\frac{1}{pq-1}}(pq-1)^{\frac{q+1}{pq-1}}} \quad \text{and} \quad \ell_v \le \frac{q^{\frac{p+1}{pq-1}}(p+1)^{\frac{p+2}{pq-1}}}{(q+1)^{\frac{1}{pq-1}}(pq-1)^{\frac{p+1}{pq-1}}} \theta^{-\frac{p+1}{pq-1}}, \quad (4.23)
$$

where  $\theta$  was defined in (3.1).

*Proof.* First of all we observe that, by exploiting  $(2.2)$  and using the monotonicity of the components, we easily obtain the inequalities

$$
u'(r) \le -v^q(r) \Theta(r)
$$

and

$$
v'(r) \le -u^p(r) \Theta(r) ,
$$

for all  $r > 0$ . Integrating the former from r to  $+\infty$ , we infer that

$$
\ell_u - u(r) \le -\int_r^{+\infty} v^q \Theta \, ds \qquad \Longrightarrow \qquad -u(r) \le -\int_r^{+\infty} v^q \Theta \, ds,
$$

which substituted into the latter yields

$$
v'(r) \le -\left(\int_r^{+\infty} v^q \Theta \, ds\right)^p \Theta(r) \qquad \forall r > 0. \tag{4.24}
$$

Upon multiplying both sides of  $(4.24)$  by  $v<sup>q</sup>$ , note that such inequality can be rewritten as

$$
\frac{\left(v^{q+1}\right)'(r)}{q+1} \le -\left(\int_r^{+\infty} v^q \Theta \, ds\right)^p v^q(r) \Theta(r) \qquad \forall r > 0. \tag{4.25}
$$

Hence, by further integrating (4.25) from r to  $+\infty$  we end up with

$$
\frac{\ell_v^{q+1} - v^{q+1}(r)}{q+1} \le -\frac{\left(\int_r^{+\infty} v^q \Theta ds\right)^{p+1}}{p+1} \qquad \forall r > 0,
$$

which implies

$$
v(r) \ge \left(\frac{q+1}{p+1}\right)^{\frac{1}{q+1}} \left(\int_r^{+\infty} v^q \Theta \, ds\right)^{\frac{p+1}{q+1}} \qquad \forall r > 0,
$$

and this inequality can equivalently be rewritten as

$$
v^{q}(r)\Theta(r)\left(\int_{r}^{+\infty}v^{q}\Theta ds\right)^{-q\frac{p+1}{q+1}}\ge\left(\frac{q+1}{p+1}\right)^{\frac{q}{q+1}}\Theta(r)\qquad\forall r>0.
$$
\n(4.26)

On the other hand, if we integrate  $(4.26)$  from 0 to r, we find

$$
\frac{q+1}{pq-1}\left[\left(\int_r^{+\infty}v^q\,\Theta\,ds\right)^{-\frac{pq-1}{q+1}} - \left(\int_0^{+\infty}v^q\,\Theta\,ds\right)^{-\frac{pq-1}{q+1}}\right] \ge \left(\frac{q+1}{p+1}\right)^{\frac{q}{q+1}}\int_0^r\,\Theta\,ds \qquad \forall r>0,
$$

that is, upon dropping the rightmost term on the left-hand side and rearranging factors,

$$
\left(\int_{r}^{+\infty} v^q \Theta ds\right)^{\frac{pq-1}{q+1}} \le \frac{(q+1)^{\frac{1}{q+1}} (p+1)^{\frac{q}{q+1}}}{pq-1} \left(\int_{0}^{r} \Theta ds\right)^{-1} \qquad \forall r > 0.
$$

Because  $v(r) > \ell_v$ , the above estimate entails

$$
\ell_v^{q\frac{pq-1}{q+1}}\left(\int_r^{+\infty}\Theta\,ds\right)^{\frac{pq-1}{q+1}} \le \frac{(q+1)^{\frac{1}{q+1}}\,(p+1)^{\frac{q}{q+1}}}{pq-1}\left(\int_0^r\Theta\,ds\right)^{-1} \qquad \forall r>0\,,
$$

namely

$$
\ell_v \le \frac{(q+1)^{\frac{1}{q(pq-1)}} (p+1)^{\frac{1}{pq-1}}}{(pq-1)^{\frac{q+1}{q(pq-1)}}} \left( \int_0^r \Theta \, ds \right)^{-\frac{q+1}{q(pq-1)}} \left( \theta - \int_0^r \Theta \, ds \right)^{-\frac{1}{q}} \qquad \forall r > 0. \tag{4.27}
$$

A straightforward optimization argument over r ensures that the minimum of the right-hand side is attained if and only if

$$
\int_0^r \Theta ds = \frac{q+1}{q(p+1)} \theta,
$$

so by substituting such value into (4.27), and carrying out some algebraic simplifications, we deduce the claimed bound on  $\ell_v$  in (4.23). The analogous bound on  $\ell_u$  is readily obtained by symmetry, i.e. inter-<br>changing the roles of u and v along with those of p and q. changing the roles of  $u$  and  $v$  along with those of  $p$  and  $q$ .

*Proof of Theorem 1.2.* Let  $\xi > 0$  be fixed. First of all we observe that, as a direct consequence of Lemma 4.2, the set of all  $\eta > 0$  for which there exists a globally positive solution to (1.6) is necessarily an interval, which we call I. Proposition 3.4 guarantees that I is nonempty, and by virtue of Proposition 4.1 and Lemma 4.6 we can also assert that  $I$  is not a singleton. Moreover, we claim that

$$
\xi \ge (\eta - \theta \xi^p)_+^q \theta \qquad \text{and} \qquad \eta \ge (\xi - \theta \eta^q)_+^p \theta, \tag{4.28}
$$

which readily ensure that  $I$  is in addition bounded and bounded away from zero. In order to obtain (4.28) we notice that, by monotonicity,

$$
u(r) = \xi - \int_0^r \frac{1}{\psi^{n-1}(s)} \left( \int_0^s v^q \, \psi^{n-1} \, dt \right) ds \ge \xi - \eta^q \int_0^r \Theta \, ds \ge \xi - \theta \eta^q \tag{4.29}
$$

and

$$
v(r) = \eta - \int_0^r \frac{1}{\psi^{n-1}(s)} \left( \int_0^s u^p \, \psi^{n-1} \, dt \right) ds \ge \eta - \xi^p \int_0^r \Theta \, ds \ge \eta - \theta \xi^p \tag{4.30}
$$

for all  $r > 0$ , so the desired bounds follow by plugging  $(4.30)$  into  $(4.29)$  and vice versa, using the positivity of the components and eventually letting  $r \to +\infty$ . Hence, we can denote by  $\eta_m(\xi)$  and  $\eta_M(\xi)$ the strictly positive and finite infimum and supremum of  $I$ , respectively. An immediate application of Lemma 4.5 shows that they are actually a minimum and a maximum, that is, both the pairs  $(\xi, \eta_m(\xi))$ and  $(\xi, \eta_M(\xi))$  give rise to globally positive solutions to (1.6), *i.e.* I is in addition closed and thus (1.11) is necessary and sufficient for a global solution to exist. Still as a consequence of Proposition 4.1, Lemma 4.6 and the definitions of  $\eta_m$  and  $\eta_M$ , it is plain that (1.12a) and (1.12c) must hold. On the other hand, the validity of  $(1.12b)$  follows from  $(1.12a)$ ,  $(1.12c)$  and Lemma 2.7: indeed, if  $(u_m, v_m)$  and  $(u_M, v_M)$ are the globally positive solutions to (1.6) starting from  $(\xi, \eta_m(\xi))$  and  $(\xi, \eta_M(\xi))$ , respectively, and  $(u, v)$ is the one starting from  $(\xi, \eta)$ , for any  $\eta \in (\eta_m(\xi), \eta_M(\xi))$ , we have that

$$
\ell_u - \ell_{u_M} > \xi - \xi = 0
$$
 and  $\ell_v - \ell_{v_m} > \eta - \eta_m > 0$ ,

that is both  $\ell_u$  and  $\ell_v$  are strictly positive.

Let us now prove the claimed properties of the functions  $\xi \mapsto \eta_m(\xi)$  and  $\xi \mapsto \eta_M(\xi)$ , which by definition take values in  $(0, +\infty)$  and comply with  $(1.9)$ . To this aim, we can argue similarly to the proof of Theorem 1.1. Assume by contradiction that there exist  $\xi_1 > \xi_2 > 0$  such that  $\eta_m(\xi_1) < \eta_m(\xi_2)$ , and let  $(u_1, v_1)$  and  $(u_2, v_2)$  denote the corresponding solutions to  $(1.6)$  starting from  $(\xi_1, \eta_m(\xi_1))$  and  $(\xi_2, \eta_m(\xi_2))$ , respectively. Then, Lemma 2.7 would entail

$$
\ell_{v_2} - \ell_{v_1} > \eta_m(\xi_2) - \eta_m(\xi_1) > 0,
$$

which is absurd since we already know that  $\ell_{v_2} = \ell_{v_1} = 0$ . Hence, the function  $\xi \mapsto \eta_m(\xi)$  is nondecreasing. Via an analogous argument, we infer that also  $\xi \mapsto \eta_M(\xi)$  is nondecreasing (in this case one has to use the monotonicity of  $u_1 - u_2$ ). Upon reversing the roles of p, q and  $\xi, \eta$ , we notice that it is well defined a function  $\xi_M$  :  $(0, +\infty) \to (0, +\infty)$  that to every  $\eta > 0$  associates the only value  $\xi \equiv \xi_M(\eta) > 0$  such that  $(\xi, \eta)$  gives rise to a globally positive solution to  $(1.6)$  satisfying  $\ell_v = 0$ . By construction, we have that  $\xi_M(\eta_m(\xi)) = \xi$  and  $\eta_m(\xi_M(\eta)) = \eta$  for all  $\xi, \eta > 0$ , which shows that  $\eta_m$  is a bijection of  $(0, +\infty)$ into itself with  $\xi_M = \eta_m^{-1}$ . Therefore, due to its monotonicity, it is necessarily strictly increasing and continuous. A completely analogous reasoning proves that the same properties hold for  $\eta_M$ .

Finally, as for (1.10), it is enough to observe that Lemma 2.7 and Proposition 4.7 yield

$$
\eta_M(\xi) - \eta_m(\xi) < \ell_{v_M} - \ell_{v_m} \leq C \qquad \forall \xi > 0 \,,
$$

where  $C > 0$  is, for instance, the same constant appearing in the rightmost bound of formula (4.23).  $\Box$ 

## 5. Rigidity of finite-energy solutions: proof of Theorem 1.3

Now, our goal is to show that globally positive solutions cannot have a finite energy, unless  $\mathbb{M}^n \equiv$  $\mathbb{R}^n$  and  $q, p$  are critical. Before, we a need real-analysis lemma, which will be useful especially in the stochastically incomplete case.

**Lemma 5.1.** Let  $f \in C^1([1, +\infty))$  satisfy  $f, f' > 0$  and  $\lim_{r \to +\infty} f(r) = +\infty$ . Let  $\alpha, \epsilon > 0$ . Then

$$
\limsup_{r \to +\infty} f^{\epsilon}(r) \int_{r}^{+\infty} \left(\frac{f}{f'}\right)^{\alpha} ds = +\infty.
$$
\n(5.1)

Proof. First of all, we set

$$
g(t) := f^{-1}(t) \qquad \forall t \in [f(1), +\infty)
$$

and apply the change of variables  $\tau = f(s)$  in the integral in formula (5.1), obtaining:

$$
\int_{r}^{+\infty} \left(\frac{f}{f'}\right)^{\alpha} ds = \int_{f(r)}^{+\infty} \frac{\tau^{\alpha}}{\left[f'(f^{-1}(\tau))\right]^{\alpha+1}} d\tau = \int_{f(r)}^{+\infty} \tau^{\alpha} \left[g'(\tau)\right]^{\alpha+1} d\tau \ge f^{\alpha}(r) \int_{f(r)}^{+\infty} \left[g'(\tau)\right]^{\alpha+1} d\tau. \tag{5.2}
$$

Assume by contradiction that  $(5.1)$  does not hold, namely that there exists  $C > 0$  such that

$$
f^{\epsilon}(r) \int_{r}^{+\infty} \left(\frac{f}{f'}\right)^{\alpha} ds \leq C \qquad \forall r \in [1, +\infty).
$$

Upon setting  $r = f^{-1}(t)$  and taking advantage of  $(5.2)$ , we would thus infer that

$$
t^{\alpha+\epsilon} \int_{t}^{+\infty} \left[g'(\tau)\right]^{\alpha+1} d\tau \leq C \quad \forall t \in [f(1), +\infty),
$$

which in particular implies, by Hölder's inequality,

$$
\int_{t}^{2t} g'(\tau) d\tau \leq t^{\frac{\alpha}{\alpha+1}} \left( \int_{t}^{2t} \left[ g'(\tau) \right]^{\alpha+1} d\tau \right)^{\frac{1}{\alpha+1}} \leq C^{\frac{1}{\alpha+1}} t^{-\frac{\epsilon}{\alpha+1}} \qquad \forall t \in [f(1), +\infty). \tag{5.3}
$$

Finally, we apply (5.3) with the choices  $t \equiv t_k := f(1) \cdot 2^k$ , for all  $k \in \mathbb{N}$ . This yields

$$
\int_{t_k}^{t_{k+1}} g'(\tau) d\tau \leq \frac{C^{\frac{1}{\alpha+1}}}{f(1)^{\frac{\epsilon}{\alpha+1}}} 2^{-\frac{\epsilon}{\alpha+1}k} \quad \forall k \in \mathbb{N},
$$

whence, by adding up,

$$
\int_{f(1)}^{+\infty} g'(\tau) d\tau \leq \frac{C^{\frac{1}{\alpha+1}}}{f(1)^{\frac{\epsilon}{\alpha+1}}} \cdot \frac{2^{\frac{\epsilon}{\alpha+1}}}{2^{\frac{\epsilon}{\alpha+1}} - 1},
$$

that is  $g' \in L^1([f(1), +\infty))$ , which is inconsistent with the fact that g is surjective onto  $[1, +\infty)$ .

Proof of Theorem 1.3. Having in mind Remark 1.1, in order to prove the thesis it is enough to establish that, as soon as

$$
\mathbb{M}^n \not\equiv \mathbb{R}^n \qquad \text{or} \qquad \frac{1}{p+1} + \frac{1}{q+1} < \frac{n-2}{n},\tag{5.4}
$$

the globally positive solution  $(u, v)$  satisfies

$$
\int_0^{+\infty} u'v' \,\psi^{n-1} \, dr = \int_0^{+\infty} u^{p+1} \,\psi^{n-1} \, dr = \int_0^{+\infty} v^{q+1} \,\psi^{n-1} \, dr = +\infty \,. \tag{5.5}
$$

To this aim, as a consequence of Propositions 2.5 and 2.6, it is readily seen that under (5.4) there exist constants  $r_0, K_0 > 0$  such that

$$
\psi^{n-1}(r)\left(\frac{u(r)v'(r)}{p+1} + \frac{u'(r)v(r)}{q+1}\right) \le -K_0 \qquad \forall r \in [r_0, +\infty),\tag{5.6}
$$

since requiring  $\mathbb{M}^n \neq \mathbb{R}^n$  amounts asking that  $\psi'' > 0$  in some interval. On the other hand, upon multiplying the first and the second equation in  $(1.6)$  by v and u, respectively, and integrating by parts, we end up with the identities

$$
\int_0^r u'v' \psi^{n-1} ds - \psi^{n-1}(r)u'(r)v(r) = \int_0^r v^{q+1} \psi^{n-1} ds \qquad \forall r > 0
$$
\n(5.7)

and

$$
\int_0^r u'v' \psi^{n-1} ds - \psi^{n-1}(r)u(r)v'(r) = \int_0^r u^{p+1} \psi^{n-1} ds \qquad \forall r > 0.
$$
 (5.8)

In particular, recalling that  $u' < 0$  and  $v' < 0$ , we easily obtain

$$
\int_0^{+\infty} u'v' \psi^{n-1} dr \le \int_0^{+\infty} v^{q+1} \psi^{n-1} dr \quad \text{and} \quad \int_0^{+\infty} u'v' \psi^{n-1} dr \le \int_0^{+\infty} u^{p+1} \psi^{n-1} dr,
$$

which means that  $(5.5)$  is in fact equivalent to

$$
\int_0^{+\infty} u'v' \,\psi^{n-1} \, dr = +\infty \,. \tag{5.9}
$$

The proof that, under (5.4), then (5.9) holds, will be our main focus from now on. To reach it, we will distinguish between the stochastically complete and incomplete case.

i) Let  $\Theta \notin L^1(\mathbb{R}^+)$ . Thanks to Corollary 2.3, we know that

$$
\lim_{r \to +\infty} u(r) = 0 \quad \text{and} \quad \lim_{r \to +\infty} v(r) = 0. \tag{5.10}
$$

Assume by contradiction that

$$
\int_0^{+\infty} u'v' \,\psi^{n-1} \, dr < +\infty \,. \tag{5.11}
$$

As a consequence, by virtue of (5.7) and (5.8) we can deduce that both the limits

$$
L_1 := \lim_{r \to +\infty} \psi^{n-1}(r)u'(r)v(r)
$$
 and  $L_2 := \lim_{r \to +\infty} \psi^{n-1}(r)u(r)v'(r)$ 

exist, and they are clearly nonpositive. Moreover, upon letting  $r \to +\infty$  in (5.6), we infer that

$$
\frac{L_2}{p+1} + \frac{L_1}{q+1} \le -K_0 \,,
$$

which means that either  $L_2 < 0$  or  $L_2 = 0$  and  $L_1 < 0$ . In the former case, there exist constants  $r_1, K_1 > 0$ such that

$$
\psi^{n-1}(r)u(r)v'(r) \le -K_1 \qquad \forall r \in [r_1, +\infty),\tag{5.12}
$$

that is

$$
\psi^{n-1}(r)u'(r)v'(r) \ge -K_1 \frac{u'(r)}{u(r)} \qquad \forall r \in [r_1, +\infty).
$$

Upon integrating such inequality from  $r_1$  to any  $r > r_1$ , we end up with

$$
\int_{r_1}^r u'v' \psi^{n-1} ds \ge K_1 \log\left(\frac{u(r_1)}{u(r)}\right) \qquad \forall r \in [r_1, +\infty),
$$

which is clearly in contradiction with  $(5.11)$ , recalling the left limit in  $(5.10)$ . In the latter case one argues analogously, using the right limit in (5.10) instead.

ii) Let  $\Theta \in L^1(\mathbb{R}^+)$ . From Proposition 4.1, we know that at least one between  $\ell_u$  and  $\ell_v$  is strictly positive. If both  $\ell_u, \ell_v > 0$ , then from (2.2) and the monotonicity of u and v it readily follows that

$$
-u'(r) \ge \ell_v^q \frac{\int_0^r \psi^{n-1} ds}{\psi^{n-1}(r)} \quad \text{and} \quad -v'(r) \ge \ell_u^p \frac{\int_0^r \psi^{n-1} ds}{\psi^{n-1}(r)} \qquad \forall r > 0,
$$
\n(5.13)

thus (5.9) holds provided we can show that

$$
\int_0^{+\infty} \frac{\left(\int_0^r \psi^{n-1} ds\right)^2}{\psi^{n-1}(r)} dr = +\infty.
$$
 (5.14)

This is in fact a simple consequence of Lemma 5.1 with the choices  $f(r) = \int_0^r \psi^{n-1} ds$  and  $\alpha = \epsilon = 1$ , since

$$
\int_0^{+\infty} \frac{\left(\int_0^r \psi^{n-1} \, ds\right)^2}{\psi^{n-1}(r)} \, dr \ge \int_r^{+\infty} \frac{f^2}{f'} \, ds \ge f(r) \int_r^{+\infty} \frac{f}{f'} \, ds
$$

for all  $r \ge 1$ , whence (5.14) follows upon letting  $r \to +\infty$  along a sequence that attains the lim sup of the rightmost side. Let us therefore focus on the case where  $\ell_u = 0$  and  $\ell_v > 0$  (if instead  $\ell_v = 0$  and  $\ell_u > 0$ the argument is completely symmetric). The left inequality in (5.13) still holds, and its integration from r to  $+\infty$  yields

$$
u(r) \geq \ell_v^q \int_r^{+\infty} \frac{\int_0^s \psi^{n-1} dt}{\psi^{n-1}(s)} ds \qquad \forall r > 0.
$$
 (5.15)

On the other hand, by using this information along with the monotonicity of  $u$  in the right identity of (2.2), we also deduce that

$$
-v'(r) \ge u^p(r) \frac{\int_0^r \psi^{n-1} ds}{\psi^{n-1}(r)} \ge \ell_v^{pq} \left( \int_r^{+\infty} \frac{\int_0^s \psi^{n-1} dt}{\psi^{n-1}(s)} ds \right)^p \frac{\int_0^r \psi^{n-1} ds}{\psi^{n-1}(r)} \qquad \forall r > 0,
$$
\n(5.16)

so by multiplying  $(5.15)$  and  $(5.16)$  we obtain

$$
\psi^{n-1}(r)u(r)v'(r) \le -\ell_v^{(p+1)q} \left[ \left( \int_0^r \psi^{n-1} ds \right)^{\frac{1}{p+1}} \int_r^{+\infty} \frac{\int_0^s \psi^{n-1} dt}{\psi^{n-1}(s)} ds \right]^{p+1} \qquad \forall r > 0.
$$

Thanks to Lemma 5.1 applied to the same f as above,  $\alpha = 1$  and  $\epsilon = \frac{1}{p+1}$ , we infer that the liminf of the right-hand side is  $-\infty$ , hence also

$$
\liminf_{r \to +\infty} \psi^{n-1}(r)u(r)v'(r) = -\infty.
$$
\n(5.17)

Suppose by contradiction that  $(5.11)$  holds. Then, from  $(5.8)$  we deduce again that the limit  $L_u$  exists, and in this case it must necessarily be equal  $-\infty$  due to (5.17). This entails the validity of (5.12) for other suitable constants  $r_1, K_1 > 0$ , which is however inconsistent with (5.11) as shown in i) (recall that u vanishes at infinity by assumption).  $\square$ 

### 6. Generalizations: proof of Corollary 1.5

Finally, we show that all of our main results can be extended to a class of Riemannian models slightly wider than the Cartan-Hadamard one.

Proof of Corollary 1.5. First of all, we observe that the preliminary results of Subsection 2.1 hold regardless of the Cartan-Hadamard assumption. Moreover, a straightforward computation shows that requiring the function V to be convex, that is  $V'' \geq 0$  on  $(0, +\infty)$ , is equivalent to

$$
\left(1 + \frac{1}{p+1} + \frac{1}{q+1}\right)\psi^{n-1}(r) - 2(n-1)\left(\int_0^r \psi^{n-1} ds\right) \frac{\psi'(r)}{\psi(r)} \le 0 \qquad \forall r \in (0, +\infty). \tag{6.1}
$$

On the other hand, from the proof of Proposition 2.5 it is clear that  $(6.1)$  is precisely what we need to assert that  $P_{(u,v)}$  is monotone non-increasing, and  $P_{(u,v)}(r) \leq 0$  for every  $r \in (0, R_{\xi,\eta})$ , for any local solution to  $(1.6)$  starting from  $(\xi, \eta) \in (0, +\infty)^2$ . It is not difficult to verify that all the proofs of Section 3 and the proof of Theorem 1.1 solely rely on this property, *i.e.* we never use directly the fact that  $\psi'' \geq 0$ . As concerns Theorems 1.2 and 1.3, let us notice in addition that, if  $\mathbb{M}^n$  is a noncompact model manifold

such that V is convex and  $(5.4)$  holds, then  $V'' > 0$  somewhere. Indeed, if by contradiction  $V''(r) = 0$  for every  $r \in (0, +\infty)$ , from the definition of V we would end up with the identity

$$
\left(\int_0^r \psi^{n-1} ds\right)^{\frac{pq-1}{2(p+1)(q+1)}} = cr \qquad \forall r \in [0, +\infty)
$$

for some  $c > 0$ , that is

$$
\psi(r) = \tilde{c} r^{\frac{1}{n-1} \left[ \frac{2(p+1)(q+1)}{pq-1} - 1 \right]} \qquad \forall r \in [0, +\infty)
$$

for another constant  $\tilde{c} > 0$ . However, since  $\psi'(0) = 1$ , this is possible if and only if  $\tilde{c} = 1$  and

$$
\frac{1}{n-1} \left[ \frac{2(p+1)(q+1)}{pq-1} - 1 \right] = 1 \qquad \Longleftrightarrow \qquad \frac{1}{p+1} + \frac{1}{q+1} = \frac{n-2}{n},
$$

which also entails  $\psi(r) = r$ , that is the exponents are critical and  $\mathbb{M}^n \equiv \mathbb{R}^n$ , a contradiction. Hence, under  $(5.4)$  we can infer that  $\mathcal{V}''$  must be positive somewhere, which implies in turn that inequality  $(6.1)$ is strict in an interval, so (5.6) does hold (recall Proposition 2.6) and the proof of Theorem 1.3 can be carried out exactly as above. The fact that the volume of  $\mathbb{M}^n$  is infinite, which is used when Lemma 5.1 is invoked, is an immediate consequence of the convexity of  $V$  (and it is in any case always true if  $\mathbb{M}^n$  is stochastically incomplete). The same holds for Theorem 1.2, as the stochastic-incompleteness assumption yields  $\mathbb{M}^n \neq \mathbb{R}^n$ , so (4.1) is again satisfied and from there on the proof can be repeated identically.  $\Box$ 

Acknowledgments. M.M. was supported by the PRIN Project "Direct and Inverse Problems for Partial Differential Equations: Theoretical Aspects and Applications" (grant no. 201758MTR2, MIUR, Italy). The authors are partially supported by the INdAM-GNAMPA group.

#### **REFERENCES**

- [1] Z. M. Balogh and A. Kristály. Sharp isoperimetric and Sobolev inequalities in spaces with nonnegative Ricci curvature. Math. Ann., 385, 1747–1773, 2023.
- [2] E. Berchio, A. Ferrero, and G. Grillo. Stability and qualitative properties of radial solutions of the Lane-Emden-Fowler equation on Riemannian models. J. Math. Pures Appl. (9), 102(1):1–35, 2014.
- [3] M. Bonforte, F. Gazzola, G. Grillo, and J. L. Vázquez. Classification of radial solutions to the Emden-Fowler equation on the hyperbolic space. Calc. Var. Partial Differential Equations, 46(1-2):375–401, 2013.
- [4] J. Busca and R. Manásevich. A Liouville-type theorem for Lane-Emden systems. Indiana Univ. Math. J., 51(1):37-51, 2002.
- [5] G. Catino and D. D. Monticelli. Semilinear elliptic equations on manifolds with nonnegative Ricci curvature. Preprint arXiv: <https://arxiv.org/pdf/2203.03345.pdf>, 2022.
- [6] W. Chen, C. Li, and B. Ou. Classification of solutions for a system of integral equations. Comm. Partial Differential Equations, 30(1-3):59–65, 2005.
- [7] M. Fogagnolo, A. Malchiodi and L. Mazzieri. A note on the critical Laplace Equation and Ricci curvature. Preprint arXiv: <https://arxiv.org/pdf/2203.04678.pdf>, 2022.
- [8] B. Gidas and J. Spruck. Global and local behavior of positive solutions of nonlinear elliptic equations. Comm. Pure Appl. Math. 34 (4), 525–598, 1981.
- [9] D. Gilbarg and N. S. Trudinger. Elliptic partial differential equations of second order. Classics in Mathematics. Springer-Verlag, Berlin, 2001. Reprint of the 1998 edition.
- [10] R. E. Greene and H. Wu. Function theory on manifolds which possess a pole, volume 699 of Lecture Notes in Mathematics. Springer, Berlin, 1979.
- [11] A. Grigor'yan. Analytic and geometric background of recurrence and non-explosion of the Brownian motion on Riemannian manifolds. Bull. Am. Math. Soc., New Ser., 36(2):135–249, 1999.
- [12] G. Grillo, K. Ishige, and M. Muratori. Nonlinear characterizations of stochastic completeness. J. Math. Pures Appl. (9), 139:63–82, 2020.
- [13] G. Grillo, K. Ishige, M. Muratori, and F. Punzo. A general nonlinear characterization of stochastic incompleteness, Preprint arXiv: <https://arxiv.org/abs/2301.07942>, 2023.
- [14] G. Grillo, M. Muratori, and J. L. Vázquez. The porous medium equation on Riemannian manifolds with negative curvature. The large-time behaviour. Adv. Math., 314:328–377, 2017.
- [15] J. Hulshof and R. C. A. M. van der Vorst. Asymptotic behaviour of ground states. Proc. Am. Math. Soc., 124(8):2423– 2431, 1996.

- [16] T. Kawakami and M. Muratori. Nonexistence of radial optimal functions for the Sobolev inequality on Cartan-Hadamard manifolds. In Geometric properties for parabolic and elliptic PDEs, volume 47 of Springer INdAM Ser., pages 183–203. Springer, Cham, [2021]  $\odot$ 2021.
- [17] P.-L. Lions. The concentration-compactness principle in the calculus of variations. The limit case. I. Rev. Mat. Iberoam., 1(1):145–201, 1985.
- [18] G. Mancini and K. Sandeep. On a semilinear elliptic equation in  $\mathbb{H}^n$ . Ann. Sc. Norm. Super. Pisa Cl. Sci. (5), 7(4):635–671, 2008.
- [19] E. Mitidieri. A Rellich type identity and applications. Comm. Partial Differential Equations, 18(1-2):125–151, 1993.
- [20] M. Muratori and N. Soave. Some rigidity results for Sobolev inequalities and related PDEs on Cartan-Hadamard manifolds. To appear in Ann. Sc. Norm. Super. Pisa Cl. Sci., [https://doi.org/10.2422%2F2036-2145.202105\\_071](https://doi.org/10.2422%2F2036-2145.202105_071), preprint arXiv: <https://arxiv.org/abs/2103.08240>, 2021.
- [21] P. Poláčik, P. Quittner, and P. Souplet. Singularity and decay estimates in superlinear problems via Liouville-type theorems. I. Elliptic equations and systems. Duke Math. J., 139(3):555–579, 2007.
- [22] P. Quittner and P. Souplet. Superlinear parabolic problems. Birkhäuser Advanced Texts: Basler Lehrbücher. [Birkhäuser Advanced Texts: Basel Textbooks]. Birkhäuser Verlag, Basel, 2007. Blow-up, global existence and steady states.
- [23] W. Reichel and H. Zou. Non-existence results for semilinear cooperative elliptic systems via moving spheres. J. Differ. Equations, 161(1):219–243, 2000.
- [24] J. Serrin and H. Zou. Non-existence of positive solutions of Lane-Emden systems. Differential Integral Equations, 9(4):635–653, 1996.
- [25] J. Serrin and H. Zou. Existence of positive entire solutions of elliptic Hamiltonian systems. Comm. Partial Differential Equations, 23(3-4):577–599, 1998.
- [26] J. Serrin and H. Zou. Existence of positive solutions of the Lane-Emden system. Atti Sem. Mat. Fis. Univ. Modena, 46(suppl.):369–380, 1998. Dedicated to Prof. C. Vinti (Italian) (Perugia, 1996).
- [27] P. Souplet. The proof of the Lane-Emden conjecture in four space dimensions. Adv. Math., 221(5):1409–1427, 2009.
- [28] X. J. Wang. Sharp constant in a Sobolev inequality. Nonlinear Anal., 20(3):261–268, 1993.

MATTEO MURATORI Dipartimento di Matematica, Politecnico di Milano Piazza Leonardo da Vinci 32, 20133 Milano, Italy Email address: matteo.muratori@polimi.it

Nicola Soave

Dipartimento di Matematica "Giuseppe Peano", Universita degli Studi di Torino ` Via Carlo Alberto 10, 10123 Torino, Italy Email address: nicola.soave@unito.it