

Substitution Voice Rehabilitation After Open Partial Horizontal Laryngectomy Through the Proprioceptive Elastic Method (PROEL): A Preliminary Study

*Marco Fantini, *Michela Gallia, †Gabriella Borrelli, ‡Nicole Pizzorni, §Andrea Ricci Maccarini, ¶Alfonso Borragan Torre, †Antonio Schindler, *¶Giovanni Succo, and *Erika Crosetti, *Candiolo, †¶Turin, ‡Milan, and §Cesena, Italy, and ¶Santander, Spain

Summary: Objective. The aim was to investigate the efficacy of the Proprioceptive Elastic Method (PROEL) in the rehabilitation of the substitution voice after open partial horizontal laryngectomy (OPHL).

Study design. Prospective outcome study.

Methods. Fifteen patients surgically treated by OPHL type II or type III for laryngeal cancer were recruited (experimental group). Each patient underwent a specific program of voice rehabilitation based on the PROEL method with the same speech and language pathologist. Acoustic-aerodynamic analysis: maximum phonation time (MPT); spectrographic classification (Titze's modified classification), perceptual analysis (INFVo rating scale) and self-assessments (SECEL questionnaire) were performed before the treatment (T0), after 3 months of rehabilitation (T1), and at the end of the 6-month rehabilitation program (T2). A control sample of other 15 patients who underwent OPHL type II or type III and who underwent a standard perioperative rehabilitation was randomly extracted from an historical database and compared to the experimental group.

Results. Significant voice improvements between T0-T1 and T2 were found for acoustic, aerodynamic, perceptual, and self-assessments analysis in the experimental group. Significant differences were found between the experimental group at T2 and the control sample for aerodynamic, self-assessment, and perceptual analysis.

Conclusions. The results of the present study support PROEL method as an effective approach for substitution voice rehabilitation after OPHL type II and III. Randomized controlled trials on larger groups of patients are needed in future in order to compare PROEL with other rehabilitative approaches.

Key Words: Communication—Laryngeal cancer—Laryngectomy—Rehabilitation—Voice.

INTRODUCTION

Surgical and nonsurgical treatments for laryngeal cancer can lead to voice impairment, with a severe impact on oral communication. This impairment can play a critical role affecting communication-related quality of life (QOL). People with verbal communication disabilities face demanding challenges in areas such as maintaining social roles, identity and accessing daily services.¹

Several treatment options are available today for the management of laryngeal cancer, including surgical and nonsurgical approaches. Among these, surgical function-sparing approaches allow the preservation of the main laryngeal functions and avoid a permanent tracheostoma, which causes cosmetic disability and results in a decreased QOL.²⁻⁴

Open Partial Horizontal Laryngectomies (OPHLs) represent a group of surgical procedures that have been recently classified with a systematic nomenclature by Succo et al.⁵ According to the OPHL classification system, three types of OPHLs can be described: type I (formerly defined horizontal supraglottic laryngectomy), type II (previously called supracricoid laryngectomy), and type III (also named supratracheal laryngectomy). While type I OPHL spares the glottic plane and shows little impact on voice quality; type II and III OPHLs sacrifice both vocal folds, resulting in the creation of a neoglottis. Voice after type II and III OPHL is usually highly deteriorated, with a hoarse and breathy quality. A comparative outcome study by Schindler et al showed similar results in functional outcomes after type II and type III OPHLs.⁶

The Proprioceptive Elastic Method (PROEL) is a multi-dimensional approach to voice and dysphonia developed by Alfonso Borragan.⁷ The method uses pressure, vibration, temperature, and stretching sensory stimuli in order to communicate with the phonatory organ in an intuitive and easy way. Proprioception and elasticity are obtained through voice hydration procedures, instable-balance postures, laryngeal manipulations, and semioccluded vocal tract postures. Particular importance is given to the reduction of vocal risk factors such as vocal overload, inflammatory conditions (eg, smoking and laryngopharyngeal reflux) and to vocal fold surface hydration procedures (nasal hydrotherapy). In order to eliminate tension, stress and muscular

Accepted for publication April 21, 2020.

Conflict of Interest: There are no competing interests for this article.

Financial Disclosures: The author Marco Fantini was supported by the AIRC fellowship for Italy research grant.

From the *Head and Neck Oncology Service, Candiolo Cancer Institute, FPO IRCCS, Candiolo, Italy; †Otorhinolaryngology Service, San Luigi Gonzaga Hospital, University of Turin, Turin, Italy; ‡Department of Biomedical and Clinical Sciences "L. Sacco", University of Milan, Milan, Italy; §Otorhinolaryngology Service, Bufalini Hospital, Cesena, Italy; ¶Centro de Foniatria y Logopedia, Santander, Spain; and ¶Oncology Department, University of Turin, Turin, Italy.

Address correspondence and reprint requests to Marco Fantini, Head and Neck Oncology Service, Candiolo Cancer Institute, FPO IRCCS, SP142, km 3,95, 10060 Candiolo, Italy. E-mail: marcofantini8811@hotmail.it

Journal of Voice, Vol. ■■■, No. ■■■, pp. ■■■–■■■
0892-1997

© 2020 The Voice Foundation. Published by Elsevier Inc. All rights reserved.

<https://doi.org/10.1016/j.jvoice.2020.04.025>

stiffness, body stretching maneuvers and facilitating instable postures are carried out (eg, the so-called leaning Pisa Tower position, where the patient stands on tiptoe with his/her body leaning upward while looking up at the ceiling). Laryngeal manipulations and semioccluded vocal tract exercises (eg, semioccluded ventilation mask postures) promote an easier and resonant phonation. Recently, the PROEL method showed efficacy in the treatment of a group of patients affected by functional dysphonia.⁸ Because voice quality is one of the major critical points regarding functional results after type II and type III OPHLs, postsurgical voice rehabilitation is crucial in order to reach a satisfying voice related QOL. Nevertheless, little information about voice rehabilitation programs and strategies after partial laryngectomies can be found in scientific literature.⁹⁻¹³

The aim of the present study was to investigate the efficacy of the PROEL method in the rehabilitation of substitution voice after type II and type III OPHL.

MATERIALS AND METHODS

This prospective outcome study was carried out according to the Declaration of Helsinki. All subjects enrolled in the study gave their informed consent; experimental data concerning the PROEL-based rehabilitation program were collected prospectively; a final comparison with an historical control sample was carried out in order to compare the results obtained with the PROEL method to the results historically obtained with a standard rehabilitative approach.

Population

Fifteen patients (12 males and 3 females) who underwent type II or type III OPHL at the Head and Neck Department of the FPO-IRCCS Candiolo Institute, at San Luigi Gonzaga Hospital and at Martini Hospital (Italy) were recruited during the last 2 years and enrolled in the experimental group.

Inclusion criteria were: OPHL type II or type III, no evidence of disease at the last follow-up, preservation of respiration, speech and oral feeding (absence of percutaneous endoscopic gastrostomy or naso-gastric tube), absence of the tracheostoma, no salvage total laryngectomy performed, and >6 months after surgery. Each patient underwent standard rehabilitation of voice and speech during hospitalization and no further rehabilitation after discharge. The standard voice and speech rehabilitation program consists of a supraglottic voice training aiming at a basic understanding and activation of the neoglottis. It focuses on apnea training; push-pull exercises and short vocalizations with hard onsets. The mean age of subjects recruited was 63.09 ± 9.92 years. According to the modern nomenclature,⁵ 3 patients underwent type IIa OPHL (20%); 5 patients underwent type IIa + ARY OPHL (33.33%); 2 patients underwent type IIb OPHL + ARY (13.33%); 3 patients underwent type IIIa OPHL (20%), and 2 patients underwent type IIIa OPHL + CAU (13.33%). Mean distance from surgery was 33.55 ± 40.83 months (6-98 months).

A control sample composed of other 15 patients (13 males and 2 females) who underwent type II or type III OPHL at the Head and Neck Department of the FPO-IRCCS Candiolo Institute, at San Luigi Gonzaga Hospital and at Martini Hospital in the last 10 years was randomly extracted through a random number generator from an historical database of more than 500 patients. All the patients included in the historical database satisfied the same inclusion criteria cited for the experimental sample. Functional outcomes were previously collected following the same procedure used for the experimental sample, at least 6 months after surgery. Each patient underwent a standard rehabilitation of voice and speech during hospitalization and no further rehabilitation after discharge. The mean age of the patients of the control sample was 65 ± 7.49 years. Two patients underwent type IIa OPHL (13.3%); 5 patients underwent type IIa + ARY OPHL (33.3%); 1 patient underwent a type IIb OPHL (6.7%); 4 patients underwent type IIIa OPHL (26.7%), and 3 patients underwent type IIIa OPHL + CAU (20%). Mean distance from surgery was 35.86 ± 27.12 months (6-75 months). Sociodemographic and clinical features of the two groups of patients are shown in Table 1.

Procedures

Each patient of the experimental group underwent a rehabilitation program following the standard principles of the PROEL method as described by Alfonso Borrigan^{7,8}: (1) control of vocal risk factors; (2) vocal proprioceptive awareness; (3) elimination of the mechanisms of stress, tension, and muscular stiffness; (4) Projection and resonance of the voice; (5) research into the feeling of freedom and well-being. Each patient underwent a 45-60 min PROEL session per week for the first 3 months. After each session, the therapist gave the patient concrete instructions to carry out a series of exercises for 5 minutes twice a day. For the last 3 months, the sessions took place every 2 weeks, with the aim of consolidating the results. All the therapeutic programs were conducted by the same experienced speech language pathologist (SLP).

The patients' voices were assessed by a trained speech and language pathologist before the PROEL-based voice therapy (T0), after 3 months of rehabilitation (T1) and at the end (6 months) of the speech therapy program (T2). Concerning self-assessments, each participant completed the Italian version of the Self-Evaluation of Communication Experiences after Laryngeal Cancer (I-SECEL).^{14,15} The I-SECEL questionnaire specifically assesses communication dysfunction after laryngectomy and their effects on patients' daily living activities. The questionnaire is composed of 34 items divided into 3 subscales: General (5 items), Environmental (14 items), and Attitudinal (15 items). Total scores range from 0 to 102, the general subscale from 0 to 15, the environmental subscale from 0 to 42, and the attitudinal subscale from 0 to 45. The higher the score, the greater the perception of communication dysfunction. The maximum

TABLE 1.
Sociodemographic and Clinical Factors in the Two Samples of Patients

	Experimental Group	Control Group	P Value
N. patients	15	15	ns
Age (years)	63.09 ± 9.92	65 ± 7.49	ns
Sex (n. and %)			ns
Male	12 (80%)	13 (86.7%)	
Female	3 (20%)	2 (13.3%)	
Distance from surgery (months)	33.55 ± 40.83	35.86 ± 27.12	ns
Type of surgery			ns
OPHL IIa	3 (20%)	2 (13.33%)	
OPHL IIa + ARY	5 (33.33%)	5 (33.33%)	
OPHL IIb		1 (6.7%)	
OPHL IIb + ARY	2 (13.33%)		
OPHL IIIa	3 (20%)	4 (26.7%)	
OPHL IIIb	2 (13.33%)	3(20%)	

phonation time (MPT) was measured through window selection on the production of 3 sustained /a/; the longest phonation time was recorded. Perceptual evaluation was carried out with the INFVo rating scale, a perceptual scale specifically developed for substitution voice, assessing overall quality impression and intelligibility (I), additive and unnecessary noise (N), speech fluency (F), and presence of voiced segments (Vo). Each parameter is scored on a visual analog scale from 0 (minimally deviant) to 10 (maximally deviant substitution voicing) and then is reported to a 4 categories ordinal scale.¹⁶ The perceptual evaluation was performed by listening to a recorded 56-word and 99-syllable passage. A spectrography of the sustained vowel /a/ was performed. Patients' voices were classified into four categories on the basis of the spectrogram analysis, according to the proposed modified Titze's classification.¹⁷ The following categories were used: (1) type 1 voices, periodic without strong modulations or subharmonics; (2) type 2 voices, with strong modulations, bifurcations, or subharmonics; (3) type 3 voices, smearing of energy across harmonics with visible fundamental frequency and 1 or 2 harmonics; (4) type 4 voices, aperiodic. Voices were recorded with a microphone Samson Meteor Mic (Samson Technologies, Hauppauge, NY) placed at a distance of 30 cm from the mouth of the patient, connected via USB to a MacBook Pro computer (Apple, Cupertino, CA) running PRAAT software (Version 5.3.57 for Mac, Boersma & Weenick, University of Amsterdam, Amsterdam, the Netherlands). The audio signals were digitized on 16 bit at sampling frequency of 48 kHz.

Voice recordings were assessed randomly and independently by two raters blinded to session assignment, speech and language pathologists, who underwent specific training. Both SLPs had at least 2 years of experience with perceptual assessment of substitution voice. In case of disagreement between the raters, they jointly reassessed the parameter until a consensus was reached.

Multiparametric voice assessments of the experimental group at T2 were then compared with the acoustic,

aerodynamic, perceptual, and self-assessment outcomes of the control sample.

Statistical analysis

Statistical analysis was carried out with GraphPad Prism software (Version 7.0, Apple, Cupertino, CA). Means and standard deviations (SDs) for acoustic, aerodynamic, self-assessment, and perceptual analysis were calculated. The normality of the distributions was assessed with the Kolmogorov-Smirnov test. Friedman tests with Dunn's post hoc corrections for multiple comparisons and ANOVA tests with Tukey's post hoc corrections for multiple comparisons were used to detect statistical differences between the variables measured at T0, T1, and T2, as appropriate. Unpaired *t* tests and Mann-Whitney tests were used to compare the outcomes of the experimental group and the control sample, as appropriate. An alpha of 0.05 was considered for the statistical procedures.

RESULTS

No significant differences were found concerning age, gender distribution, type of surgery and months from surgery distribution between the experimental group and the control sample, as shown in [Table 1](#).

Acoustic-aerodynamic analysis

Means and SDs for MPT in the experimental group were 6.67 ± 3.05 seconds at T0; 8.72 ± 2.76 seconds at T1 and 9.98 ± 3.2 seconds at T2, as shown in [Figure 1](#). Means and SDs for Titze's modified spectrographic class were 3.91 ± 0.30 at T0; 3.64 ± 0.50 at T1, and 3.09 ± 0.83 at T2, as shown in [Figure 2](#). For MPT, significant differences were found between T0-T1 ($P = 0.022$); T1-T2 (0.03) and T0-T2 ($P = 0.011$). Concerning Titze's modified spectrographic class, significant differences were found between T0 and T2 ($P = 0.015$).




FIGURE 1. MPT mean values and SD in the experimental group at T0, T1, T2, and in the control group.

Mean values for MPT and Titze's modified spectrographic class for the control sample were 6.60 ± 3.16 seconds and 3.2 ± 0.67 as shown in Figures 1 and 2, respectively. Significant differences between experimental T2 mean values and control mean values were found for MPT ($P = 0.012$).

Perceptual analysis

Mean and SD for the INFVo rating scale in the experimental group are shown in Table 2. The INFVo perceptual rating scale showed significant differences in all the subscales. In particular, post hoc tests showed significant differences between T0 and T2 for the subscales I ($P = 0.0001$), N ($P = 0.0001$), F ($P = 0.0004$), Vo ($P = 0.0002$); between T0 and T1 for the subscales F ($P = 0.0017$), Vo ($P = 0.0426$), and between T1 and T2 for the subscale F ($P = 0.0084$).

Means and SD for the INFVo rating scale of the control sample were I = 2.06 ± 0.53 ; N = 2.32 ± 0.48 ; F = 1.94 ± 0.37 ; Vo = 2.34 ± 0.58 . Significant differences between the




FIGURE 2. Titze's modified spectrographic class mean values and SD in the experimental group at T0, T1, T2, and in the control group.

experimental group at T2 and the control sample were found for the subscales I ($P = 0.0489$); N ($P = 0.0078$); F ($P = 0.0004$); Vo ($P = 0.0001$). Graphical results as shown in Figure 3.

Self-assessments

Means and SDs for the SECEL questionnaire are shown in Table 3. Significant differences were found for all the subscales. In particular, post hoc tests showed significant differences between T0 and T2 for the subscales T ($P = 0.0400$), G ($P = 0.0290$); between T0 and T1 for the subscales T ($P = 0.0013$), E ($P = 0.0040$), and A ($P = 0.0218$).

Means and SDs for the SECEL questionnaire in the control sample were T: 35.33 ± 11.92 ; G: 9.93 ± 2.37 ; E: 17.53 ± 6.95 ; A: 7.87 ± 5.67 . A significant difference between the experimental group at T2 and the control sample were found for the subscale G ($P = 0.0001$). Graphical results are shown in Figure 4.

TABLE 2.
Mean Values, Standard Deviations, Test, and *P* Value of INFVo Subscales at T0, T1, and T2

Subscale	T0 (Mean and SD)	T1 (Mean and SD)	T2 (Mean and SD)	Test	<i>P</i> Value
I	2.20 ± 0.65	1.58 ± 0.64	1.25 ± 0.61	Friedman test + Dunn's multiple comparisons test	0.0001
N	2.07 ± 0.63	1.69 ± 0.61	1.33 ± 0.64		
F	1.96 ± 0.49	1.55 ± 0.44	1.22 ± 0.51		
Vo	1.52 ± 1.22	0.73 ± 0.95	0.43 ± 0.73		

TABLE 3.
Mean Values and Standard Deviations, Test, and *P* Value of I-SECEL Subscales at T0, T1, and T2

Subscale	T0 (Mean and SD)	T1 (Mean and SD)	T2 (Mean and SD)	Test	<i>P</i> Value
T	39.36 ± 12.81	23.72 ± 8.32	28.09 ± 13.56	RM one way ANOVA + Tukey's multiple comparisons test	0.0012
G	5.63 ± 2.69	4.27 ± 2.28	3.36 ± 1.96		
E	21.63 ± 7.59	14.63 ± 6.5	18.00 ± 7.94		
A	12.09 ± 7.07	4.81 ± 3.19	6.73 ± 5.20		




FIGURE 3. INFVo rating scale mean values and SD in the experimental group at T0, T1, T2, and in the control group.




FIGURE 4. SECEL questionnaire mean values and SD in the experimental group at T0, T1, T2, and in the control group.

DISCUSSION

The current study investigated voice quality and voice-related QOL improvements after a specific voice rehabilitation program based on the PROEL method in patients who underwent type II and type III OPHLs. Multiparametric voice assessments showed significant improvements regarding acoustic-aerodynamic analysis, perceptual evaluations and self-assessments after the rehabilitation program. One of the strengths of the study is the multiparametric voice assessment showing similar trends in voice improvement in aerodynamic, acoustic, and perceptual analysis at T0, T1, and T2. Self-assessments on the contrary improved at T1 compared to T0 but did not improve further at T2 for some subscales.

MPT significantly improved between T0-T1 and T0-T2, suggesting an increased valving activity of the neoglottis, while the significant reduction of mean Titze's spectrographic class suggested an improvement of the mean acoustic quality of the substitution voice produced by the neoglottis. Perceptually, the patient's voices improved significantly for all the INFVo subscales: overall quality impression and intelligibility (I), additive and unnecessary noise (N), speech fluency (F), and presence of voiced segments (Vo). A significant reduction in SECEL questionnaire

scores was detected too, suggesting an improvement in voice-related QOL.

The comparison between the experimental group after a PROEL-based voice rehabilitation and the control sample (with no further rehabilitation after hospitalization) showed overall better results for the experimental group and some significant differences concerning aerodynamic measures (MPT), self-assessments and perceptual analysis, suggesting the efficacy of a PROEL-based rehabilitation program after hospitalization for the vocal function of patients who underwent type II and III OPHLs.

Looking individually at the results of the experimental group, the overall impression of the authors is that patients who underwent an OPHL procedure longer ago are likely to have little improvement compared to the ones who underwent surgery more recently. Nonetheless, the small size of the sample and the lack of a multivariate analysis does not allow to generalize such statement at the moment.

Concerning PROEL application to voice rehabilitation, a previous study by Lucchini *et al* suggested the method's efficacy in the treatment of functional dysphonia.⁸ No experience about the application of the PROEL method for substitution voice rehabilitation after OPHL has been described so far. Long-term functional outcomes after OPHLs—in particular, type II OPHLs—have been investigated widely. In this surgery, both vocal folds are sacrificed and the voice is produced by the vibration of the arytenoid mucosa against the epiglottis/tongue base. The site of the mucosal wave is normally observed between the anterior part of the body of one or both arytenoids and the tongue base or epiglottis.¹⁸ Thus, the vocal signal after type II OPHL shows substantially poor volitionally induced valving activity and resistance to airflow during voicing,¹⁹⁻²² resulting in a strained, deep and asexuated voice (difficult to modulate and to raise) and speech that is composed of reduced phrase groups, because the patients rapidly become short of breath.^{5,23-25} Functional results of type III OPHLs have recently been analyzed,²⁶ showing similar outcomes to type II OPHLs,⁶ with severe voice impairment but well preserved oral communication, and with almost all patients showing a good attitude toward their communication dysfunction. Concerning technological communication systems, a recent study by Crosetti *et al*²⁷ investigated telephonic voice intelligibility after various laryngeal cancer treatment options. Type II and III OPHLs showed the poorest intelligibility rates comparing to transoral laser microsurgery procedures, radiotherapy and type I OPHL. Concerning voice-related QOL, most authors have reported a moderate reduction in voice-related QOL after type II and III OPHL; however, the reports are somewhat contradictory.^{18,21,24,25,28}

Regarding substitution voice rehabilitation strategies after open partial laryngectomy, a recent study by Palmer *et al*¹³ tested the Expiratory Muscle Strength Training after supracricoid laryngectomy, obtaining some interesting results on some objective and subjective respiratory parameters like the peak cough flow and the dyspnea index. The

improvement of respiratory parameters could have a positive impact on some vocal aerodynamic parameters like the MPT, thus representing an interesting goal for rehabilitation after OPHL. Anyway, the study has some considerable limitations, like the very small sample size and the lack of a control group. In the present study, the significant improvement of MPT registered after the PROEL-based rehabilitation approach could be likely related to the effect of respiratory muscles training obtained through instable balance postures and water resistance therapy.^{7,8}

Other studies investigated the efficacy of some semioccluded vocal tract postures on voice rehabilitation after OPHL. Fouquet et al¹² found an efficacy of the hand-over-mouth exercise on voice perceptual quality and neoglottis closure and vibratory patterns after 2 minutes of training. Silveira et al¹¹ tested a bilabial fricative exercise combined with the vowel /e/. The authors found a vocal improvement concerning both perceptual and endoscopic findings after 4 minutes spent performing the proposed exercise. Even if the described studies have great limitations represented by the small sample sizes, they both suggest a possible positive effect of semioccluded vocal tract postures on substitution voice quality. This aspect has to be taken into account, since one of the basic principles of the PROEL method, represented by the proprioception of the vocal tract, is reached through the use of semioccluded vocal tract exercises performed through semioccluded ventilation masks.^{29,30}

Another aspect that is worth of consideration is represented by the great attention given by the PROEL method to vocal folds hydration and lubrication. As shown by a recent systematic review by Alves et al³¹ most of the recent literature about voice hydration is of good quality evidence and suggests that both systemic and surface hydration procedures should be encouraged in voice hygiene programs. Even if OPHLs require the removal of the true vocal folds, the vibrating residual mucosa presumably benefits from a condition of optimal hydration and lubrication.

The main limitations of the present study were the small number of recruited patients and the lack of a randomized controlled trial study design, even if a random selection of the control patients from an historical database was performed in order to reduce the selection bias. Future research should include larger number of patients and should focus on the various components of the PROEL method in order to investigate their effectiveness separately and to determine the weight of their contribution to the final functional result. Future studies should also aim at comparing the PROEL method to other rehabilitative approaches with randomized clinical trials.

CONCLUSIONS

Because voice and oral communication can be severely deteriorated after oncologic laryngeal surgery, it is very important for clinicians to guide patients through effective and rewarding rehabilitative programs. The results of the present study seem promising, as they outline some significant

positive changes in patients' voices after a PROEL-based rehabilitation program, suggesting PROEL as a likely effective approach for voice rehabilitation after OPHL.

Acknowledgments

Sincere thanks to Alfonso Borragan Torre and his team for their valuable suggestions.

REFERENCES

1. Yorkston KM, Bourgeois MS, Baylor CR. Communication and aging. *Phys Med Rehabil Clin N Am*. 2010;21:309–319. <https://doi.org/10.1016/j.pmr.2009.12.011>.
2. Mosconi P, Cifani S, Crispino S, et al. Head and Neck Cancer Italian Working Group. The performance of Sf-36 health survey in patients with laryngeal cancer. *Head Neck*. 2000;22:175–182. [https://doi.org/10.1002/\(sici\)1097-0347\(200003\)22:2<175::aid-hed10>3.0.co;2-v](https://doi.org/10.1002/(sici)1097-0347(200003)22:2<175::aid-hed10>3.0.co;2-v).
3. Weinstein GS, El-Sawy MM, Ruiz C, et al. Laryngeal preservation with supracricoid partial laryngectomy results in improved quality of life when compared with total laryngectomy. *Laryngoscope*. 2001;111:191–199. <https://doi.org/10.1097/00005537-200102000-00001>.
4. Olthoff A, Steuer-Vogt MK, Licht K, et al. Quality of life after treatment for laryngeal carcinomas. *ORL J Otorhinolaryngol Relat Spec*. 2006;68:253–258. <https://doi.org/10.1159/000093094>. Epub 2006 May 4.
5. Succo G, Peretti G, Piazza C, et al. Open partial horizontal laryngectomies: a proposal for classification by the working committee on nomenclature of the European Laryngological Society. *Eur Arch Otorhinolaryngol*. 2014;271:2489–2496. <https://doi.org/10.1007/s00405-014-3024-4>.
6. Schindler A, Pizzorni N, Fantini M, et al. Long-term functional results after open partial horizontal laryngectomy type IIa and type IIIa: a comparison study. *Head Neck*. 2016;38(Suppl 1):E1427–E1435. <https://doi.org/10.1007/s00405-014-3024-4>.
7. Borragan A, Lucchini E, Agudo M, et al. Il Metodo Propriocettivo Elastico (PROEL) nella terapia vocale. *Acta Phon Lat*. 2008;30:18–50.
8. Lucchini E, Ricci Maccarini A, Bissoni E, et al. Voice improvement in patients with functional dysphonia treated with the proprioceptive-elastic (PROEL) method. *J Voice*. 2018;32:209–215. <https://doi.org/10.1016/j.jvoice.2017.05.018>.
9. Schindler A, Pizzorni N, Mozzanica F, et al. Functional outcomes after supracricoid laryngectomy: what do we not know and what do we need to know? *Eur Arch Otorhinolaryngol*. 2016;273:3459–3475. <https://doi.org/10.1007/s00405-015-3822-3>. Epub 2015 Nov 6. Review.
10. Figueiredo IC, Vendramini SHF, Lourenção LG, et al. Profile and speech-language rehabilitation of patients with laryngeal cancer. *Codas*. 2019;31: e20180060. <https://doi.org/10.1590/2317-1782/20182018060>.
11. Silveira HSL, Simões-Zenari M, Kulcsar MA, et al. Combined vocal exercises for rehabilitation after supracricoid laryngectomy: evaluation of different execution times. *J Voice*. 2018;32:723–728. <https://doi.org/10.1016/j.jvoice.2017.09.009>.
12. Fouquet ML, Vieira TPG, Murata CJM, et al. Hands-over-mouth exercise in supracricoid horizontal partial laryngectomy: initial study. *Pro Fono*. 2012;17:346–350. <https://doi.org/10.1590/S1516-80342012000300018>.
13. Palmer AD, Bolognone RK, Thomsen S, et al. The safety and efficacy of expiratory muscle strength training for rehabilitation after supracricoid partial laryngectomy: a pilot investigation. *Ann Otol Rhinol Laryngol*. 2018 3489418812901. <https://doi.org/10.1177/0003489418812901>.
14. Schindler A, Mozzanica F, Brignoli F, et al. Reliability and validity of the Italian self-evaluation of communication experiences after laryngeal cancer questionnaire. *Head Neck*. 2013;35:1606–1615. <https://doi.org/10.1002/hed.23198>.
15. Blood G. Development and assessment of a scale addressing communication needs of patients with laryngectomies. *Am J Speech Lang Pathol*. 1993;82–87. <https://doi.org/10.1044/1058-0360.0203.82>.
16. Moerman M, Martens JP, Crevier-Buchman L, et al. The INFVo perceptual rating scale for substitution voicing: development and reliability. *Eur Arch Otorhinolaryngol*. 2006;263:435–439. <https://doi.org/10.1007/s00405-005-1033-z>. Epub 2006 Jan 11.

17. Sprecher A, Olszewski A, Jiang JJ, et al. Updating signal typing in voice: addition of type 4 signals. *J Acoust Soc Am*. 2010;127:3710–3716. <https://doi.org/10.1121/1.3397477>.
18. Weinstein GS, El-Sawy MM, Ruiz C, et al. Laryngeal preservation with supracricoid partial laryngectomy results in improved quality of life when compared with total laryngectomy. *Laryngoscope*. 2001;111:191–199. <https://doi.org/10.1097/00005537-200102000-00001>.
19. Saito K, Araki K, Ogawa K, et al. Laryngeal function after supracricoid laryngectomy. *Otolaryngol Head Neck Surg*. 2009;140:487–492. <https://doi.org/10.1016/j.otohns.2008.12.036>.
20. Webster KT, Samlan RA, Jones B, et al. Supracricoid partial laryngectomy: swallowing, voice, and speech outcomes. *Ann Otol Rhinol Laryngol*. 2010;119:10–16. <https://doi.org/10.1177/000348941011900102>.
21. So YK, Yun YS, Baek CH, et al. Son YI speech outcome of supracricoid partial laryngectomy: comparison with total laryngectomy and anatomic considerations. *Otolaryngol Head Neck Surg*. 2009;141:770–775. <https://doi.org/10.1016/j.otohns.2009.08.028>.
22. Laudadio P, Presutti L, Dall'olio D, et al. Supracricoid laryngectomies: long-term oncological and functional results. *Acta Otolaryngol*. 2006;126:640–649. <https://doi.org/10.1080/00016480500469024>.
23. Bron L, Pasche P, Brossard E, et al. Functional analysis after supracricoid partial laryngectomy with cricothyroidopiglotomy. *Laryngoscope*. 2002;112(7 Pt 1):1289–1293. <https://doi.org/10.1097/00005537-200207000-00027>.
24. Dworkin JP, Meleca RJ, Zacharek MA, et al. Voice and deglutition functions after the supracricoid and total laryngectomy procedures for advanced stage laryngeal carcinoma. *Otolaryngol Head Neck Surg*. 2003;129:311–320. [https://doi.org/10.1016/s0194-5998\(03\)01314-7](https://doi.org/10.1016/s0194-5998(03)01314-7).
25. Torrejano G, Guimarães I. Voice quality after supracricoid laryngectomy and total laryngectomy with insertion of voice prosthesis. *J Voice*. 2009;23:240–246. <https://doi.org/10.1016/j.jvoice.2007.08.005>.
26. Schindler A, Fantini M, Pizzorni N, et al. Swallowing, voice, and quality of life after supratracheal laryngectomy: preliminary long-term results. *Head Neck*. 2015;37:557–566. <https://doi.org/10.1002/hed.23636>.
27. Crosetti E, Fantini M, Arrigoni G, et al. Telephonic voice intelligibility after laryngeal cancer treatment: is therapeutic approach significant? *Eur Arch Otorhinolaryngol*. 2017;274:337–346. <https://doi.org/10.1007/s00405-016-4217-9>.
28. Schindler A, Mozzanica F, Ginocchio D, et al. Voice-related quality of life in patients after total and partial laryngectomy. *Auris Nasus Larynx*. 2012;39:77–83. <https://doi.org/10.1016/j.anl.2011.03.009>.
29. Fantini M, Succo G, Crosetti E, et al. Voice quality after a semi-occluded vocal tract exercise with a ventilation mask in contemporary commercial singers: acoustic analysis and self-assessments. *J Voice*. 2017;31:336–341. <https://doi.org/10.1016/j.jvoice.2016.05.019>.
30. Frisancho K, Salfate L, Lizana K, et al. Immediate effects of the semi-occluded ventilation mask on subjects diagnosed with functional dysphonia and subjects with normal voices. *J Voice*. 2018. <https://doi.org/10.1016/j.jvoice.2018.10.004>. pii: S0892-1997(18)30371-0.
31. Alves M, Krüger E, Pillay B, et al. The effect of hydration on voice quality in adults: a systematic review. *J Voice*. 2019;33:125. <https://doi.org/10.1016/j.jvoice.2017.10.001>. e13-125.e28.